

Brain Peptides and Glial Growth.

II. Identification of Cells That Secrete Glia-promoting Factors

Dana Giulian and Douglas G. Young

Program of Neuroscience, Department of Neurology, Baylor College of Medicine, Houston, Texas 77030

Abstract. Glia-promoting factors (GPFs) are brain peptides which stimulate growth of specific macroglial populations *in vitro*. To identify the cellular sources of GPFs, we examined enriched brain cell cultures and cell lines derived from the nervous system for the production of growth factors. Ameboid microglia secreted astroglia-stimulating peptides, while growing neurons were the best source of the oligodendroglia-

stimulating factors. These secretion products co-purified by gel filtration, anion exchange chromatography, and reverse-phase high performance liquid chromatography with GPFs isolated from goldfish and rat brain. Our findings suggest that glial growth in the central nervous system is regulated in part by a signaled release of peptides from specific secretory cells.

As described in the preceding report, peptides recovered from the central nervous system stimulated the proliferation of oligodendroglia or astroglia in culture (10). Elevated concentrations of some of these glia-promoting factors (GPFs)¹ were found during periods of glial proliferation in the developing rat brain, in the regenerating goldfish visual system, or after injury to the adult rat brain (10). Since our findings suggested that secretion of growth-stimulating peptides might be associated with gliogenesis, we sought to identify the cells which were responsible for the production of GPF-like factors. In this report, we isolated GPFs from cells in culture using gel filtration, ion-exchange chromatography, and reverse-phase high-performance liquid chromatography (HPLC). Astroglia-stimulating GPFs were secreted by microglia, and oligodendroglia-stimulating GPFs were produced by neurons. Proliferation of glia in the brain may, therefore, be regulated by the release of peptides from specific secretory cells.

Materials and Methods

Tissue Preparation and Cell Cultures

Optic tecta obtained from common goldfish (Ozark Fisheries, Ozark, MO) or cerebral cortices from newborn albino rat (Holtzman Co., Madison, WI) were isolated as described previously (5, 10, 11). Pooled tissues were dispersed by mild sonication in phosphate-buffered saline (PBS) and centrifuged for 30 min at 15,000 *g* (Microfuge) at 4°C. The supernatant was filtered (0.45 μ m; Millipore/Continental Water Systems, Bedford, MA) and applied to a P-10 column (100 \times 0.9 cm, Bio-Rad Laboratories, Richmond, CA). The column was eluted with sterile PBS (pH 7.4) and 700- μ l fractions of eluate were collected in sterile plastic collecting trays. The cell lines C-1300 (CCL #147), C6 (CCL #107), GH3 (CCL #821), N2A (CCL #131), and 3T3 (CCL #92) were obtained from American Type Culture Collection, Rockville, MD. The B65 neuronal cell line

¹ *Abbreviations used in this paper:* Dil-ac-LDL, acetylated low density lipoprotein dioctadecyl-1-3,3,3',3'-tetramethyl-indo-carbocyanine; GC, galactocerebroside; GFAP, glial fibrillary acidic protein; GPF, glia-promoting factor; HPLC, high performance liquid chromatography.

from rat brain tumor was a gift from Dr. Yasuko Tomozawa (Baylor College of Medicine, Houston, TX; reference 22). Astroglia were isolated by the method described by McCarthy and de Vellis (19) with 5 mM L-leucine methyl ester used to destroy contaminating microglia (24). Ganglion cells were isolated from the goldfish retina by microdissection as described previously (4). The fluorescent compound, true blue (Sigma Chemical Co., St. Louis, MO), was used as a retrograde tracer (13) to confirm the identity of isolated ganglion cells (Fig. 1).

Ameboid microglia were obtained from mixed glial cell cultures grown for 1 wk in plastic flasks containing chemically defined medium described by Bottenstein and Sato (2) with 10% fetal calf serum. The flasks of 1-wk-old cultures were agitated for 15 h on a rotary shaker (180 rpm) at 37°C, and the suspended cells were transferred to new plastic flasks. Microglia selectively adhered to plastic within 3 h at 37°C, in defined medium containing 10% fetal calf serum (6). These adhering cells were re-suspended in a Ca²⁺, Mg²⁺-free PBS with 5 mM EDTA, and transferred to a plastic flask containing defined medium with 10% fetal calf serum. The population of cells adhering after a second 3-h period consisted of 95 \pm 3% nonspecific esterase-positive microglia. These isolated ameboid microglia, proliferated *in vitro*, were capable of phagocytosing 5- μ m latex beads and contained the acetylated low density lipoprotein receptor (Fig. 2) (6, 21) as well as the macrophage surface antigens, MAC-1 and MAC-3 (23). The fluorescent probe for acetylated low density lipoprotein, dioctadecyl-1-3,3,3',3'-tetramethyl-indo-carbocyanine (Dil-ac-LDL) was a gift from Dr. David Via of the Department of Medicine, Baylor College of Medicine, Houston, TX.

Glial Cell Assays

Preparation of glial cultures and identification of specific cell populations by indirect immunofluorescence were carried out as described in the previous report (10, 11). Galactocerebroside (GC) served as a marker for oligodendroglia and glial fibrillary acidic protein (GFAP) was used to identify astroglia. Bioassays to test for GPF activities used glia isolated from the cerebral cortex of newborn rat and plated on poly-L-lysine-coated glass coverslips in 35-mm plastic dishes containing 1.5 ml of defined medium with 10% fetal calf serum. After 48 h, these cultures were washed three times with defined medium (10). Between 10 and 100 μ l of partially purified GPFs in PBS were added to glial cultures in 1.5 ml of defined medium. The final concentrations of the partially purified GPFs ranged from 0.05 to 10 μ g/ml of culture medium (1). Matching control cultures were prepared with equivalent volumes (10–150 μ l) of PBS. After an incubation period of 72 h, cultures were stained for GC⁽⁺⁾ or GFAP⁽⁺⁾ cells. The mean cell number was determined from 10 randomly selected fields (0.314 mm²) of each coverslip when viewed by a microscope with epifluorescence (Nikon). Data were expressed as a fold of increase in the mean cell




Figure 1. Photomicrographs of isolated ganglion cell layer recovered by microdissection from the goldfish retina (4). The preparation as seen by phase-contrast microscopy consists of unmyelinated axons and ganglion cell bodies (A and C). These cells filled with a fluorescent label, true blue, 72 h after retrograde transport in vivo (B and D). Previous study (4), using morphological and biochemical markers, has shown this dissection to be free of major glial contaminants. Bar, 20 μ m.




Figure 2. Photomicrographs of enriched microglia preparations. Ameboid microglia were isolated by adhesion to plastic, as described in Materials and Methods. As shown here, these isolated cells contain the acetylated low density lipoprotein receptor and bind the fluorescent probe Dil-ac-LDL (B). Bar, 20 μ m.

number over cultures incubated with matching aliquots of PBS controls. A unit of biological activity for a specific GPF was defined as the fold increase above the mean control cell number per microgram protein added to a milliliter of defined culture medium for 72 h.

Column Chromatography

GPFs separated by gel filtration were further purified using ion-exchange chromatography. Pooled fractions with GPF₁ or GPF₂ were applied to DEAE-

5PW (Bio-Rad Laboratories, Richmond, CA) and eluted with a gradient of NaCl in 20 mM of sodium phosphate buffer (pH 7.4). In the case of GPF₁, the peak of oligodendroglia-stimulating activity was found routinely in fractions 17 through 25, while GPF₂ eluted in fractions 8 through 12 (Fig. 3). A combination of gel filtration and ion-exchange chromatography gave a final purification of ~2,000- to 3,000-fold for GPF₁ or GPF₂ (Table I). Between 80 and 90% of total biological activity for either GPF₁ or GPF₂ was recovered from the ion-exchange column.

GPFs partially purified by gel filtration were also separated on a C3 column




Figure 3. Separation of GPFs by ion-exchange chromatography. GPF₁ and GPF₂, partially purified by gel filtration, were eluted from a DEAE-5PW column with a 0.02-M sodium phosphate buffer (pH 7.2) and a gradient of sodium chloride. The total protein concentration in the 700- μ l fractions was determined by the fluorescamine method. The biological assay was carried out as described in Materials and Methods using three cultures per data point. (Upper panel) Recovery of active GPF₁; (lower panel) recovery of active GPF₂.

(Beckman Instruments, Inc., Palo Alto, CA) by reverse-phase HPLC using an acetonitrile gradient (Fisher Chemical Co., Pittsburgh, PA) in distilled, deionized water containing 10 mM trifluoroacetic acid (Pierce Chemical Co., Rockford, IL). Two peaks of biological activity were recovered from fractions containing partially purified GPF₁, one peak eluting between 37 and 40% and the second between 46 and 49% of acetonitrile (Fig. 4). A single peak of activity for GPF₂ was eluted with a 25% concentration of acetonitrile (Fig. 4). The two-step purification using gel filtration and reverse-phase HPLC showed ~8,500-fold purification for GPF₁ and 2,000-fold for GPF₂ (Table I). After lyophilization and renaturation of samples, the recovery of biological activity ranged from 10 to 30% of GPF₁ and from 30 to 50% for GPF₂.

For sieving HPLC, we used two BioSil TSK-125 columns (Bio-Rad Laboratories) with an eluting buffer of 100 mM sodium sulfate and 20 mM potassium phosphate (pH 6.8).

Results

GPFs have been identified as peptide growth factors present

during gliogenesis in the central nervous system (10). Since the secretion of these peptides might represent an important mechanism for controlling glial growth, we sought to determine the cellular sources of GPFs in the brain. This type of investigation required us to distinguish GPFs from other factors that stimulated cell growth by several different biochemical criteria.

Identification of Cells That Promote Macroglial Growth

We screened cell lines derived from tumors of the nervous system or enriched cell preparations from newborn rat brain for the presence of glia-stimulating activities. Microglia, isolated by selective adhesion, were nearly homogenous with

Table I. Specific Activity and Degree of Purification of GPF₁ and GPF₂

Techniques	GPF ₁		GPF ₂	
	Specific activity	Fold purification	Specific activity	Fold purification
	<i>U/μg per ml</i>		<i>U/μg per ml</i>	
Tectal supernatant	0.007 ± 0.003	1	0.008 ± 0.001	1
P10	2.40 ± 0.63	350	2.00 ± 0.29	250
P10 + DEAE	19.00 ± 2.29	2,700	15.63 ± 2.03	1,950
P10 + C3	61.00 ± 3.60	8,700	20.00 ± 2.00	2,500

A unit of biological activity represents a 100% increase in the mean numbers of oligodendroglia or astroglia per mm² when compared to control preparations treated with an equivalent volume of PBS. The increase in cell number is calculated from dose-response curves involving at least four cultures for each factor concentration assayed. Specific activity is expressed as U/μg protein in 1 ml of culture medium. Data are presented as mean values ± SEM.

95% of cells containing acetylated lipoprotein receptors (Fig. 2). Astroglial cultures, prepared using McCarthy's method (19), provided 98% GFAP⁽⁺⁾ cells. We also examined the neuronal cell lines B65, N-2A, and C-1300, as well as the C6 glial cell line. The fibroblast cell line 3T3 and the pituitary cell line GH3 provided nonneural control material.

Cell lines were harvested from 35-mm plastic dishes and dispersed by sonication. These sonicates were incubated with newborn rat brain cultures and after 48 h, glia populations were identified by immunofluorescence. Significant increases in the number of oligodendroglia were found in cultures incubated with sonicates from the neuronal lines N-2A and C-1300, but not with sonicates from microglia or the C6, GH3, or the 3T3 cell lines (Fig. 5). By comparison, the microglial sonicates stimulated the appearance of GFAP⁽⁺⁾ astroglia (Fig. 5). We next explored the possibility that the glia-stimulators found in cell sonicates were similar to GPFs isolated from brain.

Microglia Secrete Astroglia-stimulating GPFs

Putative astroglia-stimulating factors from cell preparations were fractionated by gel filtration. Once again, only microglial sonicates contained significant biological activity. Two of the microglial factors co-purified by gel filtration with fish GPF₂ and GPF₄ (Fig. 6).

As suggested earlier, a signaled secretion of growth factors might help to control glial proliferation in the brain. Our initial attempts to detect microglial secretion of GPF-like substances proved unsuccessful. We found, however, that a 24-h incubation of microglia with fixed *Staphylococcus aureus*, a known macrophage activator (3), significantly increased the amount of astroglia-stimulating activity released into culture medium. This growth activity, fractionated by gel filtration, contained factors similar to GPF₂ (9 kD) and GPF₄ (3 kD) (Fig. 7). We confirmed the identity of the 9-kD factor secreted by microglia as GPF₂ by co-purification with goldfish and rat brain factors using anion exchange chromatography (Fig. 8) and reverse-phase HPLC (Fig. 9). The similarities in apparent molecular masses, biological activities, and co-elution profiles suggested that microglia secreted GPF₂ in the central nervous system (20).




Figure 4. Separation of GPFs by reverse-phase chromatography. GPF₁ and GPF₂, partially purified by gel filtration, were eluted from a reverse-phase C3 column with 0.01 M trifluoroacetic acid and a gradient of acetonitrile (20–70%). The total protein concentrations in each of the 700-μl fractions were determined by the fluorescamine method. Biological assay was carried out as described in Materials and Methods using three cultures per data point. (*Upper panel*) Two peaks of biologically active GPF₁ were recovered; (*lower panel*) the major peak of biologically active GPF₂ was eluted with 25–30% acetonitrile.

Neuronal Cell Lines Secrete Oligodendroglia-stimulating GPFs

Neuronal cell lines contained factors that stimulated the growth of oligodendroglia in culture (Fig. 5). Sonicates from




Figure 5. Whole cell sonicates (25 μ l) containing \sim 100 μ g protein from glial cell line (C6), microglia (MIC), a fibroblast cell line (3T3), a pituitary tumor cell line (GH3), and two neuronal cell lines (N-2A and C-1300) were incubated with primary glial cell cultures grown in defined media for 48 h. The fold increase in GC⁽⁺⁾ oligodendroglia or GFAP⁽⁺⁾ astroglia were calculated from increases in mean cell numbers/mm² when compared to control cultures. These controls were incubated with 25 μ l of PBS for 48 h. As shown, sonicates from the neuronal cell lines increased the number of oligodendroglia found in culture while microglia sonicates contained astroglia-stimulating activity.




Figure 6. Cell production of astroglia-stimulating GPFs. Soluble protein from goldfish optic tectum or cell sonicates were separated by gel filtration and assayed for the presence of astroglia-stimulating factors. Microglia contained biological activity which co-eluted with GPF₂ (9 kD) and GPF₄ (3 kD) found in goldfish brain. Such biological activity was not detected in astroglia, the glial cell line, or the neuronal cell lines, N-2A and C-1300.




Figure 7. Secretion of astroglia-stimulating factors. Media conditioned for 24 h by microglia, astroglia, glial cell line C6, and the neuronal cell line C-1300, were assayed for the presence of GPFs. All cell cultures were activated by the addition of a suspension of fixed *Staphylococcus aureus* (20 μ l per 1.5 ml culture medium). The conditioned medium was collected after 24 h, and fractionated by gel filtration. Only microglia secreted detectable levels of GPF₂- and GPF₄-like factors. Microglia also released an 18-kD astroglia-stimulating peptide (not found in collected fractions) which has been identified as Interleukin-1 (Guilian D., T. J. Baker, and L. B. Lachman, unpublished data).

the cell lines C-1300, B65, and N-2A, when fractionated by gel filtration, showed GPF₁- and GPF₃-like activities; such activities were not detected in the C6 glial cell line, the fibroblast cell line 3T3, astroglia, or microglia (Fig. 10). We also examined media conditioned by these cell preparations for the presence of secreted oligodendroglia-stimulating peptides. Only the neuronal cell lines released factors which appeared similar to GPF₁ and GPF₃ (Fig. 11). The 15-kD peptide secreted by C-1300 co-eluted with authentic GPF₁ from goldfish optic tectum and from newborn rat brain using anion exchange chromatography (Fig. 12) and reverse-phase HPLC (Fig. 13).

Purification of GPF₁

As a final step to confirm the identities of the oligodendroglia-stimulating factors, we purified GPF₁ by combining the techniques of gel filtration, anion exchange chromatography, reverse-phase HPLC, and sieving HPLC. 200 mg of fish brain (400 optic tecta) yielded <10 ng of peptide with an estimated 100,000-fold purification (Fig. 14). This highly purified GPF₁




Figure 8. Co-purification of GPF₂ from fish, rat, and microglia using anion exchange chromatography confirms the similarity of these factors from different sources. Conditions for chromatography are described in Fig. 3.




Figure 9. Co-purification of GPF₂ from rat brain and from microglia using reverse-phase HPLC. Conditions for chromatography are described in Fig. 4.

appeared as a single peak of biological activity when separated by sieving HPLC with an apparent molecular mass of 15 kD. Using the same chromatographic techniques, we also isolated GPF₁, which was secreted by the C-1300 cell line. As shown in Fig. 15, highly purified GPF₁ from the neuronal cell line co-eluted by sieving HPLC with fish material. The co-purification of these GPFs suggest a high degree of structural homology between the oligodendroglia-stimulating peptides




Figure 10. Presence of oligodendroglia-stimulating GPFs in different cellular sources. Cell sonicates were separated by gel filtration and monitored for GPF activity. Only the neuronal cell lines, B65 and N-2A, showed detectable levels of GPF₁- and GPF₃-like activity.




Figure 11. Secretion of GPF₁- and GPF₃-like factors by neuronal cell lines. 10 ml of defined media conditioned by cells for 24 h were concentrated by ultrafiltration (YM-2 filter, Amicon Corp., Danvers, MA) and separated by gel filtration. As shown, neuronal cell lines (C-1300, N-2A) but not astroglia, microglia, or the cell lines C6 and 3T3, released oligodendroglia-stimulating factors.




Figure 12. Co-purification of GPF₁ from fish optic tectum, rat brain, and the C-1300 cell line using anion exchange chromatography as described in Fig. 3.




Figure 13. Co-purification of GPF₁ from rat cerebral cortex and from neuronal cell line C-1300 by reverse-phase HPLC. Conditions for isolation as described in Fig. 4. Two peaks of biological activity are recovered from either source at 36–39% and 45–47% acetonitrile.




Figure 14. Dose-response curves showing purification of GPF₁. Supernatants of goldfish optic tecta were separated by gel filtration (P-10), anion exchange chromatography (DEAE), reverse-phase HPLC (C3), and sieving HPLC (TSK) as described in Materials and Methods. Specific biological activity suggests a 100,000-fold purification of GPF₁, when compared to tectal supernatants. The yield for 400 optic tecta was <10 ng of GPF₁.




Figure 15. Highly purified GPF₁ isolated from goldfish brain or secreted by the neural cell line C-1300 were separated using sieving HPLC. These peptides co-eluted with an apparent molecular mass of 15 kD. Goldfish GPF was isolated as described in Fig. 14. 6 liters of conditioned media served as starting material for the C-1300 factor. This medium was concentrated by ultrafiltration (YM-2, Amicon Corp.) and then processed by the same techniques used for goldfish material. Molecular weight markers: (a) 68 kD; (b) 18 kD; (c) 12 kD; (d) 6 kD.

found in fish optic tectum and in a mammalian neuronal cell line.

Regenerating Neurons Produce GPF₁ and GPF₃

As noted before (10), brain levels of GPF₁ and GPF₃ were elevated during regeneration or development of the central nervous system. Since GPF₁ and GPF₃ were secreted by neuronal cell lines, we speculated that neurons might be a major source of oligodendroglia-stimulating peptides in vivo.




Figure 16. GPF production in neurons. Sonicates of 50 ganglion cell layer dissections (~1.5 mg protein) were separated by gel filtration. Oligodendroglia-stimulating factors GPF₁ and GPF₃ were found in retinal ganglion cells isolated from the regenerating goldfish visual system 12 d after axotomy (upper panel). No astroglia-stimulating activity was detected in the same preparation (lower panel).

As shown in Fig. 1, ganglion cells isolated from the goldfish retina by microdissection (4) consisted of cell bodies and unmyelinated axons. Sonicates of this glial-free preparation (4) contained significant levels of GPF₁ and GPF₃ (Fig. 16) with no evidence of astroglia-stimulating activity. Moreover, the concentrations of GPF₁ and GPF₃ were markedly elevated in neurons undergoing axonal regeneration (Fig. 17). We concluded that growing neurons found in the developing rat brain and the regenerating goldfish visual system were the likely sources of oligodendroglia-stimulating factors.

Discussion

We report here on the isolation of GPF₁, an oligodendroglia-promoting factor, by several different chromatographic methods. Picomolar concentrations of fish GPF₁, after a 100,000-fold purification, stimulated oligodendroglial proliferation *in vitro*. Based upon the specificity of its biological activity, GPF₁ appears to represent a new class of brain growth factor. The stability of this factor after lyophilization, exposure to acetonitrile, and SDS PAGE (Giulian, D., unpublished data) increases the likelihood of obtaining bulk quantities of homogenous peptide. The neuronal cell lines as sources of GPF₁ may provide sufficient material needed for such large-scale purifications.

Microglia are the principal phagocytic cells of the brain and consist of two forms (12, 18, 20): the ameboid cell, morphologically similar to the macrophage, is found in developing and injured brain, while the ramified cell is associated with the normal adult central nervous system (18). Investigators using histochemistry and electron microscopy (12, 18), have suggested that ameboid microglia serve as active scavenger cells and eventually differentiate into quiescent, ramified cells. Our study of enriched brain cell cultures indicate that "activated" ameboid microglia are the best secretors of the astro-




Figure 17. Comparison of GPF₁ and GPF₃ activities recovered from intact and regenerating retinal ganglion cells. Pooled fractions of biological activity recovered by gel filtration were assayed as described in Materials and Methods. Regenerating ganglion cells were isolated 12 d after axotomy. Isolated ganglion cells from intact retina provided control tissue. The greater specific biological activities found in regenerating cells suggested a stimulated production of GPFs in growing neurons.

Table II. GPFs in Brain

	Oligodendroglia-stimulating Factors	Astroglia-stimulating factors
Peptide factors	GPF ₁ (15 kD) GPF ₃ (6 kD)	GPF ₂ (9 kD) GPF ₄ (3 kD)
Developing mammalian brain	+++ (Postnatal)	+++ (Prenatal)
Injured adult mammalian brain	---	+++
Regenerating goldfish visual system	+++	---+
Brain cellular source	"Growing" neurons	Ameboid microglia

A summary of biological specificities and cellular sources for GPFs recovered from the central nervous system. The oligodendroglia-stimulating peptides are produced by neurons found in developing or regenerating tissues. Astroglia-stimulating factors are secreted by ameboid microglia found in embryonic brain or at sites of central nervous system injury.

glia-stimulating factors, GPF₂ and GPF₄ (Table II). Moreover, we detect significant levels of GPF₂ and GPF₄ in such microglial-rich tissues as the cerebral cortex of embryonic rat and wound sites of brain-injured adults (12, 18). Such findings suggest that ameboid microglia serve not only as scavenger

cells in the brain but also as regulators of astroglial growth by the release of peptide factors.

Neurons are the probable source of oligodendroglia-stimulating GPFs in developing rat brain and in the regenerating goldfish visual system (Table II). As reported here, neuronal cell lines secrete GPF₁ and GPF₃ in vitro, implying such secretion also occurs in vivo. Although the conditions for eliciting production of GPF₁ and GPF₃ are unknown, regenerating neurons showed greater factor concentrations than did quiescent cells. We suggest that peptides released by neurons stimulate oligodendroglial proliferation in the neighborhood of growing axons (7, 8).

Our findings point to the existence of a regulatory network whereby peptides released from specific secretory cells control the growth and proliferation of specific glial populations (7–9, 14–17). Perhaps application of these peptides will allow manipulation of cell growth in the developing or injured brain. Moreover, the identification of events which elicit the production and secretion of GPFs may help to elucidate mechanisms that control cellular organization of the nervous system.

We thank Joan Bales, Ivey Burton, and Richard Stoldt for their technical assistance and Judy Walker for the graphic artwork.

This work was supported by grants R01-EY04915 and R01-NS2063801 from the National Institutes of Health, by a Teacher Investigator Award (NS00806), by the Basil O'Conner Starter Researcher from the March of Dimes, and by funds from the Kroc Foundation.

Received for publication 2 January 1985, and in revised form 8 November 1985.

References

1. Bohlen, P., S. Stein, W. Dairman, and S. Udenfriend. 1973. Fluorometric assay of proteins in the nanogram range. *Arch. Biochem. Biophys.* 155:213–220.
2. Bottenstein, J. E., and G. H. Sato. 1979. Growth of a rat neuroblastoma cell line in serum-free supplemented medium. *Proc. Natl. Acad. Sci. USA.* 76:514–517.
3. Davies, P. 1981. Secretory functions of mononuclear phagocytes: overview and methods for preparing conditioned supernatants. In *Methods for Studying Mononuclear Phagocytes*. D. O. Adams, P. J. Edelson, and H. Koren, editors. Academic Press, Inc., New York. 549–559.
4. Giulian, D. 1980. Isolation of retinal ganglion cells from retina. *Brain Res.* 189:135–155.
5. Giulian, D. 1984. Peptides from the regenerating central nervous system of goldfish stimulate glia. *Proc. Natl. Acad. Sci. USA.* 81:3567–3571.
6. Giulian, D., and T. Baker. 1986. Characterization of amoeboid microglia isolated from mammalian brain. *J. Neurosci.* In press.
7. Giulian, D., V. Iwanij, and H. Stuckenbrok. 1985. The response of optic tract glia during regeneration of the goldfish visual system. I. Biosynthetic activity within different glial populations after transection of retinal ganglion cell axons. *Brain Res.* 339:87–96.
8. Giulian, D., and V. Iwanij. 1985. The response of optic tract glia during regeneration of the goldfish visual system. II. Tectal factors stimulate optic tract glia. *Brain Res.* 339:97–104.
9. Giulian, D., and L. B. Lachman. 1985. Interleukin-1 stimulation of astroglial proliferation after brain injury. *Science (Wash. DC)*. 228:497–499.
10. Giulian, D., R. L. Allen, T. J. Baker, and Y. Tomozawa. 1986. Brain peptides and glial growth. I. Glia-promoting factors as regulators of gliogenesis in the developing and injured central nervous system. *J. Cell Biol.* 102:803–811.
11. Giulian, D., Y. Tomozawa, H. Hindman, and R. L. Allen. 1985. Peptides from regenerating central nervous system promote specific populations of microglia. *Proc. Natl. Acad. Sci. USA.* 82:4287–4290.
12. Hortege, R. 1932. Microglia. In *Cytology and Cellular Pathology of the Nervous System*, Vol. 2. W. Penfield, editor. Paul B. Hoeber, Inc., New York. 481–534.
13. Illert, M., N. Fritz, A. A. Aschoff, and H. Hollander. 1982. Fluorescent compounds as retrograde tracers compared with horseradish peroxidase (HRP). II. A parametric study in the peripheral motor system of the cat. *J. Neurosci. Methods.* 6:199–218.
14. Lemke, G. E., and J. P. Brookes. 1984. Identification and purification of glial growth factor. *J. Neurol.* 4:75–83.
15. Lemmon, S. K., M. C. Riley, K. A. Thomas, G. A. Hoover, T. Maciag, and R. A. Bradshaw. 1982. Bovine fibroblast growth factor: comparison of brain and pituitary preparation. *J. Cell Biol.* 95:162–169.
16. Leutz, A., and M. Schachner. 1981. Epidermal growth factor stimulates DNA-synthesis of astrocytes in primary cerebellar cultures. *Cell & Tissue Res.* 220:393–404.
17. Lim, R. 1980. Glia maturation factor. *Curr. Top. Dev. Biol.* 16:305–322.
18. Ling, E. A. 1981. The origin and nature of microglia. In *Advances in Cellular Neurobiology*, Vol. 2. S. Fedoroff and L. Hertz, editors. Academic Press, Inc., New York. 33–82.
19. McCarthy, K., and J. de Vellis. 1980. Preparation of separate astroglial and oligodendroglial cell cultures from rat cerebral tissue. *J. Cell Biol.* 85:890–902.
20. Oehmichen, M. 1983. Inflammatory cells in the central nervous system. *Prog. Neuropathol.* 5:277–325.
21. Pitas, R. E., T. L. Innerarity, J. N. Weinstein, and R. W. Mahley. 1981. Acetoacetylated lipoproteins used to distinguish fibroblasts from macrophages in vitro by fluorescence microscopy. *Arteriosclerosis.* 1:177–185.
22. Schubert, D., S. Heineman, W. Carlisle, H. Tarikas, B. Kimes, J. Patrick, J. H. Steinbach, W. Culp, and B. L. Brandt. 1974. Clonal cell lines from rat central nervous system. *Nature (Lond.)*. 249:224–226.
23. Springer, T. A., and M. K. Ho. 1982. Macrophage differentiation antigens: markers of macrophage subpopulations and tissue localization. In *Hybridomas in Cancer Diagnosis and Treatment*. M. S. Mitchell and H. F. Oettgen, editors. Raven Press, New York. 35–46.
24. Thiele, G. L., M. Kurosaka, and P. E. Lipsky. 1983. Phenotype of the accessory cell necessary for mitogen-stimulated T and B cell responses in human peripheral blood: delineation by its sensitivity to the lysosomotropic agent, L-leucine methyl ester. *J. Immunol.* 131:2282–2290.