

Meier et al. Vol. 127, No. 6, December 1994. Pages 1505–1514.

The underlined and bold portion of the corrected rat NAP57 sequence below represents new carboxy-terminal amino acids due to the insertion of an extra guanoside nucleotide at position 1464 of the NAP57 cDNA.

MADAEAAAMTFPKKKKKERKPLPEADVAEIQHAEDFLIKPESKAAQLDT	50
SQWPLLLKNFDRLNVRTTHYTPIPCGSNPLKREIGEYVRTGFINLDKPSN	100
PSSHEVVAVIRRILRVEKTGHSGTLDPKVTGCLIVCIERATRLVKSQQSA	150
GKEYGVVRLHNAIEGTAQLSRALETLTGALFQRPPLIAAVKRQLRVRTI	200
YESRVVEYDPERRLGVFWSCEAGTYIRTLCVHLLGVGGQMQLRRL	250
RSGVVGGERDHMVTMHDVLDDAQYLYDHHRDESYLRRVVFPLEKLLTSHKRL	300
VMKDSAVNAICYGAKIMLPGLLRYEDGIEVNQEVVVITTKGEAVCVAIAL	350
MTTAVISTCDHGVVAKIKRIMERDTYPRKWGLGPKASQKKQLIKQGLLD	400
KHGRPTDGT PASWTRDYVDYSDSSKKATAAEATPGPGVTADAASIVKRKR	450
DSDSDADE ATPTTPRVKKEKKKKADGEEAAEDGDGDA TRKKKKKK	500
ARAAEELSG	509