
Meier et al. Vol. 127, No. 6, December 1994. Pages 1505–1514.

The underlined and bold portion of the corrected rat NAP57 sequence below represents new carboxy-terminal amino acids due to the insertion of an extra guanoside nucleotide at position 1464 of the NAP57 cDNA.

MADAEAAMTFPKKHKKKKERKPLPEADVAEIQHAEDFLIKPESKAAQLDT	50
SQWPLLLKNFDRNLNVRTTHYTPICGSNPLKREIGEYVRTGFINLDKPSN	100
PSSHEVVAWIRRIILRVEKTGHSGTLDPKVTGCLIVCIERATRLVKSQQSA	150
GKEYVGVVRLHNAIEGTAQLSRALETLTGALFQRPPLIAAVKRQLRVRTI	200
YESRVVEYDPERRLGVFWVSCEAGTYIRTLCVHLGLLLGVGGQMQLRRV	250
RSGVVGGERDHMVTMHDVLDAYLYDHRDESYLRRVVFPLEKLLTSHKRL	300
VMKDSAVNAICYGAKIMLPGLLRYEDGIEVNQEVVITTKGEAVCVAIL	350
MTTAVISTCDHGVVAKIKRVIMERDTPRWGLGPKASQKKQLIKQGLLD	400
KHGRPTDGT PASWTRDYVDYSDSSKKATAAEATPGPGVTADAASIVKRKR	450
DSDSDADE <u>ATPTTTPRVKKKKKKKEKADGGEAAEDGDGDATRKKKKK</u>	500
<u>ARAAEELSG</u>	509