

Primary Megakaryocytes Reveal a Role for Transcription Factor NF-E2 in Integrin α IIB β 3 Signaling

Masamichi Shiraga,* Alec Ritchie,* Sallouha Aidoudi,* Veronique Baron,* David Wilcox,† Gilbert White,§ Belen Ybarrondo,|| George Murphy,¶ Andrew Leavitt,¶ and Sanford Shattil*

*Department of Vascular Biology, The Scripps Research Institute, La Jolla, California 92037; †Department of Pediatrics, Medical College of Wisconsin, Milwaukee, Wisconsin 53226; §Department of Medicine, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599; ||PharMingen, Inc., San Diego, California 92139; and ¶Department of Laboratory Medicine, The University of California at San Francisco, San Francisco, California 94143

Abstract. Platelet integrin α IIB β 3 responds to intracellular signals by binding fibrinogen and triggering cytoskeletal reorganization, but the mechanisms of α IIB β 3 signaling remain poorly understood. To better understand this process, we established conditions to study α IIB β 3 signaling in primary murine megakaryocytes. Unlike platelets, these platelet precursors are amenable to genetic manipulation. Cytokine-stimulated bone marrow cultures produced three arbitrary populations of α IIB β 3-expressing cells with increasing size and DNA ploidy: small progenitors, intermediate-size young megakaryocytes, and large mature megakaryocytes. A majority of the large megakaryocytes bound fibrinogen in response to agonists, while almost none of the smaller cells did. Fibrinogen binding to large megakaryocytes was inhibited by Sindbis virus-mediated expression of isolated β 3 integrin cytoplasmic tails. Strikingly, large megakaryocytes from mice defi-

cient in the transcription factor NF-E2 failed to bind fibrinogen in response to agonists, despite normal surface expression of α IIB β 3. Furthermore, while megakaryocytes from wild-type mice spread on immobilized fibrinogen and exhibited filopodia, lamellipodia and Rho-dependent focal adhesions and stress fibers, NF-E2-deficient megakaryocytes adhered poorly. These studies establish that agonist-induced activation of α IIB β 3 is controlled by NF-E2-regulated signaling pathways that mature late in megakaryocyte development and converge at the β 3 cytoplasmic tail. Megakaryocytes provide a physiologically relevant and tractable system for analysis of bidirectional α IIB β 3 signaling.

Key words: α IIB β 3 • integrin • megakaryocyte • NF-E2 • signaling

INTEGRIN α IIB β 3 functions in platelets as an adhesion receptor for fibrinogen and von Willebrand factor and as a signaling receptor that regulates organization of the cytoskeleton (Fox, 1996; Hartwig et al., 1996; Shattil et al., 1998). The adhesive and signaling functions of α IIB β 3 are closely intertwined. For example, ligand binding to α IIB β 3 is regulated by inside out signals that control receptor affinity and avidity (Ginsberg et al., 1992), while outside in signals triggered by integrin ligation influence platelet shape, granule secretion, and clot retraction (Schoenwaelder et al., 1994; Hartwig et al., 1996; Yuan et al., 1997; Jenkins et al., 1998; Leng et al., 1998). Until recently, most studies of α IIB β 3 were conducted using platelets or, alternatively, immortalized cell lines transfected with

α IIB β 3 (O'Toole et al., 1994; Qi et al., 1998). These studies have been useful in formulating a conceptual framework for integrin signaling that includes a complicated network of heterotrimeric G proteins, protein and lipid kinases and phosphatases, phospholipases, proteases, and small GTPases (Fox, 1996; Rittenhouse, 1996; Shattil et al., 1998). However, a detailed understanding of α IIB β 3 signaling has been impeded by the inability to manipulate gene expression in anucleate platelets and by concerns that observations in cell lines may not be entirely relevant to platelets. In this context, it is now possible to study platelets in gene-targeted mice, an advance that is beginning to yield important new insights into the structure and function of α IIB β 3 (Holmbäck et al., 1996; Offermanns et al., 1997; Aszódi et al., 1999; Hauser et al., 1999; Hodivala-Dilke et al., 1999; Law et al., 1999). Nonetheless, further analysis of α IIB β 3 signaling would be facilitated if there were a means to manipulate genes *ex vivo* in an appropriate primary cell.

Megakaryocytes, the direct precursors of platelets, represent a potential model system to achieve this goal. The

The first two authors contributed equally to this work.

Address correspondence to Dr. Sanford J. Shattil, Department of Vascular Biology, The Scripps Research Institute, 10550 North Torrey Pines Rd., VB-5, La Jolla, CA 92037. Tel.: (858) 784-7148. Fax: (858) 784-7422. E-mail: shattil@scripps.edu

development of megakaryocytes from hematopoietic stem cells requires a suitable microenvironment, the action of specific hematopoietic growth factors, including thrombopoietin (TPO)¹, and a hierarchy of transcription factors, including FOG, GATA-1, and NF-E2 (Kaushansky, 1995; Long, 1998; Vyas et al., 1999). Recombinant growth factors allow for preferential expansion and maturation of murine and human megakaryocyte progenitors *ex vivo*, enabling biochemical studies to be carried out (Mountford et al., 1999; Rojnuckarin et al., 1999). In fact, human megakaryocytes cultured under such conditions have been shown to bind fibrinogen or a ligand-mimetic anti- α IIB β 3 monoclonal antibody after acute stimulation with TPO, demonstrating the presence of inside out signaling pathways in these cells (Zauli et al., 1997a).

Therefore, we set out to determine whether primary murine megakaryocytes could be used to further characterize α IIB β 3 signaling and, in effect, bridge the gap between *in vitro* experiments with platelets and clinicopathological analyses of gene-targeted mice. The results validate this approach and establish that agonist-induced inside out signaling and fibrinogen binding to α IIB β 3 require a gene or genes regulated by the transcription factor, NF-E2. Furthermore, these studies show that protein expression can be manipulated in mature megakaryocytes using Sindbis virus vectors, enabling a molecular analysis of α IIB β 3 signaling in the appropriate cellular context.

Materials and Methods

Murine Bone Marrow Cultures

All chemicals and reagents were from Sigma Chemical Co. unless indicated otherwise. Initial studies were carried out with wild-type megakaryocytes derived from BALB/C mice. p45 NF-E2-deficient mice (a generous gift from Drs. Ramesh Shivdasani and Stuart Orkin, Harvard Medical School, Boston, MA) are in a mixed C57Bl/6-129/Sv background (Shivdasani et al., 1995). When their megakaryocytes were studied, wild-type mice with a similarly mixed background were used as controls. There was no obvious difference between wild-type megakaryocytes from these different strains with respect to the α IIB β 3 functions studied. Bone marrow cells were harvested from 8–9-wk-old mice by flushing both sets of femurs and tibias with PBS containing 2% bovine serum albumin, 0.38% trisodium citrate, and 1 U/ml DNase. Mononuclear cells were isolated over Ficoll Hypaque (1.080 g/ml; Pharmacia) after a 30-min centrifugation at 400 *g*. Low-density mononuclear cells were cultured for up to 9 d at a starting density of 10⁶ cells/ml in Iscove's Modified Dulbecco's medium (Irvine Scientific) supplemented as described (Zauli et al., 1997a), and including 10 ng/ml murine TPO (a gift from Kirin Brewery Ltd., Japan), murine IL-6 and human IL-11 (Biosource International, Inc.). Cytospins of 20,000–100,000 cells were stained with Wright-Giemsa to analyze cell morphology.

Flow Cytometry

Cell size, antigen expression and DNA ploidy were assessed by flow cytometry. After 0–9 d in culture, bone marrow-derived cells were centrifuged at 200 *g* for 10 min, washed and resuspended in PBS at 4 × 10⁶/ml. Then 50- μ l aliquots were incubated for 30 min at 4°C with one of the following antibodies: 10 μ g/ml FITC-conjugated anti- α IIB (rat anti-mouse CD41; PharMingen); biotin anti-mouse GP V (murine clone C24.25.1; a gift from V. Ramakrishnan, Cor Therapeutics, Inc., South San Francisco, CA); FITC-anti-Gr-1, a granulocyte marker (rat anti-mouse; PharMingen); phycoerythrin-anti-Sca-1, a progenitor cell marker (rat anti-mouse; PharMingen); or a 1:1,000 dilution of rabbit anti-mouse GP Iba α (from V.

Ramakrishnan). Negative controls included an irrelevant isotype-matched rat or mouse antibody (PharMingen) or normal rabbit serum, as needed. When the primary antibodies were biotin-anti-GP V or rabbit anti-GP Iba α , the cells were washed and subsequently incubated for 15 min at 4°C with 20 μ g/ml FITC-streptavidin or 14 μ g/ml FITC anti-rabbit immunoglobulin (heavy plus light chains), respectively (Biosource International). Cells were finally resuspended in PBS containing 1 μ g/ml propidium iodide and analyzed on a FACSCalibur flow cytometer (Becton Dickinson). DNA ploidy was assessed as described (Williams et al., 1998).

Inside Out Signaling in Megakaryocytes

Purified human fibrinogen (Enzyme Research Laboratories) was labeled with FITC (Shattil et al., 1987). Human fibrinogen was used as a ligand instead of murine fibrinogen because of its wider availability and its capacity to bind to and support the aggregation of activated murine platelets (Law et al., 1999). Furthermore, in preliminary studies, murine and human fibrinogen were found to be equivalent inhibitors of FITC-human fibrinogen binding to agonist-stimulated murine and human platelets. After 6 d in culture, bone marrow cells were sedimented by gravity for 60 min at 37°C in a 50-ml conical polypropylene tube. In addition to achieving modest enrichment of megakaryocytes, gravity sedimentation limited any functional damage that might occur to these fragile cells during centrifugation. When the effects of certain inhibitors were studied (see Results), they or appropriate vehicle control buffer were added at this stage for the final 20 min. Cells were then gently resuspended to 4 × 10⁶/ml in modified Tyrode's buffer containing 1 mM CaCl₂ and 1 mM MgCl₂ (O'Toole et al., 1994) and then incubated for 30 min at room temperature in the presence of FITC-fibrinogen (250 μ g/ml), specific agonists and inhibitors, 10 μ g/ml of the non-function-blocking anti- α IIB antibody, and 20 μ g/ml phycoerythrin-streptavidin (Molecular Probes). After a 10-fold dilution with buffer containing 1 μ g/ml propidium iodide, fibrinogen binding was quantified by flow cytometry (Pampori et al., 1999).

FITC-fibrinogen binding was monitored in the FL1 channel of the flow cytometer on the gated subset of viable cells (e.g., negative for propidium iodide, FL3) that expressed α IIB β 3 (FL2). Preliminary studies indicated that fibrinogen binding was inhibited \geq 90% by either 10 mM EDTA (Law et al., 1999), 5 μ M kistrin (an RGD-containing disintegrin known to block α IIB β 3 function; Adler et al., 1991), or 20 μ g/ml 1B5, a function-blocking hamster monoclonal antibody specific for murine α IIB β 3 (a gift from B. Coller and S. Smyth, Mt. Sinai Medical Center, New York, NY; Lengweiler et al., 1999). Therefore, specific fibrinogen binding was defined as binding that was inhibited by kistrin or EDTA. When it was necessary to compare the binding data between subpopulations of cells that were heterogeneous in size and α IIB β 3 density, specific FITC-fibrinogen binding was expressed as a percent of maximal binding obtained in the presence of 1 mM MnCl₂, an activator of integrins (Bazzoni and Hemler, 1998).

To determine the effect of isolated integrin cytoplasmic tails on fibrinogen binding to megakaryocytes, cDNA encoding chimeric proteins consisting of the extracellular and transmembrane domains of the Tac subunit of the human IL-2 receptor (CD25) and the human β 3 cytoplasmic tail (Tac- β 3) or the α 5 cytoplasmic tail (Tac- α 5; Chen et al., 1994; LaFlamme et al., 1994) were cloned into a Sindbis virus vector (pSinRep5; Invitrogen). As a further control, a tailless Tac construct was also prepared. Linear, capped viral RNA was prepared and pseudovirions were produced in BHK cells using a Sindbis Expression System (Invitrogen). Pseudovirion titers were such that a 1:3,000 dilution of BHK cell supernatant infected virtually all BHK cells in a 100-mm dish, as determined by surface expression of Tac using a phycoerythrin-conjugated antibody to human CD25 (PharMingen). To infect megakaryocytes, day 5 bone marrow cells were suspended in a 1:1.5 dilution of virus for 1 h and then diluted with 7 vol of serum-free medium and cultured for another 24 h. The cells were stained with FITC-fibrinogen, phycoerythrin-anti-human CD25 (PharMingen), and propidium iodide, and fibrinogen binding to large, viable megakaryocytes was quantified by flow cytometry.

Outside In Signaling in Megakaryocytes

Bone marrow-derived cells were cultured for 5 or 6 d, allowed to settle by gravity for 1 h and resuspended to 2 × 10⁵/ml in buffer containing 137 mM NaCl, 2.7 mM KCl, 3.3 mM NaH₂PO₄, 1 mM MgCl₂, and 3.8 mM Hepes, pH 7.35. Cells were then incubated on fibrinogen-coated coverslips for 45 min at 37°C in the presence or absence of 100 nM phorbol myristate acetate, the latter added to activate protein kinase C and enhance cell spread-

1. Abbreviations used in this paper: CAT, chloramphenicol acyltransferase; TPO, thrombopoietin.

ing (Leng et al., 1998). After removing nonadherent cells, the adherent cells were fixed in 3.7% formaldehyde, permeabilized with 0.2% Triton X-100 in PBS, and stained with monoclonal anti-vinculin antibody, FITC-anti-mouse IgG and rhodamine-phalloidin. Cells were then analyzed by confocal laser scanning microscopy and photomicrographs were prepared using Adobe Photoshop 5.0 (Hato et al., 1998; Leng et al., 1998).

In some experiments, megakaryocytes from day 5 cultures were infected with recombinant Sindbis virus expressing either C3 exoenzyme, an inhibitor of Rho GTPase (Aktories et al., 1992; Sindbis/C3), or chloramphenicol acyl transferase (Sindbis/CAT; kind gifts from Drs. C.S. Hahn, University of Virginia and M.A. Schwartz, Scripps Research Institute). Viral stocks were produced and titered in BHK cells as described (Bobak et al., 1997; Hahn et al., 1992). For infection, megakaryocytes were allowed to adhere to fibrinogen-coated coverslips for 30 min at 37°C and incubated with Sindbis/C3 or Sindbis/CAT for 1 h at 37°C at a multiplicity of infection of 50. After washing, the cells were incubated another 2.5 h in buffer, and 100 nM phorbol myristate acetate was added for the final 30 min to enhance spreading. After washing, adherent cells were fixed, permeabilized, stained, and examined by confocal microscopy. In parallel, some cells were infected on fibrinogen-coated plastic dishes, scraped into SDS sample buffer, and 50- μ g aliquots were subjected to SDS-PAGE under reducing conditions (Laemmli, 1970). After transfer to nitrocellulose, blots were probed with a rabbit polyclonal antibody specific for Sindbis E1 protein (a gift from J.H. Strauss, California Institute of Technology, Pasadena, CA) or with a murine monoclonal antibody to Rho (Santa Cruz Biotechnology, Santa Cruz, CA). Blots were developed for 0.5–2 min by chemiluminescence (ECL; Amersham Corp.).

Results

α Iib β 3 and Megakaryocyte Development

To obtain primary murine megakaryocytes for studies of

α Iib β 3 signaling, a low-density fraction of bone marrow was cultured in serum-free medium containing 10 ng/ml TPO, IL-6, and IL-11. Total cell number decreased progressively by 45, 73, and 89% on days 3, 6, and 9, respectively. On the other hand, Wright-Giemsa staining showed a progressive increase in the proportion of large, polyploid cells, such that by day 6 both small, immature megakaryocytes with basophilic cytoplasm and larger, mature megakaryocytes with granular, eosinophilic cytoplasm were readily apparent (Fig. 1). The vast majority of cells on day zero were small, as assessed by flow cytometric light scatter (Fig. 2 A). However, by day 3 a subpopulation of larger cells of intermediate size was evident, and by day 6 a subpopulation of large cells was present, typically amounting to 5–10% of the total. When the large cells were FACS-sorted and stained, they consisted of mature megakaryocytes (Fig. 1 G). The relative proportion of large megakaryocytes decreased slightly by day 9 in culture, presumably the result of programmed cell death (Fig. 2 A; Zauli et al., 1997b). On day 6, the DNA ploidy of the small cells was 2N, while that of the intermediate-size and large cells ranged from 2–32N and 8–128N, respectively (Fig. 2 B).

Surface antigen expression was characterized by flow cytometry (Table I). The proportion of α Iib β 3-expressing cells increased progressively with time and by day 6 it represented 21% of all small cells, 98% of the intermediate-size cells and >99% of the large cells. On day zero, 15% of the small cells were positive for α Iib β 3, due in part to the




Figure 1. Morphology of murine bone marrow cells. Bone marrow was harvested and maintained in culture as described in Materials and Methods. Wright-Giemsa-stained cytopsin preparations after zero (A), 3 (B), 6 (C), and 9 (D) d demonstrated an increase in the proportion of megakaryocytes over time. Day 6 cells were sorted on the basis of size, as determined by flow cytometric light scatter profiles (see Fig. 2 A). E, illustrates the sorted small cells; F, intermediate-size cells; and G, large cells. The objective lens was 20 \times .




Figure 2. Size and DNA ploidy of murine bone marrow cells. Cells cultured as in Fig. 1 were characterized by flow cytometry as described in Materials and Methods. A shows dot plots for forward (FSC) and side (SSC) light scatter profiles of 10,000 cells. In each plot, three arbitrary analysis gates have been drawn. The gate at the lower left includes small cells, the middle gate intermediate-size cells, and the gate at the upper right large cells. Note a progressive increase in the proportion of intermediate-size and large cells with time. In B, cells on day 6 of culture were analyzed for DNA content. Note the increase in ploidy with cell size.

presence of platelets. Throughout the culture period, a significant minority of the cells expressed Sca-1 (Table I), and some coexpressed α I**IIb** β 3, suggesting that they were committed megakaryocyte progenitors. Expression of components of the platelet GP Ib-V-IX complex appeared to lag behind expression of α I**IIb** β 3 because not every α I**IIb** β 3-expressing cell was positive for GP Ib or GP V (Table I). Moreover, some cells expressed GP Ib but not GP V, suggesting that GP V is not required for GP Ib expression. On day 6, most of the small cells were positive for the myeloid marker, Gr-1, as were 53% of the intermediate-size cells and 2% of the large cells. Taken together, these results indicate that murine megakaryocytes can undergo progressive expansion and maturation in culture, permitting a detailed analysis of α I**IIb** β 3 signaling during megakaryocyte development.

α I**IIb** β 3 Signaling in Developing Megakaryocytes

Inside out signaling in α I**IIb** β 3-expressing megakaryocyte progenitors and developing megakaryocytes was assessed on day 6 of culture by measuring the agonist-dependent

binding of a saturating concentration of FITC-fibrinogen. Large megakaryocytes bound little or no fibrinogen in the absence of an agonist (Fig. 3 A). On the other hand, specific fibrinogen binding was observed after a 30-min incubation of the cells with 10 ng/ml SDF-1 α , a CXCR4 receptor agonist (Riviere et al., 1999; Wang et al., 1998) or 100 ng/ml TPO, the agonist for c-Mpl receptors (Kaushansky, 1999). Fibrinogen binding was even greater in response to typical platelet agonists, including 10–100 μ M ADP plus epinephrine, 0.5 U/ml thrombin, or 2 mM PAR4 thrombin receptor-activating peptide (GYPGKF; Kahn et al., 1998; Fig. 3 A). Although higher concentrations of these agonists did not stimulate additional fibrinogen binding, combinations of three or more agonists markedly increased fibrinogen binding, in some cases equivalent to that obtained with MnCl₂ (Fig. 3 A). None of the agonists increased surface expression of α I**IIb** β 3.

Fibrinogen binding induced by the combination of PAR4-activating peptide, ADP, and epinephrine was inhibited >80% by preincubation of the cells with 2 μ M PGI₂ or PGE₁, compounds that inhibit platelet aggregation through receptor-mediated stimulation of adenylyl

Table I. Expression of Surface Markers during *Ex Vivo* Expansion of Megakaryocytes

	Day 0				Day 3			Day 6			Day 9		
	Small	Small	Medium	Large									
α I IIb	15.3*	5.9	50.6	99.5	21.4	98.1	99.7	27.9	72.0	99.4			
GP Ib α	1.9	5.1	50.5	90.8	11.4	78.8	84.6	17.0	56.7	92.1			
GP V	8.9	7.8	29.2	54.6	14.9	45.4	53.3	21.7	54.4	88.9			
Sca-1	12.6	18.1	45.1	32.1	7.6	27.2	11.0	13.3	21.1	22.1			
Gr-1	57.6	88.6	49.1	8.9	86.8	53.3	2.5	90.7	53.8	14.7			

*Values represent the percentage of cells expressing a given surface marker and are the average of two independent experiments, each performed in duplicate.




Figure 3. Fibrinogen binding to α IIb β 3-expressing megakaryocyte progenitors and megakaryocytes. Bone marrow cells on day 6 of culture were stimulated with the indicated agonists in the presence of FITC-fibrinogen, a non-function-blocking anti- α IIb antibody (to identify α IIb β 3-expressing cells), and propidium iodide (to exclude dead cells). Agonist concentrations were: SDF-1 α , 10 ng/ml; TPO, 100 ng/ml; ADP and Epi(nephrine), 100 μ M each; Thrombin, 0.5 U/ml; and PAR4 receptor-activating peptide, 2 mM. The data in A depict specific fibrinogen binding determined by flow cytometry and are expressed as a percent of maximum specific binding obtained in response to MnCl₂. Fibrinogen binding in B is expressed as the percent of cells that stained positive for FITC-fibrinogen. Data represent the means \pm SEM of five separate experiments.

cyclase (Hawiger et al., 1980). In addition, fibrinogen binding was inhibited 48% by 100 nM wortmannin, an inhibitor of certain PI 3-kinases, 40% by 12 μ M bisindolylmaleimide, a protein kinase C inhibitor, and 57% by 100 μ M BAPTA-AM, a chelator of cytoplasmic free Ca²⁺. In contrast, preincubation of cells with 1 mM aspirin to inhibit cyclooxygenase had no effect on fibrinogen binding, although it did inhibit the response to arachidonic acid as expected. Collectively, these results indicate that excitatory agonists can stimulate and certain inhibitory agonists can block the ligand binding function of α IIb β 3 in large megakaryocytes, just as in platelets. Furthermore, activation of α IIb β 3 is regulated in a positive fashion by signaling pathways that likely involve PI 3-kinase, protein kinase C, and free Ca²⁺, and in a negative fashion by cyclic AMP.

Fig. 3 B shows that up to 60% of large megakaryocytes bound fibrinogen in response to agonists. To place this in perspective, \geq 90% of agonist-activated blood platelets obtained from the same donor mice bound fibrinogen (not shown). The lack of responsiveness of some large megakaryocytes appeared to be due to impaired signaling to α IIb β 3 because FITC-fibrinogen binding to all of the large cells could be induced with MnCl₂. Furthermore, very few intermediate-size megakaryocytes and no small α IIb β 3-expressing cells bound fibrinogen in response to the agonists depicted in Fig. 3 or in response to as much as 500 nM phorbol myristate acetate to activate protein kinase C. The results were similar if fetal hematopoietic tissue from the livers of day 14.5 murine fetuses was used instead of bone marrow as the source of megakaryocyte progenitors (not shown; Lecine et al., 1998a). Overall, these results indicate that inside out signaling pathways responsible for activation of α IIb β 3 become fully developed only late in megakaryocyte development.

Platelet adhesion to an immobilized α IIb β 3 ligand triggers the reorganization of actin filaments, filopodial and lamellipodial extension, and cell spreading (Nachmias and Golla, 1991; Yuan et al., 1997; Hagmann et al., 1998; Leng et al., 1998). To determine if megakaryocytes can engage in such outside in signaling, day 6 bone marrow cultures were incubated on fibrinogen-coated coverslips for 45 min. By this time, most of the large megakaryocytes had become adherent, and this response was α IIb β 3-dependent because it could be blocked by 5 μ M kistrin. Adherent megakaryocytes ranged in diameter from 10 to $>$ 50 μ m, and many exhibited filopodia and lamellipodia (Fig. 4, A and C). When 100 nM phorbol myristate acetate was present, the megakaryocytes spread more extensively and most now exhibited focal adhesions and stress fibers (Fig. 4, B and D). Actin filaments were often circumferential and parallel to the cell margins, rather than perpendicular as in fibroblasts, an observation made previously with guinea pig megakaryocytes (Leven and Nachmias, 1982). These results indicate that ligand binding to α IIb β 3 can promote cytoskeletal reorganization in megakaryocytes, accompanied by changes in cell shape.

Mechanisms of α IIb β 3 Signaling in Megakaryocytes

Studies of mice deficient in the p45 subunit of NF-E2 have revealed that this transcription factor is necessary for the final phases of megakaryocyte development and platelet




Figure 4. Distribution of vinculin and F-actin in large megakaryocytes adherent to fibrinogen. Megakaryocytes from day 6 cultures were incubated for 45 min on fibrinogen-coated coverslips in the absence (A and C) or presence (B and D) of 100 nM phorbol myristate acetate. The cells were fixed, permeabilized and stained to visualize vinculin (green) and F-actin (red) by confocal microscopy. Each panel shows a 0.2- μm confocal image taken near the ventral surface of a representative cell. The arrowheads in D point to vinculin-rich focal adhesions, and the arrow points to actin stress fibers. This experiment is representative of three so performed. Bar, 10 μm .

production (Shivdasani et al., 1995; Lecine et al., 1998b). Since inside out signaling to $\alpha\text{IIb}\beta 3$ was fully developed only in mature megakaryocytes (Fig. 3), we considered whether genes regulated by NF-E2 might be required for $\alpha\text{IIb}\beta 3$ activation. Accordingly, bone marrow from NF-E2^{-/-} mice was cultured for 6 days with TPO, IL-6, and IL-11, yielding a subpopulation of large megakaryocytes similar to wild-type and NF-E2^{+/-} controls. However, unlike cells from wild-type or NF-E2^{+/-} mice, large NF-E2^{-/-} megakaryocytes displayed a virtual absence of FITC-fibrinogen binding in response to the combination of PAR4 thrombin receptor-activating peptide, epinephrine, and ADP (Fig. 5). Similar results were obtained when the cells were stimulated with as much as 500 nM phorbol myristate acetate. This impairment of fibrinogen binding was due to a defect in inside out signaling because NF-E2^{-/-} megakaryocytes expressed normal amounts of $\alpha\text{IIb}\beta 3$ on their surfaces, and they bound FITC-fibrinogen as well as wild-type or NF-E2^{+/-} megakaryocytes in response to MnCl₂ (Fig. 5). These results indicate that NF-E2, or more likely genes regulated by NF-E2, are required for agonist-induced activation of $\alpha\text{IIb}\beta 3$.

In fibroblastic cell lines, the heterologous expression of membrane-tethered $\beta 3$ integrin cytoplasmic tails (in the form of a human Tac- $\beta 3$ fusion protein) can reverse the high-affinity state of a constitutively active mutant of $\alpha\text{IIb}\beta 3$. In contrast, Tac- $\alpha 5$ tails have no such effect (Chen

et al., 1994). To determine if chimeric $\beta 3$ tails could block agonist-induced activation of $\alpha\text{IIb}\beta 3$, day 5 bone marrow cultures were incubated for 1 h with Sindbis pseudovirions expressing Tac- $\beta 3$ RNA or, as a control, pseudovirions expressing Tac- $\alpha 5$. 24 h later, flow cytometry showed that the Tac chimeras had been expressed in 5–20% of large megakaryocytes. Under these conditions, Sindbis virus infection did not affect either $\alpha\text{IIb}\beta 3$ surface expression or megakaryocyte viability. Large megakaryocytes expressing relatively low levels of Tac- $\beta 3$ or Tac- $\alpha 5$ bound fibrinogen in response to agonists almost as well as noninfected cells. In contrast, megakaryocytes expressing relatively high levels of Tac- $\beta 3$ bound very little fibrinogen, and significantly less than cells expressing high levels of Tac- $\alpha 5$ ($P < 0.01$; Fig. 6). The more modest reduction in fibrinogen binding in cells expressing high levels of Tac- $\alpha 5$ was also observed with a tailless Tac construct (not shown), suggesting that it might have been due to viral infection, per se. Since Tac- $\beta 3$ cannot form a heterodimer with αIIb but may nevertheless interact with cytoplasmic proteins that normally bind to $\alpha\text{IIb}\beta 3$ (Chen et al., 1994), these results suggest that one or more $\beta 3$ cytoplasmic tail-binding proteins mediates physiological activation of $\alpha\text{IIb}\beta 3$.

To explore mechanisms of outside in signaling in megakaryocytes, cells adherent to fibrinogen were exposed for 1 h to Sindbis virus encoding C3 exoenzyme, which ADP-ribosylates and inactivates the small GTPase, Rho (Akt-




Figure 5. Effect of NF-E2 deficiency on agonist-induced fibrinogen binding to large megakaryocytes. In A, fibrinogen binding to large megakaryocytes was assessed as in Fig. 3 and expressed as mean fluorescence intensity. The concentration of PAR4 receptor-activating peptide was 2 mM, ADP and Epi(nephriene) were 100 μ M each and MnCl₂ was 1 mM. NF-E2^{-/-} and wild-type megakaryocytes were directly compared in 4 separate experiments (means \pm SEM), and NF-E2^{+/-} megakaryocytes were also studied on one of these occasions. Note that unlike their wild-type and NF-E2^{+/-} counterparts, the NF-E2^{-/-} megakaryocytes failed to bind fibrinogen in response to agonists, but did so in response to MnCl₂. In B, α IIB β 3 expression in wild-type and NF-E2^{-/-} megakaryocytes was compared on one occasion using an anti- α IIB antibody.

ries et al., 1992; Moorman et al., 1999). After subsequent incubation in the presence of phorbol myristate acetate to enhance cell spreading, cytoskeletal organization was analyzed. Both Sindbis/C3 and a control virus, Sindbis/CAT, infected adherent cells, as evidenced by the time-dependent expression of the 61-kD viral E1 protein on Western blots of cell lysates (not shown). Infection with Sindbis/C3 was associated with a reduction in the percent of megakaryocytes that contained focal adhesions and stress fibers (Fig. 7, A and B). Whereas 70% of adherent megakaryocytes incubated with Sindbis/CAT displayed focal adhesions and stress fibers, only 30% of cells incubated with Sindbis/C3 did so (Fig. 7 C, $P < 0.001$). Furthermore, the remaining stress fibers in Sindbis/C3-treated megakaryocytes were often thinner and less densely packed than in the control cells. The effect of Sindbis/C3 on the megakaryocyte cytoskeleton could reasonably be attributed to ADP-ribosylation of Rho, because \sim 50% of this protein in cell lysates displayed a retarded electrophoretic mobility characteristic of ribosylation (Fig. 7 C, inset). In contrast to its effect on focal adhesions and stress fibers, Sindbis/C3 did not appear to impair megakaryocyte spreading or the formation of filopodia and lamellipodia. Thus, Rho is responsible for a subset of cytoskeletal responses in




Figure 6. Effects of chimeric integrin cytoplasmic tails on agonist-induced fibrinogen binding to large megakaryocytes. After 5 d in culture, megakaryocytes in suspension were infected with Sindbis virus expressing either the Tac- β 3 cytoplasmic tail or the Tac- α 5 tail, as described in Materials and Methods. 24 h later, Tac expression and FITC-fibrinogen binding to large megakaryocytes were studied by flow cytometry. Arbitrary analysis gates were placed around megakaryocytes that expressed no Tac, relatively low levels of Tac, and relatively high levels of Tac. Data represent the means \pm SEM of seven separate experiments.

megakaryocytes that is triggered, in part, by outside in signaling through α IIB β 3.

Discussion

A common strategy to characterize intracellular signaling pathways is to express wild-type and mutant gene products and observe the effects on cell function. Unfortunately, this approach cannot be used to characterize integrin signaling in platelets because of inherent difficulties in expressing recombinant proteins in these anucleate cells. Consequently, we hypothesized that primary murine megakaryocytes might serve as a relevant and tractable model system for studies of α IIB β 3 signaling. Culture of murine bone marrow in the presence of TPO, IL-6, and IL-11 led to megakaryocyte expansion and maturation, allowing functional studies of α IIB β 3 and the following conclusions to be drawn. (a) As with platelets, megakaryocytes are capable of inside out signaling to control agonist-induced fibrinogen binding to α IIB β 3. This process becomes fully functional late in megakaryocyte development, requires a gene or genes regulated by transcription factor NF-E2, and converges on the β 3 cytoplasmic tail. (b) As with platelets, the adhesion of megakaryocytes to immobilized fibrinogen triggers outside in signals through α IIB β 3 that lead to cell spreading and cytoskeletal reorganization. Adherent megakaryocytes exhibit filopodia, lamellipodia, and Rho-dependent focal adhesions and stress fibers. (c) Sindbis virus vectors can be used to express heterologous proteins in megakaryocytes, enabling α IIB β 3 signaling to be studied in a physiological context in ways not possible with platelets.




Figure 7. Effect of C3 exoenzyme on outside in signaling through α IIb β 3 in megakaryocytes. After 5 d in culture, fibrinogen-adherent megakaryocytes were infected with Sindbis virus expressing either CAT or C3 exoenzyme as described in Materials and Methods. Cells were then incubated for another 2.5 h, the last 30 min of which 100 nM phorbol myristate acetate was added to enhance spreading. After fixation and staining as in Fig. 4, the cells were analyzed by confocal microscopy. A and B show representative confocal images of cells infected with Sindbis/CAT and Sindbis/C3, respectively. Note the prominent vinculin-rich focal adhesions (arrowheads) and stress fibers (arrow) in A, but not in B. In C, 100 adherent megakaryocytes treated with each virus were scored for focal adhesions and stress fibers by two independent observers. Data are the means \pm SEM of three separate experiments. The inset shows a Western blot of cell lysates probed with an antibody against Rho. In three such experiments, the intensity of the slower migrating (ADP-ribosylated) Rho band from Sindbis/C3-infected cells averaged 50% of the total Rho. Bar, 10 μ m.

Mechanisms of Inside Out Signaling in Megakaryocytes

In both human and murine platelets, ligand binding to α IIb β 3 is regulated by signaling events that modulate integrin affinity and/or avidity (Shattil et al., 1998; Tsakiris et al., 1999). In platelets, the use of selective enzyme inhibitors has implicated protein and lipid kinases, such as PI 3-kinase and protein kinase C, in receptor-mediated activation of α IIb β 3. However, the mechanism by which effectors modulate α IIb β 3 function is unknown. A recent report concluded that human bone marrow-derived megakaryocytes stimulated with TPO adhere to immobilized fibrinogen in a manner dependent on α IIb β 3 and PI 3-kinase (Zauli et al., 1997a). The present studies establish that the binding of soluble fibrinogen to α IIb β 3 is regulated in murine megakaryocytes. Unstimulated megakaryocytes failed to bind soluble fibrinogen, indicating that α IIb β 3 is in a default low affinity/avidity state. Stimulation with agonists, including TPO and substances that interact with G protein-linked receptors (SDF-1 α , thrombin, PAR4 receptor-activating peptide, ADP, epinephrine), caused rapid activation of α IIb β 3 and fibrinogen binding to a majority of mature megakaryocytes (Fig. 3). In fact, a combination of agonists caused as much fibrinogen binding as did MnCl₂, an activator of integrins (Bazzoni and Hemler, 1998). Platelet aggregation inhibitors, such as PGI₂ and PGE₁ as well as inhibitors of PI 3-kinase and protein kinase C, blocked agonist-induced fibrinogen binding. Thus, α IIb β 3 is likely subject to similar regulatory mechanisms in mature megakaryocytes and in platelets.

A consistent finding in this study was that few intermediate-size megakaryocytes and no small α IIb β 3-expressing progenitors bound fibrinogen in response to agonists (Fig.

3 B). This failure could not be explained by a lack of α IIb β 3 expression, by prior binding of ligands to α IIb β 3 during culture, or by structural alterations in the integrin, since these cells could bind fibrinogen in response to MnCl₂. The most likely explanation for these observations is that inside out signaling pathways to α IIb β 3 become fully developed relatively late in megakaryocytopoiesis. Developmental regulation of α IIb β 3 affinity/avidity, as opposed to regulation of α IIb β 3 expression, could have beneficial *in vivo* consequences by limiting the binding of soluble adhesive ligands to α IIb β 3 on progenitors and young megakaryocytes, thereby preventing their unnecessary aggregation. At the same time, the presence of α IIb β 3 on the surface of these cells, even in a low affinity/avidity state, might enable them to adhere to fibrinogen or other ligands in the extracellular matrix (Savage et al., 1992), and to initiate outside in signals (Haimovich et al., 1993).

The most striking finding in this study was an inability of p45 NF-E2^{-/-} megakaryocytes to bind fibrinogen in response to agonists (Fig. 5). NF-E2 contains a 45-kD subunit that is restricted to hematopoietic cells and a ubiquitous 18-kD subunit (Andrews, 1998). Since p45 and p18 hetero-oligomers are required for transcriptional activation, the results with p45-deficient megakaryocytes suggest that one or more genes regulated by NF-E2 are required for full inside out signaling. This discovery would have been difficult to validate and explore in platelets because NF-E2^{-/-} mice have so few circulating platelets (Shivdasani et al., 1995). In addition to highlighting the value of the megakaryocyte model, these results suggest that the identification and characterization of NF-E2-regulated

genes should advance our understanding of the mechanisms of inside out regulation of α IIB β 3.

Recently, an *in vivo* immunoselection strategy was used to identify thromboxane synthase as a p45 NF-E2 target gene in the HEL cell megakaryoblastic cell line (Deveaux et al., 1997). In platelets, this enzyme is responsible for the conversion of prostaglandin endoperoxides to thromboxane A₂. Since thromboxane A₂ is a platelet agonist, a deficiency of thromboxane synthase in NF-E2^{-/-} megakaryocytes could, in theory, help to explain the inside out signaling defect in these cells. However, this is unlikely for several reasons. First, most investigators including ourselves have found that agonists such as thrombin, ADP, and epinephrine stimulate little or no fibrinogen binding to HEL cells or to other megakaryoblastic cell lines. Thus, although these cells express thromboxane synthase, they appear to lack critical intermediates required for inside out signaling (Jennings et al., 1996; Cichowski et al., 1999). Second, we found that large megakaryocytes from wild-type mice bound fibrinogen in response to PAR4 peptide, ADP, and epinephrine even after treatment with aspirin, which blocks cyclooxygenase and subsequent production of thromboxane A₂ (Roth et al., 1975). Third, thromboxane A₂ is not involved in the initial activation of α IIB β 3 in platelets, but rather reinforces the responses to other agonists (Thomas et al., 1998). It is now feasible, and may be preferable, to use primary megakaryocytes instead of megakaryoblastic cell lines to identify NF-E2-regulated genes involved in α IIB β 3 signaling.

Sindbis virus-mediated expression of Tac- β 3 integrin cytoplasmic tails in megakaryocytes caused dose-dependent inhibition of agonist-induced fibrinogen binding to α IIB β 3 (Fig. 6). Although expressed to the same extent as Tac- β 3, Tac- α 5 caused significantly less inhibition of agonist-induced fibrinogen binding to megakaryocytes ($P < 0.01$), and the extent of inhibition was identical to that observed with a tailless Tac construct. The marked inhibitory effect of Tac- β 3 is consistent with the suggestion that overexpressed β 3 tails may bind and titrate cytoplasmic proteins that otherwise would interact with α IIB β 3 to regulate receptor affinity or avidity (Chen et al., 1994; LaFlamme et al., 1994). Several proteins have been shown to interact with the β 3 tail in model systems and by affinity chromatography, including cytoskeletal proteins such as talin, α -actinin, filamin, and myosin, a 14-kD polypeptide called β 3-endonexin, and the cytoplasmic tail of α IIB (Otey et al., 1993; Haas and Plow, 1996; Hemler, 1998; Jenkins et al., 1998; Pfaff et al., 1998). In principle, any or all of these interactions with the β 3 tail might influence the conformation or oligomerization state of α IIB β 3 and, therefore, its ligand binding properties (Kucik et al., 1996; Kashiwagi et al., 1997; Sampath et al., 1998; Bennett et al., 1999).

Mechanisms of Outside In Signaling in Megakaryocytes

The adhesion of platelets to surfaces coated with fibrinogen or von Willebrand factor triggers tyrosine phosphorylation of numerous proteins, cell spreading, and actin rearrangements (Haimovich et al., 1993; Heemskerk et al., 1997; Yuan et al., 1997; Leng et al., 1998). In the current study, we found that murine megakaryocytes also adhere to fibrinogen via α IIB β 3 and undergo spreading and cy-

toskeletal reorganization in an α IIB β 3-dependent fashion, particularly in the presence of a costimulus like phorbol myristate acetate (Fig. 4). This response is similar in some respects to the morphological changes and actin rearrangements in fibroblasts that are triggered by the combination of cell adhesion via β 1 integrins and stimulation by growth factors (Burrige and Chrzanowska-Wodnicka, 1996). β 1 integrins (α 4 β 1 and α 5 β 1) have been demonstrated to play key roles in hematopoietic cell development, presumably by mediating outside in signaling (Verfaillie, 1998). In addition, β 1 integrins in mature human or guinea pig megakaryocytes can support attachment and spreading on fibronectin and focal adhesion formation (Berthier et al., 1998; Schick et al., 1998). In contrast, other than regulation of fibrinogen uptake into α -granules, very little is known about the function of α IIB β 3 in megakaryocytes (Handagama et al., 1993).

α IIB β 3 signaling cannot be essential for megakaryocytopoiesis or thrombopoiesis because platelet numbers are normal in mice and humans deficient in α IIB β 3 (Hodivala-Dilke et al., 1999). This is not to say, however, that α IIB β 3 plays no role in these processes. For example, proplatelet and platelet formation may involve a series of membrane protrusive events which could involve integrin-triggered cytoskeletal reorganization (Choi et al., 1995; Lecine et al., 1998b). In support of this, certain anti- α IIB β 3 antibodies can inhibit proplatelet formation by cultured megakaryocytes (Takahashi et al., 1999). The formation of filopodia, lamellipodia, and focal adhesions in fibrinogen-adherent megakaryocytes (Fig. 4) suggests that signals from α IIB β 3 may activate Rho family GTPases, including cdc42, Rac, and Rho (Hall, 1998). Rho in particular is implicated by the reduction in focal adhesions and stress fibers in megakaryocytes transduced with C3 exoenzyme (Fig. 7). Since Rho GTPases may become activated during integrin-mediated cell adhesion (Ren et al., 1999) and signal to the cytoskeleton and nucleus (Hall, 1998; Treisman et al., 1998), we suggest that signals downstream of α IIB β 3 may play some adjunctive role in megakaryocyte or platelet development. In this context, antibody or peptide blockade of ligand binding to α IIB β 3 has been shown to impair human megakaryocyte colony formation in a fibrin gel (Zauli et al., 1997a), and treatment of a human megakaryoblastic cell line with C3 exoenzyme caused an increase in DNA ploidy (Takada et al., 1996). Further studies of the role of α IIB β 3 signaling in megakaryocytes are warranted.

Retroviruses and lentiviruses are being evaluated as transfer vectors for gene therapy, and they also hold promise for studies of megakaryocyte function (Burstein et al., 1999; Miyoshi et al., 1999; Murphy and Leavitt, 1999; Wilcox et al., 1999). The present experiments show that Sindbis virus vectors provide an alternative way to achieve transient expression of recombinant proteins in terminally differentiated hematopoietic cells like megakaryocytes. Transient expression may be particularly useful in situations where long-term expression of a recombinant protein might interfere with cell growth or differentiation, as with C3 exoenzyme and Tac- β 3 (Henning et al., 1997; Lukashev et al., 1994). Taken together with new strategies to achieve megakaryocyte-specific gene expression *in vivo* (Murphy and Leavitt, 1999), several complementary approaches are now available to manipulate gene expression

in megakaryocytes, establishing these cells as an ideal system for determining the molecular basis of α IIB β 3 signaling.

The authors are grateful to Dr. Kenneth Kaushansky and his laboratory for generous help in setting up the megakaryocyte culture system. We thank Drs. Ramesh Shivdasani and Stuart Orkin for the NF-E2^{-/-} mice, Drs. Takashi Kato and Yoshihiro Tanaka for TPO, Dr. V. Ramakrishnan for the anti-GP Ib and GP V antibodies, Drs. Barry Collier and Susan Smyth for antibody 1B5, Dr. James H. Strauss for antibody to Sindbis E1, and Drs. Chang S. Hahn and Martin A. Schwartz for the Sindbis/CAT and Sindbis C3 vectors.

These studies were supported by research grants from the National Institutes of Health (HL 56595, HL 48728, HL 54476).

Submitted: 19 September 1999

Revised: 4 November 1999

Accepted: 24 November 1999

References

- Adler, M., R.A. Lazarus, M.S. Dennis, and G. Wagner. 1991. Solution structure of kistrin, a potent platelet aggregation inhibitor and GP IIb-IIIa antagonist. *Science*. 253:445-448.
- Aktories, K., C. Mohr, and G. Koch. 1992. *Clostridium botulinum* C3 ADP-ribosyltransferase. *Curr. Topics Microbiol. Immunol.* 175:115-131.
- Andrews, N.C. 1998. The NF-E2 transcription factor. *Int. J. Biochem. Cell Biol.* 30:429-432.
- Aszódi, A., A. Pfeifer, M. Ahmad, M. Glauner, X.H. Zhou, L. Ny, K.E. Andersson, B. Kehrel, S. Offermanns, and R. Fässler. 1999. The vasodilator-stimulated phosphoprotein (VASP) is involved in cGMP- and cAMP-mediated inhibition of agonist-induced platelet aggregation, but is dispensable for smooth muscle function. *EMBO (Eur. Mol. Biol. Organ.) J.* 18:37-48.
- Bazzoni, G., and M.E. Hemler. 1998. Are changes in integrin affinity and conformation overemphasized? *Trends Biochem. Sci.* 23:30-34.
- Bennett, J.S., S. Zigmund, G. Vilaire, M. Cunningham, and B. Bednar. 1999. The platelet cytoskeleton regulates the affinity of the integrin α IIB β 3 for fibrinogen. *J. Biol. Chem.* 274:25301-25307.
- Berthier, R., M. Jacquier-Sarlin, A. Schweitzer, M.R. Block, and A. Molla. 1998. Adhesion of mature polyploid megakaryocytes to fibronectin is mediated by beta 1 integrins and leads to cell damage. *Exp. Cell Res.* 242:315-327.
- Bobak, D., J. Moorman, A. Guanzon, L. Gilmer, and C. Hahn. 1997. Inactivation of the small GTPase Rho disrupts cellular attachment and induces adhesion-dependent and adhesion-independent apoptosis. *Oncogene*. 15:2179-2189.
- Burridge, K., and M. Chrzanowska-Wodnicka. 1996. Focal adhesions, contractility, and signaling. *Annu. Rev. Cell Biol.* 12:463-519.
- Burstein, S.A., A. Dubart, F. Norol, N. Debili, P. Friese, T. Downs, X. Yu, P.W. Kincade, J.L. Villeval, and W. Vainchenker. 1999. Expression of a foreign protein in human megakaryocytes and platelets by retrovirally mediated gene transfer. *Exp. Hematol.* 27:110-116.
- Chen, Y.-P., T.E. O'Toole, T. Shipley, J. Forsyth, S.E. LaFlamme, K.M. Yamada, S.J. Shattil, and M.H. Ginsberg. 1994. "Inside-out" signal transduction inhibited by isolated integrin cytoplasmic domains. *J. Biol. Chem.* 269:18307-18310.
- Choi, E.S., J.L. Nichol, M.M. Hokom, A.C. Hornkohl, and P. Hunt. 1995. Platelets generated in vitro from proplatelet-displaying human megakaryocytes are functional. *Blood*. 85:402-413.
- Cichowski, K., M.J. Orsini, and L.F. Brass. 1999. PAR1 activation initiates integrin engagement and outside-in signalling in megakaryoblastic CHRF-288 cells. *Biochim. Biophys. Acta Mol. Cell Res.* 1450:265-276.
- Deveaux, S., S. Cohen-Kaminsky, R.A. Shivdasani, N.C. Andrews, A. Filipe, I. Kuzniak, S.H. Orkin, P.H. Romeo, and V. Mignotte. 1997. p45 NF-E2 regulates expression of thromboxane synthase in megakaryocytes. *EMBO (Eur. Mol. Biol. Organ.) J.* 16:5654-5661.
- Fox, J.E.B. 1996. Platelet activation: new aspects. *Haemostasis*. 26(Suppl.): 102-131.
- Ginsberg, M.H., X. Du, and E.F. Plow. 1992. Inside-out integrin signalling. *Curr. Opin. Cell Biol.* 4:766-771.
- Haas, T.A., and E.F. Plow. 1996. The cytoplasmic domain of α IIB β 3. A ternary complex of the integrin alpha and beta subunits and a divalent cation. *J. Biol. Chem.* 271:6017-6026.
- Hagmann, J., M. Grob, A. Welman, G. van Willigen, and M.M. Burger. 1998. Recruitment of the LIM protein hic-5 to focal contacts of human platelets. *J. Cell Sci.* 111:2181-2188.
- Hahn, C.S., Y.S. Hahn, T.J. Braciale, and C.M. Rice. 1992. Infectious Sindbis virus transient expression vectors for studying antigen processing and presentation. *Proc. Natl. Acad. Sci. USA*. 89:2679-2683.
- Haimovich, B., L. Lipfert, J.S. Brugge, and S.J. Shattil. 1993. Tyrosine phosphorylation and cytoskeletal reorganization in platelets are triggered by interaction of integrin receptors with their immobilized ligands. *J. Biol. Chem.* 268:15868-15877.
- Hall, A. 1998. Rho GTPases and the actin cytoskeleton. *Science*. 279:509-514.
- Handagama, P., R.M. Scarborough, M.A. Shuman, and D.F. Bainton. 1993. Endocytosis of fibrinogen into megakaryocyte and platelet α -granules is mediated by α IIB β 3 (glycoprotein IIb-IIIa). *Blood*. 82:135-138.
- Hartwig, J.H., S. Kung, T. Kovacsovic, P.A. Janney, L.C. Cantley, T.P. Stossel, and A. Toker. 1996. D3 phosphoinositides and outside-in integrin signaling by glycoprotein IIb-IIIa mediate platelet actin assembly and filopodial extension induced by phorbol 12-myristate 13-acetate. *J. Biol. Chem.* 271:32986-32993.
- Hato, T., N. Pampori, and S.J. Shattil. 1998. Complementary roles for receptor clustering and conformational change in the adhesive and signaling functions of integrin α IIB β 3. *J. Cell Biol.* 141:1685-1695.
- Hauser, W., K.P. Nobeloch, M. Eigenthaler, S. Gambaryan, V. Krenn, J. Geiger, M. Glazova, E. Rohde, I. Horak, U. Walter, and M. Zimmer. 1999. Megakaryocyte hyperplasia and enhanced agonist-induced platelet activation in vasodilator-stimulated phosphoprotein knockout mice. *Proc. Natl. Acad. Sci. USA*. 96:8120-8125.
- Hawiger, J., S. Parkinson, and S. Timmons. 1980. Prostacyclin inhibits mobilisation of fibrinogen-binding sites on human ADP- and thrombin-treated platelets. *Nature*. 283:195-197.
- Heemskerk, J.W.M., W.M.J. Vuist, M.A.H. Feijge, C.P.M. Reutelingsperger, and T. Lindhout. 1997. Collagen but not fibrinogen surfaces induce bleb formation, exposure of phosphatidylserine, and procoagulant activity of adherent platelets: evidence for regulation by protein tyrosine kinase-dependent Ca²⁺ responses. *Blood*. 90:2615-2625.
- Hemler, M.E. 1998. Integrin associated proteins. *Curr. Opin. Cell Biol.* 10:578-585.
- Henning, S.W., R. Galandrini, A. Hall, and D.A. Cantrell. 1997. The GTPase Rho has a critical regulatory role in thymus development. *EMBO (Eur. Mol. Biol. Organ.) J.* 16:2397-2407.
- Hodivala-Dilke, K.M., K.P. McHugh, D.A. Tsakiris, H. Rayburn, D. Crowley, M. Ullman-Cullere, F.P. Ross, B.S. Collier, S. Teitelbaum, and R.O. Hynes. 1999. Beta3-integrin-deficient mice are a model for Glanzmann thrombasthenia showing placental defects and reduced survival. *J. Clin. Invest.* 103:229-238.
- Holmbäck, K., M.J.S. Danton, T.T. Suh, C.C. Daugherty, and J.L. Degen. 1996. Impaired platelet aggregation and sustained bleeding in mice lacking the fibrinogen motif bound by integrin α IIB β 3. *EMBO (Eur. Mol. Biol. Organ.) J.* 15:5760-5771.
- Jenkins, A.L., L. Nannizzi-Alaimo, D. Silver, J.R. Sellers, M.H. Ginsberg, D.A. Law, and D.R. Phillips. 1998. Tyrosine phosphorylation of the β 3 cytoplasmic domain mediates integrin-cytoskeletal interactions. *J. Biol. Chem.* 273:13878-13885.
- Jennings, L.K., S.M. Slack, C.D. Wall, and T.H. Mondoro. 1996. Immunological comparisons of integrin alpha IIB beta 3 (GPIIb-IIIa) expressed on platelets and human erythroleukemia cells: evidence for cell specific differences. *Blood Cells Mol. Dis.* 22:23-35.
- Kahn, M.L., Y.W. Zheng, W. Huang, V. Bigornia, D.W. Zeng, S. Moff, R.V. Farese, Jr., C. Tam, and S.R. Coughlin. 1998. A dual thrombin receptor system for platelet activation. *Nature*. 394:690-694.
- Kashiwagi, H., M.A. Schwartz, M.A. Eigenthaler, K.A. Davis, M.H. Ginsberg, and S.J. Shattil. 1997. Affinity modulation of platelet integrin α IIB β 3 by β 3-endonexin, a selective binding partner of the β 3 integrin cytoplasmic tail. *J. Cell Biol.* 137:1433-1443.
- Kaushansky, K. 1995. Thrombopoietin: The primary regulator of platelet production. *Blood*. 86:419-431.
- Kaushansky, K. 1999. The enigmatic megakaryocyte gradually reveals its secrets. *Bioessays*. 21:353-360.
- Kucik, D.F., M.L. Dustin, J.M. Miller, and E.J. Brown. 1996. Adhesion-activating phorbol ester increases the mobility of leukocyte integrin LFA-1 in cultured lymphocytes. *J. Clin. Invest.* 97:2139-2144.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227:680-685.
- LaFlamme, S.E., L.A. Thomas, S.S. Yamada, and K.M. Yamada. 1994. Single subunit chimeric integrins as mimics and inhibitors of endogenous integrin functions in receptor localization, cell spreading and migration, and matrix assembly. *J. Cell Biol.* 126:1287-1298.
- Law, D.A., L. Nannizzi-Alaimo, K. Ministri, P. Hughes, J. Forsyth, M. Turner, S.J. Shattil, M.H. Ginsberg, V. Tybulewicz, and D.R. Phillips. 1999. Genetic and pharmacological analyses of Syk function in α IIB β 3 signaling in platelets. *Blood*. 93:2645-2652.
- Lecine, P., V. Blank, and R. Shivdasani. 1998a. Characterization of the hematopoietic transcription factor NF-E2 in primary murine megakaryocytes. *J. Biol. Chem.* 273:7572-7578.
- Lecine, P., J.L. Villeval, P. Vyas, B. Swencki, Y. Xu, and R.A. Shivdasani. 1998b. Mice lacking transcription factor NF-E2 provide in vivo validation of the proplatelet model of thrombocytopoiesis and show a platelet production defect that is intrinsic to megakaryocytes. *Blood*. 92:1608-1616.
- Leng, L., H. Kashiwagi, X.-D. Ren, and S.J. Shattil. 1998. RhoA and the function of platelet integrin α IIB β 3. *Blood*. 91:4206-4215.
- Lengweiler, S., S.S. Smyth, M. Jirouskova, L.E. Scudder, H. Park, T. Moran, and B.S. Collier. 1999. Preparation of monoclonal antibodies to murine platelet glycoprotein IIb/IIIa (alphaIIb beta3) and other proteins from hamster-mouse interspecies hybridomas. *Biochem. Biophys. Res. Commun.* 262:167-173.

- Leven, R.M., and V.T. Nachmias. 1982. Cultured megakaryocytes: changes in the cytoskeleton after adenosine diphosphate-induced spreading. *J. Cell Biol.* 92:313–323.
- Long, M.W. 1998. Megakaryocyte differentiation events. *Semin. Hematology.* 35:192–199.
- Lukashev, M.E., D. Sheppard, and R. Pytela. 1994. Disruption of integrin function and induction of tyrosine phosphorylation by the autonomously expressed β_1 integrin cytoplasmic domain. *J. Biol. Chem.* 269:18311–18314.
- Miyoshi, H., K.A. Smith, D.E. Mosier, I.M. Verma, and B.E. Torbett. 1999. Transduction of human CD34+ cells that mediate long-term engraftment of NOD/SCID mice by HIV vectors. *Science* 283:682–686.
- Moorman, J.P., D. Luu, J. Wickham, D.A. Bobak, and C.S. Hahn. 1999. A balance of signaling by Rho family small GTPases RhoA, Rac1 and Cdc42 coordinates cytoskeletal morphology but not cell survival. *Oncogene.* 18:47–57.
- Mountford, J.C., S.K. Melford, C.M. Bunce, J. Gibbons, and S.P. Watson. 1999. Collagen or collagen-related peptide cause $[Ca^{2+}]_i$ elevation and increased tyrosine phosphorylation in human megakaryocytes. *Thromb Haemost.* 82:1153–1159.
- Murphy, G.J., and A.D. Leavitt. 1999. A model for studying megakaryocyte development and biology. *Proc. Natl. Acad. Sci. USA.* 96:3065–3070.
- Nachmias, V.T., and R. Golla. 1991. Vinculin in relation to stress fibers in spread platelets. *Cell Motil. Cytoskel.* 20:190–202.
- O'Toole, T.E., Y. Katagiri, R.J. Faull, K. Peter, R. Tamura, V. Quaranta, J.C. Loftus, S.J. Shattil, and M.H. Ginsberg. 1994. Integrin cytoplasmic domains mediate inside out signaling. *J. Cell Biol.* 124:1047–1059.
- Offermanns, S., C.F. Toombs, Y.H. Hu, and M.I. Simon. 1997. Defective platelet activation in $G\alpha_q$ -deficient mice. *Nature.* 389:183–186.
- Otey, C.A., G.B. Vasquez, K. Burridge, and B.W. Erickson. 1993. Mapping of the α -actinin binding site within the β_1 integrin cytoplasmic domain. *J. Biol. Chem.* 268:21193–21197.
- Pampori, N., T. Hato, D.G. Stupack, S. Aidoudi, D.A. Cheresch, G.R. Nemerow, and S.J. Shattil. 1999. Mechanisms and consequences of affinity modulation of integrin $\alpha_v\beta_3$ detected with a novel patch-engineered monovalent ligand. *J. Biol. Chem.* 274:21609–21616.
- Pfaff, M., S. Liu, D.J. Erle, and M.H. Ginsberg. 1998. Integrin beta cytoplasmic domains differentially bind to cytoskeletal proteins. *J. Biol. Chem.* 273:6104–6109.
- Qi, W., E. Loh, G. Vilaire, and J.S. Bennett. 1998. Regulation of alphaIIb beta3 function in human B lymphocytes. *J. Biol. Chem.* 273:15271–15278.
- Ren, X.D., W.B. Kiosses, and M.A. Schwartz. 1999. Regulation of the small GTP-binding protein Rho by cell adhesion and the cytoskeleton. *EMBO (Eur. Mol. Biol. Organ.) J.* 18:578–585.
- Rittenhouse, S.E. 1996. Phosphoinositide 3-kinase activation and platelet function. *Blood.* 88:4401–4414.
- Riviere, C., F. Subra, K. Cohen-Solal, V. Cordette-Lagarde, R. Letestu, C. Auclair, W. Vainchenker, and F. Louache. 1999. Phenotypic and functional evidence for the expression of CXCR4 receptor during megakaryocytopoiesis. *Blood.* 93:1511–1523.
- Rojnuckarin, P., J.G. Drachman, and K. Kaushansky. 1999. Thrombopoietin-induced activation of the mitogen-activated protein kinase (MAPK) pathway in normal megakaryocytes: role in endomitosis. *Blood.* 94:1273–1282.
- Roth, G.J., N. Stanford, and P.W. Majerus. 1975. Acetylation of prostaglandin synthase by aspirin. *Proc. Natl. Acad. Sci. USA.* 72:3073–3076.
- Sampath, R., P.J. Gallagher, and F.M. Pavalko. 1998. Cytoskeletal interactions with the leukocyte integrin β_2 cytoplasmic tail—activation-dependent regulation of associations with talin and α -actinin. *J. Biol. Chem.* 273:33588–33594.
- Savage, B., S.J. Shattil, and Z.M. Ruggeri. 1992. Modulation of platelet function through adhesion receptors. A dual role for glycoprotein IIb-IIIa (integrin $\alpha_{IIb}\beta_3$) mediated by fibrinogen and glycoprotein Ib-von Willebrand factor. *J. Biol. Chem.* 267:11300–11306.
- Schick, P.K., C.M. Wojenski, X. He, J. Walker, C. Marcinkiewicz, and S. Niewiarowski. 1998. Integrins involved in the adhesion of megakaryocytes to fibronectin and fibrinogen. *Blood.* 92:2650–2656.
- Schoenwaelder, S.M., S.P. Jackson, Y. Yuan, M.S. Teasdale, H.H. Salem, and C.A. Mitchell. 1994. Tyrosine kinases regulate the cytoskeletal attachment of integrin $\alpha_{IIb}\beta_3$ (platelet glycoprotein IIb/IIIa) and the cellular retraction of fibrin polymers. *J. Biol. Chem.* 269:32479–32487.
- Shattil, S.J., M. Cunningham, and J.A. Hoxie. 1987. Detection of activated platelets in whole blood using activation-dependent monoclonal antibodies and flow cytometry. *Blood.* 70:307–315.
- Shattil, S.J., H. Kashiwagi, and N. Pampori. 1998. Integrin signaling: the platelet paradigm. *Blood.* 91:2645–2657.
- Shivdasani, R.A., M.F. Rosenblatt, D. Zucker-Franklin, C.W. Jackson, P. Hunt, C.J. Saris, and S.H. Orkin. 1995. Transcription factor NF-E2 is required for platelet formation independent of the actions of thrombopoietin/MGDF in megakaryocyte development. *Cell.* 81:695–704.
- Takada, M., N. Morii, S. Kumagai, and R. Ryo. 1996. The involvement of the *rho* gene product, a small molecular weight GTP-binding protein, in ploidy of a human megakaryocytic cell line, CMK. *Exp. Hematol.* 24:524–530.
- Takahashi, R., N. Sekine, and T. Nakatake. 1999. Influence of monoclonal anti-platelet glycoprotein antibodies on in vitro human megakaryocyte colony formation and proplatelet formation. *Blood.* 93:1951–1958.
- Thomas, D.W., R.B. Mannon, P.J. Mannon, A. Latour, J.A. Oliver, M. Hoffman, O. Smithies, B.H. Koller, and T.M. Coffman. 1998. Coagulation defects and altered hemodynamic responses in mice lacking receptors for thromboxane A_2 . *J. Clin. Invest.* 102:1994–2001.
- Treisman, R., A.S. Alberts, and E. Sahai. 1998. Regulation of SRF activity by Rho family GTPases. *Cold Spring Harbor Symp. Quant. Biol.* 63:643–651.
- Tsakiris, D.A., L. Scudder, K. Hodivala-Dilke, R.O. Hynes, and B.S. Coller. 1999. Hemostasis in the mouse (*Mus musculus*): a review. *Thromb. Haemost.* 81:177–188.
- Verfaillie, C.M. 1998. Adhesion receptors as regulators of the hematopoietic process. *Blood.* 92:2609–2612.
- Vyas, P., K. Ault, C.W. Jackson, S.H. Orkin, and R.A. Shivdasani. 1999. Consequences of GATA-1 deficiency in megakaryocytes and platelets. *Blood.* 93:2867–2875.
- Wang, J.F., Z.Y. Liu, and J.E. Groopman. 1998. The α -chemokine receptor CXCR4 is expressed on the megakaryocytic lineage from progenitor to platelets and modulates migration and adhesion. *Blood.* 92:756–764.
- Wilcox, D.A., J.C. Olsen, L. Ishizawa, M. Griffith, and G.C. White II. 1999. Integrin alphaIIb promoter-targeted expression of gene products in megakaryocytes derived from retrovirus-transduced human hematopoietic cells. *Proc. Natl. Acad. Sci. USA.* 96:9654–9659.
- Williams, J.L., G.G. Pipia, N.S. Datta, and M.W. Long. 1998. Thrombopoietin requires additional megakaryocyte-active cytokines for optimal ex vivo expansion of megakaryocyte precursor cells. *Blood.* 91:4118–4126.
- Yuan, Y., S.M. Dopheide, C. Ivanidis, H.H. Salem, and S.P. Jackson. 1997. Calpain regulation of cytoskeletal signaling complexes in von Willebrand factor-stimulated platelets. Distinct roles for glycoprotein Ib-V-IX and glycoprotein IIb-IIIa (integrin alphaIIb beta3) in von Willebrand factor-induced signal transduction. *J. Biol. Chem.* 272:21847–21854.
- Zauli, G., A. Bassini, M. Vitale, D. Gibellini, C. Celeghini, E. Caramelli, S. Pierpaoli, L. Guidotti, and S. Capitani. 1997a. Thrombopoietin enhances the $\alpha_{IIb}\beta_3$ -dependent adhesion of megakaryocytic cells to fibrinogen or fibronectin through PI 3 kinase. *Blood.* 89:883–895.
- Zauli, G., M. Vitale, E. Falcieri, D. Gibellini, A. Bassini, C. Celeghini, M. Columbaro, and S. Capitani. 1997b. In vitro senescence and apoptotic cell death of human megakaryocytes. *Blood.* 90:2234–2243.