


Sesaki and Jensen Vol. 152, No. 6, March 19, 2001. Pages 1123–1134.

Due to a printing error, Fig. 5 reproduced poorly in the *Journal* and information was lost. The corrected Fig. 5 appears below:



*Figure 5. ugo1* $\Delta$  *dnm1* $\Delta$  cells contain mitochondrial tubules and maintain mtDNA. Wild-type (FY833, WT), *ugo1* $\Delta$  (YHS72), *dnm1* $\Delta$  (YHS83), and *ugo1* $\Delta$  *dnm1* $\Delta$  (YHS85) cells expressing OM45-GFP (pKC2) were grown to log phase in SRaf medium, stained with 1  $\mu$ g/ml DAPI, and viewed by DIC and fluorescence (OM45-GFP) microscopy. (B) Wild-type, *ugo1* $\Delta$ , *dnm1* $\Delta$ , and *ugo1* $\Delta$  *dnm1* $\Delta$  cells were grown to log phase in YEPD medium. Cells were collected and resuspended in YEPD medium to an OD<sub>600</sub> of 2. Cells were then diluted in 10-fold increments, and 10  $\mu$ l of each dilution was spotted onto YEPD and YEPGE media and incubated at 30°C for 2 and 6 d, respectively. N, nuclear DNA staining. Bar, 3  $\mu$ m.