

# Control of axon elongation via an SDF-1 $\alpha$ /Rho/mDia pathway in cultured cerebellar granule neurons

Yoshiki Arakawa,<sup>1,2</sup> Haruhiko Bito,<sup>1,3,4</sup> Tomoyuki Furuyashiki,<sup>1</sup> Takahiro Tsuji,<sup>1</sup> Sayaka Takemoto-Kimura,<sup>1</sup> Kazuhiro Kimura,<sup>1</sup> Kazuhiko Nozaki,<sup>2</sup> Nobuo Hashimoto,<sup>2</sup> and Shuh Narumiya<sup>1</sup>

<sup>1</sup>Department of Pharmacology and <sup>2</sup>Department of Neurosurgery, Kyoto University Faculty of Medicine, and <sup>3</sup>PRESTO-Japan Science and Technology Corporation, Sakyo-ku, Kyoto 606-8315, Japan

<sup>4</sup>Department of Neurochemistry, University of Tokyo Graduate School of Medicine, Bunkyo-ku, Tokyo 113-0033, Japan

Rho-GTPase has been implicated in axon outgrowth. However, not all of the critical steps controlled by Rho have been well characterized. Using cultured cerebellar granule neurons, we show here that stromal cell-derived factor (SDF)-1 $\alpha$ , a neural chemokine, is a physiological ligand that can turn on two distinct Rho-dependent pathways with opposite consequences. A low concentration of the ligand stimulated a Rho-dependent pathway that mediated facilitation of axon elongation. In contrast, Rho/ROCK activation achieved by a higher concentration of SDF-1 $\alpha$  caused repression of axon formation

and induced no more increase in axon length. However, even at this higher concentration a Rho-dependent axon elongating activity could be recovered upon removal of ROCK activity using Y-27632. SDF-1 $\alpha$ -induced axon elongating activity under ROCK inhibition was replicated by the dominant-active form of the mammalian homologue of the *Drosophila* gene Diaphanous (mDia)1 and counteracted by its dominant-negative form. Furthermore, RNAi knock-down of mDia1 abolished SDF-1 $\alpha$ -induced axon elongation. Together, our results support a critical role for an SDF-1 $\alpha$ /Rho/mDia1 pathway in mediating axon elongation.

## Introduction

It has been widely accepted that rearrangement of actin and microtubule cytoskeleton lies at the heart of neuronal morphogenesis (Tanaka and Sabry, 1995). Drastic changes in neuronal shape occur immediately after the exit of neuronal cells from mitotic cycles, during embryonic and postnatal development. Process formation/extension and cell body migration must be spatially and temporally orchestrated in order to achieve patterned formation of neuronal cell layers and appropriate generation of synaptic circuits (Goodman and Shatz, 1993; Tessier-Lavigne and Goodman, 1996; Van Vactor and Flanagan, 1999).

A dynamic morphological alteration is initiated during the acquisition of neuronal polarity and must continue until the completion of synaptogenesis. Recent findings indicate a critical role for the antagonism between Rac- and Rho-GTPases in these events (Narumiya et al., 1997; Hall, 1998; Luo, 2000; Dickson, 2001). A large body of work has now established that several soluble or transmembranous guidance molecule systems can exhibit either chemorepulsion or chemoattraction

toward axonal growth cones, at least in part, via coupling to either Rho or Rac, respectively. Such systems include EphrinA (Shamah et al., 2001), EphB reverse signaling (Lu et al., 2001), Semaphorin4D/PlexinB (Driessens et al., 2001; Swiercz et al., 2002), Semaphorin3A/PlexinA-Neuropilin (Jin and Strittmatter, 1997; Liu and Strittmatter, 2001; Journey et al., 2002), Netrin/DCC (Li et al., 2002), Slit/Robo (Wong et al., 2001), and neurotrophins/p75/trk (Yamashita et al., 1999; Huang and Reichardt, 2001; Ozdinler and Erzurumlu, 2001; Nusser et al., 2002). Thus, antagonistic interactions and hierarchical cascades between distinct small GTPases had been considered as most likely ways by which distinct gradient cues could be decoded into reliable maps of afferents projecting into one specific area of the central nervous system (CNS).\*

However, to date, a clear understanding regarding what specific effectors of the small GTPases contribute to each of these opposing signaling events is still missing. Furthermore, whether Rho always antagonized with Rac remains controversial (Sebok et al., 1999; Bashaw et al., 2001).

The online version of this article includes supplemental material.

Address correspondence to Shuh Narumiya, Dept. of Pharmacology, Kyoto University Faculty of Medicine, Yoshida, Sakyo-ku, Kyoto 606-8315, Japan. Tel.: 81-75-753-4392. Fax: 81-75-753-4693. E-mail: snaru@mfour.med.kyoto-u.ac.jp

Key words: mDia; Rho; axon elongation; cerebellar granule neuron; SDF-1 $\alpha$

\*Abbreviations used in this paper: CRIB, Cdc42/Rac interactive binding; CNS, central nervous system; DA, dominant-active; DN, dominant-negative; EGFP, enhanced GFP; EGL, external granule cell layer; IGL, internal granule cell layer; mDia, mammalian homologue of the *Drosophila* gene Diaphanous; P, postnatal day; PAK, p21-associated kinase; RBD, Rho-binding domain; RNAi, RNA interference; siRNA, short interfering double stranded RNA oligomer; SDF, stromal cell-derived factor.

The rodent cerebellar granule cells represent a highly advantageous resource to study in detail the molecular sequence of events controlling axonogenesis, neuronal migration, dendritogenesis, and synaptogenesis. Indeed, each of these steps that is critical for neuronal maturation and circuit formation occurs in an organized and sequential fashion in the cerebellum during the postnatal 3 wk after birth (Altman and Bayer, 1996; Hatten, 1999). Many of these steps involve dramatic changes in neuronal morphology, yet previous studies indicated that many of their features could be recapitulated, at least in part, in primary culture (Powell et al., 1997; Bito et al., 2000; Yamasaki et al., 2001; Yacubova and Komuro, 2002).


Previous investigation of molecular determinants involved in axon formation in cerebellar granule cells, from our laboratory and others, have begun to uncover the critical importance of pia-granule cell interaction (Bhatt et al., 2000) on one hand and control of actin dynamics on the other (Bito et al., 2000; Yamasaki et al., 2001). Thus, we considered the possibility that a factor supplied by the pia mater may be involved in the signaling pathway controlling axon initiation and elongation. One candidate factor we tested was stromal cell-derived factor (SDF)-1 $\alpha$ , a chemokine heavily expressed in the pia mater and chemoattractant for migration of cerebellar granule cells (Lu et al., 2001; Klein et al., 2001; Tham et al., 2001; Zhu et al., 2002). Our interest was prompted by the striking phenotype of the knockout mice lacking either SDF-1 $\alpha$  (Ma et al., 1998) or its cognate receptor CXCR4 (Ma et al., 1998; Zou et al., 1998). In both lines of mice, the formation of the cerebellar granule cell layers was abolished. This suggested that early events in cerebellar granule cell morphogenesis, such as axon initiation, axon elongation, or migration of the cell body, might be severely perturbed in the absence of SDF-1 $\alpha$  signaling.

However, whether SDF-1 $\alpha$  could stimulate axon formation via direct control of actin cytoskeletal signaling mechanisms in cerebellar granule cells has not been demonstrated. Here we show that SDF-1 $\alpha$  activates Rho but not Rac small GTPase in cerebellar granule cells. A low concentration (100 ng/ml) of SDF-1 $\alpha$  induced significant amount of axon outgrowth in a C3 exoenzyme-sensitive fashion. In contrast, at a higher concentration SDF-1 $\alpha$  rather repressed axon formation in a Rho-dependent manner. However, Y-27632 treatment was sufficient to uncover an axon elongating activity even under high dose of SDF-1 $\alpha$ . Thus, SDF-1 $\alpha$  stimulated two distinct opposing Rho effectors: a novel Rho effector, which promoted axon elongation, and ROCK to inhibit axonogenesis. The facilitatory effect of SDF-1 $\alpha$  in the presence of Y-27632 could be mimicked by overexpressing a dominant-active (DA) form of the mammalian homologue of the *Drosophila* gene Diaphanous (mDia)1, a Rho effector involved in control of actin polymerization during cell polarization and directed cell growth. This dominant activation of mDia was dependent on intact Rac activity. Furthermore, the axon outgrowth induced by SDF-1 $\alpha$  could be antagonized either by overexpression of a dominant interfering mutant or by RNAi knockdown of mDia1. Thus, we identify a novel function for mDia1 as a critical Rho effector-mediating SDF-1 $\alpha$ -dependent axon elongation in concert with Rac.

## Results

### A pial chemokine SDF-1 $\alpha$ triggers axon elongation via Rho early in culture in cerebellar granule cells

We tested whether SDF-1 $\alpha$  was able to trigger any axon growth and whether this morphological change correlated, at least in part, with an alteration in either Rho or Rac activity in cultured cerebellar granule cells, where axonogenesis is well known to precede dendritogenesis. A 12-h exposure to SDF-1 $\alpha$  at a concentration of 100 ng/ml induced a significant increase in the mean length of first appearing process compared with control (Fig. 1, A and B; Video 1, avail-




**Figure 1. SDF-1 $\alpha$  facilitates axon elongation via a Rho-dependent pathway.** (A) Facilitation of axon elongation by 12-h exposure to SDF-1 $\alpha$  (top). SDF-1 $\alpha$ -induced facilitation is blocked by C3 treatment (bottom), indicating the existence of a Rho-dependent mode of neurite extension.  $\beta$ -tubulin immunostaining was employed to completely trace the entire length of axons. (B) Axon elongating activity induced by SDF-1 $\alpha$  reveals a bell-shaped response curve, whereas axon numbers are reduced at only higher concentrations. SDF-1 $\alpha$  effect on axon length and number are both blocked by C3 treatment.  $n \approx 42$ –209. (C) 1- or 12-h exposure to SDF-1 $\alpha$  (100 or 500 ng/ml) leads to a substantial increase in the amount of GTP-bound Rho (RBD pulldown) but not that of GTP-bound Rac (unpublished data) in cerebellar granule cells. \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ . Bar, 5  $\mu$ m.

able at <http://www.jcb.org/cgi/content/full/jcb.200210149/DC1>), while having no effect on axon number (Fig. 1, A and B). This effect was abolished in the presence of C3 exoenzyme, a Rho inhibitor (Fig. 1, A and B). SDF-1 $\alpha$  promoted a bell-shaped response in axon elongation with a peak effect at 100 ng/ml (Fig. 1, A and B). However, at a larger concentration this axon outgrowth effect was abolished and indistinguishable from the control (Fig. 1 B). In contrast, SDF-1 $\alpha$  treatment significantly reduced axon number at concentrations over 250 ng/ml. Presence of C3 exoenzyme completely flattened either response (Fig. 1 B). Together, this indicated that SDF-1 $\alpha$  may promote axon elongation via Rho, at least at lower concentrations, whereas it inhibited initiation of axon at higher concentrations. The reduction in SDF-1 $\alpha$ -dependent axon growth with the maximal concentration of SDF-1 $\alpha$  was unlikely to be caused by receptor desensitization or inactivation, since we confirmed that axon numbers were now negatively affected at the same dose. Furthermore, either 1- or 12-h exposure to either 100 or 500 ng/ml SDF-1 $\alpha$  was accompanied with a similarly strong elevation in the amount of GTP-bound form of Rho (Fig. 1 C), whereas no apparent increase was discerned for GTP-bound Rac (unpublished data) as determined by pull-down assays using either a GST-fused Rho-binding domain (RBD) of Rhotekin or a GST-fused Cdc42/Rac interactive binding (CRIB) domain of p21-associated kinase (PAK) were performed (Ren et al., 1999; Tsuji et al., 2002). These results indicate the possibility that stimulation of Rho pathway with a physiological ligand such as SDF-1 $\alpha$  may mediate axon elongation.

### Antagonism between a Rho-activated axon elongating pathway and ROCK-mediated control of axonogenesis


To test whether the biphasic responsiveness could be accounted for by the existence of distinct thresholds of activation of two separate Rho effectors, we carefully reexamined the dose–response relationship between C3 concentration and resulting axon length. The sampling number was increased substantially in order to detect even small differences that might have been overlooked before. At 10  $\mu$ g/ml of C3, a significant increase in axon length and number was achieved (Fig. 2). When the dose was augmented to 30–50  $\mu$ g/ml, this facilitation in axon length was substantially diminished compared with 10  $\mu$ g/ml treatment, though a small but significant net increase was still detectable; however, no additional effect was seen on axon number (Fig. 2). Therefore, different concentrations of active Rho were likely to gear two distinct pathways, repression and facilitation of axon extension, via two distinct Rho effectors. In contrast, only one Rho effector was likely to contribute to axon number control (Fig. 2), consistent with our previous work (Bito et al., 2000). A similar C3 dose–response curve was also reported in PC-12 cells (Winton et al., 2002).

What are these two Rho effectors? The finding that SDF-1 $\alpha$  seemed to antagonize axonogenesis at higher doses in a Rho-dependent manner (Fig. 1 B) was in keeping with our previous finding that Rho/ROCK signaling critically regulates axon numbers (Bito et al., 2000). To test whether ROCK, a Rho-associated kinase (Narumiya et al., 1997; Bito et al., 2000; Ishizaki et al., 2000), was indeed mediating the SDF-1 $\alpha$  effect, we bath-applied 50  $\mu$ M Y-27632, a potent




**Figure 2. Distinct C3 dose effectiveness on axon elongation and axon number in cultured cerebellar granule neurons.** Axon elongation reveals a bell-shaped C3 responsiveness consistent with the presence of two opposing effects downstream of Rho. However, the C3 effect on axon number is saturated at doses over 10  $\mu$ g/ml.  $n \approx 115$ –424; \*\*\* $P < 0.001$ .

and selective ROCK inhibitor (Uehata et al., 1997; Ishizaki et al., 2000). As expected, treatment with Y-27632 reversed the negative SDF-1 $\alpha$  effect on axon numbers, indicating that SDF-1 $\alpha$  is indeed able to activate ROCK (Fig. 3, A and B). Surprisingly, however, exposure to SDF-1 $\alpha$  in the presence of the ROCK inhibitor now resulted in a significant net increase in axon extension (Fig. 3, A and B). Thus, SDF-1 $\alpha$  activated a novel ROCK-independent, yet C3-sensitive, effector pathway that was coupled to axon elongation.



**Figure 3. SDF-1 $\alpha$ -induced axon elongation is antagonized by ROCK.** 12-h exposure to high concentration (500 ng/ml) of SDF-1 $\alpha$  represses axonogenesis (A, top, and B, right). Addition of 50  $\mu$ M Y-27632, a specific ROCK inhibitor, blocked this SDF-1 $\alpha$ -induced inhibition of axonogenesis (A, bottom, and B, right). Under the same condition, SDF-1 $\alpha$  significantly facilitated axon elongation in ROCK-inhibited neurons (B, left), while having little effect on axon numbers (B, right).  $n \approx 82$ –74. \*\*\*Compared with untreated cells or ## compared with Y-27632-treated cells;  $P < 0.001$ . Bars, 5  $\mu$ m.

**Figure 4. A putative role for mDia1 in facilitated process outgrowth.** (A) Domain structure of wild-type and dominant mutant constructs of mDia1. (B) Coupling of elevated mDia1 activity (by overexpression of mDia1- $\Delta$ N3) with lower ROCK activity (in the presence of Y-27632) is sufficient to induce elongated axon-like processes in Swiss3T3 cells. Note the high amount of F-actin stained with phalloidin in the thin processes (arrowheads) in the thin processes (arrowheads) of the GFP-mDia1- $\Delta$ N3-expressing cells. Bars, 50  $\mu$ m.




### The Rho effector mDia1, a regulator of actin polymerization, is heavily expressed in the external granule cell layer during early postnatal development and positively regulates axon outgrowth

Recently, mDia was found to be a critical Rho effector that may predominantly act as an inducer/regulator of actin polymerization in HeLa cells and may be required for establishment of cell polarity and directed growth (Watanabe et al., 1997; Ishizaki et al., 2001; Ozaki-Kuroda et al., 2001; Tsuji et al., 2002). Indeed, we found that Swiss3T3 cells can elongate prolonged neurite-like processes best when higher mDia activity was achieved by overexpression a DA form of mDia1 (mDia1- $\Delta$ N3; Fig. 4 A; Ishizaki et al., 2001) and was coupled with lower ROCK activity in the presence of 30  $\mu$ M Y-27632 (Fig. 4

B), consistent with previous work from our laboratory (Tsuji et al., 2002).

We also found that expression of mDia1 coincided with early axonogenesis in postnatal cerebellum in mice (Fig. 5, A and B). Immunohistochemical analysis showed that mDia1 was especially abundant in postnatal day 1 (P1) cerebellum at and beneath the external granule cell layer where the earliest events in axonogenesis occurred (Fig. 5 B). In round cerebellar granule cells, mDia1 protein was already colocalized with F-actin and tubulin at spots where an axon was likely to initiate (Fig. 5 C). After axon outgrowth started, mDia1 was heavily enriched at the base of early initiating process and within its growth cones (Fig. 5 C, arrowheads) in close spatial vicinity with actin filaments and microtubules (Figs. 5 C).



**Figure 5. High expression of mDia1 in cerebellar EGL during early postnatal development.** (A) Western blot analysis of mDia1 expression in cerebellar lysates at P1, P2, P9, or in adult (A). (B) mDia1-like immunoreactivity is highly concentrated at and beneath the EGL at P1. TO-PRO nuclear stain indicates the locations of the cells. Bar, 50  $\mu$ m. (C) mDia1 is highly expressed at the neck of a nascent process (top, arrowhead) or in the growth cones (bottom, arrowheads) in cerebellar granule cells in culture at 6 h (top) and 12 h (bottom) in vitro. mDia1 heavily colocalized with F-actin (phalloidin) and microtubules (tubulin) structures. Bars, 5  $\mu$ m.

To ask directly whether mDia1 activity facilitated, at least in part, axon formation and elongation in cerebellar granule neurons, we tested whether expression of the DA-mDia1 mutant was sufficient to replicate the SDF-1 $\alpha$ -induced axon elongating activity in the absence of ROCK pathway which was blocked with Y-27632. Transfection was performed immediately after trituration and during initial plating of the neurons so that expression of exogenous protein was initiated  $\sim$ 6 h after plating (Bito et al., 2000, and this study). Neurons were fixed using PFA at 12 h after transfection to examine the effect of mDia1 activity on axon initiation and elongation. Indeed, mDia1- $\Delta$ N3-expressing cerebellar granule cells revealed a similar number of axons as GFP-expressing control cells; however, the axons were significantly longer compared with the mock-transfected neurons in the presence of Y-27632 (Fig. 6, A–D), consistent with the idea that mDia1 may work downstream of Rho to facilitate axon growth.

During transfection of mDia1- $\Delta$ N3, we noticed that the newly formed processes were abnormal in neurons untreated with Y-27632. Axon outgrowth seemed to be prematurely aborted as processes were filled with an exuberant amount of filamentous actin and  $\beta$ -tubulin (Fig. 6 B, mDia1- $\Delta$ N3),

and thus exhibited an abnormal width (Fig. 6 B and unpublished data). Basal ROCK activity, in the context of excessive mDia1 activity, might account for a prominent increase in actin polymerization (Watanabe et al., 1999), while also sustaining a tonic level of actomyosin contractility, thereby negatively acting on axon elongation.


### Rho/mDia pathway mediates SDF-1 $\alpha$ -stimulated axon elongation in cerebellar granule neurons in part via a coordination with a Rac-dependent pathway

We next examined the contribution of endogenous mDia1 on SDF-1 $\alpha$ -stimulated axon elongation by use of a dominant-negative (DN) form of mDia1, mDia1- $\Delta$ N3(Hind-III) (Fig. 4 A; Tsuji et al., 2002). This mutant was able to abolish the effect of DA-mDia1 on axon length (Fig. 7 A). DN-mDia1 also inhibited axon numbers back to baseline levels (Fig. 7 A); however, since DA-mDia1 had little effect per se (Fig. 6 D), we currently favor the simplest view that rather than acting on the axonogenesis itself, mDia1's axon elongating activity may be required in order to visualize even the smallest process. Consistently, when this DN mutant was transfected in neurons stimulated with SDF-1 $\alpha$  in the presence of Y-27632, both axon number and axon length were significantly diminished (Fig. 7, B and C). Together, these results support the notion that axon elongation induced by SDF-1 $\alpha$  may be regulated in an mDia1-dependent manner.

To obtain an independent confirmation of these results, we applied the RNA interference techniques using short interfering double stranded RNA oligomers (siRNAs) (Elbashir et al., 2001). mDia1-specific siRNA was designed and its efficiency in knocking down mDia1 protein was tested in NIH3T3 cells by Western blot analysis (Fig. 8 A). Loss of mDia1 immunoreactivity was specifically obtained in cells expressing a cotransfection marker enhanced GFP (EGFP; unpublished data). Using identical transfection procedures, EGFP-positive cerebellar granule neurons were screened to identify the neurons, which have taken up the mDia1 siRNA, and axon lengths were measured in these neurons. A significant reduction was observed both in axon numbers and in axon length (Fig. 8, B and C), compared with scramble siRNA-treated neurons, in keeping with the DN approach.


We finally wanted to examine the relevance of mDia1 activity vis-a-vis of Rac, a small GTPase classically found to mediate axon outgrowth. A DN form of Rac, RacN17, was introduced into cerebellar granule cells along with a DA-mDia1 mutant mDia1- $\Delta$ N3. Axon elongation facilitated by DA-mDia1 in the presence of Y-27632 was repressed, in the presence of RacN17, back to baseline levels (Fig. 9 A). These findings are most consistent with the idea that Rac may significantly contribute to the mDia1 effect on axon outgrowth. Thus, mDia1 may directly regulate Rac or one of its upstream regulators.


Could Rho/mDia pathway regulate Rac? We recently proposed a candidate mechanism in HeLa and Swiss3T3 cells: as ROCK is known to down-regulate Rac activity, ROCK inhibition, under some circumstances, may suffice to up-regulate Rac via mDia1 (Tsuji et al., 2002). We tested whether such a possibility may be true in cerebellar granule



**Figure 6. DA mDia1 facilitates axon elongation.** Morphology of cerebellar granule cells overexpressing GFP (A), GFP-mDia1- $\Delta$ N3 alone (B), or GFP-mDia1- $\Delta$ N3 in the presence of Y-27632 (C). When ROCK activity was reduced, expression of GFP-mDia1- $\Delta$ N3 resulted in a significantly enhanced elongation (D, left) of axons ( $n \approx 65$ –157) compared with EGFP-expressing controls (A). Overexpression of GFP-mDia1- $\Delta$ N3 alone successfully induced an axon, which, however, exhibited a significantly altered shape (enlarged width, premature stop), presumably due to an increased actin stability in the presence of intact ROCK activity (B). Basal ROCK activity, in the context of excessive mDia1 activity, might cause a prominent increase in actin polymerization, while also sustaining a tonic level of actomyosin contractility, thereby negatively acting on axon elongation. \* $P < 0.05$ ; \*\*\* $P < 0.001$ . Bars, 5  $\mu$ m.

**Figure 7. A DN mDia1 mutant interferes with SDF-1 $\alpha$ -dependent axon elongation.** (A) Coexpression of the GFP-mDia1- $\Delta$ N3(HindIII) mutant abolished the effect of FLAG-mDia1- $\Delta$ N3 expression on axon length (left).  $n \approx 33$ –80. (B and C) The effect of GFP-mDia1- $\Delta$ N3(HindIII) overexpression was examined on SDF-1 $\alpha$ -facilitated axon elongation in the presence of Y-27632. A potent inhibition on both SDF-1 $\alpha$ -dependent axon elongation (B and C, left) and axon initiation (B and C, right) was detected.  $n \approx 35$ –145. Bars, 5  $\mu$ m.






**Figure 8. mDia1 knockdown by RNAi using siRNA completely abolishes SDF-1α-dependent axon elongation.** (A) Significant reduction of mDia1 protein is achieved in NIH3T3 within 24 h by RNAi using siRNA. (B and C) mDia1 knockdown by RNAi annihilates both SDF-1α-dependent axon elongation (C, left) and axon initiation (C, right) back to baseline levels, thereby confirming the DN experiments.  $n \approx 26-140$ . \*\*\* $P < 0.001$ . Bars, 5 μm.

cells. Consistent with previous work, Y-27632 treatment in SDF-1α-stimulated neurons was sufficient to increase the GTP-bound form of Rac (Fig. 9 B).

Together, our data establish a critical role for the Rho/mDia1 pathway in mediating axon elongation in SDF-1α-stimulated cerebellar granule neurons, presumably in concert with Rac activity (Fig. 10).



**Figure 9. Expression of the morphological effects of mDia1 is mediated, at least in part, by Rac activity.** (A) Co-expression of GFP-N17Rac, a DN mutant of Rac, suppresses expression of the facilitatory effects of FLAG-mDia1-ΔN3 on either axon length (left) in the presence of Y-27632 (50 μM) back to baseline levels.  $n \approx 45-137$ . (B) Inhibition of ROCK activity in the presence of high concentration of SDF-1α (500 ng/ml) increases the GTP-bound form of Rac. \*\*\* $P < 0.001$ .

## Discussion

### Discovery of a Rho/mDia-dependent signaling pathway crucial for axon elongation

Previous reports have shown an antagonism between Rac and Rho-mediated pathways in neurite extension and control of axonal growth cones (for review see Narumiya et al., 1997; Gallo and Letourneau, 1998; Hall, 1998; Luo, 2000; Dickson, 2001; Nikolic, 2002). In the mammalian CNS, we showed that enhanced Rho/ROCK activity led to inhibition of axonogenesis, whereas inhibition of Rho/ROCK pathway induced precocious outgrowth of neurites (Bito et al., 2000). In spite of a few reports suggesting a distinct, rather facilitatory effect for Rho during neuritogenesis (Threadgill et al., 1997; Sebok et al., 1999; Bashaw et al., 2001), it has remained unclear whether these experimental results truly reflected a genuine primary effect mediated by Rho.

In this study, we have formally demonstrated that Rho may in fact mediate both stimulation and inhibition of axon outgrowth downstream of the same ligand SDF-1α in the context of early postnatal cerebellar granule cells. Although a classically recognized Rho/ROCK pathway repressed axon formation, a Rho/mDia pathway was identified which rather potentially facilitated axon elongation. How ROCK can domi-




Figure 10. **Proposed model for the role of mDia in SDF-1 $\alpha$ -dependent axon outgrowth in cultured cerebellar granule neurons.** Usually Rho and Rac antagonistically controls axon outgrowth. During the very early stages in the primary culture, ROCK action predominates to favor suppression of precocious outgrowth of axons from the cerebellar granule cells. However, as these neurons express a high amount of mDia1 gradual decline in ROCK activity facilitates expression of Rho-dependent mDia1 activity and a subsequent recruitment of a signaling complex, which in concert with the Rac-dependent signaling cascade may help the transition from inhibition to stimulation of axon outgrowth and elongation. Thus, the SDF-1 $\alpha$ /Rho/mDia1 pathway may play a critical role in defining and modulating the balance between the Rho- and Rac-based signaling pathways during axon outgrowth.

nantly antagonize the expression of the axon elongation machinery, while another Rho effector mDia is facilitating this event, is currently being investigated. One possibility that is consistent with our finding is that ROCK and mDia1 may locally respond to different levels of GTP-bound Rho. Our finding of a multiplicity in Rho-mediated sites of action is reminiscent of the recent findings suggesting that Rac also acts on multiple steps of axon outgrowth during *Drosophila* development (Ng et al., 2002).

How then does mDia, an adaptor protein containing multiple formin homology domains, exert its effect on axon elongation? A clear picture is still missing, since only a few binding partners of mDia such as profilin (Watanabe et al., 1997), IRSp53 (Fujiwara et al., 2000), or mDia-interacting protein (Satoh and Tominaga, 2002) have been reported. Interestingly, in *Saccharomyces cerevisiae* the Diaphanous homologue Bni1p was shown to be critically involved in controlling assembly of actin cables required for establishment of cell polarity and directed growth (Ozaki-Kuroda et al., 2001; Pruyne et al., 2002; Sagot et al., 2002). Furthermore, recently a DA mutant of mDia1 was also shown to affect orientation of microtubules in HeLa cells, thus implying that mDia may perhaps spatially coordinate actin polymerization and the stability/rate of assembly of microtubule polymers (Ishizaki et al., 2001; Palazzo et al., 2001).

Intriguingly, in our study Rac activity was shown to be required to support an mDia-based change in neuronal morphology. Indeed, a high amount of basal Rac activity is found in the early postnatal cerebellum (unpublished data). In vivo and in vitro studies examining the consequence of defects in Rac/Pak/Cdk5/p35 signaling have provided numerous lines of evidence in support for a role for Rac-mediated control of cytoskeletal dynamics during axon outgrowth (Luo et al., 1994, 1996; Nikolic et al., 1996, 1998; Kaufmann et al., 1998; Hing et al., 1999; Newsome et al., 2000; Zukerberg et al., 2000; Ng et al., 2002). Since we detected an increase in Rho but not Rac activity upon SDF-1 $\alpha$  stimulation, we presume that Rac may play a rather permissive role in mediating the Rho/mDia-dependent events, although we cannot exclude the possibility that mDia may also directly regulate Rac or one of its upstream regulators (Fig. 9; Tsuji et al., 2002).

### A critical role for a pial chemokine SDF-1 $\alpha$ as a guidance molecule regulating axon outgrowth

Recently, SDF-1 $\alpha$  (Tashiro et al., 1993; Nagasawa et al., 1994) has attracted a lot of attention as a prototypical chemokine that acts as a mitogen and participates in positional control of a large variety of highly differentiated cell types. Gene targeting of either the ligand molecule SDF-1 $\alpha$  or its receptor CXCR4 resulted in a strikingly similar phenotype in the respective knockout mice, thereby demonstrating a crucial role for SDF-1 $\alpha$ /CXCR4 signaling in the development of appropriate cell layer formation in the cerebellum, the hippocampal dentate gyrus, the thymus, the spleen, vascular endothelium, and the intestinal epithelium (Nagasawa et al., 1996; Ma et al., 1998; Tachibana et al., 1998; Zou et al., 1998; Lu et al., 2002). In the cerebellum, absence of SDF-1 $\alpha$ /CXCR4 signaling resulted in premature invasion of many proliferating granule cells into the cerebellar anlage, and aberrant onset of migration from the external granule cell layer (EGL) to the internal granule cell layer (IGL) (Ma et al., 1998; Zou et al., 1998). Complementary experiments demonstrated that SDF-1 $\alpha$  acted as a chemoattractant for migration of both EGL- and IGL-derived cerebellar granule cells in culture (Klein et al., 2001; Lu et al., 2001; Zhu et al., 2002). Together, these lines of evidence are consistent with the idea that neuronal migration and patterning may heavily involve chemokine and G protein-coupled receptor signaling.

However, it was not known which exact step of neuronal morphogenesis was directly targeted by SDF-1 $\alpha$ , nor was it clear what intracellular signaling it activated in order to fulfill its role as a neural guidance molecule. Here we found that a large part of the initial morphogenetic and cytoskeletal action of SDF-1 $\alpha$  might be specifically mediated by the Rho signaling pathway, independent of its potency as a mitogen, in a pertussis toxin-insensitive manner (unpublished data). Previously, it was suspected that SDF-1 $\alpha$  might exhibit its growth promoting or chemotactic effect on neurons via pertussis-sensitive Gi/o heterotrimeric G proteins (Klein et al., 2001) and/or via PLC-mediated activation of intracellular calcium stores (Klein et al., 2001; Lu et al., 2002). The possible contribution of these signaling pathways to the neurite elongation still needs to be properly addressed in our cultured cerebellar granule neurons.



## Control of actin dynamics via multiple effectors of small GTPases: potential antagonism and coordination between distinct signaling pathways

Our study demonstrates that the Rho-dependent signaling cascade implicated in the developmental control of neural hardwiring may consist of two apparently opposing streams of signals: an inhibitory pathway mediated by ROCK (Bito et al., 2000) and a facilitatory pathway mediated by mDia1 (this study). Coordination of these two pathways may be suited to sequentially express the unique property of SDF-1 $\alpha$  either as a repellent or as an attractant. These findings shed light on the new possibility that Rho- and Rac-dependent pathways may interact not just in an antagonistic but also in a cooperative way in the regulation of the timing and the extent of axon elongation in cultured cerebellar granule cells (Fig. 10). Our study raises the possibility that the stimulation of Rho/mDia-mediated pathway by SDF-1 $\alpha$  might provide a modulatory link between the Rho- and Rac-based modes.

Together, these findings underscore the significance of ROCK/mDia antagonism during early CNS development. An appropriate balance and coordination between these two signaling systems may be key to controlling the initial timing and extent of axon outgrowth as a function of the strength of stimuli exhibited by the gradient of external guidance cues such as the pial chemokine SDF-1 $\alpha$ .

## Materials and methods

### Primary culture of mouse cerebellar granule cells

Mouse cerebellar granule cell cultures were as described in Bito et al. (2000). A detailed protocol can be obtained from the authors upon request.

### Materials and reagents

Y-27632, a ROCK inhibitor, was supplied by Dr. Masayoshi Uehata (Mitsubishi Pharma Co., Osaka, Japan). Botulinum C3 exoenzyme was purified as described (Morii and Narumiya, 1995). SDF-1 $\alpha$  was purchased from PeproTech EC. GST-Rhotekin RBD and GST-PAK CRIB proteins were prepared as described (Ren et al., 1999; Tsuji et al., 2002). pEGFP-C1 was from CLONTECH Laboratories, Inc. pEGFP-mDia1 $\Delta$ N3, pFL-mDia1 $\Delta$ N3, pEGFP-mDia1 $\Delta$ N3 (HindIII), and pEGFP-N17Rac were as described (Watanabe et al., 1999; Tsuji et al., 2002). All other reagents were of analytical grade unless noted otherwise.

### Transient transfection of plasmid cDNA constructs and pharmacological assays

Transfection was performed in the following way as modified from Bito et al. (2000). At the end of cerebellar granule cell preparation, a cell pellet was recovered from a 1,000 rpm, 10-min centrifugation at 4°C and resuspended with 20% FCS/HANKS(-) at densities of  $5 \times 10^5$  cells. The cell mixture (80  $\mu$ l) was plated onto 12-mm round Matrigel-coated (Becton Dickinson) coverslips (Assistent) placed in a 24-well plate. Each coverslip had been coated with 80  $\mu$ l Matrigel, 4 h before use. Cells were allowed to settle down for 30 min at RT and were incubated for 30 min at 37°C in a CO<sub>2</sub> incubator after each coverslip was fed with 0.5 ml of Medium A (Earle's MEM supplemented with 10% FCS [Hyclone], 1 $\times$  B-27 supplement [Invitrogen], insulin [25 mg/L; Sigma-Aldrich], glucose [5 g/L], transferrin [100 mg/L; Calbiochem], and 2 mM glutamine). Then, for each coverslip subjected to transfection 100  $\mu$ l OptiMEM (Invitrogen) containing a total amount of 1  $\mu$ g of plasmids (0.5  $\mu$ g of each plasmid for cotransfection) and 1  $\mu$ l of Lipofectamine 2000 (Invitrogen) was added drop by drop. The plate was incubated in a CO<sub>2</sub> incubator for 3 h and replaced with 0.5 ml of Medium A (same as above) per well. When necessary, either Rho or ROCK activity was inhibited in transfected cells by adding Y-27632 (final concentration 50  $\mu$ M) or C3 (final concentration 50  $\mu$ g/ml), respectively. Cells were put back into a CO<sub>2</sub> incubator and cultured for another 12 h.

To examine the effect of various pharmacological reagents on various aspects of axon outgrowth, cerebellar granule cells were plated onto 12-mm round Matrigel-coated coverslips placed in a 24-well plate as de-

scribed above. Cells were allowed to settle down for 30 min at RT. After each coverslip was fed with 0.5 ml of Medium A, they were incubated for another 3.5 h at 37°C in a CO<sub>2</sub> incubator and then replaced with Medium A supplemented with either the vehicle or SDF-1 $\alpha$  (500 ng/ml), Y-27632 (50  $\mu$ M), SDF-1 $\alpha$  (10, 100, 250, 500, and 1,000 ng/ml) plus Y-27632 (50  $\mu$ M), SDF-1 $\alpha$  (500 ng/ml) plus C3 (50  $\mu$ g/ml), and indicated concentrations of C3 exoenzyme. The cells were put back into a CO<sub>2</sub> incubator and cultured for another 12 h.

Pharmacological assays were performed in transfected cells in a similar way except that the cells were incubated in a CO<sub>2</sub> incubator for 3 h after the end of transfection procedure and then replaced with 0.5 ml of Medium A containing similar amounts of drugs as described above.

For transfection, Swiss 3T3 cells were plated on a cover glass at a density of  $2 \times 10^4$  cells per 22-mm dish and cultured in DME containing 10% FCS for 24 h. The cells were washed with PBS(-) once and incubated with the indicated amounts of pEGFP-C1 or pEGFP-mDia1 $\Delta$ N3 mixed with 1  $\mu$ l of Lipofectamine 2000 for 3 h in 1 ml of OPTI-MEM. The medium was replaced with DME containing 10% FCS, and the cells were cultured for 18 h. The cells were then cultured in serum-free DME for 24 h and treated with 30  $\mu$ M Y-27632 for the last 30 min. The cells were fixed and stained as described below.

### Immunofluorescence microscopy and measurements of axon parameters

Immunocytochemistry and GFP-imaging was performed essentially as described (Bito et al., 1996, 2000) using a LSM 510 system (Carl Zeiss Micro-Imaging, Inc.). Primary and secondary antibodies were as follows: rabbit polyclonal anti-p140mDia1 antibody (AP50) (Watanabe et al., 1997), anti-FLAG mAb (M2; Eastman Kodak Co.), rat antitubulin mAb (Chemicon), anti- $\beta$ -tubulin mAb (TUB.2.1) (Sigma-Aldrich); anti-GFP monoclonal and polyclonal (Molecular Probes); and Alexa Fluor 488-, 594-, and 633-conjugated goat anti-rabbit, anti-rat, and anti-mouse secondary (Molecular Probes). Alexa Fluor 594-, and 633-phalloidin (1:1,000; Molecular Probes) were used to stain the F-actin. In all image analyses, no background subtraction was performed, and all pseudocolor representations were assembled using Adobe Photoshop® version 6.0 for illustrative purpose only.

Quantitation of axon length and number was performed on cerebellar granule neurons cultured in the presence of 10% FCS using parameters similar to those described previously (Bito et al., 2000). Axons were distinguished from the filopodia by the presence of microtubules as verified by  $\beta$ -tubulin staining. As established previously (Bito et al., 2000),  $\beta$ -tubulin immunoreactivity was missing in virtually all processes with less than 3  $\mu$ m in length. Because the average axon number was below two (i.e., exact number was usually one or zero) during the experimental time window in most of our assays in this study, we restricted our measurement of axon length to the longer first process. Since an axon was equal or larger than 3  $\mu$ m in length, its contour could be unambiguously traced from the cell soma up to the very tip of the processes; all turning points within the longest process was defined and the sum of the cumulated distances between these points was considered as axon length. The calculation was performed off-line using the software plug-in of the LSM 510 system. Measurements in transfected neurons were performed identically except that transfected neurons were first identified using an antitag immunostaining (with an anti-GFP or an anti-FLAG antibody) to maximize the probability of detection; when directly compared, FLAG- and GFP-tagged constructs yielded similar results and phenotypes. Statistical analyses were performed using Prism 3.02 (Graphpad Software). All data are indicated as means  $\pm$  SEM. Unpaired *t* test with Welch's correction was employed to determine statistical significance and *p* values below 0.05 was considered as significant.

### Videomicroscopy

Cerebellar granule cells labeled with PKH-26 (Sigma-Aldrich) were seeded at a density of  $10^5$  per dish in a 35-mm glass-bottom dish (Matsunami Glass) and cultured for 3 h. The dish was transferred to a temperature-controlled CO<sub>2</sub> incubator (Carl Zeiss MicroImaging, Inc.) attached to the microscope stage. SDF-1 $\alpha$  (100 ng/ml, a final concentration) was then added, and the cell movement was monitored at 37°C in 5% CO<sub>2</sub> for 60 min using a confocal laser scanning unit (LSM 510-V2.5; Carl Zeiss MicroImaging, Inc.). One optical section was acquired every 3 min, and the video image were constructed from these sequential images.

### Immunohistochemistry

P1 or adult brain was taken out and immediately frozen in TissueTek O.C.T. Compound (Sakura) on dry ice. Cryostat sections of 20- $\mu$ m width were obtained and subjected to indirect immunofluorescence staining using an anti-p140mDia1 primary (AP50; Watanabe et al., 1997) and Alexa

Fluor 488-conjugated goat anti-rabbit secondary antibodies. TO-PRO-3 iodide (Molecular Probes) was used for nuclear staining. Specimens were examined on Axioplan or LSM 510 confocal imaging system (Carl Zeiss MicroImaging, Inc.). Stacked optical sections were merged using Maximum Projection Software (Carl Zeiss MicroImaging, Inc.).

### Western blot analysis

Total brain or cerebellar tissues were collected from ICR mice of indicated ages. Thin slices were cut in parasagittal directions in ice-cold HANKS and lysed in the lysis buffer containing 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, and 1× Complete (Protease Inhibitor Cocktail Tablets) (Roche Diagnostics). The samples were centrifuged at 100,000 g for 30 min, and the supernatant was collected as the cell lysates. Protein concentration of the lysates was determined by the Lowry method. For Western blot analysis using the total cell lysates, one fifth volume of the 5× Laemmli sample buffer was added to the lysates. The mixtures were boiled for 5 min and subjected to SDS-PAGE and Western blot analysis using anti-p140mDia1 antibody (AP50).

### Rho and Rac pull-down assays

Cerebellar granule cells were plated at a density of  $5 \times 10^5$  cells per 10-cm dish. After treatment with Y-27632, C3 exoenzyme, and/or SDF-1 $\alpha$ , the cells were washed once with ice-cold HANKS and lysed in the pull-down lysis buffer containing 50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 2 mM MgCl<sub>2</sub>, 10% glycerol, 1% NP-40, 1× Complete (Roche Diagnostics), 1 mM sodium fluoride, 1 mM EGTA, and 1 mM PMSF. The samples were centrifuged at 12,000 g for 30 min, and the supernatants were saved as the cell lysates. For cerebellar tissue lysates, cerebella were collected from P1 and adult ICR mice, and thin slices were cut in parasagittal directions in ice-cold HANKS and lysed in the pull-down lysis buffer. The tissue lysates were centrifuged at 100,000 g for 30 min, and the supernatant was collected as the cell lysates. 200  $\mu$ g of cell lysates was incubated with 20  $\mu$ g GST-Rhotekin RBD or GST-PAK CRIB immobilized on glutathione-Sepharose 4B beads (Amersham Biosciences) for 1 h at 4°C. Beads were washed three times with the pull-down buffer, and GTP-bound RhoA and GTP-bound Rac1 were detected by Western blot with an anti-RhoA (Santa Cruz Biotechnology, Inc.) and an anti-Rac1 antibody (Upstate Biotechnology). Protein concentration was determined by the Lowry method.

### Protein knockdown of mDia1 by RNAi using siRNA

siRNA corresponding to mDia1 mRNA sequences were designed as recommended with 5'-phosphate, 3'-hydroxyl, and two-base overhangs on each strand; it was chemically synthesized and annealed for duplex siRNA formation by Dharmacon Research, Inc. The following gene-specific sequences were used successfully: si-mDia1 (K2) sense 5'-GCUGGUCAGACCAUGGAU-3' and antisense 5'-CGACCAGUCUCGGUACCU-3'. Transfection of mDia1 siRNA for NIH-3T3 cells ( $10^4$  cells/ml) was performed with Lipofectamine 2000 in 6-well plates. Per well, 2  $\mu$ l lipofectamine 2000 diluted in 100  $\mu$ l Opti-MEM was applied to a pre-mix consisting of siRNA (20  $\mu$ M, 4  $\mu$ l)/Opti-MEM (100  $\mu$ l) and incubated for 30 min. The whole mixture was added to the medium, which then was changed to DME 3 h after transfection. Cells were incubated for 24–48 h before analysis of knock-down mDia1 by Western blot as described above.

Cotransfection of pEGFP-C1 and siRNA to cerebellar granule cells ( $5 \times 10^5$  cells/ml) were performed with Lipofectamine 2000. Briefly, for  $10^6$  cells 2  $\mu$ l Lipofectamine 2000 diluted in 100  $\mu$ l Opti-MEM was applied to a pEGFP-C1 (1  $\mu$ g)/siRNA (20  $\mu$ M, 8  $\mu$ l)/Opti-MEM (200  $\mu$ l) mixture and incubated for 30 min. The entire mixture was then added to the cell-containing medium, which was refed with the original culture medium 3 h after transfection. Cells were incubated for 12 h in solution and then finally plated onto 12-mm round Matrigel-coated coverslips placed in a 24-well plate. After another 12 h of culture on the coverslips, measurements of axon parameters were performed as described above.

### Online supplemental material

Video 1, showing facilitation of axon elongation by a 100 ng/ml SDF-1 $\alpha$  treatment in cultured cerebellar granule neurons, is available online at <http://www.jcb.org/cgi/content/full/jcb.200210149/DC1>.

We thank K. Nonomura, H. Nose, and T. Arai for assistance, T. Ishizaki and N. Watanabe for discussion, and S. Arakawa for support.

This work was supported in part by grants in aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to S. Narumiya, K. Kimura, and H. Bito) and grants from PRESTO-Japan Science and Technology Corporation, the Tanabe Medical Frontier Conference, the

Cell Science Research Foundation, and Human Frontier Science Program (to H. Bito). T. Furuyashiki is a postdoctoral fellow and Y. Arakawa and S. Takemoto-Kimura are predoctoral fellows from the Japan Society for Promotion of Science.

Submitted: 28 October 2002

Revised: 10 March 2003

Accepted: 10 March 2003

## References

- Altman, J., and S.A. Bayer. 1996. The third stage of cerebellar development: maturation of the cerebellar system. *In* Development of the Cerebellar System. CRC Press, Boca Raton, FL. 324–469.
- Bashaw, G.J., H. Hu, C.D. Nobes, and C.S. Goodman. 2001. A novel Dbl family RhoGEF promotes Rho-dependent axon attraction to the central nervous system midline in *Drosophila* and overcomes Robo repulsion. *J. Cell Biol.* 155:1117–1122.
- Bhatt, R.S., T. Tomoda, Y. Fang, and M.E. Hatten. 2000. Discoidin domain receptor 1 functions in axon extension of cerebellar granule neurons. *Genes Dev.* 14:2216–2228.
- Bito, H., K. Deisseroth, and R.W. Tsien. 1996. CREB phosphorylation and dephosphorylation: a Ca<sup>2+</sup>- and stimulus duration-dependent switch for hippocampal gene expression. *Cell.* 87:1203–1214.
- Bito, H., T. Furuyashiki, H. Ishihara, Y. Shibasaki, K. Ohashi, K. Mizuno, M. Maekawa, M. Ishizaki, and S. Narumiya. 2000. A critical role for a Rho-associated kinase, p160ROCK, in determining axon outgrowth in mammalian CNS neurons. *Neuron.* 26:431–441.
- Dickson, B.J. 2001. Rho GTPases in growth cone guidance. *Curr. Opin. Neurobiol.* 11:103–110.
- Driessens, M.H., H. Hu, C.D. Nobes, A. Self, I. Jordens, C.S. Goodman, and A. Hall. 2001. Plexin-B semaphorin receptors interact directly with active Rac and regulate the actin cytoskeleton by activating Rho. *Curr. Biol.* 11:339–344.
- Elbashir, S.M., J. Harborth, W. Lendeckel, A. Yalcin, K. Weber, and T. Tuschl. 2001. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature.* 411:494–498.
- Fujiwara, T., A. Mammoto, Y. Kim, and Y. Takai. 2000. Rho small G-protein-dependent binding of mDia to an Src homology 3 domain-containing IRSp53/BAIAP2. *Biochem. Biophys. Res. Commun.* 271:626–629.
- Gallo, G., and P.C. Letourneau. 1998. Axon guidance: GTPases help axons reach their targets. *Curr. Biol.* 8:R80–R82.
- Goodman, C.S., and C.J. Shatz. 1993. Developmental mechanisms that generate precise patterns of neuronal connectivity. *Cell.* 72:77–98.
- Hall, A. 1998. Rho GTPases and the actin cytoskeleton. *Science.* 279:509–514.
- Hatten, M.E. 1999. Central nervous system neuronal migration. *Annu. Rev. Neurosci.* 22:511–539.
- Hing, H., J. Xiao, N. Harden, L. Lim, and S.L. Zipursky. 1999. Pak functions downstream of Dock to regulate photoreceptor axon guidance in *Drosophila*. *Cell.* 97:853–863.
- Huang, E.J., and L.F. Reichardt. 2001. Neurotrophins: roles in neuronal development and function. *Annu. Rev. Neurosci.* 24:677–736.
- Ishizaki, T., M. Uehata, I. Tamechika, J. Keel, K. Nonomura, M. Maekawa, and S. Narumiya. 2000. Pharmacological properties of Y-27632, a specific inhibitor of rho-associated kinases. *Mol. Pharmacol.* 57:976–983.
- Ishizaki, T., Y. Morishima, M. Okamoto, T. Furuyashiki, T. Kato, and S. Narumiya. 2001. Coordination of microtubules and the actin cytoskeleton by the Rho effector mDia1. *Nat. Cell Biol.* 3:8–14.
- Jin, Z., and S.M. Strittmatter. 1997. Rac1 mediates collapsin-1-induced growth cone collapse. *J. Neurosci.* 17:6256–6263.
- Jurney, W.M., G. Gallo, P.C. Letourneau, and S.C. McLoon. 2002. Rac1-mediated endocytosis during Ephrin-A2- and Semaphorin 3A-induced growth cone collapse. *J. Neurosci.* 22:6019–6028.
- Kaufmann, N., Z.P. Wills, and D. Van Vactor. 1998. *Drosophila* Rac1 controls motor axon guidance. *Development.* 125:453–461.
- Klein, R.S., J.B. Rubin, H.D. Gibson, E.N. DeHaan, X. Alvarez-Hernandez, R.A. Segal, and A.D. Luster. 2001. SDF-1 alpha induces chemotaxis and enhances Sonic hedgehog-induced proliferation of cerebellar granule cells. *Development.* 128:1971–1981.
- Li, X., E. Saint-Cyr-Proulx, K. Aktories, and N. Lamarche-Vane. 2002. Rac1 and Cdc42 but not RhoA or Rho kinase activities are required for neurite outgrowth induced by the Netrin-1 receptor DCC (deleted in colorectal cancer) in N1E-115 neuroblastoma cells. *J. Biol. Chem.* 277:15207–15214.

- Liu, B.P., and S.M. Strittmatter. 2001. Semaphorin-mediated axonal guidance via Rho-related G proteins. *Curr. Opin. Cell Biol.* 13:619–626.
- Lu, M., E.A. Grove, and R.J. Miller. 2002. Abnormal development of the hippocampal dentate gyrus in mice lacking the CXCR4 chemokine receptor. *Proc. Natl. Acad. Sci. USA.* 99:7090–7095.
- Lu, Q., E.E. Sun, R.S. Klein, and J.G. Flanagan. 2001. Ephrin-B reverse signaling is mediated by a novel PDZ-RGS protein and selectively inhibits G protein-coupled chemoattraction. *Cell.* 105:69–79.
- Luo, L. 2000. Rho GTPases in neuronal morphogenesis. *Nat. Rev. Neurosci.* 1:173–180.
- Luo, L., Y.J. Liao, L.Y. Jan, and Y.N. Jan. 1994. Distinct morphogenetic functions of similar small GTPases: *Drosophila* Drac1 is involved in axonal outgrowth and myoblast fusion. *Genes Dev.* 8:1787–1802.
- Luo, L., T.K. Hensch, L. Ackerman, S. Barbel, L.Y. Jan, and Y.N. Jan. 1996. Differential effects of the Rac GTPase on Purkinje cell axons and dendritic trunks and spines. *Nature.* 379:837–840.
- Ma, Q., D. Jones, P.R. Borghesani, R.A. Segal, T. Nagasawa, T. Kishimoto, and R.T. Bronson. 1998. Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc. Natl. Acad. Sci. USA.* 95:9448–9453.
- Morii, N., and S. Narumiya. 1995. Preparation of native and recombinant *Clostridium botulinum* C3 ADP-ribosyltransferase and identification of Rho proteins by ADP-ribosylation. *Methods Enzymol.* 256:196–206.
- Nagasawa, T., H. Kikutani, and T. Kishimoto. 1994. Molecular cloning and structure of a pre-B-cell growth-stimulating factor. *Proc. Natl. Acad. Sci. USA.* 91:2305–2309.
- Nagasawa, T., S. Hirota, K. Tachibana, N. Takakura, S. Nishikawa, Y. Kitamura, N. Yoshida, H. Kikutani, and T. Kishimoto. 1996. Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature.* 382:635–638.
- Narumiya, S., T. Ishizaki, and N. Watanabe. 1997. Rho effectors and reorganization of actin cytoskeleton. *FEBS Lett.* 410:68–72.
- Newsome, T.P., S. Schmidt, G. Dietzl, K. Keleman, B. Asling, A. Debant, and B.J. Dickson. 2000. Trio combines with dock to regulate Pak activity during photoreceptor axon pathfinding in *Drosophila*. *Cell.* 101:283–294.
- Ng, J., T. Nardine, M. Harms, J. Tzu, A. Goldstein, Y. Sun, G. Dietzl, B.J. Dickson, and L. Luo. 2002. Rac GTPases control axon growth, guidance and branching. *Nature.* 416:442–447.
- Nikolic, M. 2002. The role of Rho GTPases and associated kinases in regulating neurite outgrowth. *Int. J. Biochem. Cell Biol.* 34:731–745.
- Nikolic, M., H. Dudek, Y.T. Kwon, Y.F. Ramos, and L.H. Tsai. 1996. The cdk5/p35 kinase is essential for neurite outgrowth during neuronal differentiation. *Genes Dev.* 10:816–825.
- Nikolic, M., M.M. Chou, W. Lu, B.J. Mayer, and L.H. Tsai. 1998. The p35/Cdk5 kinase is a neuron-specific Rac effector that inhibits Pak1 activity. *Nature.* 395:194–198.
- Nusser, N., E. Gosmanova, Y. Zheng, and G. Tigyi. 2002. NGF signals through TrkA, PI3 kinase, and Rac1 to inactivate RhoA during the initiation of neuronal differentiation of PC12 cells. *J. Biol. Chem.* 277:35840–35846.
- Ozaki-Kuroda, K., Y. Yamamoto, H. Nohara, M. Kinoshita, T. Fujiwara, K. Irie, and Y. Takai. 2001. Dynamic localization and function of Bni1p at the sites of directed growth in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 21:827–839.
- Ozdinler, P.H., and R.S. Erzurumlu. 2001. Regulation of neurotrophin-induced axonal responses via Rho GTPases. *J. Comp. Neurol.* 438:377–387.
- Palazzo, A.F., T.A. Cook, A.S. Alberts, and G.G. Gundersen. 2001. mDia mediates Rho-regulated formation and orientation of stable microtubules. *Nat. Cell Biol.* 3:723–729.
- Powell, S.K., R.J. Rivas, E. Rodriguez-Boulan, and M.E. Hatten. 1997. Development of polarity in cerebellar granule neurons. *J. Neurobiol.* 32:223–236.
- Pruyne, D., M. Evangelista, C. Yang, E. Bi, S. Zigmond, A. Bretscher, and C. Boone. 2002. Role of formins in actin assembly: nucleation and barbed-end association. *Science.* 297:612–615.
- Ren, X.D., W.B. Kiosses, and M.A. Schwartz. 1999. Regulation of the small GTP-binding protein Rho by cell adhesion and the cytoskeleton. *EMBO J.* 18:578–585.
- Sagot, I., A.A. Rodal, J. Moseley, B.L. Goode, and D. Pellman. 2002. An actin nucleation mechanism mediated by Bni1 and Profilin. *Nat. Cell Biol.* 4:626–631.
- Satoh, S., and T. Tominaga. 2002. mDia-interacting protein acts downstream of Rho-mDia and modifies Src activation and stress fiber formation. *J. Biol. Chem.* 276:39290–39294.
- Sebok, A., N. Nusser, B. Debrenceni, Z. Guo, M.F. Santos, J. Szeberenyi, and G. Tigyi. 1999. Different roles for RhoA during neurite initiation, elongation, and regeneration in PC12 cells. *J. Neurochem.* 73:949–960.
- Shamah, S.M., M.Z. Lin, J.L. Goldberg, S. Estrach, M. Sahin, L. Hu, M. Bazalakov, R.L. Neve, G. Corfas, A. Debant, and M.E. Greenberg. 2001. EphA receptors regulate growth cone dynamics through the novel guanine nucleotide exchange factor ephexin. *Cell.* 105:233–244.
- Swiercz, J.M., R. Kuner, J. Behrens, and S. Offermanns. 2002. Plexin-B1 directly interacts with PDZ-RhoGEF/LARG to regulate RhoA and growth cone morphology. *Neuron.* 35:51–63.
- Tachibana, K., S. Hirota, H. Iizasa, H. Yoshida, K. Kawabata, Y. Kataoka, Y. Kitamura, K. Matsushima, N. Yoshida, S. Nishikawa, et al. 1998. The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature.* 393:591–594.
- Tanaka, E., and J. Sabry. 1995. Making the connection: cytoskeletal rearrangements during growth cone guidance. *Cell.* 83:171–176.
- Tashiro, K., H. Tada, R. Heilker, M. Shirozu, T. Nakano, and T. Honjo. 1993. Signal sequence trap: a cloning strategy for secreted proteins and type I membrane proteins. *Science.* 261:600–603.
- Tessier-Lavigne, M., and C.S. Goodman. 1996. The molecular biology of axon guidance. *Science.* 274:1123–1133.
- Tham, T.N., F. Lazarini, I.A. Franceschini, F. Lachapelle, A. Amara, and M. Dubois-Dalq. 2001. Developmental pattern of expression of the alpha chemokine stromal cell-derived factor 1 in the rat central nervous system. *Eur. J. Neurosci.* 13:845–856.
- Threadgill, R., K. Bobb, and A. Ghosh. 1997. Regulation of dendritic growth and remodeling by Rho, Rac, and Cdc42. *Neuron.* 19:625–634.
- Tsuji, T., T. Ishizaki, M. Okamoto, C. Higashida, K. Kimura, T. Furuyashiki, Y. Arakawa, R.B. Birge, T. Nakamoto, H. Hirai, and S. Narumiya. 2002. ROCK and mDia1 antagonize in Rho-dependent Rac activation in Swiss 3T3 fibroblasts. *J. Cell Biol.* 157:819–830.
- Uehata, M., T. Ishizaki, H. Satoh, T. Ono, T. Kawahara, T. Morishita, H. Tamakawa, K. Yamagami, J. Inui, M. Maekawa, and S. Narumiya. 1997. Calcium sensitization of smooth muscle mediated by a Rho-associated protein kinase in hypertension. *Nature.* 389:990–994.
- Van Vactor, D., and J.G. Flanagan. 1999. The middle and the end: slit brings guidance and branching together in axon pathway selection. *Neuron.* 22:649–652.
- Watanabe, N., P. Madaule, T. Reid, T. Ishizaki, G. Watanabe, A. Kakizuka, Y. Saito, K. Nakao, B.M. Jockusch, and S. Narumiya. 1997. p140mDia, a mammalian homolog of *Drosophila* diaphanous, is a target protein for Rho small GTPase and is a ligand for profilin. *EMBO J.* 16:3044–3056.
- Watanabe, N., T. Kato, A. Fujita, T. Ishizaki, and S. Narumiya. 1999. Cooperation between mDia1 and ROCK in Rho-induced actin reorganization. *Nat. Cell Biol.* 1:136–143.
- Winton, M.J., C.I. Dubreuil, D. Lasko, N. Leclerc, and L. McKerracher. 2002. Characterization of new cell permeable C3-like proteins that inactivate Rho and stimulate neurite outgrowth on inhibitory substrates. *J. Biol. Chem.* 277:32820–32829.
- Wong, K., X.R. Ren, Y.Z. Huang, Y. Xie, G. Liu, H. Saito, H. Tang, L. Wen, S.M. Brady-Kalnay, L. Mei, et al. 2001. Signal transduction in neuronal migration: roles of GTPase activating proteins and the small GTPase Cdc42 in the Slit-Robo pathway. *Cell.* 107:209–221.
- Yacubova, E., and H. Komuro. 2002. Intrinsic program for migration of cerebellar granule cells in vitro. *J. Neurosci.* 22:5966–5981.
- Yamasaki, T., K. Kawaji, K. Ono, H. Bito, T. Hirano, N. Osumi, and M. Kengaku. 2001. Pax6 regulates granule cell polarization during parallel fiber formation in the developing cerebellum. *Development.* 128:3133–3144.
- Yamashita, T., K.L. Tucker, and Y.A. Barde. 1999. Neurotrophin binding to the p75 receptor modulates Rho activity and axonal outgrowth. *Neuron.* 24:585–593.
- Zhu, Y., T. Yu, X.C. Zhang, T. Nagasawa, J.Y. Wu, and Y. Rao. 2002. Role of the chemokine SDF-1 as the meningeal attractant for embryonic cerebellar neurons. *Nat. Neurosci.* 5:719–720.
- Zou, Y.R., A.H. Kottmann, M. Kuroda, I. Taniuchi, and D.R. Littman. 1998. Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature.* 393:595–599.
- Zukerberg, L.R., G.N. Patrick, M. Nikolic, S. Humbert, C.L. Wu, L.M. Lanier, F.B. Gertler, M. Vidal, R.A. Van Etten, and L.H. Tsai. 2000. Cables links Cdk5 and c-Abl and facilitates Cdk5 tyrosine phosphorylation, kinase up-regulation, and neurite outgrowth. *Neuron.* 26:633–646.