

PROTEIN SYNTHESIS AND RNA SYNTHESIS DURING MITOSIS IN ANIMAL CELLS

CAROL G. KONRAD, Ph.D.

From the Department of Zoology, University of California, Berkeley

ABSTRACT

Protein synthesis and RNA synthesis during mitosis were studied by autoradiography on mammalian tissue culture cells. Protein synthesis was followed by incubating hamster epithelial and human amnion cells for 10 or 15 minutes with phenylalanine- C^{14} . To study RNA synthesis the hamster cells were incubated for 10 minutes with uridine- C^{14} . Comparisons of the synthetic capacity of the interphase and mitotic cells were then made using whole cell grain counts. The rate of RNA synthesis decreased during prophase and reached a low of 13 to 16 per cent of the average interphase rate during metaphase-anaphase. Protein synthesis in the hamster cells showed a 42 per cent increase during prophase with a subsequent return to the average interphase value during metaphase-anaphase. The human amnion cells showed no significant change at prophase but there was a 52 to 56 per cent drop in phenylalanine incorporation at metaphase-anaphase as compared to the average interphase rate. Colcemide was used on the hamster cells to study the effect of a prolonged mitotic condition on protein and RNA synthesis. Under this condition, uridine incorporation was extremely low whereas phenylalanine incorporation was still relatively high. The drastic reduction of RNA synthesis observed under mitotic conditions is believed to be due to the coiled condition of the chromosomes. The lack of a comparable reduction in protein synthesis during mitosis is interpreted as evidence for the presence in these cells of a relatively stable messenger RNA.

INTRODUCTION

Studies of the growth pattern during the life cycle of various protista have yielded conflicting evidence on the metabolic status of the dividing cell. Some investigators find no evidence of disruption in the general interphase growth pattern during the time of cell division (Adolph and Bayne-Jones, 1932; Knaysi, 1940; Mitchison, 1957, 1958; Abbo and Pardee, 1960; Williamson and Scopes, 1961). Other data, however, show marked reduction in growth rate during division (Bayne-Jones and Adolph, 1932; Knaysi, 1940; Zeuthen, 1953; Lindgren and Haddad, 1954; Prescott, 1955; Maruyama, 1956; Christensson, 1959; Hamburger

and Zeuthen, 1960; Satir and Zeuthen, 1961) and, in some cases, there is evidence for a loss of mass at the time of division (Kimball, Caspersson, Svensson, and Carlson, 1959; Sandritter, Schiemer, Kraus, and Dörrien, 1960; Woodard, Gelber, and Swift, 1961).

Changes in cell morphology that occur during mitosis are so drastic that on this basis alone it might be predicted that there would be a change from the interphase pattern of metabolism. The chromosomes become condensed; the nuclear membrane breaks down; the nucleolus disappears; and the spindle and its related structures form, oc-

cupping a large fraction of the cell and thus disorganizing the interphase cytoplasmic structure. Fine structure cytology reveals that the endoplasmic reticulum is disorganized (Porter, 1954; Porter and Machado, 1960) and mitochondria are often pushed to the periphery (Porter, 1954; Porter and Machado, 1960; Gross, 1957; Kurosumi, 1958).

Uptake of $P^{32}O_4$ and $S^{35}O_4$ has been studied in several cell types and found to decline during division (Zeuthen, 1951; Taylor and Taylor, 1953; Mazia and Prescott, 1954; Taylor, 1958; Hamburger and Zeuthen, 1960; de Terra, 1960). Much work has been done recently on the uptake of specific precursors of protein, DNA, and RNA during the life cycle of the cell (Siskin, 1959; Feinendegen, Bond, Shreeve, and Painter, 1960; Taylor, 1959, 1960; Woodard, Rasch, and Swift, 1961; Baserga, 1962 *a*, 1962 *b*; Mitchison and Wilbur, 1962; Prescott and Bender, 1962). In general, mitosis is found to be a period of depressed metabolic activity.

Except for the few instances when DNA synthesis extends into prophase or begins in telophase, most cells synthesize DNA well before division (Taylor, 1957). The work of Taylor, Woods, and Hughes (1957) with colchicine, and of Bibring (cited by Bucher and Mazia, 1960) with mercaptoethanol-treated cells supports the conclusion that condensed chromosomes do not synthesize DNA.

Taylor (1960) and Prescott and Bender (1962) report that RNA synthesis completely stops when the chromosomes condense in mitosis. Das (1963) finds that although condensed chromosomes no longer synthesize RNA, it is only after the nucleolus breaks down that RNA synthesis comes to a complete halt. Prescott and Bender, however, state that RNA synthesis completely stops in prophase shortly before the nucleoli disappear.

The following experiments were undertaken to study the metabolic status of some mammalian tissue culture cells during the course of mitosis in regard to protein and RNA synthesis. Rapidly growing hamster and human amnion cells were pulse labeled with phenylalanine- C^{14} or uridine- C^{14} and the incorporation was quantitatively evaluated by autoradiography. Comparisons of this work and similar studies carried out by other investigators are made in the discussion.

MATERIALS AND METHODS

Two mammalian tissue culture lines were used in the following experiments. Most of the work was done on hamster epithelial cells isolated in 1958 in the laboratory of Dr. George Yerganian and carried since 1960 by the Naval Biological Laboratory, School of Public Health, University of California, Berkeley, California. The other cells were human amnion, isolated at the Virus Laboratory, University of California, Berkeley, California (Zitcer and Dunnebacke, 1957).

Culture medium for the hamster cells was composed of 80 per cent medium NCTC 109 (McQuilken, Evans, and Earle, 1957), 10 per cent fetal bovine serum and 10 per cent penicillin (100 units/ml)-streptomycin (50 μ g/ml) solution; the human amnion cells were grown in a medium composed of 10 per cent of a 10 \times concentrate of medium 199 (Morgan, Morton, and Parker, 1950) made up in 1 \times Hanks' balanced salt solution, 5 per cent lamb serum, 75 per cent Hanks' balanced salt solution, and 10 per cent of the penicillin-streptomycin solution. All stock cultures were grown in 100-ml stoppered glass medicine bottles and kept at 37°C.

Cells to be used for the experiments were trypsinized (0.1 per cent trypsin-Versene solution in Puck's saline) from rapidly growing stock cultures and harvested by centrifugation. Approximately 3×10^4 cells in 0.2 ml of medium were pipetted onto 22-mm² glass coverslips and placed in individual 35 \times 10-mm covered plastic Petri dishes. This volume of cells was allowed to remain on the coverslip for at least 4 hours to allow attachment of the cells to the coverslip before adding more medium to the dish to a final volume of 2 ml. The coverslip cultures were incubated in a 5 per cent CO₂ atmosphere at 37°C in a constant high humidity incubator.

Coverslip cultures used in experiments were allowed to grow until they were in the period of exponential growth. Insofar as possible, the growth conditions were constant for all cultures. Checks made on generation time, mitotic time, chromosome number, and appearance revealed no significant change during the time these experiments were carried on.

To study protein synthesis, the cells were fed DL-phenylalanine- C^{14} (California Corp. for Biochemical Research, Los Angeles, 4 to 12 mC/mm); uridine- C^{14} (New England Nuclear Corp., Boston, 12 to 16 mC/mm) was used to study RNA synthesis. Both precursors were added directly to the medium in which the cells had already been growing; phenylalanine to a final concentration of 20 μ C/ml and uridine to 10 μ C/ml. In some experiments colcemid (5×10^{-6} per cent) was used to study the effect of a

prolonged metaphase condition on the uptake of phenylalanine and uridine. It, too, was made up in medium in which the cells had previously been growing.

Generation time and length of time in the various stages of mitosis were determined for both cell lines (Edwards, Koch, Youcis, Freese, Laite, and Donaldson, 1960). The length of time in the labeled precursor was adjusted to a time no greater than the time in any one mitotic stage: 10 minutes for the hamster cells and 15 minutes for the human amnion cells. Thus the cells in a given phase, when fixed, were in either that phase and/or the preceding phase during labeling.

Immediately after the cells were exposed to the labeled precursor, the medium was removed and the coverslips washed in unlabeled medium and then in several changes of Hanks' saline and fixed in methanol for at least 15 minutes. The coverslips were then attached to microscope slides with Euparal, cell side up, and allowed to dry for 3 days.

Slides labeled with uridine- C^{14} were then treated with deoxyribonuclease (Worthington Biochemical Corp.) for 3 hours at 37°C, at a concentration of 0.2 mg/ml, made up in 0.003 M $MgSO_4$ and adjusted to pH 6.7 with dilute NaOH. Enzyme-treated slides were rinsed in H_2O and again dried before further handling.

Kodak AR-10 stripping film was then applied to the slides (Doniach and Pelc, 1950) and the slides were exposed for 1 to 3 weeks. The resulting autoradiograms were stained with either toluidine blue (0.05 per cent toluidine blue in Walpole's buffer pH 4) or with Azure Eosin Giemsa (procedure of T. McClure, University of California Medical Center, San Francisco, California). After rinsing, the slides were air dried.

All grain counts were done visually at a magnification of 1000. Grain counts were made over single cells and corrected for background by determining the area of the cell with the aid of an ocular grid and counting grains in a cell-free area of known size. All counts over cells of experimental slides were significantly different from background. A minimum of 20 cells in each mitotic phase was counted on each slide. Checks made on previously counted cells gave an error of no more than 5 per cent.

Tritium-labeled precursors were used in earlier experiments but carbon-14 was substituted in the experiments described here because of the possibility of self-absorption in the more rounded and, therefore, presumably thicker, dividing cells. Although carbon-14 has a much longer range and, therefore, overcomes to a large extent the error due to possible differences in self-absorption among the cells, there is another geometric problem characteristic of long range emitters: one should expect differences in

losses of grains in the counts among cells of varying geometry due to emissions hitting the emulsion outside the cell area. Therefore, controls were used in which the cells were labeled under the same conditions as were the experimentals but returned to unlabeled medium for some length of time greater than the division time (but less than the generation time) before being fixed. In this way, observations could be made on cells which were in mitosis at the time of fixation but which were labeled while they were in interphase.

However, for counts of control interphase cells under these conditions, correction must be made for the fact that during the time between labeling and fixation some cells had gone through mitosis, and, therefore, interphase counts include a group of daughter cells with half the amount of label of the parent cell. A formula for this correction may readily be derived by use of the formula for cumulative phase index given by Edwards *et al.* (1960, formula 1). Let x be the corrected mean interphase grain count. Then the observed mean interphase grain count is equal to:

$$\frac{p_1}{p_2} \cdot \frac{x}{2} + \frac{p_2 - p_1}{p_2} \cdot x$$

where p_2 is the fraction of cells in interphase and p_1 is the fraction of cells which have divided between the time of labeling and the time of fixation. Referring to formula 1 of Edwards *et al.*, we see that:

$$p_1 = 2(1 - 2^{-t_1/T})$$

$$p_2 = 2(1 - 2^{-t_2/T})$$

Where T = generation time (15 hours in the hamster cells and 30 hours in the human amnion cells); t_1 = time elapsed after labeling (time in unlabeled medium before fixation; note that t_1 is always less than t_2 in this experiment); t_2 = time in interphase (generation time minus mitotic time).

Thus, the corrected mean interphase grain count can be calculated if the generation time, the time elapsed between labeling and fixation, and the time in interphase are known. No correction was made for the small per cent of cells in mitosis when labeled and in interphase when fixed.

RESULTS

Uridine Incorporation

HAMSTER CELLS: A drastic reduction in the incorporation of uridine occurred during mitosis as seen from the data in Table I and Fig. 1. As

TABLE I
Results of Whole Cell Grain Counts over Hamster Cells Exposed to Uridine- C^{14} for 10 Minutes

Experiment	Interphase	Prophase	Metaphase	Anaphase	¶Telophase
Slide 1	*33.6 ± 2.1 ‡247.1 ± 15.4	18.4 ± 2.1 135.3 ± 15.4	3.4 ± 0.8 25.0 ± 5.9	3.1 ± 0.5 22.8 ± 3.7	9.5 ± 1.8 69.8 ± 13.2
Slide 2	38.8 ± 4.1 240.7 ± 25.4	21.3 ± 2.0 132.1 ± 12.4	5.2 ± 1.0 32.2 ± 6.2	4.6 ± 0.6 28.5 ± 3.7	10.7 ± 1.4 66.4 ± 8.7
Slide 3	24.6 ± 2.1 278.3 ± 23.8	9.4 ± 1.2 106.3 ± 13.6	1.2 ± 0.2 13.6 ± 2.3	2.2 ± 0.6 24.9 ± 6.8	6.8 ± 0.9 76.9 ± 10.2
§Average	255.4 ± 12.7	124.6 ± 8.0	23.6 ± 3.0	25.4 ± 2.9	71.0 ± 6.3
Corrected average	100.0 ± 6.6	56.5 ± 4.3	12.6 ± 1.7	16.4 ± 2.1	37.6 ± 3.8

* Average grain counts with standard error of the mean based on 20 cells per phase.

‡ Normalized average grain counts obtained by dividing the average grain count per phase by the mean grain count of that slide. This procedure overcomes differences among replicate slides due to differences in time of film exposure and allows them to be averaged.

§ Average of the normalized average grain counts of replicate experiments.

|| Normalized average grain count of each mitotic phase corrected for self-absorption, etc., due to the geometry of cells in that phase, by dividing by the normalized average grain count for control cells in the same phase. The resulting interphase value was then set equal to 100 to facilitate comparison. These values from Tables I, II, and III are plotted in Fig. 1.

¶ Telophase counts are the sum of counts over sister cells.




FIGURE 1 Normalized average grain counts corrected for self-absorption, etc. 95 per cent confidence limits are included for every point.

TABLE II
Results of Whole Cell Grain Counts over Hamster Cells Exposed to Phenyldanine-C¹⁴ for 10 Minutes

Experiment	Interphase	Prophase	Metaphase	Anaphase	Telophase
Slide 1	* 22.4 ± 1.1 ‡ 104.0 ± 5.1	32.0 ± 1.2 148.6 ± 5.6	20.0 ± 1.2 92.8 ± 5.6	10.6 ± 1.2 49.2 ± 5.6	22.7 ± 1.8 105.4 ± 8.4
Slide 2	55.2 ± 4.0 108.5 ± 7.9	67.9 ± 2.8 133.4 ± 5.5	45.5 ± 2.6 89.4 ± 5.1	34.6 ± 2.3 68.0 ± 4.5	51.3 ± 3.6 100.8 ± 7.1
Slide 3	50.5 ± 2.9 119.9 ± 6.9	53.7 ± 3.8 127.5 ± 9.0	35.8 ± 2.2 85.0 ± 5.2	30.3 ± 1.6 72.0 ± 3.8	40.2 ± 2.8 95.5 ± 6.6
§ Average	110.8 ± 3.9	136.5 ± 4.0	89.1 ± 3.1	63.1 ± 2.7	100.6 ± 4.3
Corrected average	100.0 ± 5.6	142.5 ± 7.1	109.4 ± 5.9	94.0 ± 6.6	123.0 ± 7.6

Geometry Control	**Hours	Uncorrected Interphase	‡‡Corrected Interphase	Prophase	Metaphase	Anaphase	Telophase
Slide 1	2.5	19.3 ± 1.5	21.7 ± 1.7 118.2 ± 9.2	21.8 ± 1.5 118.7 ± 8.2	18.7 ± 1.4 101.8 ± 7.6	14.8 ± 1.6 80.6 ± 8.7	14.8 ± 1.3 80.6 ± 7.1
Slide 2	5.0	15.4 ± 1.2	19.4 ± 1.5 115.5 ± 8.9	17.4 ± 1.5 103.6 ± 8.9	16.5 ± 1.2 98.2 ± 7.1	13.4 ± 1.4 79.8 ± 8.3	17.3 ± 1.4 103.0 ± 8.3
Slide 3	10.0	26.3 ± 1.8	41.7 ± 2.9 146.8 ± 10.2	30.2 ± 1.7 106.3 ± 6.0	22.6 ± 1.3 79.6 ± 4.6	19.9 ± 1.3 70.1 ± 4.6	27.6 ± 1.8 97.2 ± 6.3
Average		126.8 ± 5.5		109.5 ± 4.5	93.2 ± 3.8	76.8 ± 4.3	93.6 ± 4.2

* See Table I.

‡ See Table I.

§ See Table I.

|| See Table I.

¶ See Table I.

** Hours in unlabeled medium after labeling.

‡‡ Correction was made for those cells which have gone through mitosis during the time in unlabeled medium.

TABLE III
Results of Whole Cell Grain Counts over Haman Amnion Cells Exposed to Phenylalanine-C¹⁴ for 15 Minutes

Experiment	Interphase	Prophase	Metaphase	Anaphase	Telophase
Slide 1	* 119.2 ± 6.2 ‡ 144.9 ± 7.5	127.8 ± 7.0 155.4 ± 8.5	55.2 ± 3.8 67.1 ± 4.6	44.8 ± 2.6 54.5 ± 3.2	64.2 ± 4.2 78.1 ± 5.1
Slide 2	138.6 ± 8.1 127.8 ± 7.5	194.4 ± 7.6 179.2 ± 7.0	71.0 ± 4.2 65.4 ± 3.9	54.4 ± 2.7 50.1 ± 2.5	84.0 ± 5.4 77.4 ± 5.0
Slide 3	157.1 ± 11.0 139.8 ± 9.8	199.8 ± 7.9 177.8 ± 7.0	71.5 ± 4.7 63.6 ± 4.2	58.2 ± 2.9 51.8 ± 2.6	75.2 ± 5.0 66.9 ± 4.4
§ Average	137.5 ± 4.8	170.8 ± 4.3	65.4 ± 2.4	52.1 ± 1.6	74.1 ± 2.8
Corrected Average	100.0 ± 7.2	110.9 ± 6.2	47.9 ± 2.9	43.8 ± 3.2	52.5 ± 3.7

Geometry Control	**Hours	Uncorrected Interphase	††Corrected Interphase	Prophase	Metaphase	Anaphase	Telophase
Slide 1	3	124.6 ± 7.8	133.5 ± 8.4 99.9 ± 6.3	149.6 ± 7.6 111.9 ± 5.7	132.8 ± 6.4 99.3 ± 4.8	115.5 ± 7.6 86.4 ± 5.7	137.0 ± 8.3 102.5 ± 6.2

* See Table I.
 ‡ See Table I.
 § See Table I.
 || See Table I.
 ¶ See Table I.
 ** See Table II.
 †† See Table II.

TABLE IV
Results of Whole Cell Grain Counts over Hamster Cells Exposed to Phenylalanine-C¹⁴ for 1 Hour or Uridine-C¹⁴ for 10 Minutes at End of Colcemide Treatment

Phenylalanine-C ¹⁴	**Hours	Interphase	C-Mitosis	Uridine-C ¹⁴	**Hours	Interphase	C-Mitosis
Slide 1	3	114.4 ± 6.5	81.1 ± 6.3	Slide 1	5.5	52.8 ± 3.8	2.6 ± 0.5
	†	117.0 ± 6.6	83.0 ± 6.4			190.6 ± 13.7	9.4 ± 1.8
Slide 2	15	106.0 ± 9.1	48.0 ± 3.8	Slide 2	24.0	63.9 ± 7.1	2.0 ± 0.4
		137.7 ± 11.8	62.3 ± 4.9			193.9 ± 21.5	6.1 ± 1.2
Slide 3	30	118.4 ± 10.0	51.5 ± 4.6	Slide 3	39.5	66.2 ± 10.2	4.9 ± 1.5
		139.4 ± 11.8	60.6 ± 5.4			186.2 ± 28.7	13.8 ± 4.2
Slide 4	45	148.8 ± 12.8	76.7 ± 9.4				
		132.0 ± 11.4	68.0 ± 8.3				
§ Average (2, 3, and 4)		136.4 ± 6.7	§§ 63.6 ± 3.7	Average		190.2 ± 12.8	§§ 9.8 ± 1.6
Corrected (average)		100.0 ± 6.6	76.8 ± 6.2	Corrected (average)		100.0 ± 6.7	8.5 ± 1.5

* See Table I.

† See Table I.

§ See Table I.

|| See Table I.

¶ See Table I.

** Hours in colcemide.

§§ Anaphase geometry control was used as the correction for cells in C-mitosis.

early as prophase, the rate of RNA synthesis had fallen to 56 per cent of that of the average interphase value. By metaphase-anaphase, synthesis had fallen to a low of 13 to 16 per cent that of interphase. At telophase, the rate was increasing.

Phenylalanine Incorporation

HAMSTER CELLS: There is a significant rise in the incorporation of phenylalanine in prophase (42 per cent) over the average interphase rate. This period of high protein synthesis is followed by a fall at metaphase-anaphase to a rate not significantly different from the average interphase value. The results of the geometry controls indicate that the geometry of the anaphase hamster cell has a significant effect on the observed grain count of that phase. All data from these experiments are shown in Table II and Fig. 1.

HUMAN AMNION CELLS: Prophase appears to have the highest rate of protein synthesis, although the difference between it and the average interphase rate is not statistically significant when the three experiments are combined. At any rate, these cells show a significant drop in protein synthesis during metaphase-anaphase, at which time the rate is 52 to 56 per cent lower than the average interphase value. The geometry control data for the human amnion cells indicate that geometry does not significantly affect the grain counts. The data from these experiments are found in Table III and Fig. 1.

Colcemide Treatment

HAMSTER CELLS: Colcemide was employed to hold cells in a metaphase condition (C-mitosis) in order to test the effects of prolonged mitotic condition on uridine and phenylalanine incorporation in hamster cells. Phenylalanine-C¹⁴ was added to the medium for the last hour, or uridine-C¹⁴ for the last 10 minutes, of exposure (3 to 45 hours) to colcemide.

These cells cannot be held in C-mitosis indefinitely and eventually return to an abortive interphase condition with fragmented nuclei; this condition makes it possible to distinguish them from normal interphase cells. The interphase cells with fragmented nuclei accumulate steadily and are apparently incapable of reaching mitosis again. The length of time that cells remain in C-mitosis is somewhere between 3 and 15 hours, as judged from these experiments. The uptake of phenyl-

alanine by cells treated with colcemide for greater than 3 hours was 77 per cent that of the average interphase. The uptake of uridine fell to 8 per cent of that of the average interphase rate. The data from these experiments are to be found in Table IV.

DISCUSSION

RNA Synthesis

The drop in uridine uptake observed in these experiments during normal and C-mitosis in the hamster cell line was much more drastic than any drop observed for phenylalanine uptake. Unlike the reports of other investigators (Feinendegen *et al.*, 1960; Taylor, 1960; Baserga, 1962 *b*; Prescott and Bender, 1962), however, these experiments showed RNA synthesis to be significantly above zero during mitosis.

RNA synthesis declined with the time spent in the mitotic condition and reached a very low level in the extended colcemide mitosis. The DNA of condensed mitotic chromosomes thus seems to be masked in such a way as to lower its ability to prime RNA synthesis. For example, the DNA-dependent RNA polymerase may not be able to bind to the DNA. Unless the cell is completely effective in masking its primer DNA, one might expect some new RNA to be made during mitosis. Also, it might be possible for some nucleolar material, although not visible, to still function in RNA synthesis.

Protein Synthesis

The high rate of phenylalanine incorporation found in prophase for both hamster and human amnion cells might be interpreted as an indication that the cell reaches its maximum rate of protein synthesis at that time. However, Prescott and Bender (1962) find no prophase increase in hamster cells when using histidine-H³, and Baserga (1962 *a*), using leucine-C¹⁴ to label Ehrlich ascites tumor cells, finds protein synthesis to be maximal during the DNA synthetic period (S period).

The human amnion cells show a drop at metaphase-anaphase to about one-half of the interphase rate of phenylalanine incorporation. However, in the hamster cells, although there is a definite fall from the high rate at prophase, the metaphase-anaphase cells show no significant change from interphase. The hamster cell treated with colcemide is beginning to show a lowered rate of

protein synthesis, but there seems certainly to be no linear relationship between time in mitosis and drop in synthesis. Baserga (1962 *a*) and Prescott and Bender (1962) find the rate of amino acid incorporation to be lowered during mitosis, whereas Taylor (1960) finds amino acid incorporation unaffected by the division stage. It is difficult to explain the different results of Taylor, Prescott, and Bender, and those of this paper (all obtained from hamster cells). Prescott and Bender find a 75 per cent drop in histidine- H^3 incorporation during mitosis. One possible explanation of the differences between their results and those reported here involves the use of different isotopes. Due to the short range of tritium and the possible greater thickness of the mitotic cell relative to interphase cells, one may find the grain counts with tritium lower than those found using carbon-14 if no geometry correction is made.

The telophase rise in incorporation of both uridine and phenylalanine probably represents a transition period between the synthetic capacity of one cell and two cells. At some time after mitosis, the two daughter cells would be expected to show a combined rate twice that of the parent cell, but further experiments on both cell lines would be necessary to determine when this occurs.

Several differences between the patterns of RNA synthesis and protein synthesis are seen in these experiments: 1. The high rate of protein synthesis during prophase occurs at the same time that RNA synthesis has fallen drastically; 2. During mitosis, protein synthesis falls at most to one-half the average interphase value, while RNA synthesis is reduced to one-tenth; 3. Although protein synthesis would be expected eventually to cease under extended mitotic conditions, the results of the colcemide experiment indicate that, after several hours of extremely low RNA synthesis, protein synthesis is still relatively high. An interesting experiment of Goldstein, Micou, and Crocker (1960) showed that for 10 to 30 hours enucleated

human amnion tissue culture cells are able to incorporate amino acids, but are unable to incorporate detectable amounts of RNA precursors.

The similar results found for mitotic and enucleated cells support Mazia's idea that the mitotic condition is in many ways similar to the enucleate condition (Mazia and Prescott, 1954). In both mitotic and enucleate cells, RNA synthesis is greatly suppressed while protein synthesis is affected much less. In the mitotic cell this condition lasts on the order of 10 minutes, with colcemide it can be extended to 3 hours, and in the enucleate cell it continues for as long as 30 hours. Unless normally the rate of messenger RNA synthesis is a very small fraction of the total RNA synthesis—so that the small rate of RNA synthesis in mitosis could represent a large fraction of normal messenger synthesis—the messenger RNA must have a half-life equal to or greater than 10 minutes in dividing cells, 3 hours in colcemide-treated cells, and 30 hours in enucleated cells. Levinthal, Keynan, and Higa (1962) find a much less stable messenger RNA in bacteria. However, the stability of messenger RNA might well vary among different cell types.

The author wishes to thank Mrs. E. Zitcer of the Virus Laboratory, University of California, Berkeley, for the human amnion cells, and the Naval Biological Laboratory, School of Public Health, University of California, Berkeley, for the hamster cells used in these experiments. Thanks are especially due to Dr. Morgan Harris for the use of his tissue culture laboratory. Acknowledgment is given to Dr. Daniel Mazia for his continued advice and encouragement throughout the course of this work.

This study was submitted as a thesis to the Graduate Division of the University of California, Berkeley, in partial fulfillment of the requirements for the Ph.D. degree. Most of the work was done during the tenure of a Predoctoral Research Fellowship of the National Institutes of Health, United States Public Health Service (No. GPM-9708-C3).

Received for publication, February 18, 1963.

REFERENCES

- ABBO, F. E., and PARDEE, A. B., 1960, Synthesis of macromolecules in synchronously dividing bacteria, *Biochim. et Biophysica Acta*, **39**, 478.
- ADOLPH, E. F., and BAYNE-JONES, S., 1932, Growth in size of micro-organisms measured from motion pictures. II. *Bacillus megatherium*, *J. Cell. and Comp. Physiol.*, **1**, 409.
- BASERGA, R., 1962 *a*, A radioautographic study of the uptake of C^{14} -leucine by tumor cells in deoxyribonucleic acid synthesis, *Biochim. et Biophysica Acta*, **61**, 445.
- BASERGA, R., 1962 *b*, A study of nucleic acid synthesis in ascites tumor cells by two emulsion autoradiography, *J. Cell Biol.*, **12**, 633.
- BAYNE-JONES, S., and ADOLPH, E. F., 1932, Growth in size of micro-organisms measured from motion

- pictures. I. Yeast, *Saccharomyces cerevisiae*, *J. Cell. and Comp. Physiol.*, **1**, 387.
- BUCHER, N. L. R., and MAZIA, D., 1960. Deoxyribonucleic acid synthesis in relation to duplication of centers in dividing eggs of the sea urchin, *Strongylocentrotus purpuratus*, *J. Biophysic. and Biochem. Cytol.*, **7**, 651.
- CHRISTENSSON, E., 1959, Changes in free amino acids and proteins during cell growth and synchronous division in cultures of *Tetrahymena pyriformis*, *Acta Physiol. Scand.*, **45**, 339.
- DAS, N., 1963, Chromosomal and nucleolar RNA synthesis in root tips during mitosis, *Science*, **140**, 1231.
- DE TERRA, N., 1960, The effect of enucleation on restoration of the interphase rate of P³² uptake after cell division in *Stentor coeruleus*, *Exp. Cell Research*, **21**, 34.
- DONIAGH, I., and PELC, S. R., 1950, Autoradiographic technique, *Brit. J. Radiol.*, **23**, 184.
- EDWARDS, J. L., KOCH, A. L., YOUICIS, P., FREESE, H. L., LAITE, M. B., and DONALSON, J. T., 1960, Some characteristics of DNA synthesis and the mitotic cycle in Ehrlich ascites tumor cells, *J. Biophysic. and Biochem. Cytol.*, **7**, 273.
- FEINENDEGEN, L. E., BOND, V. P., SHREEVE, W. W., and PAINTER, R. B., 1960, RNA and DNA metabolism in human tissue culture cells studied with tritiated cytidine, *Exp. Cell Research*, **19**, 443.
- GOLDSTEIN, L., MICOU, J., and CROCKER, T. T., 1960, Nuclear-cytoplasmic relationships in human cells in tissue culture. IV. A study of some aspects of nucleic acid and protein metabolism in enucleate cells, *Biochim. et Biophysica Acta*, **45**, 82.
- GROSS, P. R., 1957, Labile biocolloids, cell division and the structure of the mitotic apparatus, *Tr. New York Acad. Sc.*, **20**, 154.
- HAMBURGER, K., and ZEUTHEN, E. 1960, Some characteristics of growth in normal and synchronized populations of *Tetrahymena pyriformis*, *Compt. rend. trav. lab. Carlsberg*, **32**, 1.
- KIMBALL, R. F., CASPERSSON, T. O., SVENSSON, G., and CARLSON, L., 1959, Quantitative cytochemical studies on *Paramecium aurelia*. I. Growth in total dry weight measured by the scanning interference microscope and x-ray absorption methods, *Exp. Cell Research*, **17**, 160.
- KNAYSI, G., 1940, A photomicrographic study of the rate of growth of some yeasts and bacteria, *J. Bact.*, **40**, 247.
- KUROSUMI, K., 1958, Electron microscope studies on mitosis in sea urchin blastomeres, *Protoplasma*, **49**, 116.
- LEVINTHAL, C., KEYNAN, A., and HIGA, A., 1962, Messenger RNA turnover and protein synthesis in *B. subtilis* inhibited by actinomycin D, *Proc. Nat. Acad. Sc.*, **48**, 1631.
- LINDEGREN, C. C., and HADDAD, S. A., 1954, Growth rates of individual yeast cells, *Genetica*, **27**, 45.
- MARUYAMA, Y., 1956, Biochemical aspects of the cell growth of *Escherichia coli* as studied by the method of synchronous culture, *J. Bact.*, **72**, 821.
- MAZIA, D., and PRESCOTT, D. M., 1954, Nuclear function and mitosis, *Science*, **120**, 120.
- MCQUILKIN, W. T., EVANS, V. J., and EARLE, W. R., 1957, The adaptation of additional lines of NCTC clone 929 (strain L) cells to chemically defined protein-free medium, *J. Nat. Cancer Inst.*, **19**, 885.
- MITCHISON, J. M., 1957, The growth of single cells. I. *Schizosaccharomyces pombe*, *Exp. Cell Research*, **13**, 244.
- MITCHISON, J. M., 1958, The growth of single cells. II. *Saccharomyces cerevisiae*, *Exp. Cell Research*, **15**, 214.
- MITCHISON, J. M., and WILBUR, K. M., 1962, The incorporation of protein and carbohydrate precursors during the cell cycle of a fission yeast, *Exp. Cell Research*, **26**, 144.
- MORGAN, J. F., MORTON, H. J., and PARKER, R. C., 1950, Nutrition of animal cells in tissue culture. I. Initial studies on a synthetic medium, *Proc. Soc. Exp. Biol. and Med.*, **73**, 1.
- PORTER, K. R., 1954, Changes in cell fine structure accompanying mitosis, in *Fine Structure of Cells*, Symposium 8th Congress on Cell Biology, Leiden, 1954, Groningen, The Netherlands, Noordhoff Ltd., 236-250.
- PORTER, K. R., and MACHADO, R. D., 1960, Studies on the endoplasmic reticulum. IV. Its form and distribution during mitosis in cells of onion root tip, *J. Biophysic. and Biochem. Cytol.*, **7**, 167.
- PRESCOTT, D. M., 1955, Relations between cell growth and cell division. I. Reduced weight, cell volume, protein content and nuclear volume of *Amoeba proteus* from division to division, *Exp. Cell Research*, **9**, 328.
- PRESCOTT, D. M., and BENDER, M. A., 1962, Synthesis of RNA and protein during mitosis in mammalian tissue culture cells, *Exp. Cell Research*, **26**, 260.
- SANDRITTER, W., SCHIEMER, H. G., KRAUS, H., and DÖRRIEN, U., 1960, Das Wachstum von Einzelzellen in der Gewebe Kultur. Interferenz Mikroskopische Untersuchungen an HeLa-Zellen, *J. Biophysic. and Biochem. Cytol.*, **7**, 579.
- SATIR, P., and ZEUTHEN, E., 1961, Cell cycle and the relationship of growth rate to reduced weight (RW) in the giant amoeba, *Chaos chaos L.*, *Compt. rend. trav. lab. Carlsberg*, **32**, 241.
- SISKIN, J. E., 1959, The synthesis of nucleic acids and proteins in the nuclei of *Tradescantia* root tips, *Exp. Cell Research*, **16**, 602.

- TAYLOR, J. H., 1957, The time and mode of duplication of chromosomes, *Am. Naturalist*, **91**, 209.
- TAYLOR, J. H., 1958, Incorporation of phosphorus-32 into nucleic acids and proteins during microgametogenesis of *Tulbaghia*, *Am. J. Bot.*, **45**, 123.
- TAYLOR, J. H., 1959, Autoradiographic studies of nucleic acids and proteins during meiosis in *Lilium longiflorum*, *Am. J. Bot.*, **46**, 477.
- TAYLOR, J. H., 1960, Nucleic acid synthesis in relation to the cell division cycle, *Ann. New York Acad. Sc.*, **90**, 409.
- TAYLOR, J. H., and TAYLOR, S. H., 1953, The autoradiograph—a tool for cytogeneticists, *J. Heredity*, **44**, 129.
- TAYLOR, J. H., WOODS, P. S., and HUGHES, W. L., 1957, The organization and duplication of chromosomes as revealed by autoradiographic studies using tritium labeled thymidine, *Proc. Nat. Acad. Sc.*, **43**, 122.
- WILLIAMSON, D. H., and SCOPES, A. W. 1961, Protein synthesis and nitrogen uptake in synchronously dividing cultures of *Saccharomyces cerevisiae*, *J. Inst. Brewing*, **67**, 39.
- WOODARD, J., GELBER, B., and H. SWIFT, 1961, Nucleoprotein changes during the mitotic cycle in *Paramecium aurelia*, *Exp. Cell Research*, **23**, 258.
- WOODARD, J., RASCH, E., and SWIFT, H., 1961, Nucleic acid and protein metabolism during the mitotic cycle in *Vicia faba*, *J. Biophysic. and Biochem. Cytol.* **9**, 445.
- ZEUTHEN, E., 1951, Segmentation, nuclear growth and cytoplasmic storage in eggs of echinoderms and amphibia, *Pubbl. staz. zool. Napoli*, **23**, 47.
- ZEUTHEN, E., 1953, Growth as related to the cell cycle in single cell cultures of *Tetrahymena pyriformis*, *J. Embryol. and Exp. Morphol.*, **1**, 239.
- ZITCER, E. M., and DUNNEBACKE, T. H., 1957, Transformation of cells from normal human amnion into established strains, *Cancer Research*, **17**, 1047.