


Location is everything



CK1 δ (green) localizes to the centrosome (red), where it promotes neurite outgrowth in response to Wnt-3a.

A specific isoform of casein kinase 1 (CK1) localizes to the centrosome to promote neurite extension, Greer and Rubin reveal.

The signaling protein Wnt-3a stimulates rapid neurite outgrowth from certain cell lines by activating the effector proteins Dishevelled (Dvl) 2 and 3. Dvl proteins' downstream effects can be modulated by phosphorylation. After finding that Dvl-2 and -3 were phosphorylated during Wnt-3a-induced neurite extension, Greer and Rubin screened a panel of small molecule inhibitors to identify the responsible protein kinases.

IC261, an inhibitor of the closely related CK1 isoforms δ and ϵ , blocked Wnt-3a-induced neurite outgrowth and Dvl-2/3 phosphorylation. Knocking down either kinase by RNAi also


reduced Dvl-2/3 phosphorylation, yet the two isoforms had opposing effects on neurite formation. Cells lacking CK1 δ no longer grew neurites in response to Wnt-3a, whereas CK1 ϵ depletion prompted neurite extension in the absence of Wnt ligand, suggesting that this isoform usually inhibits neurite growth.

Greer and Rubin found that, unlike CK1 ϵ , CK1 δ localized strongly to the centrosome, an organelle that controls neurite outgrowth in many types of neurons. Blocking this localization prevented Wnt-3a from inducing neurite extension. CK1 δ was targeted to the centrosome by a region in its C terminus; substituting this centrosomal localization signal for the C-terminal domain of CK1 ϵ allowed that isoform to promote neurite formation in place of CK1 δ , indicating that location is the key determinant of the isoforms' different activities.

Many Wnt signaling pathway proteins localize to centrosomes. The authors now want to investigate whether they cooperate with CK1 δ to drive neurite outgrowth and other centrosome functions such as ciliogenesis.

Greer, Y.E., and J.S. Rubin. 2011. *J. Cell Biol.* doi:10.1083/jcb.201011111.

Connecting the cytokinesis dots



Clusters of Myo2 (green) form at the equator of wild-type cells (top) but not cells lacking the IQGAP Rng2 (bottom).

Laporte et al. describe how precursors of the cytokinetic contractile ring assemble at the cell equator.

As fission yeast enter mitosis, protein clusters called cytokinesis nodes form around the cell cortex at the future site of cell division. The nodes eventually coalesce into a contractile ring to separate daughter cells, but little is known about the nodes' initial assembly, except that their localization at the cell cortex is determined by Mid1, a homologue of the animal-cell cleavage furrow protein anillin.


Laporte et al. analyzed the composition of cytokinesis nodes in different mutant backgrounds to define two protein modules that

assemble around Mid1 at the plasma membrane. The first module consisted of the myosin light chain Cdc4 and the actin-bundling IQGAP protein Rng2, which together recruited the myosin motor protein Myo2. The second module incorporated the membrane-bending F-BAR protein Cdc15. Both modules recruited the actin-nucleating formin Cdc12 into the nodes, Laporte et al. found.

The researchers then used a single-molecule imaging technique called SHREC to examine how these components are organized within each cytokinesis node. Myo2's head domain points into the cell interior, an orientation that may help it capture and pull actin filaments nucleated from neighboring nodes as the separate clusters condense into a full contractile ring. Senior author Jian-Qiu Wu now wants to examine how the organization of the node proteins changes as they form the contractile ring. He's particularly interested in whether Myo2 polymerizes into filaments, as has been observed in the contractile ring of other organisms.

Laporte, D., et al. 2011. *J. Cell Biol.* doi:10.1083/jcb.201008171.

Osteoblasts are bone idle without Frizzled-9



Compared to wild-type (left), a vertebra from a Fzd9-null mouse (right) shows reduced calcein staining (green), indicating a decreased rate of bone formation.

The Wnt receptor Frizzled-9 (Fzd9) promotes bone formation, Albers et al. report, providing a potential new target for the treatment of osteoporosis.

Adult bones are maintained by a balance of bone-forming osteoblasts and bone-resorbing osteoclasts. Although Wnt signaling affects this balance in mice and humans, the Wnt receptors involved remain unknown. Albers et al. found that the Wnt receptor Fzd9 was upregulated during osteoblast differentiation and that mice lacking Fzd9 had fragile bones due to low rates of bone formation.

Fzd9-null osteoblasts differentiated normally, but they failed to mineralize their extracellular matrix. The loss of Fzd9 disrupted

a non-canonical branch of the Wnt signaling pathway, resulting in reduced levels of the transcription factor STAT1, which was, in turn, required for the expression of several interferon-regulated genes. One of these genes encoded a ubiquitin-like molecule called Isg15. Though little is known about Isg15's function, restoring its expression in Fzd9-null osteoblasts boosted matrix mineralization, whereas mice lacking Isg15 had similar bone defects to Fzd9-knockout animals.

Mice lacking one copy of Fzd9 also had low bone mass, suggesting that insufficient Fzd9 may cause the reduced bone density seen in Williams-Beuren syndrome patients, who have a hemizygous deletion of the chromosomal region that includes the *FZD9* gene. Senior author Thorsten Schinke now wants to investigate whether boosting Fzd9 expression has the opposite effect to Fzd9 depletion and can stimulate bone formation. If so, Fzd9 would be an attractive drug target for treating a variety of bone-loss disorders.

Albers, J., et al. 2011. *J. Cell Biol.* doi:10.1083/jcb.201008012.