

Casp8p41 generated by HIV protease kills CD4 T cells through direct Bak activation

Amy M. Sainski, Haiming Dai, Sekar Natesampillai, Yuan-Ping Pang, Gary D. Bren, Nathan W. Cummins, Cristina Correia, X. Wei Meng, James E. Tarara, Marina Ramirez-Alvarado, David J. Katzmann, Christina Ochsenbauer, John C. Kappes, Scott H. Kaufmann, and Andrew D. Badley

Vol. 206 No. 7, September 29, 2014. Pages 867–876.

Due to an error during the production process, an incorrect version of Fig. 3 was published. Specifically, incorrect images were included for the top two panels of part I, and several figure parts contained minor formatting and labeling errors. A corrected version of the figure is shown below.




Figure 3. Casp8p41 facilitates Bak oligomerization and Bak-mediated MOMP. (A) Dextran-FITC-loaded liposomes composed of MOM lipids were incubated with 50 nM His₆-BakΔTM in the absence or presence of 200 nM GST, GST-Casp8p41, or GST-Casp8p41 V150E/L157K at 37°C for 30 min. Bim BH3 peptide served as a positive control. After bismaleimido-hexane cross-linking, samples were subjected to SDS-PAGE and blotted for Bak. (B) After His₆-BakΔTM was incubated with GST or GST-Casp8p41 in buffer containing 1% (wt/vol) CHAPS, fractions from size exclusion chromatography were blotted for Bak and GST. Representative of *n* = 3. (C) FITC-dextran release over time from FITC-dextran-loaded liposomes treated as in A. Representative of *n* = 3. (D) Summarized results from three independent experiments at 300 s of FITC-dextran release from liposomes. (E) FITC-dextran release from liposomes treated with 50 nM His₆-BakΔTM in the presence of 200 nM GST-Casp8p41, GST-Casp8p41 V150E/L157K (EK), or GST. 100% release in D and E was determined by treating liposomes with Triton X-100. Error bars indicate ± 1 SD from three independent experiments (*, *P* < 0.05; ***, *P* < 0.001). (F–J) After mitochondria (F) from wt (G), Bax^{-/-}Bak^{-/-} double knockout (H), Bax^{-/-} (I), or Bak^{-/-} (J) MEFs were incubated with 200 (1), 500 (2), or 1,000 nM (3) of purified protein for 1 h, sedimented, and washed, the supernatants and pellets were blotted for cytochrome c (Cyto C) and, as a control, the mitochondrial matrix protein Hsp60. Bim BH3 (200 nM), which induces CytoC release in the presence of either Bak or Bax, served as a positive control. Representative of *n* = 3.

The html and pdf versions of this article have been corrected. The error remains only in the print version.