

# Septins recognize micron-scale membrane curvature

Damián Lobato-Márquez and Serge Mostowy


Section of Microbiology, Medical Research Council Centre for Molecular Bacteriology and Infection, Imperial College London, London SW7 2AZ, England, UK

How cells recognize membrane curvature is not fully understood. In this issue, Bridges et al. (2016. *J. Cell Biol.* <http://dx.doi.org/10.1083/jcb.201512029>) discover that septins, a component of the cytoskeleton, recognize membrane curvature at the micron scale, a common morphological hallmark of eukaryotic cellular processes.

Eukaryotic cells have dedicated proteins that sense membranes, depending on their curvature (Antony, 2011). Sensors of membrane curvature are important because they organize a wide variety of cellular functions, including vesicle trafficking and organelle shaping (McMahon and Gallop, 2005). Curvature-sensing proteins, for example, the Bin-Amphiphysin-Rvs (BAR) domain-containing proteins, have been mostly described to work at the nanometer scale (Zimmerberg and Kozlov, 2006). In contrast, a clear mechanism of sensing membrane curvature at the micron scale in eukaryotic cells has not been described. In this issue, Bridges et al. discover that septins, a poorly understood component of the cytoskeleton, recognize plasma membrane curvature at the micron scale and serve as landmarks for eukaryotic cells to know their local shape.

Septins are an evolutionarily conserved family of GTP-binding proteins that assemble into nonpolar filaments and rings (John et al., 2007; Sirajuddin et al., 2007; Bertin et al., 2008). Septins have been implicated in diverse membrane organization events where micron-scale curvature takes place (Saarikangas and Barral, 2011; Mostowy and Cossart, 2012), including the cytokinetic furrow, the annulus of spermatozoa, the base of cellular protrusions (e.g., cilium and dendritic spines), and the phagocytic cup surrounding invasive bacterial pathogens (Fig. 1). However, the precise role of septin-membrane interactions remains elusive. It was first suggested in 1999 that the interaction of human septins with phosphatidylinositol 4,5-bisphosphate (PI(4,5)P<sub>2</sub>) is important for septin localization (Zhang et al., 1999). More recently, work using recombinant septins from budding yeast *Saccharomyces cerevisiae* assembled on PI(4,5)P<sub>2</sub> lipid monolayers showed that septins interact with membrane to facilitate filament assembly (Bridges et al., 2014). Membrane-facilitated septin assembly has also been observed using phospholipid liposomes, and in this case septins were also shown to induce membrane tubulation (Tanaka-Takiguchi et al., 2009). Given that (a) septins can interact with membrane, (b) septin assembly is membrane facilitated, and (c) septin assemblies are associated with a variety of membrane organization events from yeast to mammals, Bridges et al. (2016) hypothesized that septins serve as a mechanism to recognize membrane curvature.

Correspondence to Serge Mostowy: s.mostowy@imperial.ac.uk



**Figure 1. Morphological hallmarks of eukaryotic cells characterized by micron-scale membrane curvature and septin assembly.** Septins have been implicated in membrane organization events where micron-scale curvature takes place. (A) A septin ring acts as a scaffold for cytokinesis proteins and forms a diffusion barrier at the cytokinetic furrow of a dividing cell. (B) A septin ring forms a diffusion barrier at the annulus of a mammalian spermatozoon, which separates the anterior and posterior tail. (C) A septin ring forms a diffusion barrier at the base of a cilium to separate the ciliary membrane from the plasma membrane. (D) In neurons, a septin-dependent diffusion barrier can localize at the base of dendritic spine necks. (E) During phagocytosis, a cup is formed at the plasma membrane; septin rings assemble at the base of the phagocytic cup to regulate entry.

nanometer scale (e.g., BAR domain-containing proteins). However, it is likely that septins do more than recognize membrane, and the precise role of septins in membrane recognition remains unknown. The highly conserved structural and biochemical properties of septins suggest they organize, stabilize, and functionalize membrane domains (Caudron and Barral, 2009; Kusumi et al., 2012; Bridges and Gladfelter, 2015). Although we are far from knowing the full repertoire of septin function, this new work by Bridges et al. (2016) reminds us that understanding how membranes can specify septin assembly is essential to understand the role of septins in eukaryotic cells.

### Acknowledgments

Work in the S. Mostowy laboratory is supported by a Wellcome Trust Research Career Development Fellowship (WT097411MA) and the Lister Institute of Preventive Medicine.

The authors declare no competing financial interests.

Submitted: 17 March 2016

Accepted: 22 March 2016

### References

Antonny, B. 2011. Mechanisms of membrane curvature sensing. *Annu. Rev. Biochem.* 80:101–123. <http://dx.doi.org/10.1146/annurev-biochem-052809-155121>

Bertin, A., M.A. McMurray, P. Grob, S.S. Park, G. Garcia III, I. Patanwala, H.L. Ng, T. Alber, J. Thorner, and E. Nogales. 2008. *Saccharomyces cerevisiae* septins: supramolecular organization of heterooligomers and the mechanism of filament assembly. *Proc. Natl. Acad. Sci. USA.* 105:8274–8279. <http://dx.doi.org/10.1073/pnas.0803330105>

Bridges, A.A., and A.S. Gladfelter. 2015. Septin form and function at the cell cortex. *J. Biol. Chem.* 290:17173–17180. <http://dx.doi.org/10.1074/jbc.R114.634444>

Bridges, A.A., H. Zhang, S.B. Mehta, P. Occhipinti, T. Tani, and A.S. Gladfelter. 2014. Septin assemblies form by diffusion-driven annealing on membranes. *Proc. Natl. Acad. Sci. USA.* 111:2146–2151. <http://dx.doi.org/10.1073/pnas.1314138111>

Bridges, A.A., M.S. Jentzsch, P.W. Oakes, P. Occhipinti, and A.S. Gladfelter. 2016. Micron-scale plasma membrane curvature is recognized by the septin cytoskeleton. *J. Cell Biol.* <http://dx.doi.org/10.1083/jcb.201512029>

Caudron, F., and Y. Barral. 2009. Septins and the lateral compartmentalization of eukaryotic membranes. *Dev. Cell.* 16:493–506. <http://dx.doi.org/10.1016/j.devcel.2009.04.003>

John, C.M., R.K. Hite, C.S. Weirich, D.J. Fitzgerald, H. Jawhari, M. Faty, D. Schläpfer, R. Kroschewski, F.K. Winkler, T. Walz, et al. 2007. The *Caenorhabditis elegans* septin complex is nonpolar. *EMBO J.* 26:3296–3307. <http://dx.doi.org/10.1038/sj.emboj.7601775>

Kusumi, A., T.K. Fujiwara, R. Chadda, M. Xie, T.A. Tsunoyama, Z. Kalay, R.S. Kasai, and K.G. Suzuki. 2012. Dynamic organizing principles of the plasma membrane that regulate signal transduction: commemorating the fortieth anniversary of Singer and Nicolson's fluid-mosaic model. *Annu. Rev. Cell Dev. Biol.* 28:215–250. <http://dx.doi.org/10.1146/annurev-cellbio-100809-151736>

McMahon, H.T., and J.L. Gallop. 2005. Membrane curvature and mechanisms of dynamic cell membrane remodelling. *Nature.* 438:590–596. <http://dx.doi.org/10.1038/nature04396>

Mostowy, S., and P. Cossart. 2012. Septins: the fourth component of the cytoskeleton. *Nat. Rev. Mol. Cell Biol.* 13:183–194.

Saarikangas, J., and Y. Barral. 2011. The emerging functions of septins in metazoans. *EMBO Rep.* 12:1118–1126. <http://dx.doi.org/10.1038/embor.2011.193>

Sirajuddin, M., M. Farkasovsky, F. Hauer, D. Kühlmann, I.G. Macara, M. Weyand, H. Stark, and A. Wittinghofer. 2007. Structural insight into filament formation by mammalian septins. *Nature.* 449:311–315. <http://dx.doi.org/10.1038/nature06052>

Tanaka-Takiguchi, Y., M. Kinoshita, and K. Takiguchi. 2009. Septin-mediated uniform bracing of phospholipid membranes. *Curr. Biol.* 19:140–145. <http://dx.doi.org/10.1016/j.cub.2008.12.030>

Zhang, J., C. Kong, H. Xie, P.S. McPherson, S. Grinstein, and W.S. Trimble. 1999. Phosphatidylinositol polyphosphate binding to the mammalian septin H5 is modulated by GTP. *Curr. Biol.* 9:1458–1467. [http://dx.doi.org/10.1016/S0960-9822\(00\)80115-3](http://dx.doi.org/10.1016/S0960-9822(00)80115-3)

Zimmerberg, J., and M.M. Kozlov. 2006. How proteins produce cellular membrane curvature. *Nat. Rev. Mol. Cell Biol.* 7:9–19. <http://dx.doi.org/10.1038/nrm1784>