

Ragulator—a multifaceted regulator of lysosomal signaling and trafficking

Alexandria Colaço and Marja Jäätelä

Cell Death and Metabolism, Center for Autophagy, Recycling and Disease, Danish Cancer Society Research Center, Copenhagen, Denmark

The lysosomal Ragulator complex regulates cell metabolism and growth by coordinating the activities of metabolic signaling pathways with nutrient availability. In this issue, Filipek et al. (2017. *J. Cell Biol.* <https://doi.org/10.1083/jcb.201703061>) and Pu et al. (2017. *J. Cell Biol.* <https://doi.org/10.1083/jcb.201703094>) introduce a role for Ragulator in growth factor- and nutrient-regulated lysosomal trafficking.

Lysosomes are emerging as central regulators of a wide range of cellular functions. Besides their established role in macromolecule degradation and recycling, they control numerous other processes such as cell adhesion, cell motility, tumor invasion, cell death, metabolic signaling, cytosolic and extracellular pH, plasma membrane repair, and bone resorption. To accomplish these versatile functions, lysosomes must localize to appropriate subcellular sites. Accordingly, they can be divided into at least two spatially distinct pools, a juxtanuclear pool in the vicinity of the microtubule organizing center, which is responsible for most of the degradative “housekeeping” functions of lysosomes, and a smaller peripheral pool close to the plasma membrane that participates in various cell type-specific functions such as cell motility, invasion, bone resorption, and plasma membrane repair (Hämälistö and Jäätelä, 2016). Lysosome positioning has been linked to lysosomal function. Thus, lysosomes have to be capable of moving in response to environmental cues. In line with this, starvation induces a perinuclear lysosomal distribution required, for example, for autophagic flux (Korolchuk et al., 2011), and conversely oncogenic receptor tyrosine kinase signaling drives lysosomes to the cell periphery to promote cell motility and invasion (Rafn et al., 2012).

The centrifugal and centripetal movements of lysosomes occur mainly along the microtubule tracks and are mediated by plus end-directed kinesin motors and minus end-directed dynein motors. The small GTPase Rab7 can tether lysosomes to both dynein and kinesin motors by interacting with the adaptors RILP and protrudin/FYCO, respectively. Additionally, the coupling of lysosomes to kinesin motors can be achieved by interactions between a recently identified biogenesis of lysosome-related organelles complex 1 (BLOC1)-related complex (BORC), the Arf-like GTPase Arl8, and a kinesin-interacting linker protein SKIP (Pu et al., 2015). The hetero-octameric BORC is tethered by a myristoyl group in the N terminus of its Myrlysin subunit to the cytosolic surface of the lysosomal membrane. Despite our increasing knowledge of the proteins

involved in the bidirectional movement of lysosomes, it has remained largely unknown how cells alter lysosomal positioning in response to changes in their environment. In this issue, Filipek et al. and Pu et al. elucidate this regulation by showing that growth factor signaling and nutrient availability enhance the outward movement of lysosomes through the weakening of the lysosomal interaction between BORC and Ragulator, thereby enhancing the recruitment of Arl8/SKIP to lysosomes and the subsequent coupling of lysosomes to kinesin motors (Fig. 1).

The lysosomal Ragulator complex, which now emerges as the regulator of lysosomal positioning, is best known as an essential activation platform for metabolic signaling. It is composed of five late endosomal/lysosomal adaptor and MAPK and mammalian target of rapamycin [mTOR] activator/regulator (LAMTOR) subunits. As the long name implies, this scaffold complex regulates both MAPK and mTOR complex 1 (mTORC1). The recently solved crystal structure reveals that Ragulator is formed by tightly packed LAMTOR2/3 and LAMTOR4/5 heterodimers wrapped and held together by LAMTOR1, which anchors the complex to the lysosomal membrane by its N-terminal myristoyl and palmitoyl groups (de Araujo et al., 2017; Fig. 1). The LAMTOR2/3 heterodimer associates with RagA/B and RagC/D GTPases and serves as a guanine nucleotide exchange factor for RagA/B, whose GTP-bound forms recruit mTORC1 to the complex (Bar-Peled et al., 2012). LAMTOR2/3 also interacts with MAPK kinase 1/MEK1, thereby facilitating the activation of MAPK3/ERK1 (Wunderlich et al., 2001). Furthermore, Ragulator interacts with Axin, a scaffold protein for liver kinase B1-mediated activation of AMP-activated protein kinase (AMPK) complex on the lysosomal surface (Zhang et al., 2014). It is presently unknown which LAMTOR subunit interacts with Axin, but the ability of Axin to inhibit the guanine nucleotide exchange factor activity of LAMTOR2/3 toward RagA/B suggests that it may also bind to LAMTOR2/3 (Zhang et al., 2014). Finally, Ragulator interacts with vacuolar H⁺-ATPase and neutral amino acid transporter SLC38A9, which serve as amino acid sensors for the switching between anabolic mTORC1 and catabolic AMPK activities (Zhang et al., 2014; Wang et al., 2015; Fig. 1).

Filipek et al. (2017) and Pu et al. (2015, 2017) identified the interaction between BORC and Ragulator by tandem affinity purification with Ragulator subunits and BLOS2 (a shared subunit of BLOC1 and BORC) as baits, respectively. Using similar analyses with other BLOC1 and BORC subunits, both

Correspondence to Marja Jäätelä: mj@cancer.dk

© 2017 Colaço and Jäätelä This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms/>). After six months it is available under a Creative Commons License [Attribution-Noncommercial-Share Alike 4.0 International license, as described at <https://creativecommons.org/licenses/by-nc-sa/4.0/>].



Nutrients / growth factors absent: Perinuclear lysosomes and catabolic signaling





Figure 1. Schematic presentation of the regulation of metabolic signaling and lysosomal movement by the lysosome-associated Ragulator complex that consists of five LAMTOR subunits (LT1–5). (Top) In the absence of nutrients and growth factors, Ragulator activates the catabolic liver kinase B1 (LKB1)–AMPK signaling pathway via Axin scaffold protein and retains lysosomes in the perinuclear area by binding to the Lyspersin subunit of BORG, thereby inhibiting the Arl8/SKIP-mediated association of BORG and kinesin motors. (Bottom) In the presence of nutrients and growth factors, Axin and BORG dissociate from the Ragulator, which leads to the inactivation of AMPK and activation of BORG/Arl8/SKIP/Kinesin-mediated antegrade movement of lysosomes away from the microtubule organizing center (MTOC). Subsequently, Ragulator is more efficient in promoting Rag GTPase-dependent mTORC1 activation and Mek1-dependent Erk1 activation. In nutrient-rich conditions, mTORC1 is further stimulated by amino acid sensed by the V-H⁺-ATPase and SLC38A9.

Nutrients / growth factors present: Peripheral lysosomes and anabolic signaling



groups verified the interaction to be specific for the BORG complex, and a subsequent yeast two-hybrid (Y2H) screen for pairwise interactions between all BORG and Ragulator subunits by Pu et al. (2017) identified a strong interaction between Lyspersin and LAMTOR2. By analyzing the interactions of truncated and mutated versions of Lyspersin, both groups came to similar conclusions regarding the areas of Lyspersin that interact with LAMTOR2 and the BORG complex. Specifically, mutations in the conserved Lyspersin residues L221 (Pu et al., 2017) and K224 (Filipek et al., 2017) abolished the association with Ragulator, but not with another BORG component (Myrlysin), whereas the conserved Lyspersin residues L349/L352 (Pu et al., 2017) and K316/K357 (Filipek et al., 2017) were required for the association with Myrlysin. Pu et al. (2017) further verified these interactions using Y2H analyses. Thus, Lyspersin associates with Ragulator and BORG via two distinct but adjacent domains in its structured C-terminal part, suggesting that it can form a physical link between these two complexes.

Having established the interaction between BORG and Ragulator, Filipek et al. (2017) and Pu et al. (2017) examined its functional consequences using Lyspersin and LAMTOR1-depleted cells. Lysosomes in cells depleted for LAMTOR1

(and lacking the entire Ragulator complex) were dispersed to the cellular periphery and had increased Arl8-BORG association, whereas lysosomes depleted for Lyspersin alone or together with LAMTOR1 clustered in the perinuclear area and failed to recruit Arl8 (Filipek et al., 2017; Pu et al., 2017). C-terminal Lyspersin fragments retaining binding to BORG fully rescued the perinuclear lysosomal phenotype of Lyspersin knockout cells, whereas C-terminal fragments binding only Ragulator and N-terminal fragments binding neither BORG nor Ragulator failed to do so (Filipek et al., 2017; Pu et al., 2017). Collectively, this comprehensive set of data indicates that Ragulator negatively regulates the BORG-Arl8 association and the Arl8-mediated outward movement of lysosomes via the LAMTOR2–Lyspersin interaction, and that the C-terminal part of Lyspersin is necessary and sufficient for BORG to recruit Arl8 to lysosomes and subsequent lysosomal dispersion to the cellular periphery.

The association of both BORG and mTORC1-activating Rag GTPases with LAMTOR2 raised the question whether BORG regulates the mTORC1-activating association between Rag GTPases and Ragulator to influence lysosomal positioning, or vice versa. These questions were addressed by Pu et

al. (2017) by examining the mTORC1 activity and lysosomal positioning after the depletion of various BORC subunits and mTORC1 inhibition, respectively. Notably, BORC silencing and subsequent perinuclear clustering of lysosomes did not affect mTORC1 activity, and neither genetic nor pharmacological mTORC1 inhibitors altered lysosomal positioning (Pu et al., 2017). Based on these results, Pu et al. (2017) concluded that BORC does not interfere with mTORC1 activation and that lysosomal positioning is independent of mTORC1 activity. However, this does not exclude the possibility that BORC and Rag GTPases compete for the same or overlapping binding site in LAMTOR2 as was suggested previously by Schweitzer et al. (2015) based on the ability of overexpressed Lyspersin to disrupt the Rag GTPase–Ragulator interaction and inhibit mTORC1 activity. In line with this, silencing the relatively little expressed Lyspersin by Pu et al. (2017) may have increased the binding of Rag GTPase–mTORC1 without having a detectable increase in already high mTORC1 activity. Additionally, inhibiting mTORC1 signaling may have failed to affect lysosomal positioning because Rag GTPases could compete with Lyspersin for LAMTOR2 binding even after the dissociation of mTORC1. Supporting this possibility, Filipek et al. (2017) demonstrated by size exclusion chromatography of the LAMTOR1 interactome that most Ragulator complexes are associated with Rag GTPases, whereas only a relatively minor subpopulation associated with endogenous BORC subunits (Filipek et al., 2017). Also, their data showing that affinity purification using a BORC subunit (BLOS1) as bait copurified the other BORC subunits and the Ragulator complex, but not Rag GTPases, supports the competitive binding of BORC and Rag GTPases to Ragulator (Filipek et al., 2017). In this context, it should be noted that LAMTOR2 also interacts with various other proteins (Fig. 1), and that the strength of its interaction with one protein can shift that with others, as previously suggested for Axin and mTORC1 (Zhang et al., 2014). Such a competitive scenario for Lyspersin and Rag GTPases could also explain the data by Pu et al. (2017) and Filipek et al. (2017) demonstrating that EGF signaling and amino acid availability, both of which are known activators of mTORC1, weaken the interaction between LAMTOR2 and Lyspersin to trigger the outward movement of lysosomes. Thus, the regulation of numerous protein interactions around the “crowded” Ragulator complex may occur in a competitive fashion to coordinate the metabolic signaling pathways and lysosomal positioning in response to changing needs of the cell (Fig. 1).

How such competitive binding could be regulated by growth factors and nutrients remains largely unknown, and the data presented by Pu et al. (2017) and Filipek et al. (2017) offer some directions to be further investigated. Performing surface hydrophobicity analyses of the LAMTOR2/3 heterodimer crystal structure, Pu et al. (2017) predicted the presence of a hydrophobic patch on a LAMTOR2 surface distal to LAMTOR3. The subsequent Y2H analyses revealed that alanine substitutions of several hydrophobic residues at this site abolished the binding of LAMTOR 2 to Lyspersin but not to LAMTOR3, suggesting that the hydrophobic surface of LAMTOR2 mediates its interaction with BORC (Pu et al., 2017). Furthermore, Filipek et al. (2017) showed that EGF, a well-established activator of the MEK1–ERK pathway, inhibited this interaction. Thus, it is tempting to speculate that the LAMTOR2/3-associated local activation of the MAPK3 could result in the phosphorylation of the hydrophobic surface of LAMTOR2, thereby reducing

hydrophobicity, releasing BORC from Ragulator, and promoting lysosomal movement toward the periphery. Such a model is fascinating in the context of metastatic cancer cells, where lysosomal peripheral distribution seems paramount to the invasive and aggressive phenotype. For instance, the ErbB2 oncogene, which can be activated by EGF, promotes the accumulation of lysosomes in cell protrusions and invasion of breast cancer cells (Rafn et al., 2012). These events are mediated by complex signaling events, including the activation of the MAPK–ERK pathway (Rafn et al., 2012). It remains, however, to be tested whether the MAPK–ERK pathway or other EGF/ErbB2-induced signaling pathways directly control the hydrophobicity of LAMTOR2 and thereby the interaction between LAMTOR2 and Lyspersin. If this is the case, it would open attractive therapeutic possibilities to control lysosomal trafficking by targeting MAPK–ERK or other kinases responsible for LAMTOR2 phosphorylation.

Overall, it is becoming evident that Ragulator plays a variety of cellular functions in regulating lysosomal biogenesis, activity, and, as illustrated in these two papers, positioning. Identification of Ragulator (via Lamtor2) as the negative regulator of anterograde directed movement and demonstration that this regulation is controlled by both amino acids and the EGF, highlights the varied levels of cellular regulation that converge on the Ragulator complex. How these proteins spatially interact with the Ragulator and how such interactions are regulated will be important in understanding how a Rag GTPase–LAMTOR2 interaction influences a Lyspersin–LAMTOR2 interaction, and will better enable us to manipulate the regulatory pathways to target lysosomal positioning for therapies ranging from the aforementioned invasive cancer to neurodegenerative disorders and infectious diseases.

Acknowledgments

We apologize to those colleagues whose work could not be cited because of strict space restrictions.

The related work in the Jäätelä laboratory is supported by the European Research Council (AdG 340751), Danish National Research Foundation (DNRF125), Danish Cancer Society (R90-A5783), and Novo Nordisk Foundation (NNF15OC0016914).

The authors declare no competing financial interests.

References

- Bar-Peled, L., L.D. Schweitzer, R. Zoncu, and D.M. Sabatini. 2012. Ragulator is a GEF for the rag GTPases that signal amino acid levels to mTORC1. *Cell*. 150:1196–1208. <https://doi.org/10.1016/j.cell.2012.07.032>
- de Araujo, M.E.G., A. Naschberger, B.G. Fürnrohr, T. Stasyk, T. Dünzendorfer-Matt, S. Lechner, S. Welti, L. Kremser, G. Shivalingaiah, M. Offenderinger, et al. 2017. Crystal structure of the human lysosomal mTORC1 scaffold complex and its impact on signaling. *Science*. 358:377–381. <https://doi.org/10.1126/science.aao1583>
- Filipek, P.A., M.E.G. de Araujo, G.F. Vogel, C.H. De Smet, D. Eberharter, M. Rebsamen, E.L. Rudashevskaya, L. Kremser, T. Yordanov, P. Tschaikner, et al. 2017. LAMTOR/Ragulator is a negative regulator of Arl8b- and BORC-dependent late endosomal positioning. *J. Cell Biol.* <https://doi.org/10.1083/jcb.201703061>
- Hämälistö, S., and M. Jäätelä. 2016. Lysosomes in cancer-living on the edge (of the cell). *Curr. Opin. Cell Biol.* 39:69–76. <https://doi.org/10.1016/j.ccb.2016.02.009>
- Korolchuk, V.I., S. Saiki, M. Lichtenberg, F.H. Siddiqi, E.A. Roberts, S. Imarisio, L. Jahreiss, S. Sarkar, M. Futter, F.M. Menzies, et al. 2011. Lysosomal positioning coordinates cellular nutrient responses. *Nat. Cell Biol.* 13:453–460. <https://doi.org/10.1038/ncb2204>
- Pu, J., C. Schindler, R. Jia, M. Jarnik, P. Backlund, and J.S. Bonifacio. 2015. BORC, a multisubunit complex that regulates lysosome positioning. *Dev. Cell*. 33:176–188. <https://doi.org/10.1016/j.devcel.2015.02.011>

- Pu, J., T. Keren-Kaplan, and J.S. Bonifacino. 2017. A Ragulator–BORC interaction controls lysosome positioning in response to amino acid availability. *J. Cell Biol.* <https://doi.org/10.1083/jcb.201703094>
- Rafn, B., C.F. Nielsen, S.H. Andersen, P. Szyniarowski, E. Corcelle-Termeau, E. Valo, N. Fehrenbacher, C.J. Olsen, M. Daugaard, C. Egebjerg, et al. 2012. ErbB2-driven breast cancer cell invasion depends on a complex signaling network activating myeloid zinc finger-1-dependent cathepsin B expression. *Mol. Cell.* 45:764–776. <https://doi.org/10.1016/j.molcel.2012.01.029>
- Schweitzer, L.D., W.C. Comb, L. Bar-Peled, and D.M. Sabatini. 2015. Disruption of the Rag-Ragulator Complex by c17orf59 Inhibits mTORC1. *Cell Reports.* 12:1445–1455. <https://doi.org/10.1016/j.celrep.2015.07.052>
- Wang, S., Z.Y. Tsun, R.L. Wolfson, K. Shen, G.A. Wyant, M.E. Plovanich, E.D. Yuan, T.D. Jones, L. Chantranupong, W. Comb, et al. 2015. Metabolism. Lysosomal amino acid transporter SLC38A9 signals arginine sufficiency to mTORC1. *Science.* 347:188–194. <https://doi.org/10.1126/science.1257132>
- Wunderlich, W., I. Fialka, D. Teis, A. Alpi, A. Pfeifer, R.G. Parton, F. Lottspeich, and L.A. Huber. 2001. A novel 14-kilodalton protein interacts with the mitogen-activated protein kinase scaffold mp1 on a late endosomal/lysosomal compartment. *J. Cell Biol.* 152:765–776. <https://doi.org/10.1083/jcb.152.4.765>
- Zhang, C.S., B. Jiang, M. Li, M. Zhu, Y. Peng, Y.L. Zhang, Y.Q. Wu, T.Y. Li, Y. Liang, Z. Lu, et al. 2014. The lysosomal v-ATPase-Ragulator complex is a common activator for AMPK and mTORC1, acting as a switch between catabolism and anabolism. *Cell Metab.* 20:526–540. <https://doi.org/10.1016/j.cmet.2014.06.014>