

SPOTLIGHT

Fine-tuning activity-dependent bulk endocytosis via kinases and phosphatases

 Ira Milosevic^{1,2}  and Michael A. Cousin^{3,4,5} 

The regulation of activity-dependent bulk endocytosis, the dominant mode of membrane retrieval in response to intense neuronal activity, is poorly understood. In this *JCB* issue, Peng et al. (2021. *J. Cell. Biol.* <https://doi.org/10.1083/jcb.202011028>) propose a novel molecular mechanism for the coordination of activity-dependent bulk endocytosis that builds on Minibrain kinase and its presynaptic substrate synaptojanin-1.

Brain function necessitates sustained synaptic transmission regardless of activity demands. The preservation of synaptic transmission depends on the efficient (re)formation of synaptic vesicles (SVs) by endocytosis after their insertion into the synaptic plasma membrane during neuronal stimulation (1). During mild and sparse stimulation, the dominant endocytosis modes are ultrafast endocytosis and clathrin-mediated endocytosis (CME; 1). Both modes appear to have a fixed rate and limited capacity, and therefore cannot adapt to high frequency stimulations that accumulate inserted SV membranes at the presynaptic terminal. Under these conditions, a different endocytosis mode is predominantly used, termed activity-dependent bulk endocytosis (ADBE). ADBE retrieves large areas of the presynaptic plasma membrane to form bulk endosomes, from which new SVs are then generated (1). This form of endocytosis is particularly common in synapses that operate with high rates of neurotransmission, e.g., ribbon synapses of sensory neurons. ADBE contributes to presynaptic plasticity, having recently been demonstrated to control neurotransmitter release probability (2). Importantly, defects in ADBE and SV endocytosis in general have profound consequences on neuronal function and

survival, with dysfunction linked to a series of neurodevelopmental disorders (3).

Considering the importance of ADBE to brain physiology and pathology, it is essential to understand the molecular machinery that controls this process and synchronizes it with other synaptic events. Amazingly, despite the fact that ADBE was described in the early 1970s, its regulation remains mysterious. Several protein kinases and phosphatases that contribute to regulation of CME and other endocytosis modes (1) may also contribute to ADBE. For example, the calcium/calmodulin-dependent phosphatase calcineurin activates ADBE, working with glycogen synthase kinase-3 to provide bidirectional control via the phosphorylation of specific substrates (4). However, many presynaptic proteins are calcineurin substrates, suggesting other protein kinases may perform complementary roles.

In a recent paper, Chang and colleagues (5) present data in support of calcineurin and Minibrain (Mnb) as coregulators of ADBE in fruit flies via bidirectional control of the phosphorylation status of synaptojanin (Synj)-1 phosphatase. The authors argue that the Synj-1 phosphorylation status coordinates the activity-dependent balance between CME versus ADBE (Fig. 1). Namely, during mild stimulation CME is promoted by Mnb, while ADBE is inhibited. During

intense stimulation, dephosphorylation of Synj-1 by calcineurin is required to activate ADBE (Fig. 1). An interesting novel aspect arises from examination of domain-specific Synj-1 mutants: its 4'-phosphatase SAC1 activity supports ADBE, while its 5'-phosphatase (5'-PPase) domain suppresses it. The Bin/Amphiphysin/Rvs domain protein endophilin-A has been implicated in ADBE (6); however, a Synj-1 mutant lacking the endophilin-A binding proline-rich domain (PRD) had no effect. Further studies may therefore be required to dissect synaptojanin-1-dependent and -independent roles of endophilin in ADBE.

Collectively, the data by Chang and colleagues consolidate the key role played by calcineurin in ADBE and identify Mnb as a new ADBE protein kinase. Intriguingly, the number of synapses performing ADBE is increased in Mnb hypomorphs, suggesting there is additional endocytic capacity that can be recruited on demand. There also appears to be bidirectional control of ADBE via Mnb, since Mnb overexpression represses this pathway. Notably, the enzyme activities of Synj-1 are regulated by Mnb- and calcineurin-dependent turnover of phosphorylation of S1029 (Fig. 1; 7, 8). In mammals, cyclin-dependent kinase 5 is suggested to control Synj-1 activity (9); therefore, it is important to confirm whether

¹Wellcome Centre for Human Genetics, Nuffield Department of Medicine, National Institute for Health Research Oxford Biomedical Research Centre, Oxford, UK;

²University of Coimbra, Multidisciplinary Institute of Ageing, Coimbra, Portugal; ³Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh, UK; ⁴Muir Maxwell Epilepsy Centre, University of Edinburgh, Edinburgh, UK; ⁵Simons Initiative for the Developing Brain, University of Edinburgh, Edinburgh, UK.

Correspondence to Ira Milosevic: ira.milosevic@well.ox.ac.uk; Michael A. Cousin: M.Cousin@ed.ac.uk.

© 2021 Milosevic and Cousin. This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms/>). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 4.0 International license, as described at <https://creativecommons.org/licenses/by-nc-sa/4.0/>).




Figure 1. Control of CME and ADBE via Minibrain kinase and calcineurin phosphatase. Synj-1 is phosphorylated by Mnb kinase on Ser1029 on its PRD. This promotes the 5'-PPase activity of Synj-1 and inhibits association with the endocytosis protein endophilin-A. These events promote CME. During intense neuronal activity, calcineurin (CaN) is activated and dephosphorylates Synj-1. This reduces 5'-PPase activity and promotes association with endophilin. The dephosphorylation also promotes ADBE via inhibition of Synj-1 5'-PPase activity. This phospho-regulation of the endophilin interaction does not impact ADBE. The SAC activity of Synj-1 is essential for ADBE and is unaffected by phosphorylation.

Synj-1 is also phosphorylated by the Mnb orthologue, dual specificity tyrosine-phosphorylation-regulated kinase (DYSRK1A), in mammals. A key test of the causality of activity-dependent phosphorylation events is whether they occur to the same stimulation intensities as the biological event. In this study, activity-dependent dephosphorylation of S1029 on Synj-1 was not demonstrated; instead, an absence of activity-dependent Mnb phosphorylation was observed. In mitigation, the authors convincingly demonstrated that Synj-1 phosphorylation increased during prolonged stimulus in the absence of calcineurin function.

This work also confirmed a key role for the phospholipid PI(4,5)P₂ in ADBE (1). Interestingly, it further revealed a hitherto undiscovered role for the SAC domain, but not the 5'-PPase domain of Synj-1 in ADBE. This latter activity is essential for other forms of endocytosis, such as CME and ultrafast endocytosis, with SAC activity required for clathrin-dependent vesicle generation from endosomes (10, 11). In addition to potential roles for Synj-1 SAC activity discussed by Chang and colleagues, a more provocative (and simplistic) explanation is

that the end product, phosphatidylinositol (PI) itself, is important for ADBE. In support, the neurons without diacylglycerol kinase (which generates the PI precursor phosphatidic acid) display SV endocytosis defects that are exacerbated during high activity (12).

A lack of accurate assays that monitor ADBE in both time and space has limited research in small nerve terminals for decades. In this work, ADBE is evoked and monitored using multiple approaches. This is important, since there is no simple method to monitor ADBE; therefore, it requires cross corroboration wherever possible. This study was greatly assisted via the use of genetically tractable model organisms, allowing precise intervention to abate the function of key proteins and enzymes *in vivo*. Yet, the trade-off is the relative imprecision of stimulation to evoke SV turnover, with prolonged periods of stimulation (and parallel inhibition of CME) required to evoke and isolate ADBE.

Since Peng et al. shed light on new aspects of ADBE regulation, further questions can now be envisioned. In particular, how localized production and degradation of

membrane phospholipids coordinate the temporal and spatial triggering of specific endocytosis modes. The essential role for calcineurin in most forms of endocytosis suggests where and when dephosphorylation events occur at the presynapse may be critical in the recruitment of discrete SV reformation pathways. Furthermore, Mnb/DYSRK1A is linked to brain pathologies, including Down's syndrome and autism-spectrum disorders, which is yet to be explored. These and other questions will no doubt drive further studies of remarkable plasticity when it comes to formation of new SVs and synaptic transmission, and how they organize and govern our brain activity.

Acknowledgments

We apologize for any references omitted due to space constraints. We thank Dr. Elizabeth Davenport for production of the Spotlight figure.

I. Milosevic is supported by funding from the John Black Foundation, John Fell Fund (H5D00100), and Horizon 2020 (MIA-Portugal/857524), while M.A. Cousin is supported by The Wellcome Trust (204954/Z/16/Z), Simons Foundation (529508), Epilepsy Research UK (P2003), and Cure Huntington's Disease Initiative (A-11210).

The authors declare no competing financial interests.

References

1. Chanaday, N.L., et al. 2019. *J. Neurosci.* <https://doi.org/10.1523/JNEUROSCI.1158-19.2019>
2. Ivanova, D., et al. 2021. *Sci. Adv.* <https://doi.org/10.1126/sciadv.abf3873>
3. Bonnycastle, K., et al. 2021. *J. Neurochem.* <https://doi.org/10.1111/jnc.15035>
4. Clayton, E.L., et al. 2010. *Nat. Neurosci.* <https://doi.org/10.1038/nn.2571>
5. Peng, Y.J., et al. 2021. *J. Cell Biol.* <https://doi.org/10.1083/jcb.202011028>
6. Kononenko, N.L., et al. 2014. *Neuron.* <https://doi.org/10.1016/j.neuron.2014.05.007>
7. Chen, C.K., et al. 2014. *Nat. Commun.* <https://doi.org/10.1038/ncomms5246>
8. Geng, J., et al. 2016. *J. Neurosci.* <https://doi.org/10.1523/JNEUROSCI.1470-16.2016>
9. Lee, S.Y., et al. 2004. *Proc. Natl. Acad. Sci. USA.* <https://doi.org/10.1073/pnas.0307813100>
10. Mani, M., et al. 2007. *Neuron.* <https://doi.org/10.1016/j.neuron.2007.10.032>
11. Watanabe, S., et al. 2018. *Neuron.* <https://doi.org/10.1016/j.neuron.2018.06.005>
12. Goldschmidt, H.L., et al. 2016. *Cell Rep.* <https://doi.org/10.1016/j.celrep.2015.12.022>