

REVIEW

Structure and function of the synaptonemal complex

Brenda I. Cesar¹ and Yumi Kim¹

A defining feature of meiosis is the synaptonemal complex (SC), a zipper-like protein structure that forms between homologous chromosomes to regulate their recombination and segregation. Historically viewed as an enigmatic electron-dense scaffold, the SC is now recognized as a dynamic signaling platform that coordinates key meiotic processes. Here, we review recent advances in understanding SC structure and function. We describe diverse complementary approaches that have expanded the catalog of SC components and their network of interactions within this architecture. We highlight striking conservation in structural organization and ancient molecular modules that couple SC structure to crossover regulation and further discuss how the SC implements feedback mechanisms controlling meiotic DNA break formation and repair capacity to ensure faithful chromosome segregation across generations.

Introduction

Sexual reproduction in eukaryotes depends on meiosis, a specialized cell division that produces haploid gametes from a diploid precursor. Unique to meiosis is its prolonged prophase I, during which homologous chromosomes pair, synapse, and exchange genetic material through crossover formation. These crossovers, together with sister chromatid cohesion, create the physical linkages needed for proper homolog alignment and segregation during meiosis I. Beyond their mechanical role, crossovers also shuffle genetic information between maternal and paternal chromosomes, generating new allele combinations that enhance genetic diversity and drive eukaryotic evolution.

In most eukaryotes, homologous chromosome synapsis is reinforced by the assembly of the synaptonemal complex (SC). First discovered over 70 years ago by electron microscopy (Fawcett, 1956; Moses, 1956), the SC is a ladder-like protein structure that forms between homologous chromosomes. Despite its absence in some protists and fungi, the SC displays remarkable conservation across diverse organisms, maintaining a characteristic tripartite architecture comprising two parallel lateral elements and a central region that bridges the two chromosome axes in a zipper-like fashion (Page and Hawley, 2004). The central region is subdivided into transverse filaments that span the width of the SC and central elements that form the midline of the complex (Fig. 1 A).

Initial insights into SC structure and function came from electron microscopy observations. The zipper-like appearance of the SC and the strong correlation between chromosome synapsis and crossover frequency suggested that this structure functions to align homologous chromosomes and promote crossover formation (Maguire, 1965). Moreover, recombination nodules, electron-dense ellipsoidal structures thought to represent protein complexes at sites of ongoing recombination, were observed

in association with the SC in *Drosophila melanogaster* females, including early recombination nodules that appear concurrent with SC assembly during zygotene and late nodules whose number and distribution closely matched those of chiasmata (Carpenter, 1975; Carpenter, 1979). Similar structures were subsequently reported in several other species (Anderson et al., 2001), establishing a direct link between the SC and recombination sites. Notably, *Drosophila* males lack both SC and meiotic recombination, further supporting this correlation (Rasmussen, 1973).

While these correlations between the presence of the SC and crossover formation seemed compelling, molecular identification of SC components and mutant analyses have substantially revised our understanding of its function. Contrary to early models that the SC creates preconditions for crossover formation by bridging homologs into close apposition (von Wettstein et al., 1984), SC assembly in most eukaryotes, including the budding yeast *Saccharomyces cerevisiae*, plants, and mammals, depends on the initiation of meiotic recombination through programmed DNA double-strand breaks (DSBs) (Alani et al., 1990; Padmore et al., 1991; Baudat et al., 2000; Romanienko and Camerini-Otero, 2000; Grelon et al., 2001). However, this requirement is not universal. In *Drosophila* females and *Caenorhabditis elegans*, the SC forms independently of DSB formation (Dernburg et al., 1998; McKim et al., 1998), yet crossovers strictly depend on SC assembly (Page and Hawley, 2001; MacQueen et al., 2002).

The relationship between the SC and crossover formation becomes even more complex when examining organisms with reduced or absent SC. In most organisms, crossovers exhibit interference, a phenomenon in which the presence of a crossover reduces the probability of additional crossovers nearby, thereby distributing crossovers more evenly along chromosomes

¹Department of Biology, Johns Hopkins University, Baltimore, MD, USA.

Correspondence to Yumi Kim: yumi.kim@jhu.edu.

© 2026 Cesar and Kim. This article is distributed under the terms as described at <https://rupress.org/pages/terms102024/>.




Figure 1. SC structure and protein components across model organisms. (A) Schematic of the SC architecture showing the lateral elements (blue), which associate with chromatin (gray), connected by the SC central region comprising transverse filaments (dark green) and central element proteins (light green). **(B)** A current compilation of known SC subunits. Proteins are organized by their localization, interaction, and functions within the SC. For axis-associated cohesins, only the meiosis-specific cohesin subunits are shown. SYCE3 and SYP-2 are grouped with transverse filament components because of their direct interactions.

(Muller, 1916). The fission yeast *Schizosaccharomyces pombe* lacks a canonical SC structure and forms crossovers with little or no interference (Olson et al., 1978; Bähler et al., 1993). *Aspergillus* species lack SC entirely yet still form crossovers, with *Aspergillus fumigatus* producing up to 30 per chromosome pair, the highest known in any eukaryote (Egel-Mitani et al., 1982; Auxier et al., 2023). Strikingly, disrupting SC assembly in the model plant *Arabidopsis thaliana* increases crossover numbers while abolishing interference (Capilla-Pérez et al., 2021; France et al., 2021). These findings reveal a complex, species-specific relationship between SC assembly and crossover formation, suggesting that the SC may primarily function to limit crossover numbers and impose interference rather than to promote crossover formation.

How can we reconcile this paradox? Recent advances have provided key insights into SC architecture and function. Continued genetic studies and biochemical purification, complemented by AlphaFold-based structural predictions revealing protein-protein interactions, have identified previously missing subunits across model organisms (Collins et al., 2014; Gómez-H et al., 2016; Hurlock et al., 2020; Zhang et al., 2020; Vrielynck

et al., 2023; Feng et al., 2025) (Fig. 1 B). Super-resolution and expansion microscopy, together with quantitative live imaging, have revealed the molecular organization and biophysical properties of the SC (Schücker et al., 2015; Cahoon et al., 2017; Rog et al., 2017; Wang et al., 2018; Xu et al., 2019; Zwertler et al., 2020; Köhler et al., 2025). In parallel, progress in biochemical reconstitution and structural biology has begun to map protein interfaces within the SC (Dunce et al., 2018; Dunce et al., 2021; Blundon et al., 2024), advancing toward in vitro reconstitution of the entire complex. Here, we provide an updated overview of the SC structure and function, beginning with the chromosome axes that form the structural foundation for SC assembly. We focus particularly on how conserved molecular modules within the SC regulate crossover formation and implement feedback mechanisms to ensure the obligate crossover essential for accurate homolog segregation.

Chromosome axis assembly

Axis assembly is one of the earliest events in meiotic prophase and is essential for all subsequent processes, including meiotic




Figure 2. Meiotic chromosome axis assembly and recombination control. (A) Assembly of the meiotic chromosome axis. Meiosis-specific cohesin complexes (light blue rings) extrude chromatin into loops and provide the structural foundation for recruiting coiled-coil proteins and meiotic HormADs. (B) Rec114–Mei4–Mer2 (RMM) complexes are recruited to the chromosome axis by HormADs and form nucleoprotein condensates, recruiting the Spo11 complex and inducing meiotic DSBs. (C) Upon synapsis, HormADs are actively removed from chromosome axes by TRIP13/Pch2, thereby dismantling the DSB-forming machinery and preventing further DSB formation. Unsynapsed chromosome axes retain HormAD proteins, which recruit ATR kinase and BRCA1 to activate checkpoint signaling.

DSB formation, homolog pairing, and SC assembly (Ur and Corbett, 2021). Once homologs synapse, these axes (also called axial elements) become the lateral elements of the SC, which are then connected by the transverse filaments that span the ~100 nm distance between paired chromosomes (Fig. 1 A). Axis formation begins when meiosis-specific cohesin complexes containing the kleisin subunit Rec8 establish sister chromatid cohesion and organize chromatin into arrays of loops anchored around a central proteinaceous core (Schalbetter et al., 2019). This cohesin-based scaffold provides a structural framework for the recruitment of conserved coiled-coil proteins, which assemble into tetrameric complexes that further oligomerize into filamentous structures (West et al., 2019) (Fig. 2 A). These include Red1 in budding yeast (Smith and Roeder, 1997), SYCP2/SYCP3 in mammals (Lammers et al., 1994; Offenberger et al., 1998), and ASY3/ASY4 in plants (Ferdous et al., 2012; Chambon et al., 2018) (Fig. 1 B). Although these proteins show little sequence homology across species, they act as functional homologs through shared closure motifs that recruit meiotic HormADs, named after their conserved HormA (Hop1, Rev7, and Mad2) domains, to the chromosome axes.

While chromosome axis components are broadly conserved across eukaryotes, some species have evolved distinct mechanisms for axis assembly. In *Drosophila*, meiotic HormADs and Red1 homologs are absent (Tromer et al., 2021). Instead, specialized cohesin-related proteins (including C(2)M, ORD, SOLO,

and SUNN) mediate sister chromatid cohesion and organize chromosome axes (Webber et al., 2004; Yan and McKee, 2013; Krishnan et al., 2014; Gyuricza et al., 2016). In contrast, *C. elegans* encodes four meiotic HormADs (HIM-3, HTP-1, HTP-2, and HTP-3) that assemble hierarchically to form the chromosome axis but lacks Red1 homologs (Zetka et al., 1999; Couteau and Zetka, 2005; Martinez-Perez and Villeneuve, 2005; Kim et al., 2014; West et al., 2019). Despite these differences in composition, axis formation is universally required for interhomolog recombination, a role mediated in large part by meiotic HormADs.

HORMAD proteins promote DSB formation and interhomolog recombination

Once recruited to chromosome axes, HormADs play essential roles in meiotic recombination (Gu et al., 2022). First, they promote DSB formation by recruiting the conserved Rec114–Mei4–Mer2 complex (known in mammals as REC114–MEI4–IHO1) to chromosome axes through direct interaction with Mer2/IHO1 (Panizza et al., 2011; Stanzione et al., 2016; Rousova et al., 2021; Laroussi et al., 2023). Recent studies further show that the Rec114–Mei4 complex and Mer2 independently form DNA-driven nucleoprotein phase-separated condensates in vitro, which may help recruit the Spo11 holoenzyme to chromatin loops tethered to the axis (Claeys Bouuaert et al., 2021) (Fig. 2 B). Interestingly, *C. elegans* lacks obvious Mer2/IHO1 homologs, while

duplicated REC114 orthologs DSB-1 and DSB-2 form complexes with the MEI4 ortholog DSB-3 that load onto chromatin without clear axis enrichment (Rosu et al., 2013; Stamper et al., 2013; Hinman et al., 2021). Nevertheless, HTP-3 is essential for DSB formation through association with MRE-11/RAD-50 (Goodyer et al., 2008).

Following DSB formation, HORMADs also help establish interhomolog bias, ensuring that recombination occurs preferentially between homologous chromosomes rather than sister chromatids. Rec8-containing cohesin creates a default sister bias by holding sister chromatids together, and genetic assays in budding yeast suggest that intersister repair accounts for up to one-third of meiotic recombination events (Goldfarb and Lichten, 2010; Kim et al., 2010). In response to meiotic DSBs, Mec1 and Tel1 (the yeast homologs of ATR and ATM kinases) phosphorylate Hop1, which in turn recruits and activates the meiosis-specific kinase Mek1 to chromosome axes (Carballo et al., 2008). Mek1 then phosphorylates Rad51 regulators, including Rad54 and Hed1, which together suppress Rad51-mediated strand invasion between sister chromatids (Niu et al., 2009; Callender et al., 2016). By inhibiting Rad51 activity through this mechanism, the meiosis-specific recombinase Dmcl becomes the primary mediator of strand invasion, favoring the use of homologous chromosomes as templates (Schwacha and Kleckner, 1997; Lao et al., 2013). Together, axis components and Dmcl counteract the intrinsic sister preference by Rec8 cohesin to generate strong interhomolog bias (Kim et al., 2010; Hong et al., 2013).

In mouse oocytes deficient for HORMAD1 or HORMAD2, irradiation-induced DSBs are rapidly repaired through intersister pathways (Shin et al., 2013; Rinaldi et al., 2017), suggesting that mammals also use HORMAD-dependent mechanisms to establish interhomolog bias. Meiotic HORMADs preferentially accumulate on unsynapsed chromosome axes and are actively removed from synapsed regions by the AAA+ ATPase TRIP13 (Wojtasz et al., 2009) (Fig. 2 C). Thus, their removal upon synapsis is thought to relieve the HORMAD-dependent barrier that prevents intersister repair. HORMAD2 recruits ATR to unsynapsed axes, where ATR signals through its effector kinase CHK1 to promote the loading of RAD51 and DMC1 recombinases onto resected DSBs, thereby facilitating interhomolog strand invasion and SC elongation (Pacheco et al., 2018; Widger et al., 2018). Although HORMAD1 and HORMAD2 themselves undergo ATR-dependent phosphorylation (Fukuda et al., 2012; Royo et al., 2013), it remains unknown whether this modification directly contributes to recombinase recruitment.

Plant meiosis employs similar axis-dependent mechanisms to promote interhomolog recombination, though with distinct regulatory features. In *Arabidopsis*, the HORMAD protein ASY1 is dispensable for meiotic DSB formation but is crucial for sustaining DMC1 activity and promoting interhomolog recombination (Sanchez-Moran et al., 2007). In the absence of DMC1 and/or ASY1, DSBs are efficiently repaired, likely using sister chromatids without resulting in chromosome fragmentation (Sanchez-Moran et al., 2007). Interhomolog bias therefore emerges from ASY1-mediated suppression of intersister repair rather than being an intrinsic property of the recombinase

(Kurzbaue et al., 2012). ASY1 achieves this by counteracting FIGL1, an AAA-ATPase that limits RAD51/DMC1-mediated interhomolog repair (Emmenecker et al., 2024). Thus, while the basic logic of axis proteins working with DMC1 to direct repair toward homologs is conserved, plants appear to rely on a balance between ASY1 and FIGL1 rather than phosphorylation cascades that suppress RAD51.

C. elegans, which lacks DMC1, provides another variation. RAD-51 foci form and disappear without delay in *him-3* and *htp-1* mutants, even though homologs are unavailable, suggesting that these HORMAD proteins promote interhomolog repair by creating a barrier between sister chromatids (Couteau et al., 2004; Martinez-Perez and Villeneuve, 2005). These evolutionary variations highlight that while the requirement for interhomolog recombination during meiosis is universal, the molecular strategies to achieve this outcome differ considerably across lineages.

Identification of SC central region components

Compared with chromosome axis components, defining the molecular composition of the SC central region has been challenging because of the rapid sequence divergence of its proteins (Hemmer and Blumenstiel, 2016). Below, we outline the complementary approaches that have enabled the identification of transverse filament proteins and, more recently, central element components.

Transverse filament proteins

The first transverse filament protein characterized at the molecular level was rat SYCP1 (formally known as SCP1), identified through screening of cDNA libraries with antibodies generated from purified SCs (Meuwissen et al., 1992). SYCP1 has predicted coiled-coil domains that span the width of the SC, bridging the lateral elements. Shortly thereafter, budding yeast Zip1, discovered in forward genetic screens, was shown to form the transverse filaments of the SC, sharing structural organization with SYCP1 despite limited sequence homology (Sym et al., 1993).

Transverse filament proteins in other model organisms were subsequently identified through various experimental approaches. In *Drosophila*, the *c(3)G* gene, long known from mutants lacking both SC and meiotic crossovers (Hall, 1972), was later identified as the transverse filament component (Page and Hawley, 2001). In *C. elegans*, SYP-1 was discovered through forward genetic screens (MacQueen et al., 2002), while two paralogs, SYP-5 and SYP-6, were identified recently as additional transverse filament proteins through biochemical purification (Hurlock et al., 2020; Zhang et al., 2020). In *Arabidopsis*, ZYP1a and ZYP1b were found through bioinformatic searches for proteins with predicted coiled-coil domains flanked by globular domains (Higgins et al., 2005).

Central element proteins

Central element proteins have been discovered more recently, using increasingly diverse strategies. Expression profiling and

localization studies identified the first mammalian central element proteins, SYCE1 and SYCE2 (Costa et al., 2005), followed by TEX12, which interacts with SYCE2 (Hamer et al., 2006). SYCE3 was later shown to be required for the loading of other central element proteins onto chromosomes (Schramm et al., 2011). SIX6OS1 was discovered through a human genome-wide association study that linked a nonsynonymous variant to an elevated female recombination rate (Kong et al., 2014). Subsequent functional analyses confirmed that SIX6OS1 encodes a component of the SC central element (Gómez-H et al., 2016), illustrating how population genetics can uncover novel constituents of the SC.

Beyond mammals, a combination of forward genetic screens, RNA interference screens, yeast two-hybrid assays, and computational approaches has expanded the repertoire of central element proteins across model organisms. Early examples include Ecm11 and Gmc2 in budding yeast (Humphryes et al., 2013); Cona and Corolla in *Drosophila* (Page et al., 2008; Collins et al., 2014); and SYP-2, SYP-3, and SYP-4 in *C. elegans* (Colaiácovo et al., 2003; Smolikov et al., 2007; Smolikov et al., 2009). Biochemical purification coupled with mass spectrometry has proven powerful for identifying previously missing SC components, including paralogs with partially redundant functions and moonlighting proteins with canonical roles outside meiosis. This approach identified two paralogous Skp1-related proteins, SKR-1 and SKR-2, as structural subunits of the SC in *C. elegans* (Blundon et al., 2024). Remarkably, six *C. elegans* SC subunits form a long-sought soluble protein complex in vitro, suggesting that the entire set of SC subunits is now known in this organism (Blundon et al., 2024). Complementary approaches, including transcriptomics-based candidate screening and proximity labeling, have further expanded the inventory of central element proteins, leading to the discovery of SCEP1, SCEP2, and SCEP3 in *A. thaliana* (Vrielynck et al., 2023; Feng et al., 2025).

A current compilation of known SC subunits across model organisms is shown in Fig. 1 B. These findings underscore that our knowledge of the molecular composition of the SC remains incomplete, even in well-studied model organisms. Nevertheless, as the catalog of SC components and their interaction networks have expanded, conserved structural and functional features have begun to emerge from comparative analyses across species.

Molecular features that govern SC assembly

Structural organization of transverse filaments

One of the most conserved structural features of the SC is its ~100 nm width (von Wettstein et al., 1984), which is defined by the length of the coiled-coil domains of transverse filament proteins (Sym and Roeder, 1995; Ollinger et al., 2005; Duncce et al., 2018). Across diverse organisms, these proteins assemble in a head-to-head configuration with their N termini at the midline and their C termini oriented toward the chromosome axes (Liu et al., 1996; Dong and Roeder, 2000; Anderson et al., 2005; Schild-Prüfert et al., 2011; Hurlock et al., 2020; Billmyre et al., 2023) (Fig. 1 A). When overexpressed in yeast or somatic

cells, transverse filament proteins form higher-order structures known as polycomplexes that recapitulate ultrastructural features of the SC, indicating an inherent propensity for self-assembly (Sym and Roeder, 1995; Ollinger et al., 2005). Biochemical analyses have further revealed that SYCP1 forms a tetramer through N-terminal interactions in vitro, forming the interface for head-to-head assembly that subsequently polymerizes into a zipper-like lattice (Duncce et al., 2018). The functional importance of this interface was recently validated as mutations that disrupt SYCP1 head-to-head assembly completely abolish synapsis in mouse spermatocytes (Billmyre et al., 2023). Similarly, studies in *Drosophila* have demonstrated that the N terminus of C(3)G is essential for SC assembly, while the C-terminal region mediates its interaction with the lateral elements (Jeffress et al., 2007). In *C. elegans*, the N-terminus of SYP-1 is acetylated in a NatB-dependent manner (Gao et al., 2016), likely co-translationally, a modification that may license folding or interactions required for SC assembly.

Despite these conserved structural principles, transverse filament proteins vary considerably across lineages. In the cnidarian *Hydra*, SYCP1 retains short sequence motifs with sufficient similarity to mammalian SYCP1, allowing it to cluster together in phylogenetic analyses (Fraune et al., 2012). However, other lineages show extensive sequence divergence while preserving the characteristic length and coiled-coil organization (Kursel et al., 2021). Interestingly, *C. elegans* uses two shorter transverse filament proteins, SYP-1 and SYP-5/6, each roughly half the length of transverse filament proteins in other species (MacQueen et al., 2002; Hurlock et al., 2020; Zhang et al., 2020). These proteins work together to span the SC width, demonstrating how divergent molecular mechanisms can generate a structurally conserved SC.

The conserved module linking SC initiation to meiotic recombination

In most eukaryotes, synapsis initiation is tightly coupled to DSB repair, ensuring that SC assembly occurs specifically between homologous chromosomes. In budding yeast and mammals, this coupling relies on a conserved central element module that initiates SC assembly from recombination sites through its interaction with Zip4, a scaffolding subunit of the meiosis-specific Zip2-Zip4-Spo16 (ZZS) complex that is essential for crossover formation (De Muyt et al., 2018). In budding yeast, the central element proteins Ecm11 and Gmc2 are recruited to DSB repair sites independently of Zip1, where they nucleate SC assembly (Humphryes et al., 2013). Recent work shows that this recruitment requires a direct interaction between Ecm11 and Zip4 (Pyatnitskaya et al., 2022). Intriguingly, Ecm11 and Gmc2 are predicted to share structural similarities with mammalian TEX12 and SYCE2 (Pyatnitskaya et al., 2022). Moreover, TEX12 interacts with TEX11, the mammalian homolog of Zip4, reinforcing the idea that the mechanism linking SC initiation to crossover recombination is conserved in mammals. Together, these findings illustrate how central element proteins, despite extensive sequence divergence, retain structural features and interactions that enable synapsis initiation at recombination intermediates.

An analogous mechanism has recently been described in plants, where the central element component SCEP3 loads independently of ZYP1 and helps coordinate SC assembly with crossover formation (Feng et al., 2025; Seear et al., 2025). However, SC assembly proceeds normally in mutants lacking ZIP4, even though it is essential for crossover formation (Chelysheva et al., 2007). These observations suggest that plants may initiate SC assembly through a pathway distinct from budding yeast and mammals. By contrast, the absence of ZYS components in *Drosophila* and *C. elegans* correlates with their recombination-independent mode of SC assembly, suggesting the lack of selective pressure for coupling synapsis initiation to crossover designation.

Molecular interfaces required for SC elongation across diverse species

Once nucleated at initial sites of homolog interaction, the SC central region rapidly extends along chromosome lengths through its inherent self-assembly properties and affinity for chromosome axes. This elongation mechanism varies across organisms, with a detailed understanding in yeast, mammals, and *C. elegans*, but significant gaps remain in other model organisms.

In budding yeast, SC elongation occurs as Zip1 is recruited to recombination intermediates through the Ecm11–Gmc2 complex (Humphryes et al., 2013). The C terminus of Zip1 interacts with the chromosome axis protein Red1, and this interface formation requires small ubiquitin-like modifier (SUMO) and SUMO-interacting motifs on both proteins (Cheng et al., 2006; Hooker and Roeder, 2006; Eichinger and Jentsch, 2010). Ecm11 also undergoes SUMOylation in a Gmc2 and Zip1 N terminus-dependent manner, and this modification is essential for Zip1 polymerization along chromosomes (Humphryes et al., 2013; Leung et al., 2015). Therefore, SUMOylation is expected to increase as Zip1 assembly proceeds, creating a positive feedback loop that promotes robust SC elongation (Leung et al., 2015).

By contrast, mammalian SC elongation relies on a hierarchical assembly process driven by the intrinsic self-assembly properties of its components, which has been primarily defined in vitro by biochemical and structural analyses. SYCP1 provides the structural foundation for this process and is recruited to chromosomes via its C-terminal interaction with chromosome axes (Winkel et al., 2009). SYCE3 then binds to the tetrameric SYCP1 core and remodels its interface into 2:1 heterotrimers (Crichton et al., 2023), generating an integrated SYCP1–SYCE3 lattice that serves as a platform for recruiting SYCE2–TEX12 and SYCE1–SIX6OS1 subcomplexes (Hamer et al., 2006; Schramm et al., 2011; Gómez-H et al., 2016; Sánchez-Sáez et al., 2020).

Once assembled on the SYCP1–SYCE3 scaffold, these additional central element complexes drive SC elongation through their polymerization. In vitro, SYCE2 and TEX12 form a 2:2 complex that self-assembles into filamentous structures (Davies et al., 2012; Duncie et al., 2021). Through these properties, SYCE2–TEX12 complexes can convert local synapsis initiation sites into extended synapsis along chromosome lengths (Bolcun-Filas et al., 2007; Hamer et al., 2008). Meanwhile, SYCE1 and

SIX6OS1 form a 1:1 complex through two essential binding interfaces and provide structural stability for the SC (Gómez-H et al., 2016; Dunne and Davies, 2019; Sánchez-Sáez et al., 2020). Together, SYCP1 and SYCE3 form a bilayered central structure at the SC midline, with the other central element proteins positioned between the layers (Hernández-Hernández et al., 2016). Through these coordinated activities, the central element proteins convert the initial SYCP1 lattice into a mature SC extended along the entire chromosome length (Fig. 3 A).

Unlike the hierarchical assembly of the SC in yeast and mammals, SC formation in *C. elegans* relies on the mutual interdependence among all SC subunits for both stability and chromosomal loading (Fig. 3 B). SC proteins are expressed before meiotic entry and assemble into polycomplexes, which adopt spherical shapes and fuse upon contact, exhibiting liquid-like properties (Rog et al., 2017). Although the molecular basis for the condensation of SC materials remains unknown, these polycomplexes serve as a reservoir for chromosomal loading and SC elongation. Synapsis initiates at special chromosomal regions called pairing centers, which serve as signaling hubs to mediate homology recognition and SC elongation (MacQueen et al., 2005). Extension of the SC central region between chromosome axes requires electrostatic interactions between the meiotic HORMAD protein HIM-3 and the acidic disordered C-terminal tails of SYP-5/6 (Gold et al., 2025; Gordon et al., 2025). Recent work has further demonstrated that SKR-1 and SKR-2, best known for their role in the Skp1–Cul1–F-box (SCF) ubiquitin ligase, moonlight as structural components of the SC via their SCF-forming interfaces (Blundon et al., 2024). This finding illustrates how a conserved cell cycle regulator is co-opted to interact with SC components and become an integral part of the SC.

While studies in *C. elegans* have provided insight into material properties of the SC, the mechanisms of SC elongation in *Drosophila* and plants still remain unclear. In *Drosophila*, SC assembly initiates at centromeres enriched for meiosis-specific cohesin complexes, where SC proteins are first recruited (Takeo et al., 2011; Tanneti et al., 2011). Although Cona and Corolla interact to promote C(3)G polymerization (Page et al., 2008; Collins et al., 2014), the molecular interfaces within the *Drosophila* SC remain largely uncharacterized. Super-resolution microscopy shows that the *Drosophila* SC adopts a bilayered architecture similar to that of mammals, with Cona and Corolla forming parallel tracks along chromosomes (Cahoon et al., 2017).

In *A. thaliana*, a hierarchical model of SC assembly is emerging. SCEP3 acts at the top of this hierarchy, recruiting other SC subunits and initiating SC formation (Feng et al., 2025; Seear et al., 2025). SCEP1 and SCEP2 interact with each other and are required for ZYP1 loading (Vrielynck et al., 2023). Despite these advances, elucidating the mechanisms of SC elongation will likely require identifying additional SC subunits and mapping their interaction networks and assembly dependencies.

The role of the SC in crossover regulation

Early cytological studies initially portrayed the SC as a passive structural scaffold for meiotic recombination. However, the SC




Figure 3. **SC architecture in mouse and *C. elegans*.** (A) Schematic representation of the mouse SC region. The molecular organization of SC central region components and AlphaFold 3-predicted structure of the SYCE-SIX6OS1 complex are shown on the right. The predicted aligned error (PAE) plot is also shown. (B) Schematic representation of the *C. elegans* SC central region. Right panel shows the molecular organization of SC central region components and AlphaFold 3-predicted structure of the SYP-3/SYP-4/SKR-1 complex with corresponding PAE plot. Red arrows indicate kinks that disrupt coiled-coils in both models.

is now recognized to play complex and seemingly contradictory roles in both promoting and limiting crossover formation. Here, we synthesize recent evidence to provide an updated overview of how the SC regulates crossover number and distribution.

The SC promotes the formation of class I crossovers

The pro-crossover function of the SC was first established by pioneering genetic studies in *D. melanogaster*. Mutations in *c(3)G* result in virtually no crossing-over, demonstrating an absolute

requirement for the SC (Hall, 1972). In budding yeast, however, *zip1* mutants retain some crossovers but show delayed crossover maturation and loss of crossover interference (Sym and Roeder, 1994; Storlazzi et al., 1996). The remaining crossovers in *zip1* mutants depend on structure-specific endonucleases Mus81-Mms4 (de los Santos et al., 2003), indicating that the SC is not absolutely required for all crossovers.

These findings reveal two mechanistically distinct crossover pathways. The class I pathway generates 80–90% of crossovers in most species and depends on the SC and a set of conserved ZMM proteins (named after budding yeast Zip, Mer, and Msh proteins) that stabilize crossover-specific recombination intermediates and produce crossovers exhibiting interference (Pyatnitskaya et al., 2019). In contrast, the class II pathway produces crossovers independently of the SC and lacks interference (Börner et al., 2004). Class II crossovers often increase when the class I pathway is compromised, providing a backup mechanism that helps generate chiasmata to ensure faithful chromosome segregation (Anderson et al., 2014).

How does the SC promote crossovers? Key steps in the class I pathway, including the formation of single-end invasion intermediates and double Holliday junctions (dHJs), occur in tight temporal coordination with SC morphogenesis (Börner et al., 2004) (Fig. 4, A and B). By sequestering and organizing ZMM proteins, the SC creates protective microenvironments around crossover-designated sites as recombination progresses. Super-resolution imaging in *C. elegans* shows that these sites become encased by SC central region proteins in late pachytene, forming specialized compartments that shield recombination intermediates from premature dismantling (Woglar and Villeneuve, 2018). Consistent with this protective role, recent work in budding yeast reveals a reciprocal feedback mechanism in which Zip1 stabilizes dHJ-associated ZMM complexes, which in turn reinforce the local SC central region (Henggeler et al., 2025; Tang et al., 2025). This aligns with evidence that the SC is a dynamic structure with continuous turnover and incorporation of its subunits, which modulate SC stability in response to recombination events (Voelkel-Meiman et al., 2012; Nadarajan et al., 2017; Pattabiraman et al., 2017). Importantly, this dHJ-stabilizing function of the SC does not require a fully continuous tripartite structure. Instead, local ensembles of Zip1 and ZMM proteins are sufficient to protect dHJs until they are resolved into crossovers (Voelkel-Meiman et al., 2016; Tang et al., 2025), consistent with an SC-independent role of Zip1 in meiotic recombination (Storlazzi et al., 1996). Moreover, SC disassembly is tightly coupled to dHJ resolution, ensuring that these compartments persist throughout crossover maturation (Henggeler et al., 2025). These findings suggest that the SC actively orchestrates multiple steps in crossover recombination, extending far beyond merely juxtaposing homologous chromosomes.

An ancient module linking the SC central region to RING-domain E3 ligases

A key mechanism by which the SC promotes crossover formation is through the recruitment of the Zip3 RING-domain E3 ligase family, which is essential for class I crossover formation across nearly all eukaryotes. These conserved ligases (Zip3 in

yeast; HEI10, RNF212, and RNF212B in mammals; Vilya in *Drosophila*; ZHP-1/2 and ZHP-3/4 in *C. elegans*; and HEI10 in plants) localize to discrete foci along the SC, accumulating at sites designated for crossover formation (Agarwal and Roeder, 2000; Jantsch et al., 2004; Chelysheva et al., 2012; Reynolds et al., 2013; Qiao et al., 2014; Lake et al., 2015; Zhang et al., 2018; Condezo et al., 2024; Ito et al., 2025). A growing body of work suggests that the SC interacts with these RING ligases through unexpectedly conserved molecular mechanisms. In budding yeast, the Zip1 N terminus, which includes conserved phenylalanine residues, is required for Zip3 recruitment to the SC polycomplex and recombination initiation sites (Voelkel-Meiman et al., 2019). However, Zip3 forms discrete foci along chromosomes without Zip1 (Agarwal and Roeder, 2000), suggesting that the Zip1 N terminus might stabilize Zip3 at crossover-designated sites rather than mediating its initial chromosomal loading.

This mechanism appears to be conserved, as phenylalanine-rich motifs are also present in SC central region proteins across taxa. In *C. elegans*, phenylalanine residues in the C-terminal tail of SYP-4 are required for recruitment of the ZHP-3/ZHP-4 complex to the SC, which is essential for crossover regulation but dispensable for SC assembly (Neves et al., 2025). Similar motifs are present in the C-terminal tail of mammalian SIX6OS1 and in *Drosophila* Corolla. Based on their shared functions and conserved motifs, SYP-4, SIX6OS1, and Corolla have recently been proposed as homologs (Neves et al., 2025; Williams et al., 2025), a hypothesis supported by AlphaFold structural predictions that reveal similarities in their overall coiled-coil organization and the positioning of kinks that disrupt coiled-coils (Fig. 3). While a direct interaction between Zip1 and Zip3 has recently been demonstrated in budding yeast (Sharmin et al., 2025, Preprint), experimental validation in other organisms remains an important future direction.

Remarkably, this ancient module extends to plants. In *Arabidopsis*, SCEP3 contains conserved phenylalanine repeats within its N-terminal disordered region and plays analogous roles in recruiting SC proteins to the recombination sites and initiating SC elongation (Feng et al., 2025; Seear et al., 2025). SCEP3 first localizes to recombination sites along chromosome axes and later relocates to the SC central region as synapsis proceeds (Seear et al., 2025). Interestingly, SCEP3 does not appear to interact directly with HEI10, leaving unresolved how it interfaces with crossover-promoting factors. Nevertheless, the striking conservation of this molecular module across eukaryotic lineages suggests that coupling between the SC central region and RING-domain E3 ligases is a fundamental feature of meiosis.

The role of the SC in crossover patterning

Studies across model organisms show that mutations disrupting SC assembly or integrity severely impair crossover interference, resulting in more randomly distributed crossovers (Sym and Roeder, 1994; Libuda et al., 2013; Capilla-Pérez et al., 2021; France et al., 2021). This evidence suggests that the SC acts as a structural scaffold or signaling conduit that transmits information about a designated crossover, thereby inhibiting the designation of additional crossovers nearby. However, the precise mechanism for this chromosome-wide signaling remains contentious.




Figure 4. SC morphogenesis is tightly linked to steps in homologous recombination. (A) Temporal coordination of SC morphogenesis through meiotic prophase I. Meiosis-specific cohesins load in pre-meiotic S phase and organize chromatin into arrays of loops. In leptotene, chromosome axes begin to form, and DSBs are induced. During zygotene, homologous chromosomes begin to synapse at recombination sites. In pachytene, the full SC is assembled and protects crossover-specific recombination intermediates. **(B)** Molecular steps of the class I crossover pathway in budding yeast. Spo11 catalyzes DSB formation, followed by the Mre11–Rad50–Xrs2 (MRX) complex and Exo1-mediated resection. The resulting ssDNA is initially coated by RPA and subsequently replaced by Rad51 and Dmc1 to promote strand invasion. Formation and stabilization of HJs requires ZMM proteins (light red). The resolution of dHJ triggers SC disassembly.

Recent quantitative imaging and mathematical modeling in *Arabidopsis* have led to a diffusion-based coarsening model for crossover interference (Morgan et al., 2021). In this model, the SC harbors a constrained, diffusible pool of HEI10, which initially appears as numerous small foci at early recombination nodules. According to this model, these foci undergo coarsening, analogous to Ostwald ripening in phase-separated systems, where larger, evenly dispersed foci grow at the expense of smaller ones through diffusion along the SC (Fig. 5 A). The redistribution dynamics of HEI10 is thus proposed as the primary driver of both crossover designation and interference. This framework accurately predicts crossover numbers and spacing and explains the sensitivity of crossover number to HEI10 dosage (Ziolkowski et al., 2017; Morgan et al., 2021).

This view is further supported by work in *C. elegans* showing that the ZHP-3/4 complex forms biomolecular condensates that diffuse along the SC (Zhang et al., 2025). The SC thus serves as a

one-dimensional track that compartmentalizes diffusion, allowing each chromosome pair to undergo independent coarsening and, thereby, establish its own crossover pattern (Zhang et al., 2018). Intriguingly, modeling in *Arabidopsis zyp1* mutants suggests that the coarsening process can still occur in the nucleoplasm, with competition for HEI10 across chromosomes rather than being strictly limited to individual bivalents (Durand et al., 2022; Fozard et al., 2023). This implies that crossover patterning can emerge from generic coarsening dynamics of HEI10 condensates and does not intrinsically require an SC scaffold. Together, these findings suggest that the SC functions primarily as a spatial constraint that compartmentalizes HEI10 diffusion to individual chromosome pairs, rather than being essential for the coarsening process itself. While this model explains many aspects of crossover patterning, several observations across organisms suggest that additional mechanisms must be at play.




Figure 5. **Models of crossover patterning.** (A) Coarsening model. HEI10 (magenta) initially localizes uniformly along the SC. Through diffusion-mediated coarsening, HEI10 accumulates at a subset of sites at the expense of neighboring foci, driving crossover designation and interference. (B) Beam-film model. Chromosome axes act as elastic beams that store mechanical stress. Crossover designation (yellow star) generates local “cracks” that relieve tension; this relief propagates bidirectionally (black arrows), preventing nearby precursors from reaching the stress threshold required for crossover designation. Synapsis initiates from crossover designated sites, and these initiation sites themselves exhibit interference. (C) Multistep model. This model integrates coarsening and beam-film frameworks. Stress-mediated, axis-based processes establish an initial crossover pattern, while diffusion-mediated coarsening along the SC subsequently refines and stabilizes designated sites.

If diffusion-based coarsening were the primary mechanism for crossover designation and patterning, a fully continuous SC would be required to propagate inhibitory signals along chromosomes at the time of designation. However, in budding yeast, crossover designation is established early, even before the completion of SC assembly (Allers and Lichten, 2001; Börner et al., 2004; Fung et al., 2004). These observations directly challenge a strictly SC-dependent interference mechanism and support alternative models. One such framework, the beam-film model, posits that mechanical stress along chromosome axes distributes crossovers by locally relieving tension after each crossover event (Kleckner et al., 2004), thereby providing an early, SC-independent mechanism for crossover patterning (Fig. 5 B).

Providing molecular support for this model, topoisomerase II (Top2) has been identified as a key mediator of crossover interference in budding yeast (Zhang et al., 2014b). Interference strength inversely correlates with negative supercoils (Tan et al., 2022), and Top2 degradation by the ubiquitin ligase Ufd2 modulates interference strength by tuning DNA supercoiling (Tan et al., 2026). Because this torsional stress mechanism operates along the chromosome axis rather than the SC, these findings support that the role of the SC is in implementing crossover patterning decisions that were established earlier (Börner et al.,

2004). Consistent with this view, SC nucleation sites in *Sordaria* are themselves evenly spaced by an interference-like mechanism (Zhang et al., 2014a). Furthermore, crossover frequencies covary with axis length across chromosomes within individual nuclei (Wang et al., 2019), linking patterning outcomes to the physical properties of the axis rather than the SC. Together, these observations suggest that the SC functions downstream of crossover designation, stabilizing committed sites rather than generating the pattern de novo.

Additional challenges to a simple coarsening model come from mammals, where evidence points to more complex spatiotemporal dynamics among the three RING-domain ligases. In mouse spermatocytes, RNF212B and RNF212 mark early recombination intermediates, while HEI10 appears later at maturing crossovers (Reynolds et al., 2013; Qiao et al., 2014; Condezo et al., 2024; Ito et al., 2025). However, in oocytes, RNF212 and RNF212B foci remain abundant throughout pachytene, challenging the underlying assumption that successful crossover designation relies on their depletion from noncrossover sites (Ito et al., 2025).

Recent super-resolution microscopy and real-time imaging of recombination factors in *C. elegans* indicate that crossover patterning arises from a dynamic, multilayered regulation process rather than a single decision point (Čavka et al., 2025, Preprint).

An initial selection process appears to occur shortly after DSB formation and restricts crossovers to a small subset of DSB sites that already exhibit interference, likely through axis-based cues such as mechanical stress. This early patterning is then followed by a refinement phase that coincides with the first signs of ZHP-3 coarsening along the SC. This multistep model reconciles the timing paradox of crossover designation: axis-based mechanisms establish the initial pattern, while SC-dependent diffusion-based processes subsequently refine and stabilize early-designated sites (Fig. 5 C). Together, these findings highlight that crossover designation involves multilayered regulatory inputs acting at distinct stages, with different organisms potentially weighing these mechanisms to varying degrees.

The SC coordinates feedback control of DSB formation and quality surveillance

The assembly status of the SC is closely monitored by cells to assess homolog engagement and coordinate meiotic progression. Mutants defective in SC formation exhibit elevated DSB levels, indicating that the SC provides negative feedback to limit further DSB formation (Kauppi et al., 2013; Thacker et al., 2014; Lee et al., 2021). This feedback operates in a chromosome-autonomous manner, with each chromosome pair sensing its own homolog engagement status and independently down-regulating DSB formation (Mu et al., 2020). The molecular basis of this feedback control has been well elucidated in yeast and mammals. As described earlier, axis-associated *HORMADs* are actively removed from synapsed chromosomes by TRIP13 (Pch2 in budding yeast) (Joshi et al., 2009; Wojtasz et al., 2009). Because *HORMADs* provide an axial platform for recruiting the SPO11 machinery via the REC114–MEI4–IHO1 complex in mammals (Rec114–Mei4–Mer2 complex in budding yeast) (Panizza et al., 2011; Stanzione et al., 2016; Rousová et al., 2021; Laroussi et al., 2023), their removal from synapsed regions dismantles the DSB-forming machinery, rendering synapsed regions incompetent for further DSB formation (Fig. 2 C). Through this feedback, *HORMAD* dynamics prevents excessive DSB formation and ensures that crossover formation proceeds appropriately for accurate chromosome segregation.

In mammals, *HORMADs* are also essential for recruiting ATR and BRCA1 to unsynapsed chromatin (Fig. 2 C), where ATR phosphorylates histone H2AX and activates checkpoint responses that ultimately eliminate oocytes with persistent asynapsis or recombination defects (Kouznetsova et al., 2009; Shin et al., 2010; Daniel et al., 2011; Wojtasz et al., 2012; Cloutier et al., 2015; Jiao et al., 2025). Additionally, ATR regulates *HORMAD1/2* phosphorylation and is required for proper localization of BRCA1 and ATR cofactors to unsynapsed axes, contributing to meiotic silencing of unsynapsed chromatin (Royo et al., 2013). Thus, *HORMADs* function as sensors of synapsis status, linking SC assembly to both suppression of DSB formation and activation of surveillance pathways that safeguard genome integrity and prevent aneuploidy in gametes.

A distinct feedback mechanism operates in *C. elegans*, where DSBs occur in the context of a fully assembled SC and *HORMADs*

remain axis-associated even after synapsis, inverting the logic seen in yeast and mammals. Here, the meiotic kinase CHK-2 serves as the master regulator whose activity sustains DSB competence by maintaining chromatin association of DSB proteins and is prolonged by *HORMADs* when synapsis or crossover formation fails (Rosu et al., 2013; Stamper et al., 2013; Kim et al., 2015; Hinman et al., 2021). Crossover designation stabilizes the SC central region by recruiting the Polo-like kinase PLK-2, which attenuates further DSB formation through phosphorylation of SYP-4 (Nadarajan et al., 2017; Pattabiraman et al., 2017). Chromosomes that fail to designate crossovers do not recruit PLK-2 and undergo desynapsis (Machovina et al., 2016), maintaining them in a DSB-competent state. Importantly, removing SC components from nuclei with designated crossovers reactivates CHK-2 and resumes DSB formation, demonstrating continuous surveillance (Castellano-Pozo et al., 2020). Moreover, nuclei with persistent unsynapsed chromosomes are ultimately eliminated by apoptosis, providing a final failsafe (Bhalla and Dernburg, 2005). Thus, while the molecular players differ, *C. elegans* shares with yeast and mammals the fundamental principle of using SC-associated signals to coordinate DSB formation with crossover status and eliminate defective meiotic cells.

Concluding remarks

Once viewed as an enigmatic electron-dense scaffold, the SC is now recognized as a dynamic regulatory platform that coordinates multiple essential meiotic processes. In most eukaryotes, SC assembly is coupled with meiotic recombination, positioning it as a signaling hub for self-reinforcing feedback loops. The SC recruits and organizes ZMM proteins to stabilize crossover-specific recombination intermediates, which in turn reinforce the SC structure. Additionally, the SC provides negative feedback that suppresses additional DSB formation on synapsed chromosomes, preventing excess DSB formation and enabling quality control. Together, these features illustrate how a supramolecular chromosome structure integrates its assembly with sophisticated DNA metabolism.

Studies across diverse model organisms reveal several recurring themes. Despite extensive sequence divergence among SC components, the fundamental architectural principles and core regulatory modules of the SC are far more conserved than previously appreciated. Features such as the ~100 nm SC width, head-to-head assembly of transverse filaments, and the coupling between the SC central region and ZMM proteins highlight how divergent strategies achieve similar structural and functional outcomes. A particularly striking example is the conserved interface of the SC with the Zip3 family RING-domain E3 ligases (Neves et al., 2025; Williams et al., 2025; Seear et al., 2025). This ancient module suggests that the regulatory link between the SC and the class I crossover pathway is under strong evolutionary constraint.

Central to the regulatory mechanisms of the SC are the *HORMAD* proteins, which couple homolog synapsis to recombination control and quality surveillance. Prior to SC assembly, *HORMADs* on chromosome axes establish platforms for DSB formation and enforce interhomolog bias. Upon synapsis, their

removal by TRIP13/Pch2 dismantles the DSB-forming machinery and relieves the barrier to intersister repair. HORADs that persist on asynaptic chromosomes trigger checkpoint responses that arrest or eliminate defective meiocytes, thereby preventing the production of aneuploid gametes (Wojtasz et al., 2009; Rinaldi et al., 2017; Jiao et al., 2025). These interconnected mechanisms create a robust system that balances DSB formation with the capacity for DNA repair.

Despite substantial progress, fundamental questions about the SC remain. How complete is the SC parts list, and what components are still missing? Why do the primary sequences of SC components change so rapidly on evolutionary timescales? How are individual SC subunits produced in the correct stoichiometry and coordinated to form functional complexes? What molecular mechanisms drive SC polymerization and its dynamic remodeling during recombination? Developing in vitro reconstitution systems for SC assembly will be crucial for answering these questions. The recent discovery of Skp1 proteins as SC subunits in *C. elegans* further raises intriguing questions about the prevalence and functional significance of such co-option. Resolving these questions will advance our understanding of how the SC integrates structural organization, crossover recombination, and quality control to safeguard genome integrity across generations.

Acknowledgments

We acknowledge support from the National Institutes of Health to B.I. Cesar (F31GM153163) and Y. Kim (R35GM124895).

Author contributions: Brenda I. Cesar: conceptualization, funding acquisition, resources, visualization, and writing—original draft, review, and editing. Yumi Kim: writing—original draft, review, and editing.

Disclosures: The authors declare no competing interests exist.

Submitted: 30 November 2025

Revised: 13 February 2026

Accepted: 9 March 2026

References

- Agarwal, S., and S.G. Roeder. 2000. Zip3 provides a link between recombination enzymes and synaptonemal complex proteins. *Cell*. 102:245–255. [https://doi.org/10.1016/s0092-8674\(00\)00029-5](https://doi.org/10.1016/s0092-8674(00)00029-5)
- Alani, E., R. Padmore, and N. Kleckner. 1990. Analysis of wild-type and rad50 mutants of yeast suggests an intimate relationship between meiotic chromosome synapsis and recombination. *Cell*. 61:419–436. [https://doi.org/10.1016/0092-8674\(90\)90524-i](https://doi.org/10.1016/0092-8674(90)90524-i)
- Allers, T., and M. Lichten. 2001. Differential timing and control of non-crossover and crossover recombination during meiosis. *Cell*. 106:47–57. [https://doi.org/10.1016/s0092-8674\(01\)00416-0](https://doi.org/10.1016/s0092-8674(01)00416-0)
- Anderson, L.K., K.D. Hooker, and S.M. Stack. 2001. The distribution of early recombination nodules on zygotene bivalents from plants. *Genetics*. 159:1259–1269. <https://doi.org/10.1093/genetics/159.3.1259>
- Anderson, L.K., S.M. Royer, S.L. Page, K.S. McKim, A. Lai, M.A. Lilly, and R.S. Hawley. 2005. Juxtaposition of C(2)M and the transverse filament protein C(3)G within the central region of *Drosophila*. *Proc. Natl. Acad. Sci. USA*. 102:4482–4487. <https://doi.org/10.1073/pnas.0500172102>
- Anderson, L.K., L.D. Lohmiller, X. Tang, D.B. Hammond, L. Javernick, L. Shearer, S. Basu-Roy, O.C. Martin, and M. Falque. 2014. Combined

- fluorescent and electron microscopic imaging unveils the specific properties of two classes of meiotic crossovers. *Proc. Natl. Acad. Sci. USA*. 111:13415–13420. <https://doi.org/10.1073/pnas.1406846111>
- Auxier, B., A.J.M. Debets, F.A. Stanford, J. Rhodes, F.M. Becker, F. Reyes Marquez, R. Nijland, P.S. Dyer, M.C. Fisher, J. van den Heuvel, and E. Snelders. 2023. The human fungal pathogen *Aspergillus fumigatus* can produce the highest known number of meiotic crossovers. *PLoS Biol.* 21:e3002278. <https://doi.org/10.1371/journal.pbio.3002278>
- Bähler, J., T. Wyler, J. Loidl, and J. Kohli. 1993. Unusual nuclear structures in meiotic prophase of fission yeast: A cytological analysis. *J. Cell Biol.* 121:241–256. <https://doi.org/10.1083/jcb.121.2.241>
- Baudat, F., K. Manova, J.P. Yuen, M. Jasin, and S. Keeney. 2000. Chromosome synapsis defects and sexually dimorphic meiotic progression in mice lacking Spo11. *Mol. Cell*. 6:989–998. [https://doi.org/10.1016/s1097-2765\(00\)00098-8](https://doi.org/10.1016/s1097-2765(00)00098-8)
- Bhalla, N., and A.F. Dernburg. 2005. A conserved checkpoint monitors meiotic chromosome synapsis in *Caenorhabditis elegans*. *Science*. 310:1683–1686. <https://doi.org/10.1126/science.1117468>
- Billmyre, K.K., E.A. Kesler, D. Tsuchiya, T.J. Corbin, K. Weaver, A. Moran, Z. Yu, L. Adams, K. Delventhal, M. Durmin, et al. 2023. SYCP1 head-to-head assembly is required for chromosome synapsis in mouse meiosis. *Sci. Adv.* 9:eadi1562. <https://doi.org/10.1126/sciadv.adi1562>
- Blundon, J.M., B.I. Cesar, J.W. Bae, I. Čavka, J. Haversat, J. Ries, S. Köhler, and Y. Kim. 2024. Skp1 proteins are structural components of the synaptonemal complex in *C. elegans*. *Sci. Adv.* 10:eadi4876. <https://doi.org/10.1126/sciadv.adl4876>
- Bolcun-Filas, E., Y. Costa, R. Speed, M. Taggart, R. Benavente, D.G. De Rooij, and H.J. Cooke. 2007. SYCE2 is required for synaptonemal complex assembly, double strand break repair, and homologous recombination. *J. Cell Biol.* 176:741–747. <https://doi.org/10.1083/jcb.200610027>
- Börner, G.V., N. Kleckner, and N. Hunter. 2004. Crossover/noncrossover differentiation, synaptonemal complex formation, and regulatory surveillance at the leptotene/zygotene transition of meiosis. *Cell*. 117:29–45. [https://doi.org/10.1016/s0092-8674\(04\)00292-2](https://doi.org/10.1016/s0092-8674(04)00292-2)
- Cahoon, C.K., Z. Yu, Y. Wang, F. Guo, J.R. Unruh, B.D. Slaughter, and R.S. Hawley. 2017. Superresolution expansion microscopy reveals the three-dimensional organization of the synaptonemal complex. *Proc. Natl. Acad. Sci. USA*. 114:E6857–E6866. <https://doi.org/10.1073/pnas.1705623114>
- Callender, T.L., R. Laureau, L. Wan, X. Chen, R. Sandhu, S. Laljee, S. Zhou, R.T. Suhandynata, E. Prugar, W.A. Gaines, et al. 2016. Mek1 down regulates Rad51 activity during yeast meiosis by phosphorylation of Hed1. *PLoS Genet.* 12:e1006226. <https://doi.org/10.1371/journal.pgen.1006226>
- Capilla-Pérez, L., S. Durand, A. Hurel, Q. Lian, A. Chambon, C. Taochy, V. Solier, M. Grelon, and R. Mercier. 2021. The synaptonemal complex imposes crossover interference and heterochiasmy in. *Proc. Natl. Acad. Sci. USA*. 118:e2023613118. <https://doi.org/10.1073/pnas.2023613118>
- Carballo, J.A., A.L. Johnson, S.G. Sedgwick, and R.S. Cha. 2008. Phosphorylation of the axial element protein Hop1 by Mec1/Tel1 ensures meiotic interhomolog recombination. *Cell*. 132:758–770. <https://doi.org/10.1016/j.cell.2008.01.035>
- Carpenter, A.T. 1975. Electron microscopy of meiosis in *Drosophila melanogaster* females: II. The recombination nodule—a recombination-associated structure at pachytene? *Proc. Natl. Acad. Sci. USA*. 72:3186–3189. <https://doi.org/10.1073/pnas.72.8.3186>
- Carpenter, A.T. 1979. Synaptonemal complex and recombination nodules in wild-type *Drosophila melanogaster* females. *Genetics*. 92:511–541. <https://doi.org/10.1093/genetics/92.2.511>
- Castellano-Pozo, M., S. Pacheco, G. Sioutas, A.L. Jaso-Tamame, M.H. Dore, M.M. Karimi, and E. Martinez-Perez. 2020. Surveillance of cohesin-supported chromosome structure controls meiotic progression. *Nat. Commun.* 11:4345. <https://doi.org/10.1038/s41467-020-18219-9>
- Čavka, I., A. Woglar, Y.-L. Wu, E.B. Durmus, L. Sloat, A. Gros, C.P. López, F. Hecht, A.M. Villeneuve, J. Ries, and S. Köhler. 2025. Multi-step implementation of meiotic crossover patterning. *bioRxiv*. <https://doi.org/10.1101/2025.11.12.687980> (Preprint posted November 13, 2025).
- Chambon, A., A. West, D. Vezon, C. Horlow, A. De Muyt, L. Chelysheva, A. Ronceret, A. Darbyshire, K. Osman, S. Heckmann, et al. 2018. Identification of ASYNAPTIC4, a component of the meiotic chromosome axis. *Plant Physiol.* 178:233–246. <https://doi.org/10.1104/pp.17.01725>
- Chelysheva, L., G. Gendrot, D. Vezon, M.-P. Doutriaux, R. Mercier, and M. Grelon. 2007. Zip4/Spo22 is required for class I CO formation but not for synapsis completion in *Arabidopsis thaliana*. *PLoS Genet.* 3:e83. <https://doi.org/10.1371/journal.pgen.0030083>
- Chelysheva, L., D. Vezon, A. Chambon, G. Gendrot, L. Pereira, A. Lemhendi, N. Vrielynck, S. Le Guin, M. Novatchkova, and M. Grelon. 2012.

- The *Arabidopsis* HEI10 is a new ZMM protein related to Zip3. *PLoS Genet.* 8:e1002799. <https://doi.org/10.1371/journal.pgen.1002799>
- Cheng, C.-H., Y.-H. Lo, S.-S. Liang, S.-C. Ti, F.-M. Lin, C.-H. Yeh, H.-Y. Huang, and T.-F. Wang. 2006. SUMO modifications control assembly of synaptonemal complex and polycomplex in meiosis of *Saccharomyces cerevisiae*. *Genes Dev.* 20:2067–2081. <https://doi.org/10.1101/gad.1430406>
- Clayes Bouuaert, C., S. Pu, J. Wang, C. Oger, D. Daccache, W. Xie, D.J. Patel, and S. Keeney. 2021. DNA-driven condensation assembles the meiotic DNA break machinery. *Nature.* 592:144–149. <https://doi.org/10.1038/s41586-021-03374-w>
- Cloutier, J.M., S.K. Mahadevaiah, E. Ellnati, A. Nussenzweig, A. Tóth, and J.M.A. Turner. 2015. Histone H2AFX links meiotic chromosome assembly to prophase I oocyte loss in mammals. *PLoS Genet.* 11:e1005462. <https://doi.org/10.1371/journal.pgen.1005462>
- Colaiácovo, M.P., A.J. MacQueen, E. Martínez-Pérez, K. McDonald, A. Adamo, A. La Volpe, and A.M. Villeneuve. 2003. Synaptonemal complex assembly in *C. elegans* is dispensable for loading strand-exchange proteins but critical for proper completion of recombination. *Dev. Cell.* 5:463–474. [https://doi.org/10.1016/s1534-5807\(03\)00232-6](https://doi.org/10.1016/s1534-5807(03)00232-6)
- Collins, K.A., J.R. Unruh, B.D. Slaughter, Z. Yu, C.M. Lake, R.J. Nielsen, K.S. Box, D.E. Miller, J.P. Blumenstiel, A.G. Perera, et al. 2014. Corolla is a novel protein that contributes to the architecture of the synaptonemal complex of *Drosophila*. *Genetics.* 198:219–228. <https://doi.org/10.1534/genetics.114.165290>
- Condezo, Y.B., R. Sainz-Urruela, L. Gomez-H, D. Salas-Lloret, N. Felipe-Medina, R. Bradley, I.D. Wolff, S. Tanis, J.L. Barbero, M. Sánchez-Martín, et al. 2024. RNF212B E3 ligase is essential for crossover designation and maturation during male and female meiosis in the mouse. *Proc. Natl. Acad. Sci. USA.* 121:e2320995121. <https://doi.org/10.1073/pnas.2320995121>
- Costa, Y., R. Speed, R. Ollinger, M. Alsheimer, C.A. Semple, P. Gautier, K. Maratou, I. Novak, K. Höög, R. Benavente, and H.J. Cooke. 2005. Two novel proteins recruited by synaptonemal complex protein 1 (SYCP1) are at the centre of meiosis. *J. Cell Sci.* 118:2755–2762. <https://doi.org/10.1242/jcs.02402>
- Couteau, F., and M. Zetka. 2005. HTP-1 coordinates synaptonemal complex assembly with homolog alignment during meiosis in *C. elegans*. *Genes Dev.* 19:2744–2756. <https://doi.org/10.1101/gad.1348205>
- Couteau, F., K. Nabeshima, A. Villeneuve, and M. Zetka. 2004. A component of *C. elegans* meiotic chromosome axes at the interface of homolog alignment, synapsis, nuclear reorganization, and recombination. *Curr. Biol.* 14:585–592. <https://doi.org/10.1016/j.cub.2004.03.033>
- Crichton, J.H., J.M. Duncie, O.M. Dunne, L.J. Salmon, P.S. Devenney, J. Lawson, I.R. Adams, and O.R. Davies. 2023. Structural maturation of SYCP1-mediated meiotic chromosome synapsis by SYCE3. *Nat. Struct. Mol. Biol.* 30:188–199. <https://doi.org/10.1038/s41594-022-00909-1>
- Daniel, K., J. Lange, K. Hached, J. Fu, K. Anastasiadis, I. Roig, H.J. Cooke, A.F. Stewart, K. Wassmann, M. Jasin, et al. 2011. Meiotic homolog alignment and its quality surveillance are controlled by mouse HORMAD1. *Nat. Cell Biol.* 13:599–610. <https://doi.org/10.1038/ncb2213>
- Davies, O.R., J.D. Maman, and L. Pellegrini. 2012. Structural analysis of the human SYCE2-TEX12 complex provides molecular insights into synaptonemal complex assembly. *Open Biol.* 2:120099. <https://doi.org/10.1098/rsob.120099>
- de los Santos, T., N. Hunter, C. Lee, B. Larkin, J. Loidl, and N.M. Hollingsworth. 2003. The Mus81/Mms4 endonuclease acts independently of double-Holliday junction resolution to promote a distinct subset of crossovers during meiosis in budding yeast. *Genetics.* 164:81–94. <https://doi.org/10.1093/genetics/164.1.81>
- Dernburg, A.F., K. McDonald, G. Moulder, R. Barstead, M. Dresser, and A.M. Villeneuve. 1998. Meiotic recombination in *C. elegans* initiates by a conserved mechanism and is dispensable for homologous chromosome synapsis. *Cell.* 94:387–398. [https://doi.org/10.1016/s0092-8674\(00\)81481-6](https://doi.org/10.1016/s0092-8674(00)81481-6)
- Dong, H., and G.S. Roeder. 2000. Organization of the yeast Zip1 protein within the central region of the synaptonemal complex. *J. Cell Biol.* 148:417–426. <https://doi.org/10.1083/jcb.148.3.417>
- Duncie, J.M., O.M. Dunne, M. Ratcliff, C. Millán, S. Madgwick, I. Usón, and O.R. Davies. 2018. Structural basis of meiotic chromosome synapsis through SYCP1 self-assembly. *Nat. Struct. Mol. Biol.* 25:557–569. <https://doi.org/10.1038/s41594-018-0078-9>
- Duncie, J.M., L.J. Salmon, and O.R. Davies. 2021. Structural basis of meiotic chromosome synaptic elongation through hierarchical fibrous assembly of SYCE2-TEX12. *Nat. Struct. Mol. Biol.* 28:681–693. <https://doi.org/10.1038/s41594-021-00636-z>
- Dunne, O.M., and O.R. Davies. 2019. Molecular structure of human synaptonemal complex protein SYCE1. *Chromosoma.* 128:223–236. <https://doi.org/10.1007/s00412-018-00688-z>
- Durand, S., Q. Lian, J. Jing, M. Ernst, M. Grelon, D. Zwicker, and R. Mercier. 2022. Joint control of meiotic crossover patterning by the synaptonemal complex and HEI10 dosage. *Nat. Commun.* 13:5999. <https://doi.org/10.1038/s41467-022-33472-w>
- Egel-Mitani, M., L.W. Olson, and R. Egel. 1982. Meiosis in *Aspergillus nidulans*: Another example for lacking synaptonemal complexes in the absence of crossover interference. *Hereditas.* 97:179–187. <https://doi.org/10.1111/j.1601-5223.1982.tb00761.x>
- Eichinger, C.S., and S. Jentsch. 2010. Synaptonemal complex formation and meiotic checkpoint signaling are linked to the lateral element protein Red1. *Proc. Natl. Acad. Sci. USA.* 107:11370–11375. <https://doi.org/10.1073/pnas.1004248107>
- Emmenecker, C., S. Pakzad, F. Ture, J. Guerin, A. Hurel, A. Chambon, C. Girard, R. Mercier, and R. Kumar. 2024. FIGL1 attenuates meiotic interhomolog repair and is counteracted by the RAD51 paralogue XRCC2 and the chromosome axis protein ASY1 during meiosis. *New Phytol.* 244:2442–2457. <https://doi.org/10.1111/nph.20181>
- Fawcett, D.W. 1956. The fine structure of chromosomes in the meiotic prophase of vertebrate spermatocytes. *J. Biophys. Biochem. Cytol.* 2:403–406. <https://doi.org/10.1083/jcb.2.4.403>
- Feng, C., J. Lorenz, S. Dreissig, V. Schubert, B. Wang, F. Hartmann, M. Cuacos, N. Fernández-jiménez, Z. Zhao, C. Eggeling, et al. 2025. The synaptonemal complex central element SCEP3 interlinks synapsis initiation and crossover formation in *Arabidopsis thaliana*. *Nat. Plants.* 11:1353–1366. <https://doi.org/10.1038/s41477-025-02030-9>
- Ferdous, M., J.D. Higgins, K. Osman, C. Lambing, E. Roitinger, K. Mechtler, S.J. Armstrong, R. Perry, M. Pradillo, N. Cufiádo, and F.C.H. Franklin. 2012. Inter-homolog crossing-over and synapsis in *Arabidopsis* meiosis are dependent on the chromosome axis protein AtASY3. *PLoS Genet.* 8:e1002507. <https://doi.org/10.1371/journal.pgen.1002507>
- Fozard, J.A., C. Morgan, and M. Howard. 2023. Coarsening dynamics can explain meiotic crossover patterning in both the presence and absence of the synaptonemal complex. *Elife.* 12:e79408. <https://doi.org/10.7554/eLife.79408>
- France, M.G., J. Enderle, S. Röhrig, H. Puchta, F.C.H. Franklin, and J.D. Higgins. 2021. ZYP1 is required for obligate cross-over formation and cross-over interference in. *Proc. Natl. Acad. Sci. USA.* 118:e2021671118. <https://doi.org/10.1073/pnas.2021671118>
- Fraune, J., M. Alsheimer, J.-N. Volff, K. Busch, S. Fraune, T.C.G. Bosch, and R. Benavente. 2012. Hydra meiosis reveals unexpected conservation of structural synaptonemal complex proteins across metazoans. *Proc. Natl. Acad. Sci. USA.* 109:16588–16593. <https://doi.org/10.1073/pnas.1206875109>
- Fukuda, T., F. Pratto, J.C. Schimenti, J.M.A. Turner, R.D. Camerini-Otero, and C. Höög. 2012. Phosphorylation of chromosome core components may serve as axis marks for the status of chromosomal events during mammalian meiosis. *PLoS Genet.* 8:e1002485. <https://doi.org/10.1371/journal.pgen.1002485>
- Fung, J.C., B. Rockmill, M. Odell, and G.S. Roeder. 2004. Imposition of crossover interference through the nonrandom distribution of synapsis initiation complexes. *Cell.* 116:795–802. [https://doi.org/10.1016/s0092-8674\(04\)00249-1](https://doi.org/10.1016/s0092-8674(04)00249-1)
- Gao, J., C. Barroso, P. Zhang, H.-M. Kim, S. Li, L. Labrador, J. Lightfoot, M.V. Gerashchenko, V.M. Labunskyy, M.-Q. Dong, et al. 2016. N-terminal acetylation promotes synaptonemal complex assembly in *C. elegans*. *Genes Dev.* 30:2404–2416. <https://doi.org/10.1101/gad.277350.116>
- Gold, A.L., M.E. Hurlock, A.M. Guevara, L.Y.Z. Isenberg, and Y. Kim. 2025. Identification of the polo-like kinase substrate required for homologous synapsis. *J. Cell Biol.* 224:e202408092. <https://doi.org/10.1083/jcb.202408092>
- Goldfarb, T., and M. Lichten. 2010. Frequent and efficient use of the sister chromatid for DNA double-strand break repair during budding yeast meiosis. *PLoS Biol.* 8:e1000520. <https://doi.org/10.1371/journal.pbio.1000520>
- Gómez-H, L., N. Felipe-Medina, M. Sánchez-Martín, O.R. Davies, I. Ramos, I. García-Tuñón, D.G. de Rooij, I. Dereli, A. Tóth, J.L. Barbero, et al. 2016. C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is essential for mouse fertility. *Nat. Commun.* 7:13298. <https://doi.org/10.1038/ncomms13298>
- Goodyer, W., S. Kaitna, F. Couteau, J.D. Ward, S.J. Boulton, and M. Zetka. 2008. HTP-3 links DSB formation with homolog pairing and crossing over during *C. elegans* meiosis. *Dev. Cell.* 14:263–274. <https://doi.org/10.1016/j.devcel.2007.11.016>

- Gordon, S.G., A.A. Rodriguez, Y. Gu, K.D. Corbett, C.F. Lee, and O. Rog. 2025. The synaptonemal complex aligns meiotic chromosomes by wetting. *Sci. Adv.* 11:eadt5675. <https://doi.org/10.1126/sciadv.adt5675>
- Grelon, M., D. Vezon, G. Gendrot, and G. Pelletier. 2001. AtSPO11-1 is necessary for efficient meiotic recombination in plants. *EMBO J.* 20:589–600. <https://doi.org/10.1093/emboj/20.3.589>
- Gu, Y., A. Desai, and K.D. Corbett. 2022. Evolutionary dynamics and molecular mechanisms of HORMA domain protein signaling. *Annu. Rev. Biochem.* 91:541–569. <https://doi.org/10.1146/annurev-biochem-090920-103246>
- Gyuricza, M.R., K.B. Manheimer, V. Apte, B. Krishnan, E.F. Joyce, B.D. McKee, and K.S. McKim. 2016. Dynamic and stable cohesins regulate synaptonemal complex assembly and chromosome segregation. *Curr. Biol.* 26:1688–1698. <https://doi.org/10.1016/j.cub.2016.05.006>
- Hall, J.C. 1972. Chromosome segregation influenced by two alleles of the meiotic mutant c(3)G in *Drosophila melanogaster*. *Genetics.* 71:367–400. <https://doi.org/10.1093/genetics/71.3.367>
- Hamer, G., K. Gell, A. Kouznetsova, I. Novak, R. Benavente, and C. Höög. 2006. Characterization of a novel meiosis-specific protein within the central element of the synaptonemal complex. *J. Cell Sci.* 119:4025–4032. <https://doi.org/10.1242/jcs.03182>
- Hamer, G., H. Wang, E. Bolcun-Filas, H.J. Cooke, R. Benavente, and C. Höög. 2008. Progression of meiotic recombination requires structural maturation of the central element of the synaptonemal complex. *J. Cell Sci.* 121:2445–2451. <https://doi.org/10.1242/jcs.033233>
- Hemmer, L.W., and J.P. Blumenstiel. 2016. Holding it together: Rapid evolution and positive selection in the synaptonemal complex of *Drosophila*. *BMC Evol. Biol.* 16:91. <https://doi.org/10.1186/s12862-016-0670-8>
- Henggeler, A., L. Orlić, D. Velikov, and J. Matos. 2025. Holliday junction-ZMM protein feedback enables meiotic crossover assurance. *Nature.* 647:766–775. <https://doi.org/10.1038/s41586-025-09559-x>
- Hernández-Hernández, A., S. Masich, T. Fukuda, A. Kouznetsova, S. Sandin, B. Daneholt, and C. Höög. 2016. The central element of the synaptonemal complex in mice is organized as a bilayered junction structure. *J. Cell Sci.* 129:2239–2249. <https://doi.org/10.1242/jcs.182477>
- Higgins, J.D., E. Sanchez-Moran, S.J. Armstrong, G.H. Jones, and F.C.H. Franklin. 2005. The *Arabidopsis* synaptonemal complex protein ZYP1 is required for chromosome synapsis and normal fidelity of crossing over. *Genes Dev.* 19:2488–2500. <https://doi.org/10.1101/gad.354705>
- Hinman, A.W., H.-Y. Yeh, B. Roelens, K. Yamaya, A. Woglar, H.-M.G. Bourbon, P. Chi, and A.M. Villeneuve. 2021. DSB-3 reveals conservation and divergence among protein complexes promoting meiotic double-strand breaks. *Proc. Natl. Acad. Sci. USA.* 118:e2109306118. <https://doi.org/10.1073/pnas.2109306118>
- Hong, S., Y. Sung, M. Yu, M. Lee, N. Kleckner, and K.P. Kim. 2013. The logic and mechanism of homologous recombination partner choice. *Mol. Cell.* 51:440–453. <https://doi.org/10.1016/j.molcel.2013.08.008>
- Hooker, G.W., and G.S. Roeder. 2006. A role for SUMO in meiotic chromosome synapsis. *Curr. Biol.* 16:1238–1243. <https://doi.org/10.1016/j.cub.2006.04.045>
- Humphries, N., W.-K. Leung, B. Argunhan, Y. Terentyev, M. Dvorackova, and H. Tsubouchi. 2013. The Ecm11-Gmc2 complex promotes synaptonemal complex formation through assembly of transverse filaments in budding yeast. *PLoS Genet.* 9:e1003194. <https://doi.org/10.1371/journal.pgen.1003194>
- Hurlock, M.E., I. Čavka, L.E. Kursel, J. Haversat, M. Wooten, Z. Nizami, R. Turniansky, P. Hoess, J. Ries, J.G. Gall, et al. 2020. Identification of novel synaptonemal complex components in *C. elegans*. *J. Cell Biol.* 219:e201910043. <https://doi.org/10.1083/jcb.201910043>
- Ito, M., Y. Yun, D.S. Kulkarni, S. Lee, S. Sandhu, B. Nuñez, L. Hu, K. Lee, N. Lim, R.M. Hirota, et al. 2025. Distinct and interdependent functions of three RING proteins regulate recombination during mammalian meiosis. *Proc. Natl. Acad. Sci. USA.* 122:e2412961121. <https://doi.org/10.1073/pnas.2412961121>
- Jantsch, V., P. Pasierbek, M.M. Mueller, D. Schweizer, M. Jantsch, and J. Loidl. 2004. Targeted gene knockout reveals a role in meiotic recombination for ZHP-3, a Zip3-related protein in *Caenorhabditis elegans*. *Mol. Cell Biol.* 24:7998–8006. <https://doi.org/10.1128/MCB.24.18.7998-8006.2004>
- Jeffress, J.K., S.L. Page, S.K. Royer, E.D. Belden, J.P. Blumenstiel, L.K. Anderson, and R.S. Hawley. 2007. The formation of the central element of the synaptonemal complex may occur by multiple mechanisms: The roles of the N- and C-terminal domains of the *Drosophila* C(3)G protein in mediating synapsis and recombination. *Genetics.* 177:2445–2456. <https://doi.org/10.1534/genetics.107.078717>
- Jiao, X., Z. Liang, J. Li, L. Bai, J. Xu, Y. Liu, and L.-Y. Lu. 2025. Aberrant activation of chromosome asynapsis checkpoint triggers oocyte elimination. *Nat. Commun.* 16:2260. <https://doi.org/10.1038/s41467-025-57702-z>
- Joshi, N., A. Barot, C. Jamison, and G.V. Börner. 2009. Pch2 links chromosome axis remodeling at future crossover sites and crossover distribution during yeast meiosis. *PLoS Genet.* 5:e1000557. <https://doi.org/10.1371/journal.pgen.1000557>
- Kauppi, L., M. Barchi, J. Lange, F. Baudat, M. Jasin, and S. Keeney. 2013. Numerical constraints and feedback control of double-strand breaks in mouse meiosis. *Genes Dev.* 27:873–886. <https://doi.org/10.1101/gad.213652.113>
- Kim, K.P., B.M. Weiner, L. Zhang, A. Jordan, J. Dekker, and N. Kleckner. 2010. Sister cohesion and structural axis components mediate homolog bias of meiotic recombination. *Cell.* 143:924–937. <https://doi.org/10.1016/j.cell.2010.11.015>
- Kim, Y., S.C. Rosenberg, C.L. Kugel, N. Kostow, O. Rog, V. Davydov, T.Y. Su, A.F. Dernburg, and K.D. Corbett. 2014. The chromosome axis controls meiotic events through a hierarchical assembly of HORMA domain proteins. *Dev. Cell.* 31:487–502. <https://doi.org/10.1016/j.devcel.2014.09.013>
- Kim, Y., N. Kostow, and A.F. Dernburg. 2015. The chromosome axis mediates feedback control of CHK-2 to ensure crossover formation in *C. elegans*. *Dev. Cell.* 35:247–261. <https://doi.org/10.1016/j.devcel.2015.09.021>
- Kleckner, N., D. Zickler, G.H. Jones, J. Dekker, R. Padmore, J. Henle, and J. Hutchinson. 2004. A mechanical basis for chromosome function. *Proc. Natl. Acad. Sci. USA.* 101:12592–12597. <https://doi.org/10.1073/pnas.0402724101>
- Köhler, S., M. Wojcik, K. Xu, and A.F. Dernburg. 2025. Dynamic molecular architecture of the synaptonemal complex. *Sci. Adv.* 11:eadq9374. <https://doi.org/10.1126/sciadv.adq9374>
- Kong, A., G. Thorleifsson, M.L. Frigge, G. Masson, D.F. Gudbjartsson, R. Villemoes, E. Magnusdottir, S.B. Olafsdottir, U. Thorsteinsdottir, and K. Stefansson. 2014. Common and low-frequency variants associated with genome-wide recombination rate. *Nat. Genet.* 46:11–16. <https://doi.org/10.1038/ng.2833>
- Kouznetsova, A., H. Wang, M. Bellani, R.D. Camerini-Otero, R. Jessberger, and C. Höög. 2009. BRCA1-mediated chromatin silencing is limited to oocytes with a small number of asynapsed chromosomes. *J. Cell Sci.* 122:2446–2452. <https://doi.org/10.1242/jcs.049353>
- Krishnan, B., S.E. Thomas, R. Yan, H. Yamada, I.B. Zhulin, and B.D. McKee. 2014. Sisters unbound is required for meiotic centromeric cohesion in *Drosophila melanogaster*. *Genetics.* 198:947–965. <https://doi.org/10.1534/genetics.114.166009>
- Kursel, L.E., H.D. Cope, and O. Rog. 2021. Unconventional conservation reveals structure-function relationships in the synaptonemal complex. *Elife.* 10:e72061. <https://doi.org/10.7554/eLife.72061>
- Kurzbaue, M.-T., C. Uanschou, D. Chen, and P. Schlögelhofer. 2012. The recombinases DMCI and RAD51 are functionally and spatially separated during meiosis in *Arabidopsis*. *Plant Cell.* 24:2058–2070. <https://doi.org/10.1105/tpc.112.098459>
- Lake, C.M., R.J. Nielsen, F. Guo, J.R. Unruh, B.D. Slaughter, and R.S. Hawley. 2015. Vilya, a component of the recombination nodule, is required for meiotic double-strand break formation in *Drosophila*. *Elife.* 4:e08287. <https://doi.org/10.7554/eLife.08287>
- Lammers, J.H., H.H. Offenberger, M. van Aalderen, A.C. Vink, A.J. Dietrich, and C. Heyting. 1994. The gene encoding a major component of the lateral elements of synaptonemal complexes of the rat is related to X-linked lymphocyte-regulated genes. *Mol. Cell Biol.* 14:1137–1146. <https://doi.org/10.1128/mcb.14.2.1137-1146.1994>
- Lao, J.P., V. Cloud, C.-C. Huang, J. Grubb, D. Thacker, C.-Y. Lee, M.E. Dresser, N. Hunter, and D.K. Bishop. 2013. Meiotic crossover control by concerted action of Rad51-Dmcl1 in homolog template bias and robust homeostatic regulation. *PLoS Genet.* 9:e1003978. <https://doi.org/10.1371/journal.pgen.1003978>
- Laroussi, H., A.B. Juarez-Martinez, A. Le Roy, E. Boeri Erba, F. Gabel, B. de Massy, and J. Kadlec. 2023. Characterization of the REC114-ME14-IHO1 complex regulating meiotic DNA double-strand break formation. *EMBO J.* 42:e113866. <https://doi.org/10.15252/embj.2023113866>
- Lee, M.-S., M.T. Higashide, H. Choi, K. Li, S. Hong, K. Lee, A. Shinohara, M. Shinohara, and K.P. Kim. 2021. The synaptonemal complex central region modulates crossover pathways and feedback control of meiotic double-strand break formation. *Nucleic Acids Res.* 49:7537–7553. <https://doi.org/10.1093/nar/gkab566>

- Leung, W.-K., N. Humphryes, N. Afshar, B. Argunhan, Y. Terentyev, T. Tsubouchi, and H. Tsubouchi. 2015. The synaptonemal complex is assembled by a polySUMOylation-driven feedback mechanism in yeast. *J. Cell Biol.* 211:785–793. <https://doi.org/10.1083/jcb.201506103>
- Libuda, D.E., S. Uzawa, B.J. Meyer, and A.M. Villeneuve. 2013. Meiotic chromosome structures constrain and respond to designation of crossover sites. *Nature.* 502:703–706. <https://doi.org/10.1038/nature12577>
- Liu, J.G., L. Yuan, E. Brundell, B. Björkroth, B. Daneholt, and C. Höög. 1996. Localization of the N-terminus of SCP1 to the central element of the synaptonemal complex and evidence for direct interactions between the N-termini of SCP1 molecules organized head-to-head. *Exp. Cell Res.* 226: 11–19. <https://doi.org/10.1006/excr.1996.0197>
- Machovina, T.S., R. Mainpal, A. Daryabeigi, O. McGovern, D. Paouneskou, S. Labella, M. Zetka, V. Jantsch, and J.L. Yanowitz. 2016. A surveillance system ensures crossover formation in *C. elegans*. *Curr. Biol.* 26: 2873–2884. <https://doi.org/10.1016/j.cub.2016.09.007>
- MacQueen, A.J., M.P. Colaiácovo, K. McDonald, and A.M. Villeneuve. 2002. Synapsis-dependent and -independent mechanisms stabilize homolog pairing during meiotic prophase in *C. elegans*. *Genes Dev.* 16:2428–2442. <https://doi.org/10.1101/gad.1011602>
- MacQueen, A.J., C.M. Phillips, N. Bhalla, P. Weiser, A.M. Villeneuve, and A.F. Dernburg. 2005. Chromosome sites play dual roles to establish homologous synapsis during meiosis in *C. elegans*. *Cell.* 123:1037–1050. <https://doi.org/10.1016/j.cell.2005.09.034>
- Maguire, M.P. 1965. The relationship of crossover frequency to synaptic extent at pachytene in maize. *Genetics.* 51:23–40. <https://doi.org/10.1093/genetics/51.1.23>
- Martinez-Perez, E., and A.M. Villeneuve. 2005. HTP-1-dependent constraints coordinate homolog pairing and synapsis and promote chiasma formation during *C. elegans* meiosis. *Genes Dev.* 19:2727–2743. <https://doi.org/10.1101/gad.1338505>
- McKim, K.S., B.L. Green-Marroquin, J.J. Sekelsky, G. Chin, C. Steinberg, R. Khodosh, and R.S. Hawley. 1998. Meiotic synapsis in the absence of recombination. *Science.* 279:876–878. <https://doi.org/10.1126/science.279.5352.876>
- Meuwissen, R.L., H.H. Offenberg, A.J. Dietrich, A. Riesewijk, M. van Iersel, and C. Heyting. 1992. A coiled-coil related protein specific for synapsed regions of meiotic prophase chromosomes. *EMBO J.* 11:5091–5100. <https://doi.org/10.1002/j.1460-2075.1992.tb05616.x>
- Morgan, C., J.A. Fozard, M. Hartley, I.R. Henderson, K. Bomblies, and M. Howard. 2021. Diffusion-mediated HEI10 coarsening can explain meiotic crossover positioning in *Arabidopsis*. *Nat Commun.* 12:4674. <https://doi.org/10.1038/s41467-021-24827-w>
- Moses, M.J. 1956. Chromosomal structures in crayfish spermatocytes. *J. Biophys. Biochem. Cytol.* 2:215–218. <https://doi.org/10.1083/jcb.2.2.215>
- Mu, X., H. Murakami, N. Mohibullah, and S. Keeney. 2020. Chromosome-autonomous feedback-down-regulates meiotic DNA break competence upon synaptonemal complex formation. *Genes Dev.* 34:1605–1618. <https://doi.org/10.1101/gad.342873.120>
- Muller, H.J. 1916. The mechanism of crossing-over. *Am. Nat.* 50:193–221. <https://doi.org/10.1086/279534>
- De Muyt, A., A. Pyatnitskaya, J. Andréani, L. Ranjha, C. Ramus, R. Laureau, A. Fernandez-Vega, D. Holoch, E. Girard, J. Govin, et al. 2018. A meiotic XPF-ERCC1-like complex recognizes joint molecule recombination intermediates to promote crossover formation. *Genes Dev.* 32:283–296. <https://doi.org/10.1101/gad.308510.117>
- Nadarajan, S., T.J. Lambert, E. Altendorfer, J. Gao, M.D. Blower, J.C. Waters, and M.P. Colaiácovo. 2017. Polo-like kinase-dependent phosphorylation of the synaptonemal complex protein SYP-4 regulates double-strand break formation through a negative feedback loop. *Elife.* 6:e23437. <https://doi.org/10.7554/eLife.23437>
- Neves, A.R.R., I. Čavka, T. Rausch, and S. Köhler. 2025. Crossovers are regulated by a conserved and disordered synaptonemal complex domain. *Nucleic Acids Res.* 53:gkaf095. <https://doi.org/10.1093/nar/gkaf095>
- Niu, H., L. Wan, V. Busygina, Y. Kwon, J.A. Allen, X. Li, R.C. Kunz, K. Kubota, B. Wang, P. Sung, et al. 2009. Regulation of meiotic recombination via Mek1-mediated Rad54 phosphorylation. *Mol. Cell.* 36:393–404. <https://doi.org/10.1016/j.molcel.2009.09.029>
- Offenberg, H.H., J.A. Schalk, R.L. Meuwissen, M. van Aalderen, H.A. Kester, A.J. Dietrich, and C. Heyting. 1998. SCP2: A major protein component of the axial elements of synaptonemal complexes of the rat. *Nucleic Acids Res.* 26:2572–2579. <https://doi.org/10.1093/nar/26.11.2572>
- Ollinger, R., M. Alsheimer, and R. Benavente. 2005. Mammalian protein SCP1 forms synaptonemal complex-like structures in the absence of meiotic chromosomes. *Mol. Biol. Cell.* 16:212–217. <https://doi.org/10.1091/mbc.e04-09-0771>
- Olson, L.W., U. Edén, M. Egel-Mitani, and R. Egel. 1978. Asynaptic meiosis in fission yeast? *Hereditas.* 89:189–199. <https://doi.org/10.1111/j.1601-5223.1978.tb01275.x>
- Pacheco, S., A. Maldonado-Linares, M. Marcet-Ortega, C. Rojas, A. Martínez-Marchal, J. Fuentes-Lazaro, J. Lange, M. Jasin, S. Keeney, O. Fernández-Capetillo, et al. 2018. ATR is required to complete meiotic recombination in mice. *Nat. Commun.* 9:2622. <https://doi.org/10.1038/s41467-018-04851-z>
- Padmore, R., L. Cao, and N. Kleckner. 1991. Temporal comparison of recombination and synaptonemal complex formation during meiosis in *S. cerevisiae*. *Cell.* 66:1239–1256. [https://doi.org/10.1016/0092-8674\(91\)90046-2](https://doi.org/10.1016/0092-8674(91)90046-2)
- Page, S.L., and R.S. Hawley. 2001. c(3)G encodes a *Drosophila* synaptonemal complex protein. *Genes Dev.* 15:3130–3143. <https://doi.org/10.1101/gad.935001>
- Page, S.L., and R.S. Hawley. 2004. The genetics and molecular biology of the synaptonemal complex. *Annu. Rev. Cell Dev. Biol.* 20:525–558. <https://doi.org/10.1146/annurev.cellbio.19.111301.155141>
- Page, S.L., R.S. Khetani, C.M. Lake, R.J. Nielsen, J.K. Jeffress, W.D. Warren, S.E. Bickel, and R.S. Hawley. 2008. Corona is required for higher-order assembly of transverse filaments into full-length synaptonemal complex in *Drosophila* oocytes. *PLoS Genet.* 4:e1000194. <https://doi.org/10.1371/journal.pgen.1000194>
- Panizza, S., M.A. Mendoza, M. Berlinger, L. Huang, A. Nicolas, K. Shirahige, and F. Klein. 2011. Spo11-accessory proteins link double-strand break sites to the chromosome axis in early meiotic recombination. *Cell.* 146: 372–383. <https://doi.org/10.1016/j.cell.2011.07.003>
- Pattabiraman, D., B. Roelens, A. Woglar, and A.M. Villeneuve. 2017. Meiotic recombination modulates the structure and dynamics of the synaptonemal complex during *C. elegans* meiosis. *PLoS Genet.* 13:e1006670. <https://doi.org/10.1371/journal.pgen.1006670>
- Pyatnitskaya, A., V. Borde, and A. De Muyt. 2019. Crossing and zipping: Molecular duties of the ZMM proteins in meiosis. *Chromosoma.* 128: 181–198. <https://doi.org/10.1007/s00412-019-00714-8>
- Pyatnitskaya, A., J. Andreani, R. Guérois, A. De Muyt, and V. Borde. 2022. The Zip4 protein directly couples meiotic crossover formation to synaptonemal complex assembly. *Genes Dev.* 36:53–69. <https://doi.org/10.1101/gad.348973.121>
- Qiao, H., H.B.D. Prasada Rao, Y. Yang, J.H. Fong, J.M. Cloutier, D.C. Deacon, K.E. Nagel, R.K. Swartz, E. Strong, J.K. Holloway, et al. 2014. Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. *Nat Genet.* 46:194–199. <https://doi.org/10.1038/ng.2858>
- Rasmussen, S.W. 1973. Ultrastructural studies of spermatogenesis in *Drosophila melanogaster* Meigen. *Z. Zellforsch. Mikrosk. Anat.* 140:125–144. <https://doi.org/10.1007/BF00307062>
- Reynolds, A., H. Qiao, Y. Yang, J.K. Chen, N. Jackson, K. Biswas, J.K. Holloway, F. Baudat, B. de Massy, J. Wang, et al. 2013. RNF212 is a dosage-sensitive regulator of crossing-over during mammalian meiosis. *Nat. Genet.* 45: 269–278. <https://doi.org/10.1038/ng.2541>
- Rinaldi, V.D., E. Bolcun-Filas, H. Kogo, H. Kurahashi, and J.C. Schimenti. 2017. The DNA damage checkpoint eliminates mouse oocytes with chromosome synapsis failure. *Mol. Cell.* 67:1026–1036.e2. <https://doi.org/10.1016/j.molcel.2017.07.027>
- Rog, O., S. Köhler, and A. Dernburg. 2017. The synaptonemal complex has liquid crystalline properties and spatially regulates meiotic recombination factors. *Elife.* 6:e21455. <https://doi.org/10.7554/eLife.21455>
- Romanienko, P.J., and R.D. Camerini-Otero. 2000. The mouse Spo11 gene is required for meiotic chromosome synapsis. *Mol. Cell.* 6:975–987. [https://doi.org/10.1016/s1097-2765\(00\)00097-6](https://doi.org/10.1016/s1097-2765(00)00097-6)
- Rosu, S., K.A. Zawadzki, E.L. Stamper, D.E. Libuda, A.L. Reese, A.F. Dernburg, and A.M. Villeneuve. 2013. The *C. elegans* DSB-2 protein reveals a regulatory network that controls competence for meiotic DSB formation and promotes crossover assurance. *PLoS Genet.* 9:e1003674. <https://doi.org/10.1371/journal.pgen.1003674>
- Rousová, D., V. Nivsarkar, V. Altmannova, V.B. Raina, S.K. Funk, D. Liedtke, P. Janning, F. Müller, H. Reichle, G. Vader, and J.R. Weir. 2021. Novel mechanistic insights into the role of Mer2 as the keystone of meiotic DNA break formation. *Elife.* 10:e72330. <https://doi.org/10.7554/eLife.72330>
- Royo, H., H. Prosser, Y. Ruzankina, S.K. Mahadevaiah, J.M. Cloutier, M. Baumann, T. Fukuda, C. Höög, A. Tóth, D.G. de Rooij, et al. 2013. ATR acts stage specifically to regulate multiple aspects of mammalian

- meiotic silencing. *Genes Dev.* 27:1484–1494. <https://doi.org/10.1101/gad.219477.113>
- Sanchez-Moran, E., J.-L. Santos, G.H. Jones, and F.C.H. Franklin. 2007. ASY1 mediates AtDMC1-dependent interhomolog recombination during meiosis in *Arabidopsis*. *Genes Dev.* 21:2220–2233. <https://doi.org/10.1101/gad.439007>
- Sánchez-Sáez, F., L. Gómez-H, O.M. Dunne, C. Gallego-Páramo, N. Felipe-Medina, M. Sánchez-Martín, E. Llano, A.M. Pendas, and O.R. Davies. 2020. Meiotic chromosome synapsis depends on multivalent SYCE1-SIX6OS1 interactions that are disrupted in cases of human infertility. *Sci. Adv.* 6:eabb1660. <https://doi.org/10.1126/sciadv.abb1660>
- Schalbetter, S.A., G. Fudenberg, J. Baxter, K.S. Pollard, and M.J. Neale. 2019. Principles of meiotic chromosome assembly revealed in *S. cerevisiae*. *Nat. Commun.* 10:4795. <https://doi.org/10.1038/s41467-019-12629-0>
- Schild-Prüfert, K., T.T. Saito, S. Smolnikov, Y. Gu, M. Hincapie, D.E. Hill, M. Vidal, K. McDonald, and M.P. Colaiácovo. 2011. Organization of the synaptonemal complex during meiosis in *Caenorhabditis elegans*. *Genetics*. 189:411–421. <https://doi.org/10.1534/genetics.111.132431>
- Schramm, S., J. Fraune, R. Naumann, A. Hernandez-Hernandez, C. Höög, H.J. Cooke, M. Alsheimer, and R. Benavente. 2011. A novel mouse synaptonemal complex protein is essential for loading of central element proteins, recombination, and fertility. *PLoS Genet.* 7:e1002088. <https://doi.org/10.1371/journal.pgen.1002088>
- Schücker, K., T. Holm, C. Franke, M. Sauer, and R. Benavente. 2015. Elucidation of synaptonemal complex organization by super-resolution imaging with isotropic resolution. *Proc. Natl. Acad. Sci. USA.* 112:2029–2033. <https://doi.org/10.1073/pnas.1414841112>
- Schwacha, A., and N. Kleckner. 1997. Interhomolog bias during meiotic recombination: Meiotic functions promote a highly differentiated interhomolog-only pathway. *Cell.* 90:1123–1135. [https://doi.org/10.1016/S0092-8674\(00\)80378-5](https://doi.org/10.1016/S0092-8674(00)80378-5)
- Seear, P.J., H.J.A. Dowling, M. Szymańska-Lejman, W. Dziegielewski, S. DeBilio, F.C.H. Franklin, K.D. Corbett, O.R. Davies, P.A. Ziolkowski, and J.D. Higgins. 2025. SCEP3 initiates synapsis and implements crossover interference in *Arabidopsis*. *Nat. Plants.* 11:2531–2547. <https://doi.org/10.1038/s41477-025-02155-x>
- Sharmin, S., K. Voelkel-Meiman, A.J. Poppel, and A.J. MacQueen. 2025. Crossover formation and coordinated assembly of synaptonemal complex relies on a direct interaction between Zip1 and Zip3. *bioRxiv*. <https://doi.org/10.1101/2025.11.21.689840> (Preprint posted November 22, 2025).
- Shin, Y.-H., Y. Choi, S.U. Erdin, S.A. Yatsenko, M. Kloc, F. Yang, P.J. Wang, M.L. Meistrich, and A. Rajkovic. 2010. Hormad1 mutation disrupts synaptonemal complex formation, recombination, and chromosome segregation in mammalian meiosis. *PLoS Genet.* 6:e1001190. <https://doi.org/10.1371/journal.pgen.1001190>
- Shin, Y.-H., M.M. McGuire, and A. Rajkovic. 2013. Mouse HORMAD1 is a meiosis I checkpoint protein that modulates DNA double-strand break repair during female meiosis. *Biol. Reprod.* 89:29. <https://doi.org/10.1095/biolreprod.112.106773>
- Smith, A.V., and G.S. Roeder. 1997. The yeast Red1 protein localizes to the cores of meiotic chromosomes. *J. Cell Biol.* 136:957–967. <https://doi.org/10.1083/jcb.136.5.957>
- Smolnikov, S., A. Eizinger, K. Schild-Prüfert, A. Hurlburt, K. McDonald, J. Engebrecht, A.M. Villeneuve, and M.P. Colaiácovo. 2007. SYP-3 restricts synaptonemal complex assembly to bridge paired chromosome axes during meiosis in *Caenorhabditis elegans*. *Genetics*. 176:2015–2025. <https://doi.org/10.1534/genetics.107.072413>
- Smolnikov, S., K. Schild-Prüfert, and M.P. Colaiácovo. 2009. A yeast two-hybrid screen for SYP-3 interactors identifies SYP-4, a component required for synaptonemal complex assembly and chiasma formation in *Caenorhabditis elegans* meiosis. *PLoS Genet.* 5:e1000669. <https://doi.org/10.1371/journal.pgen.1000669>
- Stamper, E.L., S.E. Rodenbusch, S. Rosu, J. Ahringer, A.M. Villeneuve, and A.F. Dernburg. 2013. Identification of DSB-1, a protein required for initiation of meiotic recombination in *Caenorhabditis elegans*, illuminates a crossover assurance checkpoint. *PLoS Genet.* 9:e1003679. <https://doi.org/10.1371/journal.pgen.1003679>
- Stanzione, M., M. Baumann, F. Papanikos, I. Dereli, J. Lange, A. Ramlal, D. Tränkner, H. Shibuya, B. de Massy, Y. Watanabe, et al. 2016. Meiotic DNA break formation requires the unsynapsed chromosome axis-binding protein IHO1 (CCDC36) in mice. *Nat. Cell Biol.* 18:1208–1220. <https://doi.org/10.1038/ncb3417>
- Storlazzi, A., L. Xu, A. Schwacha, and N. Kleckner. 1996. Synaptonemal complex (SC) component Zip1 plays a role in meiotic recombination independent of SC polymerization along the chromosomes. *Proc. Natl. Acad. Sci. USA.* 93:9043–9048. <https://doi.org/10.1073/pnas.93.17.9043>
- Sym, M., and G.S. Roeder. 1994. Crossover interference is abolished in the absence of a synaptonemal complex protein. *Cell.* 79:283–292. [https://doi.org/10.1016/0092-8674\(94\)90197-x](https://doi.org/10.1016/0092-8674(94)90197-x)
- Sym, M., and G.S. Roeder. 1995. Zip1-induced changes in synaptonemal complex structure and polycomplex assembly. *J. Cell Biol.* 128:455–466. <https://doi.org/10.1083/jcb.128.4.455>
- Sym, M., J.A. Engebrecht, and G.S. Roeder. 1993. ZIP1 is a synaptonemal complex protein required for meiotic chromosome synapsis. *Cell.* 72:365–378. [https://doi.org/10.1016/0092-8674\(93\)90114-6](https://doi.org/10.1016/0092-8674(93)90114-6)
- Takeo, S., C.M. Lake, E. Morais-de-Sá, C.E. Sunkel, and R.S. Hawley. 2011. Synaptonemal complex-dependent centromeric clustering and the initiation of synapsis in *Drosophila* oocytes. *Curr. Biol.* 21:1845–1851. <https://doi.org/10.1016/j.cub.2011.09.044>
- Tan, T., Y. Tan, Y. Wang, X. Yang, B. Zhai, S. Zhang, X. Yang, H. Nie, J. Gao, J. Zhou, et al. 2022. Negative supercoils regulate meiotic crossover patterns in budding yeast. *Nucleic Acids Res.* 50:10418–10435. <https://doi.org/10.1093/nar/gkac786>
- Tan, T., Y. Zhao, Y. Chen, Y. Mi, J. Zeng, P. Du, T. Han, Y. Liu, N. Li, J. Kong, et al. 2026. Ufd2p promotes efficient crossover formation by destabilizing Top2p during meiosis. *Proc. Natl. Acad. Sci. USA.* 123:e2517398123. <https://doi.org/10.1073/pnas.2517398123>
- Tang, S., S. Hariri, R. Bohn, J.E. McCarthy, J. Koo, M. Pourhosseinzadeh, E. Nguyen, N. Liu, C. Ma, H. Lu, et al. 2025. Protecting double Holliday junctions ensures crossing over during meiosis. *Nature.* 647:776–785. <https://doi.org/10.1038/s41586-025-09555-1>
- Tanneti, N.S., K. Landy, E.F. Joyce, and K.S. McKim. 2011. A pathway for synapsis initiation during zygotene in *Drosophila* oocytes. *Curr. Biol.* 21:1852–1857. <https://doi.org/10.1016/j.cub.2011.10.005>
- Thacker, D., N. Mohibullah, X. Zhu, and S. Keeney. 2014. Homologue engagement controls meiotic DNA break number and distribution. *Nature.* 510:241–246. <https://doi.org/10.1038/nature13120>
- Tromer, E.C., T.A. Wemyss, P. Ludzia, R.F. Waller, and B. Akiyoshi. 2021. Repurposing of synaptonemal complex proteins for kinetochores in *Kinetoplastida*. *Open Biol.* 11:210049. <https://doi.org/10.1098/rsob.210049>
- Ur, S.N., and K.D. Corbett. 2021. Architecture and dynamics of meiotic chromosomes. *Annu. Rev. Genet.* 55:497–526. <https://doi.org/10.1146/annurev-genet-071719-020235>
- Voelkel-Meiman, K., S.S. Moustafa, P. LeFrançois, A.M. Villeneuve, and A.J. MacQueen. 2012. Full-length synaptonemal complex grows continuously during meiotic prophase in budding yeast. *PLoS Genet.* 8:e1002993. <https://doi.org/10.1371/journal.pgen.1002993>
- Voelkel-Meiman, K., S.-Y. Cheng, S.J. Morehouse, and A.J. MacQueen. 2016. Synaptonemal complex proteins of budding yeast define reciprocal roles in MutS γ -mediated crossover formation. *Genetics*. 203:1091–1103. <https://doi.org/10.1534/genetics.115.182923>
- Voelkel-Meiman, K., S.-Y. Cheng, M. Parziale, S.J. Morehouse, A. Feil, O.R. Davies, A. de Muyt, V. Borde, and A.J. MacQueen. 2019. Crossover recombination and synapsis are linked by adjacent regions within the N terminus of the Zip1 synaptonemal complex protein. *PLoS Genet.* 15:e1008201. <https://doi.org/10.1371/journal.pgen.1008201>
- Vrielynck, N., M. Peuch, S. Durand, Q. Lian, A. Chambon, A. Hurel, J. Guérin, R. Guérois, R. Mercier, M. Grelon, and C. Mézard. 2023. SCEP1 and SCEP2 are two new components of the synaptonemal complex central element. *Nat. Plants.* 9:2016–2030. <https://doi.org/10.1038/s41477-023-01558-y>
- Wang, Y., Z. Yu, C.K. Cahoon, T. Parmely, N. Thomas, J.R. Unruh, B.D. Slaughter, and R.S. Hawley. 2018. Combined expansion microscopy with structured illumination microscopy for analyzing protein complexes. *Nat. Protoc.* 13:1869–1895. <https://doi.org/10.1038/s41596-018-0023-8>
- Wang, S., C. Veller, F. Sun, A. Ruiz-Herrera, Y. Shang, H. Liu, D. Zickler, Z. Chen, N. Kleckner, and L. Zhang. 2019. Per-nucleus crossover covariation and implications for evolution. *Cell.* 177:326–338.e16. <https://doi.org/10.1016/j.cell.2019.02.021>
- Webber, H.A., L. Howard, and S.E. Bickel. 2004. The cohesion protein ORD is required for homologue bias during meiotic recombination. *J. Cell Biol.* 164:819–829. <https://doi.org/10.1083/jcb.200310077>
- West, A.M., S.C. Rosenberg, S.N. Ur, M.K. Lehmer, Q. Ye, G. Hagemann, I. Caballero, I. Usón, A.J. MacQueen, F. Herzog, and K.D. Corbett. 2019. A conserved filamentous assembly underlies the structure of the meiotic chromosome axis. *Elife.* 8:e40372. <https://doi.org/10.7554/eLife.40372>

- von Wettstein, D., S.W. Rasmussen, and P.B. Holm. 1984. The synaptonemal complex in genetic segregation. *Annu. Rev. Genet.* 18:331–413. <https://doi.org/10.1146/annurev.ge.18.120184.001555>
- Widger, A., S.K. Mahadevaiah, J. Lange, E. ElInati, J. Zohren, T. Hirota, S. Pacheco, A. Maldonado-Linares, M. Stanzione, O. Ojarikre, et al. 2018. ATR is a multifunctional regulator of male mouse meiosis. *Nat. Commun.* 9:2621. <https://doi.org/10.1038/s41467-018-04850-0>
- Williams, S., G. McKown, Z. Yu, J. Gardner, C. Staber, M.C. Gibson, and R.S. Hawley. 2025. The synaptonemal complex component corolla regulates meiotic crossover formation in *Drosophila melanogaster*. *Chromosoma.* 134:10. <https://doi.org/10.1007/s00412-025-00839-z>
- Winkel, K., M. Alsheimer, R. Ollinger, and R. Benavente. 2009. Protein SYCP2 provides a link between transverse filaments and lateral elements of mammalian synaptonemal complexes. *Chromosoma.* 118:259–267. <https://doi.org/10.1007/s00412-008-0194-0>
- Woglar, A., and A.M. Villeneuve. 2018. Dynamic architecture of DNA repair complexes and the synaptonemal complex at sites of meiotic recombination. *Cell.* 173:1678–1691.e16. <https://doi.org/10.1016/j.cell.2018.03.066>
- Wojtasz, L., K. Daniel, I. Roig, E. Bolcun-Filas, H. Xu, V. Boonsanay, C.R. Eckmann, H.J. Cooke, M. Jasin, S. Keeney, et al. 2009. Mouse HORMAD1 and HORMAD2, two conserved meiotic chromosomal proteins, are depleted from synapsed chromosome axes with the help of TRIP13 AAA-ATPase. *PLoS Genet.* 5:e1000702. <https://doi.org/10.1371/journal.pgen.1000702>
- Wojtasz, L., J.M. Cloutier, M. Baumann, K. Daniel, J. Varga, J. Fu, K. Anastasiadis, A.F. Stewart, A. Reményi, J.M.A. Turner, and A. Tóth. 2012. Meiotic DNA double-strand breaks and chromosome asynapsis in mice are monitored by distinct HORMAD2-independent and -dependent mechanisms. *Genes Dev.* 26:958–973. <https://doi.org/10.1101/gad.187559.112>
- Xu, H., Z. Tong, Q. Ye, T. Sun, Z. Hong, L. Zhang, A. Bortnick, S. Cho, P. Beuzer, J. Axelrod, et al. 2019. Molecular organization of mammalian meiotic chromosome axis revealed by expansion STORM microscopy. *Proc. Natl. Acad. Sci. USA.* 116:18423–18428. <https://doi.org/10.1073/pnas.1902440116>
- Yan, R., and B.D. McKee. 2013. The cohesion protein SOLO associates with SMC1 and is required for synapsis, recombination, homolog bias and cohesion and pairing of centromeres in *Drosophila* meiosis. *PLoS Genet.* 9:e1003637. <https://doi.org/10.1371/journal.pgen.1003637>
- Zetka, M.C., I. Kawasaki, S. Strome, and F. Müller. 1999. Synapsis and chiasma formation in *Caenorhabditis elegans* require HIM-3, a meiotic chromosome core component that functions in chromosome segregation. *Genes Dev.* 13:2258–2270. <https://doi.org/10.1101/gad.13.17.2258>
- Zhang, L., E. Espagne, A. de Muyt, D. Zickler, and N.E. Kleckner. 2014a. Interference-mediated synaptonemal complex formation with embedded crossover designation. *Proc. Natl. Acad. Sci. USA.* 111:E5059–E5068. <https://doi.org/10.1073/pnas.1416411111>
- Zhang, L., S. Wang, S. Yin, S. Hong, K.P. Kim, and N. Kleckner. 2014b. Topoisomerase II mediates meiotic crossover interference. *Nature.* 511:551–556. <https://doi.org/10.1038/nature13442>
- Zhang, L., S. Köhler, R. Rillo-Bohn, and A.F. Dernburg. 2018. A compartmentalized signaling network mediates crossover control in meiosis. *Elife.* 7:e30789. <https://doi.org/10.7554/eLife.30789>
- Zhang, Z., S. Xie, R. Wang, S. Guo, Q. Zhao, H. Nie, Y. Liu, F. Zhang, M. Chen, L. Liu, et al. 2020. Multivalent weak interactions between assembly units drive synaptonemal complex formation. *J. Cell Biol.* 219:e201910086. <https://doi.org/10.1083/jcb.201910086>
- Zhang, L., W. Stauffer, C. Liu, H. Shao, N. Abuzahriyeh, R. Jiang, D. Zwicker, X. Liu, X. Yao, and A.F. Dernburg. 2025. Crossover patterning through condensation and coarsening of pro-crossover factors. *Nat. Cell Biol.* 27:1161–1174. <https://doi.org/10.1038/s41556-025-01688-9>
- Ziolkowski, P.A., C.J. Underwood, C. Lambing, M. Martinez-Garcia, E.J. Lawrence, L. Ziolkowska, C. Griffin, K. Choi, F.C.H. Franklin, R.A. Martienssen, and I.R. Henderson. 2017. Natural variation and dosage of the HEI10 meiotic E3 ligase control crossover recombination. *Genes Dev.* 31:306–317. <https://doi.org/10.1101/gad.295501.116>
- Zwettler, F.U., M.-C. Spindler, S. Reinhard, T. Klein, A. Kurz, R. Benavente, and M. Sauer. 2020. Tracking down the molecular architecture of the synaptonemal complex by expansion microscopy. *Nat. Commun.* 11:3222. <https://doi.org/10.1038/s41467-020-17017-7>