

TUBULAR CONNECTIONS BETWEEN DICTYOSOMES AND FORMING SECRETORY VESICLES IN PLANT GOLGI APPARATUS

HILTON H. MOLLENHAUER and D. JAMES MORRÉ. From the Charles F. Kettering Research Laboratory, Yellow Springs, Ohio, and the Department of Botany and Plant Pathology, Purdue University, Lafayette, Indiana

The Golgi apparatus of most plant and animal cells occurs as a system of dictyosomes (4), often dispersed throughout a large portion of the cytoplasm. Each dictyosome consists of a stack of cisternae (usually 5 or 6) and associated vesicles. The central region of a typical cisterna is sac- or platelike and about $\frac{1}{2} \mu$ in diameter. Tubular proliferations extend from this central structure, uniting at intervals to form an anastomosing network of tubules (reference 1 and Figs. 1 and 2). In plant cells, two vesicle types consistently form part of the tubular network (reference 1 and Figs. 1 and 2). These vesicles represent regions of specialization different from the usual secretion granules or vesicles reported for maize root tip (5), slime-secreting cells of insectivores (8), and root hairs (9).

Secretion vesicles are generally lost during dictyosome isolation (6), but their relationship to the cisternal tubules can be visualized from sectioned material. Due to the usual cup-shape of the dictyosomes, tangential sections through an entire cisterna are rare. However, partial tangential sections are numerous and were used to clarify the mode of secretion vesicle attachment to the cisternal tubules.

Tangentially sectioned dictyosomes from glutaraldehyde-fixed, OsO_4 -postfixed maize root tip have the same tubular cisternal form as isolated dictyosomes. The secretion product is contained in forming vesicles attached to the cisternae by one or more tubules (Figs. 3 and 4).

To clarify the tubular nature of plant dictyosomes and to preclude artifacts solely due to

glutaraldehyde, we include results with tissues fixed in KMnO_4 (Figs. 5 and 6), confirming the above observation. The oft published (5, 10) cross-sectional view of a KMnO_4 -fixed dictyosome is shown on the right in Fig. 6. Only in tangential section is the tubular nature of the structure clearly revealed (at the left of Fig. 6). Thus, the tubular character of the plant dictyosome is not restricted to cold-stored onion (1) and storage tissues (radish of Fig. 1 and cauliflower of Fig. 2) in which the dictyosomes might be expected to be relatively inactive. It is also present in dictyosomes of actively secreting cells. Preliminary observations on other tissues suggest that the tubular nature of dictyosome cisternae is widespread and not necessarily restricted to plant cells. That cisternal tubules are preserved by KMnO_4 fixation demonstrates that they are not artifacts attributable to glutaraldehyde stabilization.

A tubular attachment for 2 forms of secretion vesicles of cap cells of maize root tip has been demonstrated (Figs. 3 to 6). During formation of secretion vesicles, the cisternal tubules provide a channel for transfer of materials and possibly serve to segregate the activities of the cisternal lumen from those of the forming vesicles.

Contribution No. 219 from the Charles F. Kettering Research Laboratory, Yellow Springs, Ohio. Work supported in part by a grant from the National Science Foundation, GB-1084. Purdue University AES Journal Paper No. 2702.

Received for publication 22 December 1965.





FIGURE 1 Isolated radish dictyosome stabilized with 0.1% glutaraldehyde (7) and stained with phosphotungstic acid (1). The component cisternae have separated, revealing a central region consisting of discs or flattened sacs about $\frac{1}{2} \mu$ in diameter. $\times 73,000$.

FIGURE 2 Isolated dictyosome of cauliflower Golgi apparatus stabilized with glutaraldehyde and stained as described for Fig. 1. Tubular proliferations often extend a considerable distance from the central region. Arrow points to a "shaggy" vesicle (1). $\times 42,000$.



FIGURES 3 and 4 Micrographs from tangential sections through maize root cap dictyosomes, showing the tubular interconnections between dictyosomes and forming secretion vesicles. Tissue fixed in glutaraldehyde, postfixed in OsO_4 , and embedded in No. 1 epoxy resin mixture of Mollenhauer (2). Fig. 3. Arrow points to a "shaggy" vesicle (1). $\times 32,000$. Fig. 4. $\times 41,000$.

FIGURES 5 and 6 Micrographs from tangential sections through maize root tip dictyosomes, showing the tubular nature of the cisternae and the tubular connections to the forming secretion vesicles. Tissue fixed in 2% aqueous KMnO_4 and embedded in No. 1 epoxy resin mixture. Fig. 5. Annular region of root cap (3). Approximately $\times 45,000$. Fig. 6. Root cap mantle region (3). $\times 40,000$.

REFERENCES

1. CUNNINGHAM, W. P., MORRÉ, D. J., and MOLLENHAUER, H. H., Structure of isolated plant Golgi apparatus revealed by negative staining, *J. Cell Biol.*, 1966, **28**, 169.
2. MOLLENHAUER, H. H., Plastic embedding mixtures for use in electron microscopy, *Stain Technol.*, 1964, **39**, 111.
3. MOLLENHAUER, H. H., Transition forms of Golgi apparatus secretion vesicles, *J. Ultrastruct. Research*, 1965, **12**, 439.
4. MOLLENHAUER, H. H., and MORRÉ, D. J., Interassociation of dictyosomes to form plant cell Golgi apparatus, *J. Cell Biol.*, 1965, **27**, 68A.
5. MOLLENHAUER, H. H., WHALEY, W. G., LEECH, J. H., A function of the Golgi apparatus in outer root cap cells, *J. Ultrastruct. Research*, 1961, **5**, 193.
6. MORRÉ, D. J., and MOLLENHAUER, H. H., Isolation of the Golgi apparatus from plant cells, *J. Cell Biol.*, 1964, **23**, 295.
7. MORRÉ, D. J., MOLLENHAUER, H. H., and CHAMBERS, J. E., Glutaraldehyde stabilization as an aid to Golgi apparatus isolation, *Exp. Cell Research*, 1965, **38**, 672.
8. SCHNEPF, E., Physiologie und Morphologie sekretorischer Pflanzenzellen, in Wissenschaftliche Konferenz der Gesellschaft Deutscher Naturforscher und Ärzte, 2. Sekretion und Exkretion, Friedrichroda, 1964, (K. E. Wohlfarth-Bottermann, editor), Berlin, Springer-Verlag, 1965, 72.
9. SIEVERS, A., Funktion des Golgi-Apparatus in pflanzlichen und tierischen Zellen, in Wissenschaftliche Konferenz der Gesellschaft Deutscher Naturforscher und Ärzte, 2. Sekretion und Exkretion, Friedrichroda, 1964, (K. E. Wohlfarth-Bottermann, editor) Berlin, Springer-Verlag, 1965, 89.
10. WHALEY, W. G., KEPHART, J. E., and MOLLENHAUER, H. H., The dynamics of cytoplasmic membranes during development, in 22nd Symposium of the Society for Study, Development, and Growth, Cellular Membranes in Development, Storrs, Connecticut, 1963, **22**, (M. Locke, editor), New York, Academic Press Inc., 1964, 135.