

SORTING OUT OF NORMAL AND VIRUS-TRANSFORMED CELLS IN CELLULAR AGGREGATES

HOWARD GERSHMAN, JUDITH DRUMM, and LLOYD CULP

From the Departments of Biochemistry and Microbiology, Case Western Reserve University Medical School, Cleveland, Ohio 44106

ABSTRACT

The sorting-out behavior (self-segregation of two cell types from mixtures of the two) of five different established cell lines was studied. Eight of the ten possible binary combinations of these lines, cultured as cellular aggregates, were examined. Mouse BALB/c 3T3 cells sorted out internally to the corresponding malignant SV40 virus-transformed 3T3 cells. The transformed 3T3 line (SVT-2) did not sort out from a revertant line selected from SVT-2 cells by resistance to concanavalin A (con A). The revertant cells sorted out externally to the parent BALB/c 3T3 cells, although segregation was generally incomplete. BALB/c 3T3 cells did not sort out from another contact-inhibited line of 3T3 cells derived from Swiss albino mice (Swiss 3T3).

Both BALB/c 3T3 and Swiss 3T3 cells sorted out from cells of the contact-inhibited hamster line, NIL B. Instead of a two-layered sphere, however, a three-layered structure was observed with most of the NIL B cells external to the 3T3 cells, and a few NIL B cells comprising the center of the sphere. On the other hand, NIL B cells did not consistently sort out from either the SVT-2 or con A cells.

In general, sorting out between pairs of these five lines are slower and less complete than is generally observed between the more extensively studied chick embryonic tissue cells, suggesting that the cultured cells may be more closely related in their adhesive properties. The internal segregation of BALB/c 3T3 cells relative to SVT-2 cells is consistent with the hypothesis that transformed cells are less adhesive than their nontransformed counterparts.

The interaction of normal and malignant cells in culture has been the subject of considerable study. Much of this work has dealt with cells cultured on artificial substrata such as glass or plastic (e.g., 2, 4, 20, 21, 29, 32), especially that work utilizing established cell lines and their virus-transformed counterparts. While the study of such established lines has greatly contributed to our knowledge of how growth and movement are controlled, the presence of a nonphysiological substratum may interfere with the application of this knowledge to

in vivo situations. In particular, the study of adhesive interactions between normal and transformed cells in monolayer culture should be viewed with caution. Several workers have pointed out that in such situations the competition of cell-adhesions with cell-dish adhesions must be considered (6, 26, 33) rather than limiting the analysis to cell-to-cell adhesions. In order to avoid this particular complication, we have studied the interaction of virus-transformed and nontransformed cells cultured as cellular aggregates or

masses, in which cells are maintained in suspension in agitated liquid medium. Our purpose was to compare the properties of a nontransformed: transformed: revertant family of cell lines and also to compare the properties of a number of lines that exhibit varying degrees of contact inhibition.

The property of cells in these aggregates that was examined in this study was their ability to sort out or self-segregate. When cells of two suitable tissue types are intermixed, they sort out with the cells of one type forming a sphere surrounding the cells of the second type. Sorting out has been demonstrated for embryonic tissues from a number of vertebrate species, including mouse (5, 13, 22), chick (23, 28), human (8), and amphibian (31). Steinberg (27, 28) has proposed that the equilibrium configuration achieved after sorting out is the result of quantitative differences in adhesiveness between the two cell types. Although universal agreement on all the details of this hypothesis is still lacking, it is generally accepted that sorting-out behavior is a reflection of an adhesive interaction between the cells. For the purposes of discussion in this report, we will also tentatively consider that these adhesive interactions represent different strengths of intercellular adhesion.

MATERIALS AND METHODS

Chemicals and Radiochemicals

All chemicals were purchased from commercial sources and were either biological grade or else the highest grade available.

[Methyl-³H]thymidine was purchased from New England Nuclear (Boston, Mass.) or Amersham/Searle Corp. (Arlington Hts., Ill.) and had a spec act of 18–20 Ci/mmol.

Growth of Cell Lines

The growth of mouse fibroblast BALB/c 3T3 (1) and SV40-transformed BALB/c 3T3 (designated SVT-2) (1), con A revertant (10, 11, 12), hamster NIL B (14), and Swiss albino mice (Swiss 3T3) cells (1) has been described previously (18). Cells were found to be free of *Mycoplasma* contamination by an autoradiographic assay (18, 11).

Cells were routinely subcultured using trypsin-(ethylenedinitrilo)tetraacetic acid (EDTA)¹ solutions (0.05% Difco trypsin 1:250 (Difco Laboratories, Detroit, Mich.) and 0.5 mM EDTA). For experimental use, cells

¹ *Abbreviations used in this paper:* EDTA, (ethylenedinitrilo)tetraacetic acid; EGTA, [ethylenebis(oxyethylenetrilo)]tetraacetic acid; and PBS, phosphate-buffered saline (Dulbecco's A solution).

were inoculated into 100-mm plastic culture dishes in 15-ml medium and grown as subconfluent cultures for 3–4 days (three to five generations). The medium was changed 24 h after plating and every 48 h thereafter. Cells were removed from the dishes with [ethylenebis(oxyethylenetrilo)]tetraacetic acid (EGTA) (0.5 mM in Ca⁺⁺-Mg⁺⁺-free phosphate-buffered saline (Dulbecco's A solution) (PBS) and used either to inoculate dishes or to make aggregates. Thus, cells used in these experiments were grown as subconfluent cultures (not more than 25% confluent) and were not harvested with trypsin. These cells reached the following densities if allowed to grow to saturation with daily changes of medium: BALB/c 3T3, 10 × 10⁴ cells/cm²; SVT-2, 100 × 10⁴ cells/cm²; con A, 20 × 10⁴ cells/cm²; Swiss 3T3, 5 × 10⁴ cells/cm²; NIL B, 15–20 × 10⁴ cells/cm².

Aggregates were formed by a technique previously reported (18). Briefly, this involves mixing together 1.5 × 10⁶ cells of each type, centrifuging the suspension at low speed, and after a 1.5-h incubation period, cutting the pellet into cubical fragments about 0.4 mm on a side. The aggregates are cultured in hanging drops for 24 h and on a gyratory shaker for an additional 1.5–3.5 days.

Labeling of Cells with [³H]Thymidine

Cells were radiolabeled by the addition of [³H]thymidine to the culture medium 3–4 days before the experiment. 24 h before the experiment, the radioactive medium was replaced by nonradioactive medium. If this chase period was omitted, the nonradioactive cells in the aggregates became radioactive within 1–2 days. This may have occurred by transfer of nucleotides between cells, since this process has already been suggested as a possible explanation for the contact-mediated "metabolic co-operation" observed in culture (30).

In early experiments cells were labeled by the addition of 2.6 μCi/ml [³H]thymidine to the medium. This concentration was reduced to 0.1 μCi/ml for some of the experiments reported here. In each case in which the higher concentration was used the experiment was also repeated with the lower concentrations. No differences were found between these duplicate experiments. When 0.1 μCi/ml [³H]thymidine was added, however, total labeling per cell was decreased, and autoradiography was done by the method of Weingrad et al.² Histology and autoradiography were performed as previously described (18).

Determination of Plating Efficiency

EGTA-suspended cells were washed and either inoculated into dishes or formed into aggregates. The aggregates (pellet fragments) were cultured in hanging drops for one day and then in small vials on a gyratory shaker

² Weingrad, D., R. M. Sade, C. Haudenschild, and J. Folkman. Manuscript in preparation.

(180–200 gyrations per min) for an additional 2 days. The cells inoculated into dishes were cultured in a 37° C CO₂-gassed incubator for the same 3 days.

After this period the aggregates and dishes were rinsed once in PBS and treated with trypsin (0.05% Difco 1:250 plus 0.5 mM EDTA) in PBS for 10 min at 37°C. The single cell suspensions were counted and diluted into complete medium to the appropriate concentration. Cells were then seeded into 60-mm plastic dishes at 250, 500, or 1,000 cells per dish and cultured for 2 weeks, with changes of medium every 3–4 days. Colonies were counted after staining with Wright's solution.

In early experiments dish-maintained cells were inoculated at varying densities. In later experiments they were inoculated at the following densities, which were just sufficient to produce confluent cultures within 24 h: BALB/c 3T3, 1.8×10^6 cells/60-mm dish; SVT-2, 5×10^6 cells/60-mm dish; NIL B, 3×10^6 cells/60-mm dish.

RESULTS

Plating Efficiency Experiments

In order to assess the viability of cells maintained as aggregates, the ability of these cells to form colonies relative to cells maintained as monolayers was studied. In one set of experiments, cells cultured as aggregates were compared to parallel cultures in which cells were maintained on dishes at low to moderate density. The results showed a moderate amount of variability, not only between cell lines but also between experiments with the same line. Relative plating efficiencies, i.e., plating efficiency of cells in aggregates divided by the

plating efficiency of cells in dishes, varied from a low of 44% to a high of 100%. Absolute plating efficiencies varied from 10 to 40 colonies per 100 cells plated.

A second series of experiments was performed using some of these lines in which the cell density of monolayer cultures was very high and therefore more comparable to the cell density in aggregates. The cells were seeded into 60-mm dishes at a density just sufficient to produce confluent monolayers when the cells had attached and spread. These data are presented in Table I. Both the absolute and relative plating efficiencies observed ranged from a low of 14.3 to a high of 32.9 colonies per 100 cells plated. These experiments suggest that the viability of cells maintained in aggregates for 3 days is similar to their viability when maintained in a confluent state in dishes for a similar period of time.

Sorting Out of Mouse-Mouse Cell

Line Mixtures

BALB/c 3T3 cells and SV-40-transformed 3T3 cells were mixed and formed into aggregates in reciprocal combinations (³H]thymidine-labeled 3T3 cells plus unlabeled SVT-2 cells on the one hand, and [³H]thymidine-labeled SVT-2 cells and unlabeled BALB/c 3T3 cells, on the other). The results were visualized by sectioning and autoradiography as described in Materials and Methods. One of these autoradiographs is shown in Fig. 1 a.

TABLE I
Plating Efficiencies of the Cell Lines Cultured in Dishes as Confluent Monolayers and in Aggregates

Cell line	Plating efficiency of aggregate-maintained cells			Plating efficiency of dish-maintained cells				Relative plating efficiency*			
	Cells seeded per 60-mm dish			Cells seeded per 60-mm dish				Cells seeded per 60-mm dish			
	250	500	1,000	250	500	1,000	250	500	1,000		
	<i>Colonies/100 cells</i>			<i>Colonies/100 cells</i>				<i>%</i>			
BALB/c 3T3	19.0	14.5	9.3 (14.3)‡	32.2	19.6	12.6 (21.4)	59	74	74	(69)	
SVT-2	19.0	18.4	24.4 (20.6)	27.2	20.0	12.2 (19.8)	90	92	155	(112)	
NIL B	22.4	20.8	18.0 (20.4)	40.4	35.8	22.7 (32.9)	55	58	80	(64)	

* $\frac{\text{No. colonies formed from aggregate-maintained cells}}{\text{No. colonies formed from dish-maintained cells}} \times 100$.

‡ Nos. in parentheses indicate mean values.




FIGURE 1 (a) Autoradiograph of a histological section of an aggregate of [^3H]thymidine-labeled BALB/c 3T3 cells and unlabeled SVT-2 cells. The aggregate was cultured for 2.5 days. While more than half of the cells visible in this section have silver grains over their nuclei (see Fig. 1 b), most of the grains are not in the focal plane of this photograph. This autoradiograph was exposed for a shorter time than usual in order to reveal as much nuclear detail as possible. The dark staining of the peripheral cell layer is common in these aggregates, although the cause is not known. The bar in this photograph represents $50\ \mu\text{m}$. Magnification, $\times 200$. (b) Camera lucida drawing of the nuclei of the cells in the aggregate section shown in Fig. 1 a. The [^3H]thymidine-labeled nuclei are indicated by filled ovals while unlabeled nuclei are indicated by open ovals. The presence or absence of silver grains was determined with a $100\times$ oil immersion objective. 10 grains located over the nucleus above background was the minimum used to identify a cell as being labeled. Sorting out is nearly complete in this experiment; only a few unlabeled cells remain in the center of the aggregate, while the outer rim is virtually devoid of labeled cells. Not all of the nuclei indicated in this figure are visible in Fig. 1 a, since some are above or below the focal plane of the photograph.

A zero-time control was also performed for this and all subsequent experiments. In each case, aggregates fixed immediately after cutting were found to be random mixtures of the two cell types. After 2.5 days of culture the aggregates had formed two concentric spheres, with the SVT-2 cells on the outside and the BALB/c 3T3 cells internally. This experiment was performed four separate times, each in reciprocal combination for a total of eight separate experiments. The results of these eight experiments indicated that SVT-2 cells sorted out externally to BALB/c 3T3 cells.

Sorting out was usually but not always complete after 2 days, although the pattern was always clear. In those experiments carried out for 4.5 days, however, segregation was virtually complete, i.e., a complete shell of SVT-2 cells, containing no 3T3 cells, surrounded an inner core of BALB/c 3T3 cells. These experiments, as well as the following sorting-out experiments, are summarized in Table II. The reciprocal combination of Figs. 1 a and b is shown in Figs. 2 a and b. In these eight experiments, the ratio of BALB/c 3T3 to SVT-2 cells was either 1:2, 1:1, or 2:1. No differences in

TABLE II
Sorting out of Binary Combinations of BALB/c 3T3, SVT-2, con A, NIL B, and Swiss 3T3 Cells

Cell combination	No. of exp	Culture period	Result
		<i>days</i>	
BALB/c 3T3 + SVT-2	8	0	No sorting
	5	2.5	SVT-2 external to BALB/c 3T3
	4	4.5	SVT-2 external to BALB/c 3T3
	1	2.5*	SVT-2 external to BALB/c 3T3
BALB/c 3T3 + BALB/c 3T3	2	0	No sorting
	2	2.5	No sorting
SVT-2 + SVT-2	2	0	No sorting
	2	2.5	No sorting
SVT-2 + con A	4	0	No sorting
	4	2.5	No sorting
BALB/c 3T3 + con A	3	0	No sorting
	3	2.5	Con A external to BALB/c 3T3‡
BALB/c 3T3 + Swiss 3T3	4	0	No sorting
	4	2.5	No sorting
BALB/c 3T3 + NIL B	4	0	No sorting
	4	2.5	NIL B external and internal to BALB/c 3T3§
SVT-2 + NIL B	4	0	No sorting
	4	2.5	SVT-2 external to NIL B
Con A + NIL B	2	0	No sorting
	2	2.5	No sorting
Swiss 3T3 + NIL B	4	0	No sorting
	4	2.5	NIL B external and internal to Swiss 3T3§

* In this experiment the BALB/c 3T3 cells were taken from confluent culture. In all other experiments in this table, cells were taken from subconfluent culture.

‡ con A was external to BALB/c 3T3 in two of three experiments. In the third experiment no sorting out was observed.

§ Trilayered spheres with NIL B cells external, 3T3 cells in the middle, and more NIL B cells internal. See text for details.

sorting-out behavior were observed as this ratio was varied.

In another pair of experiments, BALB/c 3T3 cells taken from either subconfluent (as the experi-

ments above) or confluent cultures sorted out internally to SVT-2 cells after 2.5 days. As a control, BALB/c 3T3 cells labeled with [³H]thymidine were mixed with unlabeled

BALB/c 3T3 cells. No sorting out was seen after 2.5 days. Similarly, [³H]thymidine-labeled SVT-2 cells did not sort out from unlabeled SVT-2 cells in control experiments.

When SVT-2 cells were mixed with con A revertant cells (variants of SVT-2 which have regained contact inhibition of growth), no sorting out was observed. Aggregates fixed 2.5 days after seeding showed the same random mixture of labeled and unlabeled cells observed in aggregates fixed immediately after mixing. The criteria for lack of sorting out were the following: (a) examination of the outer four cell layers in the aggregate showed the same percentage of labeled versus unlabeled cells as in the aggregate interior; (b) the

outermost layer of cells contained both labeled and unlabeled cells; and (c) no discernible clumping of either labeled or unlabeled cells was seen in the interior ("patchy sorting" as described by Wiseman [34]).

When BALB/c 3T3 cells and con A revertant cells were mixed together, sorting out was observed in two of three experiments. In these two cases a tendency for con A cells to be peripheral to BALB/c 3T3 cells was observed, although sorting out was incomplete. One of these aggregate sections is shown in Figs. 3 a and b. The word "tendency" is used to mean that the distribution of labeled and unlabeled cells was not random; but neither was there a complete shell of con A cells




FIGURE 2 (a) Autoradiograph of a histological section of a portion of an aggregate of unlabeled BALB/c 3T3 cells and [³H]thymidine-labeled SVT-2 cells. The aggregate was cultured for 2.5 days. This experiment is the reciprocal combination of the one illustrated in Fig. 1. The outer few cell layers consist of labeled SVT-2 cells, while the interior consists of unlabeled BALB/c 3T3 cells. The bar represents 20 μ m. Magnification, \times 525. (b) The same field as Fig. 2 a, but photographed with phase contrast. The bar in this photograph represents 20 μ m. Magnification, \times 525.




FIGURE 3 (a) Autoradiograph of a histological section of an aggregate of [^3H]thymidine-labeled con A revertant cells and unlabeled BALB/c 3T3 cells. The aggregate was cultured for 2.5 days. Note the inner core of unlabeled BALB/c 3T3 cells. The bar in this photograph represents 50 μm . Magnification, $\times 200$. (b) Camera lucida drawing showing the nuclei of the cells in the histological section shown in Fig. 3 a. Note that sorting out is incomplete, since both con A cells and BALB/c 3T3 cells are found surrounding a small inner core of BALB/c 3T3 cells.

surrounding the BALB/c 3T3 cells. Instead, most of the cells in the outer few cell layers were con A cells, while most of the internal cells were BALB/c 3T3. In the third experiment in which this combination of cell lines was used, no discernible segregation was seen. Mixtures of BALB/c 3T3 and Swiss 3T3 cells in four experiments remained randomly mixed after 2.5 days of culture.

Sorting Out of Mouse-Hamster Cell Line Mixtures

Hamster NIL B cells were mixed with mouse BALB/c 3T3 cells in four experiments and were found to sort out. Instead of a simple two-layered sphere-within-a-sphere, however, a more complex pattern was observed. Some sections (the smallest ones) consisted entirely of NIL B cells. The next larger sections consisted of an outer rim of NIL B

cells surrounding an inner core of 3T3 cells. In about 25% of the histological sections (nearly always the largest ones), a three-layered structure was observed, with NIL B cells surrounding BALB/c 3T3 cells and with a small clump of NIL B cells in the center. One of these sections is shown in Figs. 4 a and b. It appears, then, that NIL B cells sort out both externally and internally to 3T3 cells. The three types of sections observed represent, from smallest to largest sections, tangential, partially tangential, and full-diameter sections through a three-layered sphere. This was confirmed in a few aggregates followed in serial sections. NIL B cells were also mixed with SVT-2 cells in four experiments. Although only a little sorting out was observed, a tendency for the SVT-2 cells to be external to the NIL B cells was noted. No trilayered structures, such as were seen in NIL B-BALB/c 3T3 mixtures, were observed.

Mixtures of NIL B and con A revertant cells also showed little or no sorting out after 2.5 days of culture. Mixtures of NIL B cells and Swiss 3T3 cells sorted out in exactly the same way as the NIL B and BALB/c 3T3 mixtures, i.e., an outer rim of NIL B cells surrounded an inner shell of Swiss 3T3 cells which in turn surrounded a small core of NIL B cells.

DISCUSSION

Previous studies of Gershman and Drumm (18) have indicated that cells of these established lines

can be maintained in aggregates in a healthy state. These data include: (a) histological criteria; (b) nearly quantitative recovery of cells from aggregates; and (c) comparable levels of [³H]thymidine incorporation into aggregate-maintained cells and confluent monolayer-maintained cells. In this report another criterion of cell viability, plating efficiency, indicates that cells in aggregates are about as viable as cells in confluent monolayers.

Probably the most significant result of the sorting out experiments is the finding that SVT-2 cells sort out externally to BALB/c 3T3 cells.




FIGURE 4 (a) Autoradiograph of a histological section of an aggregate of [³H]thymidine-labeled NIL B cells and unlabeled BALB/c 3T3 cells. The autoradiograph was purposely overexposed and then stained very lightly for photographic purposes. The unlabeled BALB/c 3T3 cells are consequently nearly invisible. The bar in this photograph represents 50 μ m. Magnification, \times 200. (b) Camera lucida drawing of the autoradiograph in Fig. 4 a, showing the location of the labeled NIL B nuclei (filled ovals) and the unlabeled BALB/c 3T3 nuclei (open ovals). Note that the cells are densely packed and that the aggregate profile is smooth and regular. The NIL B cells have sorted out into two populations, one internal to the unlabeled BALB/c 3T3 cells, and one external to the 3T3 cells. Sorting out is nearly complete in this aggregate, especially between the outer NIL B population and the BALB/c 3T3 cells. The cells at the surface of the aggregate appear to be packed somewhat more densely than the interior cells. This is a common feature of these aggregates which can also be seen in Fig. 1.

Interpreted according to the differential adhesiveness hypothesis of Steinberg (28), this means that SVT-2 cells are less adhesive than BALB/c 3T3 cells. This interpretation is consistent with the notion (first set forth by Coman [9]) that malignant cells are less adhesive than their nonmalignant counterparts, and also with the concept of Carter (6, 7) that cells with decreased adhesiveness are more mobile. Increased mobility of SVT-2 cells compared to BALB/c 3T3 cells has been previously observed in aggregates by Gershman and Drumm (18).

The results of sorting-out experiments between BALB/c 3T3, SVT-2, and con A revertant cells, if interpreted according to Steinberg's (28) hypothesis, suggest that these lines can be ranked in order of decreasing adhesiveness: BALB/c 3T3 > con A > SVT-2. The failure of con A cells to completely segregate from SVT-2 cells and their incomplete segregation from BALB/c 3T3 cells would then be seen as an indication that the differences in adhesiveness among these three lines is not very great, even between the extremes (BALB/c 3T3 and SVT-2). Between the intermediate cell type and one of the extremes, then, the differences are not great enough to permit clear sorting to occur during the 2.5-day culture period. It is also of interest that the three nontransformed contact-inhibited cell lines (BALB/c 3T3, Swiss 3T3, and NIL B) as well as the contact-inhibited phenotypic revertant line (con A) examined in this study showed no consistent pattern of sorting out.

NIL B cells sort out into two populations when mixed with either BALB/c 3T3 or Swiss 3T3 cells. One population sorts internally to 3T3 cells, and one moves externally. A number of explanations for these two subpopulations of NIL B cells can be envisioned including: two genetically different sublines, two subpopulations frozen in different stages of the cell cycle, normal vs. altered or damaged cells, etc. Mixtures of NIL B cells with either SVT-2 cells or con A revertant cells showed modest or no sorting out. This probably indicates that NIL B, con A, and SVT-2 cells are very similar in their sorting-out properties. The failure to sort out, or incomplete sorting out, is typical of a number of the combinations studied here. On the other hand, cells of different embryonic tissues, either mouse, chick, amphibian, or human, invariably sort out from each other rapidly and completely. This suggests that cells of the lines studied here may be more similar to each other in their surface properties than are embryonic tissue cells.

On the basis of the data presented here, a tentative ranking can be established for these five cell lines according to sorting-out properties. Because certain combinations fail to sort out even after several days of culture, it is necessary to assume overlapping ranges of values. This may imply either that cells of these established lines are more heterogeneous with regard to sorting-out properties than embryonic tissue cells, or that their sorting-out properties are very similar. A tentative scheme presenting these data graphically is shown in Fig. 5. It should be emphasized that because of the apparent similarity between these cells this figure is presented as a working hypothesis rather than a firm conclusion.

Cultured neoplastic cells have previously been examined in suspension culture, either alone, in co-culture with nonmalignant tissues (e.g., 15, 17, 19, 22, 24, 25), or in related culture systems incorporating solid substrata (3, 16, 20, 35). Moscona (22), in the only previously published report of sorting out of established cells, reports that




FIGURE 5. Proposed relationship between the adhesiveness of five established cell lines based on their sorting-out behavior. The position of the rectangles representing the cell line indicates their sorting-out behavior relative to the other lines. Two nonoverlapping bars indicate clear sorting out, with the cell line represented by the leftmost bar being external. An example of this is the SVT-2-BALB/c 3T3 combination. Near overlap or partial overlap indicates combinations in which sorting out is not complete, or not consistent, for example, the con A and BALB/c 3T3 combination. Complete failure to sort out, e.g. SVT-2 and con A, is indicated by the overlap of two rectangles. The presence of two rectangles for NIL B cells represents the observation that some NIL B cells sort out internally to 3T3 cells while others sort out externally. The relative horizontal lengths of the bars indicate the hypothesized adhesive heterogeneity of the cell lines.

mouse melanoma S91 cells sort out externally to chick embryonic cartilage and internally to chick embryonic liver. By use of an isogenic family of cell lines (BALB/c 3T3:SVT-2:con A), we have been able to extend these comparisons to more similar combinations of cells than previously attempted.

CONCLUSIONS

Mouse BALB/c 3T3 cells sort out internally to SVT-2 cells. This indicates that viral transformation of these cells results in cell surface alterations that allow the cells to recognize each other in mixed cultures. The mechanism for this recognition may operate through a change in the adhesiveness of the cells, with the SVT-2 cells being less adhesive than the BALB/c 3T3. Swiss 3T3 cells and a con A-selected revertant of SVT-2 (con A cells) fail to sort out from BALB/c 3T3 cells. On the other hand, cells of another contact-inhibited fibroblast line, NIL B, do sort out from BALB/c 3T3 cells. This suggests that the recognition phenomenon, acting during sorting out, is not specific to contact-inhibited cells.

The authors want to thank Josefina Buniel for her technical assistance.

This research was supported by American Cancer Society Grant no. IN-57-L to Case Western Reserve University and by a grant from the Cuyahoga County Unit of the American Cancer Society to Howard Gershman, and by National Cancer Institute grant no. 1-RO1-CA13513 to Lloyd Culp.

Received for publication 19 June 1975, and in revised form 26 September 1975.

REFERENCES

1. AARONSON, S. A., and G. J. TODARO. 1968. Development of 3T3-like lines from BALB/c mouse embryo cultures: transformation susceptibility to SV40. *J. Cell Physiol.* **72**:141-148.
2. ABERCROMBIE, M., J. E. M. HEAYSMAN, and H. M. KARTHAUSER. 1957. Social behavior of cells in tissue culture. III. Mutual influence of sarcoma cells and fibroblasts. *Exp. Cell Res.* **13**:276-291.
3. BARSKI, G., and J. BELEHRADEK. 1965. Étude microcinématographique du mécanisme d'invasion cancéreuse en cultures de tissu normal associé aux cellules malignes. *Exp. Cell Res.* **37**:464-480.
4. BELL, P. B. 1972. Criss-crossing, contact inhibition, and cell movement in cultures of normal and transformed 3T3 cells. *J. Cell Biol.* **55**(2, Pt. 2):16 a. (Abstr.)
5. BURDICK, M. L. 1970. Cell sorting out according to species in aggregates containing mouse and chick embryonic limb mesoblast cells. *J. Exp. Zool.* **175**:357-368.
6. CARTER, S. B. 1965. Principles of cell motility: the direction of cell movement and cancer invasion. *Nature (Lond.)* **208**:1183-1187.
7. CARTER, S. B. 1967. Haptotaxis and the mechanism of cell motility. *Nature (Lond.)* **213**:256-260.
8. CASSIMAN, J. J., and M. R. BERNFIELD. 1974. Morphogenetic properties of human embryonic cells: aggregation of dissociated cells and histogenesis in cultured aggregates. *Pediatr. Res.* **8**:184-192.
9. COMAN, D. R. 1944. Decreased mutual adhesiveness, a property of cells from squamous cell carcinomas. *Cancer Res.* **4**:625-629.
10. CULP, L. 1974. Substrate-attached glycoproteins mediating adhesion of normal and virus-transformed mouse fibroblasts. *J. Cell Biol.* **63**:71-83.
11. CULP, L. A., and P. H. BLACK. 1972. Contact-inhibited revertant cell lines isolated from Simian virus 40-transformed cells. III. Concanavalin A-selected revertant cells. *J. Virol.* **9**:611-620.
12. CULP, L. A., W. J. GRIMES, and P. H. BLACK. 1971. Contact-inhibited revertant cell lines isolated from SV40-transformed cells. I. Biologic, virologic, and chemical properties. *J. Cell Biol.* **50**:682-690.
13. DELONG, G. R., and R. L. SIDMAN. 1970. Alignment defect of reaggregating cells in cultures of developing brains of reeler mutant mice. *Dev. Biol.* **22**:584-600.
14. DIAMOND, L. 1967. Two spontaneously transformed cell lines derived from the same hamster embryo culture. *Int. J. Cancer.* **2**:143-152.
15. DODSON, E. O. 1966. Aggregation of tumor cells. *Nature (Lond.)* **209**:40-44.
16. EASTY, G. C., and D. M. EASTY. 1963. An organ culture system for the examination of tumor invasion. *Nature (Lond.)* **199**:1104-1105.
17. FOLKMAN, J., M. HOCHBERG, and D. KNIGHTON. 1974. Self-regulation of growth in three dimensions: the role of surface area limitation. *Cold Spring Harbor Conf. Cell Proliferation.* **1**:833-842.
18. GERSHMAN, H., and J. DRUMM. 1975. Mobility of normal and virus-transformed cells in cellular aggregates. *J. Cell Biol.* **67**:419-435.
19. HALPERN, B., B. PEJSACHOWICZ, H. L. LEBVRE, and G. BARSKI. 1966. Differences in patterns of aggregation of malignant and non-malignant mammalian cells. *Nature (Lond.)* **209**:157-159.
20. HEAYSMAN, J. E. M., and S. M. PEGRUM. 1973. Early contacts between normal fibroblasts and mouse sarcoma cells. *Exp. Cell Res.* **78**:479-481.
21. LUDFORD, R. J. 1932. Differences in the growth of transplantable tumors in plasma and serum culture media. *Proc. R. Soc. Lond. B. Biol. Sci.* **112**:250-263.
22. MOSCONA, A. 1957. The development *in vitro* of chimeric aggregates of dissociated embryonic chick and mouse cells. *Proc. Natl. Acad. Sci. U. S. A.* **43**:184-194.
23. MOSCONA, A., and H. MOSCONA. 1952. The dis-

- sociation and aggregation of cells from organ rudiments in the early chick embryo. *J. Anat.* **86**:287-301.
24. MOSKOWITZ, M. 1963. Aggregation of cultured mammalian cells. *Nature (Lond.)*, **200**:854-856.
 25. OPPENHEIMER, S. B., M. EDIDIN, C. W. ORR, and S. ROSEMAN. 1969. An L-glutamine requirement for intercellular adhesion. *Proc. Natl. Acad. Sci. U. S. A.* **63**:1395-1402.
 26. RUBIN, H. 1966. Fact and theory about the cell surface in carcinogenesis. In *Major Problems in Developmental Biology*. M. Locke, editor. Academic Press, Inc., New York. 315-337.
 27. STEINBERG, M. S. 1964. The problem of adhesive selectivity in cellular interactions. In *Cellular Membranes in Development*. M. Locke, editor. Academic Press, Inc., New York. 321-366.
 28. STEINBERG, M. S. 1970. Does differential adhesion govern self-assembly processes in histogenesis? Equilibrium configurations and the emergence of a hierarchy among populations of embryonic cells. *J. Exp. Zool.* **173**:395-434.
 29. STOKER, M. 1964. Regulation of growth and orientation in hamster cells transformed by polyoma virus. *Virology*. **24**:165-174.
 30. SUBAK-SHARPE, H., R. R. BURK, and J. D. PITTS. 1969. Metabolic co-operation between biochemically marked mammalian cells in tissue culture. *J. Cell Sci.* **4**:353-367.
 31. TOWNES, P. L., and J. HOLTFRETER. 1955. Directed movements and selective adhesion of embryonic amphibian cells. *J. Exp. Zool.* **128**:53-120.
 32. VASILIEV, J. M., and I. M. GELFAND. 1972. Interactions of normal and neoplastic fibroblasts with the substratum. *Ciba Found. Symp.* **14**:311-329.
 33. WESTON, J. A., and S. A. ROTH. 1969. Contact inhibition: behavioral manifestations of cellular adhesive properties *in vitro*. In *Cellular Recognition*. Smith and Good, editors. Appleton-Century-Crofts, New York. 29-37.
 34. WISEMAN, L. L. 1970. Experimental modulation of the assembly properties of embryonic cell populations. Ph.D. Thesis. Princeton University, Princeton, N.J.
 35. WOLFF, E., and N. SCHNEIDER. 1957. La culture d'un sarcome de souris sur des organes de poulet explantes *in vitro*. *Arch. Anat. Microsc. Morphol. Exp.* **46**:173-197.