

Differential and Synergistic Actions of Nerve Growth Factor and Cyclic AMP in PC12 Cells

PETER W. GUNNING, GARY E. LANDRETH, MARK A. BOTHWELL, and ERIC M. SHOOTER
*Department of Neurobiology, Stanford University School of Medicine, Stanford, California 94305; and
Department of Biochemistry, Princeton University, Princeton, New Jersey 08544*

ABSTRACT When a clonal line of rat pheochromocytoma (PC12) was exposed to β -nerve growth factor (β NGF), N^6, O^2 -dibutyryl adenosine 3':5' cyclic monophosphate (Bt_2cAMP), or a combination of the two, 10, 26, or 70% of the cell clumps, respectively, displayed neurites after 1 d. Increases in the cellular RNA concentration were also found to be additive or greater when both agents were present. Neurites induced by Bt_2cAMP alone were not maintained after replacement with β NGF. The degree of potentiated neurite outgrowth was a function of the time of simultaneous exposure to both agents. The initiation of neurite outgrowth in the presence of Bt_2cAMP was independent of RNA synthesis, in contrast to that induced by β NGF alone. We conclude that β NGF-induced initiation of morphological differentiation of these cells is not mediated by a cAMP-dependent mechanism. Consideration of Bt_2cAMP effects upon other cell lines suggests that Bt_2cAMP causes a rapid, but unstable, reorganization of the PC12 cytoskeleton, resulting in the initiation of neurite outgrowth from these cells. In contrast, β NGF alone achieves a more stable cytoskeleton reorganization by an RNA synthesis-dependent mechanism.

Nerve growth factor (NGF) is a polypeptide hormone that is involved in the development and maintenance of sympathetic and certain sensory neurons (12). Recently, a pheochromocytoma clonal cell line, PC12, showing similar NGF-responsive properties, has been isolated (5). When PC12 cells are exposed to NGF, they extend neuritic processes and cease cell division within 1 wk (5). NGF also prevents the death of these cells and stimulates their differentiation in serum-free medium (4). The induction of differentiation is RNA synthesis dependent (1). However, the regeneration of neuritic processes from cells that have been exposed to NGF for a week does not require RNA synthesis (1). These results have led to the suggestion that the synthesis of new RNA species and possibly proteins is required for the initiation of neurite outgrowth. However, once these products are made and initiation has occurred, regeneration of processes over a 24-h period can occur independent of further RNA synthesis (1). The nature and role of these gene products are unknown.

Two suggestions for the mechanisms responsible for the initiation of neurite outgrowth have been made. It has been proposed that cAMP is the second messenger in the response of PC12 cells to NGF (16–18). This proposal is based on the observations that NGF increases the cAMP content of PC12

cells during the first minutes of exposure (18); that cAMP initiates neurite outgrowth from these cells (16, 17), and that two analogs of cAMP modulate the synthesis of the same group of proteins in PC12 cells, as does NGF (2). The suggestion for an alternative mechanism comes from the finding of specific NGF receptors on the nuclear membrane of PC12 cells, and of a time- and temperature-dependent translocation of NGF (25) from its surface receptors (9, 25) to those on the nucleus (25). The association of NGF with the nucleus is necessary for both the initiation and the regeneration of neurite outgrowth from PC12 cells.¹

We have further examined the effect of N^6, O^2 -dibutyryl adenosine 3':5' cyclic monophosphate (Bt_2cAMP) upon PC12 cells. Studies at both a biochemical and morphological level indicate that NGF and cAMP act by separate mechanisms and together produce synergistic effects. In particular, initiation of neurite outgrowth from these cells in the presence of both agents is RNA- and protein-synthesis independent. It is suggested that Bt_2cAMP causes a rapid reorganization of the cytoskeleton that NGF alone achieves by an RNA synthesis-dependent mechanism.

¹ Yankner, B. A., and E. M. Shooter. Manuscript submitted.

MATERIALS AND METHODS

Cell Culture

The PC12 clone of a rat pheochromocytoma was obtained from David Schubert (Salk Institute, La Jolla, Calif.). The cells were grown in Dulbecco's modified Eagle's medium (Grand Island Biological Co., Grand Island, N. Y.), containing 10% fetal calf serum and 5% horse serum on polystyrene tissue culture dishes (NUNC, Irvine Scientific, Calif.). Cultures were maintained in a water-saturated atmosphere of 88% air/12% CO₂. Experiments were performed by replating subconfluent cells at a density of $1-2 \times 10^4$ cells/cm², and the cells were allowed to reach log phase growth (2 d). At that time, fresh medium was added, together with 5 ng/ml β NGF (prepared by the method of Smith et al. [21]) or 1 mM Bt₂cAMP (Sigma Chemical Co., St. Louis, Mo.), unless otherwise indicated. In some experiments, the fresh medium added contained 1 mg/ml bovine serum albumin (BSA, Pentex, Miles Biochemicals, Ind.), with or without fetal calf and horse serum. It has been found that, for all parameters measured in this study, maximum stimulation is attained at concentrations equal to or below 5 ng/ml for β NGF and 1 mM for Bt₂cAMP.²

The morphological differentiation of the cells was assayed by determining the percentage of cell clumps with neuritic processes longer than one cell body diameter and displaying a growth cone at their tip. Clumps, rather than single cells, were scored because addition of β NGF leads to extensive clumping of these cells (see reference 3). At least 400 clumps were examined in randomly chosen fields on each plate. All experiments were performed at least three times with similar results.

Inhibition of RNA and Protein Synthesis

Actinomycin-D (Calbiochem-Behring Corp., American Hoechst Corp., San Diego, Calif.) and cycloheximide (Sigma Chemical Co.) were added to cultures at 0.5 μ M and 10 μ g/ml, respectively. 1 h later, β NGF and Bt₂cAMP were added along with [³H]uridine (1 μ Ci/ml) or [³H]proline (2 μ Ci/ml) (Amersham Corp., Arlington Heights, Ill., sp act, 27 and 18 Ci/mmol, respectively). After 3 h, five randomly chosen fields on each plate were photographed, and the cells were immediately harvested and assayed for RNA and protein synthesis (1). Uptake into the acid-soluble compartment was unaffected by the drugs. Incorporation of [³H]uridine into RNA was inhibited >80% by 0.5 μ M actinomycin-D. Cycloheximide at 10 μ g/ml inhibited [³H]proline incorporation into protein by >97%. Uptake and incorporation were measured relative to control sister cultures that received neither drug.

DNA, RNA, and Protein Determinations

The cells were mechanically dislodged from the dishes by trituration and collected by centrifugation at 750 g_{av} for 5 min. Acid-soluble material was removed by extracting the cell pellet three times with 0.5 M perchloric acid (PCA) at 4°C. The acid-insoluble material was suspended in 1 M NaOH and incubated at 37°C for 18 h. A sample of the solution was taken for protein determination (11), using bovine serum albumin (Sigma Chemical Co.) as a standard. Concentrated PCA was added to the remainder of the solution, to a final concentration of 0.5 M PCA and left at 4°C for 1 h. The resulting precipitate was collected at 10,000 g_{av} for 10 min and washed once with 0.5 M PCA. The two supernates were combined for RNA determination (22). The pellet was extracted twice with 0.5 M PCA at 80°C for 30 min, and the combined supernates were taken for DNA determination (22). The ratios of RNA:DNA and protein:DNA were taken as a measure of the cellular RNA and protein concentrations.

RESULTS

Potentiation of Morphological Differentiation

Exposure of PC12 cells to β NGF for 24 h produces little morphological change except for a tendency to flatten out on the polystyrene substratum (Fig. 1A, and B). The cells respond to Bt₂cAMP by extending short neurites (Fig. 1C). However, β NGF in combination with Bt₂cAMP produces a morphological effect far greater than the individual effects of these agents (Fig. 1D). The combined treatment produces a morphological differentiation normally seen only after 4–5 d of exposure to β NGF.

This potentiation of neurite outgrowth depends upon the β NGF concentration. In the presence of Bt₂cAMP only, ~25% of the cell clumps possess neurites after 24 h. Addition of β NGF at increasing concentrations raises this to a maximum of 70% at 0.5–1.0 ng/ml β NGF (Fig. 2). The dose response of this potentiation by β NGF is identical to the dose-response curve for β NGF-promoted neurite regeneration from differentiated PC12 cells (3, 4).

The specificity of the Bt₂cAMP effect was evaluated by repeating the experiments described above but 2 mM Na butyrate, 1 mM Bt₂cAMP, or 1 mM AMP was substituted for Bt₂cAMP. Neurite outgrowth was not observed when these agents alone were used, and none of them potentiated the β NGF-induced response. These results indicate a specificity for the cAMP moiety of Bt₂cAMP and suggest the involvement of a cAMP-dependent mechanism. This is supported by the observation that cholera toxin, which increases cAMP levels within mammalian cells (10), produced a neurite outgrowth and β NGF potentiation similar to that observed using Bt₂cAMP (Fig. 3).

Effects on Cellular RNA Concentration

The PC12 cellular RNA concentration is increased within 24 h of exposure to β NGF, the increase being half the maximum at ~0.25 ng/ml β NGF, and the maximum between 2.5 and 5.0 ng/ml.² We, therefore, examined the effects of Bt₂cAMP on cellular RNA levels with and without β NGF. After 24 h of exposure to β NGF, in medium containing either serum or BSA, the RNA concentration increased 20% over controls (Table I). Treatment with Bt₂cAMP also increased the RNA concentration in both media. This effect was maximum at 3×10^{-4} to 10^{-3} M Bt₂cAMP. However, in medium containing serum, Bt₂cAMP produced a significantly greater stimulation than β NGF, whereas the converse was true in medium containing only BSA (Table I). The poor effect of Bt₂cAMP in medium plus BSA was due to a lack of serum rather than inhibition by BSA, because the addition of serum to medium plus BSA restored the full Bt₂cAMP stimulation (Table I). Thus, although the magnitude of β NGF stimulation is independent of the culture medium, that of Bt₂cAMP is dependent, at least in part, upon the presence of serum.

Exposure of the cells to both Bt₂cAMP and β NGF produced an increase in RNA equal to the sum of the increases separately induced by each agent when examined in medium plus serum. Using medium plus BSA, the combination produced a greater than additive increase (Table I). This result is inconsistent with a common mechanism of action for both agents, because maximally stimulating concentrations of both agents together produce an additive, or greater than additive, effect. Similar results were obtained when the PC12 cellular protein concentrations were measured (not shown).

Both the morphological and biochemical data suggest that the actions of β NGF and Bt₂cAMP are complementary and, in some situations, synergistic.

Potentiation Requires Simultaneous Presence of β NGF and Bt₂cAMP

When PC12 cells are exposed to β NGF for several days and then replated with fresh medium plus β NGF, they regenerate their neurites within 24 h (3); such cells are said to be "primed." In contrast, when cells exposed to Bt₂cAMP for several days are replated in the presence of β NGF, they do not regenerate

² Gunning, P. W., G. E. Landreth, P. Layer, M. Ignatius, and E. M. Shooter. *J. Neurosci.* In press.




FIGURE 1 Potentiation of neurite outgrowth by NGF plus Bt₂cAMP. Long phase PC12 cells were exposed to β NGF (5 ng/ml), Bt₂cAMP (1 mM), or both for 24 h, and then photographed. A, control; B, β NGF; C, Bt₂cAMP; D, β NGF plus Bt₂cAMP. Bar, 50 μ m.




FIGURE 2 Relationship of β NGF concentration to potentiation of neurite outgrowth. Log phase PC12 cells were exposed to 1 mM Bt₂cAMP plus various concentrations of β NGF for 24 h. The resulting neurite outgrowth was determined.

their neurites rapidly (G. E. Landreth, P. W. Gunning, and E. M. Shooter, unpublished observations). This suggests that the Bt₂cAMP-treated cells, although displaying significant neurite initiation, are not "primed" in a manner analogous to that produced by β NGF (4). Therefore, we investigated the stability of the Bt₂cAMP effect by pretreating the cells with Bt₂cAMP, and then adding β NGF in the continued presence or absence of Bt₂cAMP.

If cells were treated with Bt₂cAMP for 8 h, followed by exposure to β NGF for 16 h, virtually no neurite outgrowth was observed (Table II). However, the presence of β NGF plus Bt₂cAMP for the final 16 h resulted in maximum neurite outgrowth. An identical result was obtained when β NGF was present for the first 8 h, followed by both agents for the remainder of the 24-h period (Table II). Repeating this experiment with a 20-h exposure to Bt₂cAMP, followed by 4 h with β NGF alone, resulted in virtually no neurite outgrowth (Table II). After a 20-h treatment with only Bt₂cAMP, ~25% of the cell clumps had neurites. It is therefore obvious that β NGF alone cannot maintain neurites induced by Bt₂cAMP. In contrast, if β NGF and Bt₂cAMP are present for the final 4 h, a considerable potentiation of neurite outgrowth is observed. Pre-exposure to β NGF for 20 h, followed by the simultaneous presence of β NGF and Bt₂cAMP for the final 4 h, gave a

TABLE I
RNA Concentration within PC12 Cells after Exposure to β NGF and Bt₂cAMP

Additions	Medium and serum		Medium and BSA	
	ng/ml			
	5	50	5	50
β NGF				
β NGF	1.21 ^a ± 0.05 (15)	1.22 ^a ± 0.05 (16)	1.19 ^d ± 0.04 (6)	1.23 ^{d*} ± 0.05 (6)
Bt ₂ cAMP		1.36 ^b ± 0.04 (9)		1.14 ^e ± 0.03 (10)
β NGF + Bt ₂ cAMP	1.58 ^c ± 0.06 (6)	1.53 ^c ± 0.08 (3)	1.48 ^f ± 0.04 (6)	1.50 ^f ± 0.03 (6)
Bt ₂ cAMP + BSA		1.34 ± 0.01 (2)		
β NGF + Bt ₂ cAMP + BSA		1.51 ± 0.04 (2)		

Fresh medium plus serum or BSA was added to log phase PC12 cells, together with Bt₂cAMP (1 mM), β NGF (5 or 50 ng/ml), and BSA (1 mg/ml) in the appropriate plates. After 24 h, the cells were harvested and the cellular RNA concentration was measured. The values are expressed relative to control sister cultures as mean ± SD. The number of individual cultures assayed is given in parentheses. The significance of differences observed was determined from Student's *t* test for two means. Values with the same superscript were not significantly different (NS). The salient *P* values obtained were: a vs. b, *P* < 0.00005; a vs. d, NS; d vs. e, *P* < 0.02; d* vs. e, *P* < 0.001; b vs. e, *P* < 0.00005; b vs. c, *P* < 0.00005; e vs. f, *P* < 0.00005; c vs. f, NS.




FIGURE 3 Potentiation of neurite outgrowth by cholera toxin. Log phase PC12 cells were exposed for 24 h to varying concentrations of cholera toxin with (●) and without (○) β NGF (50 ng/ml). The resulting neurite outgrowth was determined.

virtually identical potentiation (Table II).

The finding that neurite outgrowth potentiation requires the simultaneous, rather than sequential, presence of both agents suggests that β NGF and Bt₂cAMP act on independent, yet complementary, cellular mechanisms. It is also apparent that the effect of Bt₂cAMP is rapidly reversible upon removal of the agent from the medium.

Resistance of Neurite Initiation to Inhibitors of Transcription and Translation

The initiation of neurite outgrowth from naive PC12 cells by β NGF is almost completely inhibited by actinomycin-D at a level that inhibits RNA synthesis by ~50% (1). Therefore, we investigated whether the initiation of neurite outgrowth in the presence of β NGF plus Bt₂cAMP was similarly susceptible. Cells were pretreated with actinomycin-D for 1 h at doses

TABLE II
Effect of Sequential, Compared with Simultaneous, Presence of Bt₂cAMP and β NGF

	Time of exposure to agent		Percent cell clumps with neurites after 24 h
	Bt ₂ cAMP	β NGF	
	h		
Control	-	-	<1
β NGF	-	0-24	10
Bt ₂ cAMP	0-24	-	26
β NGF + Bt ₂ cAMP	0-24	0-24	70
Bt ₂ cAMP pulse	0-8	8-24	4
Bt ₂ cAMP pretreated	0-24	8-24	69
β NGF pretreated	8-24	0-24	68
Bt ₂ cAMP pulse	0-20	20-24	3
Bt ₂ cAMP pretreated	0-24	20-24	46
β NGF pretreated	20-24	0-24	43

Various additions and substitutions of 1 mM Bt₂cAMP and 5 ng/ml β NGF were made over a 24 h period to log phase cells in medium containing serum. After 24 h, the percentage of cell clumps displaying neurites was determined.

ranging from 1 nM to 10 μ M, and then β NGF and Bt₂cAMP were added. After 1 h, 10% (\pm 2%) of the cells displayed neurites independent of the presence of the inhibitor at any of these concentrations (not shown). In another experiment, the cells were exposed to cycloheximide or actinomycin-D for 1 h before the addition of β NGF plus Bt₂cAMP. Neurite outgrowth examined after 3 h was not inhibited under these conditions, whereas RNA synthesis was inhibited >80% by actinomycin-D, and protein synthesis was inhibited >97% by cycloheximide (Fig. 4). Thus, initiation of neurite outgrowth induced by β NGF plus Bt₂cAMP is independent of transcription and translation.

This conclusion is supported by recent experiments performed using scanning electron microscopy, rather than phase-contrast light microscopy, to evaluate neurite outgrowth. We noted that Bt₂cAMP plus β NGF caused a significant level of neurite outgrowth within 2 min of their addition to cells.³ Such a rapid response is inconsistent with any transcriptional or translational involvement.

Effects of Cytoskeletal and DNA Synthesis Inhibitors

DNA synthesis within PC12 cells is inhibited by addition of Bt₂cAMP (M. A. Bothwell, unpublished observation). To de-

³ Bothwell, M. A. Manuscript in preparation.






FIGURE 4 Effect of inhibition of RNA and protein synthesis upon β NGF plus Bt_2cAMP -induced neurite outgrowth. Log phase PC12 cells were pre-exposed to cycloheximide (10 $\mu g/ml$) or actinomycin-D (0.5 μM) for 1 h before the addition of β NGF (5 ng/ml), Bt_2cAMP (1 mM), and [3H]uridine or [3H]proline. After 3 h, representative fields were photographed and the cells immediately harvested. The micrographs show β NGF plus Bt_2cAMP -induced morphogenesis in the presence of no inhibitor (A), cycloheximide (B), and actinomycin-D (C). Bar, 50 μm .

termine whether inhibition of DNA synthesis could cause neurite outgrowth and a potentiation of β NGF-induced morphogenesis, cells were exposed to 1 μM cytosine arabinoside (Sigma Chemical Co.) in both the presence and absence of β NGF. Although DNA synthesis was inhibited by >99%, no neurite outgrowth was observed with the inhibitor alone, and no potentiation was observed after a 24-h coexposure with β NGF. In contrast, when 0.5 $\mu g/ml$ colchicine (Sigma Chemical Co.) or 4 μM cytochalasin B (Sigma Chemical Co.) was

added to the cells, along with β NGF plus Bt_2cAMP , no neurite outgrowth was observed at any time.

DISCUSSION

The results presented in this paper lead to the following conclusions: (a) Bt_2cAMP and β NGF have a synergistic effect upon the initiation of neurite outgrowth from PC12 cells; (b) Bt_2cAMP and β NGF have additive and synergistic effects upon the RNA levels within PC12 cells; (c) β NGF cannot

maintain Bt₂cAMP-induced morphogenesis; (d) the potentiation of neurite outgrowth by Bt₂cAMP plus βNGF requires the simultaneous presence of both agents; and (e) the induction of *de novo* neurite outgrowth by Bt₂cAMP plus βNGF is independent of ongoing transcription, in contrast to that induced by βNGF alone. Therefore, it is clear that Bt₂cAMP and βNGF have complementary effects on this cell line, together producing a greatly enhanced response over that observed with either agent alone, and that they act to initiate neurite outgrowth by different mechanisms. This is made particularly clear by the observation that neurite outgrowth can be initiated by Bt₂cAMP plus βNGF, but not by βNGF alone, under conditions where RNA synthesis is largely abolished. Thus, cAMP can overcome the transcription dependency for βNGF-induced neurite outgrowth in a manner analogous to the way in which βNGF can "prime" PC12 cells for transcription-independent neurite regeneration. Because the cAMP-mediated increase in RNA levels is largely dependent on the presence of serum in the medium, whereas the NGF-mediated increase is not, it is suggested that the mechanisms leading to elevation of cellular RNA by these two agents are also distinct.

Even if βNGF is added afterward, the neurites produced by Bt₂cAMP are unstable, in the sense that they are rapidly lost on removal of Bt₂cAMP. Thus, it is unlikely that a βNGF-induced, transient rise in cAMP levels can by itself lead to neurite initiation through a cAMP-dependent step originating with cAMP as a second messenger. Instead, rapidly reversible effects of cAMP on the morphology of PC12 cells are consistent with rapidly reversible morphological and biochemical effects of cAMP described in other systems (7, 13, 14). It should also be noted that Hatanaka et al. (6) failed to detect any transient increase in the cAMP content of PC12 cells after interaction with NGF, and that Landreth et al. (8), unlike Schubert et al. (17), found no significant change in calcium efflux from NGF-treated PC12 cells.

This conclusion is supported by the observation that if βNGF is removed from the culture medium by adding anti-βNGF antibody any time during the first 3–4 h of exposure, no neurite outgrowth occurs from PC12 cells (see footnote 1). This is the time that elapses before the relevant internalization of βNGF and its receptor occurs, initiating translocation to the nucleus. Nevertheless, it should be noted that long-term stimulation of a cAMP-dependent mechanism in PC12 cells can initiate neurite outgrowth (16–18, Fig. 3), and, in this respect, cAMP mimics the action of βNGF.

Three properties of the Bt₂cAMP-induced neurite outgrowth indicate the possible intracellular sites of action. First, the morphological response is prevented by treatment with colchicine and cytochalasin B. These two agents disrupt microtubule and microfilament organization, respectively, in a wide variety of cell types (15, 23). It is likely that the Bt₂cAMP effect requires the organization of both of these cytoskeletal elements. Second, the induction of neurite outgrowth by Bt₂cAMP in the presence and absence of βNGF is rapidly reversed upon removal of Bt₂cAMP from the medium. Similar rapidly reversible effects of Bt₂cAMP upon Chinese hamster ovary cell morphology have been shown to be related to induction of microtubule and microfilament organization (7, 13, 14). Finally, the induction of neurites in the presence of Bt₂cAMP is RNA and protein synthesis independent. Similarly, Bt₂cAMP-induced changes in cell morphology and organization of the cytoskeleton in other cell lines are also independent of RNA and protein synthesis (24). We therefore conclude that Bt₂cAMP probably

causes a rapidly reversible alteration in the PC12 cell cytoskeleton that results in neurite outgrowth.

The detailed mechanism by which Bt₂cAMP alters cytoskeleton organization is unknown, but it has been reported that a microtubule-associated protein (MAP₂), which may regulate microtubule assembly, is phosphorylated by a cAMP-dependent mechanism (19, 20). It appears that the transcription-dependent event in βNGF induction of neurite outgrowth involves the production of a gene product that regulates, in a more stable manner than Bt₂cAMP, the organization of microtubules and perhaps microfilaments.

We thank P. Letourneau, R. Gunzales, and P. Layer for helpful discussions, and M. Ignatius for preparation of the micrographs.

This work was supported by grants from the National Institutes of Health (NS 04270 and NS 14632) and from the American Cancer Society (BC 325). P. W. Gunning was supported by a California Division Cancer Society Junior Fellowship J-466, and G. E. Landreth by a National Research Service Award.

Received for publication 7 October 1980, and in revised form 22 December 1980.

REFERENCES

- Burstein, D. E., and L. A. Greene. 1978. Evidence for RNA synthesis-dependent and independent pathways in stimulation of neurite outgrowth by nerve growth factor. *Proc. Natl. Acad. Sci. U. S. A.* 75:6059–6063.
- Garrels, J. E., and D. Schubert. 1979. Modulation of protein synthesis by nerve growth factor. *J. Biol. Chem.* 254:7978–7985.
- Greene, L. A. 1977. A quantitative bioassay for nerve growth factor (NGF) activity employing a clonal pheochromocytoma cell line. *Brain Res.* 133:350–353.
- Greene, L. A. 1978. Nerve growth factor prevents the death and stimulates the neuronal differentiation of clonal PC12 pheochromocytoma cells in serum-free medium. *J. Cell Biol.* 78:747–755.
- Greene, L. A., and A. S. Tischler. 1976. Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc. Natl. Acad. Sci. U. S. A.* 72:2424–2428.
- Hatanaka, H., U. Otten, and H. Thoenen. 1978. Nerve growth factor-mediated selective induction of ornithine decarboxylase in rat pheochromocytoma: a cyclic AMP-independent process. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 92:313–316.
- Hsie, A. W., and T. T. Puck. 1971. Morphological transformation of Chinese hamster cells by dibutyladenosine cyclic 3',5'-monophosphate and testosterone. *Proc. Natl. Acad. Sci. U. S. A.* 68:358–361.
- Landreth, G., P. Cohen, and E. M. Shooter. 1980. Calcium transmembrane fluxes and nerve growth factor action on a clonal cell line of rat pheochromocytoma. *Nature (Lond.)* 283:202–204.
- Landreth, G. E., and E. M. Shooter. 1980. Nerve growth factor receptors in PC12 cells: ligand-induced conversion from low to high affinity states. *Proc. Natl. Acad. Sci. U. S. A.* 77:4751–4755.
- Li, A. P., P. O'Neill, K. Kawashima, and A. W. Hsie. 1977. Correlation between changes in intracellular level of cyclic AMP, activation of cyclic AMP-dependent protein kinase, and the morphology of Chinese hamster ovary cells in culture. *Arch. Biochem. Biophys.* 182:181–187.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurements with the Folin phenol reagent. *J. Biol. Chem.* 193:265–275.
- Mobley, W., A. Server, D. Ishii, R. Riopelle, and E. Shooter. 1977. Nerve growth factor. *N. Engl. J. Med.* 297:1096–1104.
- Porter, K. R., T. T. Puck, A. W. Hsie, and D. Kelley. 1974. An electron microscope study of the effects of dibutyladenosine cyclic AMP on Chinese hamster ovary cells. *Cell.* 2:145–162.
- Puck, T. T. 1977. Cyclic AMP, the microtubule-microfilament system, and cancer. *Proc. Natl. Acad. Sci. U. S. A.* 74:4491–4495.
- Raff, E. C. 1979. The control of microtubules assembly *in vivo*. *Int. Rev. Cytol.* 59:1–96.
- Schubert, D., S. Heinemann, and Y. Kidokoro. 1977. Cholinergic metabolism and synapse formation by a rat nerve cell line. *Proc. Natl. Acad. Sci. U. S. A.* 74:2579–2583.
- Schubert, D., M. LaCorbiere, C. Whitlock, and W. Stallcup. 1978. Alterations in the surface properties of cells responsive to nerve growth factor. *Nature (Lond.)* 273:718–723.
- Schubert, D., and C. Whitlock. 1977. Alteration of cellular adhesion by nerve growth factor. *Proc. Natl. Acad. Sci. U. S. A.* 74:4055–4058.
- Shigekawa, B. L., and R. W. Olsen. 1975. Resolution of cyclic AMP stimulated protein kinase from polymerization-purified brain microtubules. *Biochem. Biophys. Res. Commun.* 63:455–462.
- Sloboda, R. D., S. A. Rudolph, J. L. Rosenbaum, and P. Greengard. 1975. Cyclic AMP-dependent endogenous phosphorylation of a microtubule-associated protein. *Proc. Natl. Acad. Sci. U. S. A.* 72:177–181.
- Smith, A. P., S. Varon, and E. M. Shooter. 1968. Multiple forms of the nerve growth factor protein and its subunits. *Biochemistry.* 7:3259–3268.
- Tsanev, R., and G. G. Markov. 1960. Substances interfering with spectrophotometric estimation of nucleic acids and their elimination by the two-wavelength method. *Biochim. Biophys. Acta.* 42:442–452.
- Wessels, N. K., B. S. Spooner, J. F. Ash, M. O. Bradley, M. A. Luduena, E. L. Taylor, J. T. Wreen, and K. M. Yamada. 1971. Microfilaments in cellular and developmental processes. *Science (Wash. D. C.)* 171:135–143.
- Willingham, M. C. 1976. Cyclic AMP and cell behavior in cultured cells. *Int. Rev. Cytol.* 44:319–363.
- Yankner, B. A., and E. M. Shooter. 1979. Nerve growth factor in the nucleus: interaction with receptors on the nuclear membrane. *Proc. Natl. Acad. Sci. U. S. A.* 76:1269–1273.