

# Intra- and Interspecific Complementation of Membrane-inexcitable Mutants of *Paramecium*

NOBUYUKI HAGA, YOSHIRO SAIMI, MIHOKO TAKAHASHI\*, and CHING KUNG  
*Laboratory of Molecular Biology and Department of Genetics, University of Wisconsin, Madison, Wisconsin 53706; and \*Institute of Biological Sciences, University of Tsukuba, Ibaraki 305, Japan*

**ABSTRACT** Membrane excitation was the basis for backward swimming of *Paramecium* facing stimulus. According to standard genetic tests, inexcitable mutants fell into three complementation groups for both *Paramecium tetraurelia* (*pwA*, *pwB*, and *pwC*) and *Paramecium caudatum* (*cnrA*, *cnrB*, and *cnrC*). Cytoplasm from a wild type transferred to a mutant through microinjection restored the excitability. Transfusions between genetically defined complementation groups of the same species effected curing, whereas transfusions between different mutants (alleles) of the same group or between sister cells of the same mutant clone did not.

Cytoplasmic transfers of all combinations among the six groups of mutants of the two species showed that any cytoplasm, except those from the same group, was able to cure. Since the pawns and the caudatum nonreversals complement one another through transfusion, they appeared to belong to six different complementation groups. The extent of curing, the amount of transfer needed to cure, and the time course of curing were characteristic of the group that received the transfusion. Variations in these parameters further suggested that the six groups represented six different genes.

Because the donor cytoplasm from either species were equally effective quantitatively in curing a given mutant, the curing factors were not species specific. These factors are discussed.

Ciliated protozoa such as *Paramecium* perform avoiding reactions when stimulated by chemicals, heat, or touch at the anterior (12). An avoiding reaction consists of transient backward swimming that results from the reversal of ciliary-beat direction. Ciliary reversal is due to an increase in intraciliary  $Ca^{++}$ , which flows in through the opened calcium channel (21). The voltage-dependent opening of the calcium channel, the essence of membrane excitation, has been investigated in great detail recently using voltage clamps (5, 6, 19).

Some ciliates can be genetically manipulated (22, 31). Mutants defective in their membranes or axonemes and, therefore, in their behavior, have been isolated and characterized (16, 17, 34). Members of one class of behavioral mutants have no avoiding reactions although they are perfectly motile. Such mutants are called pawns (after the chess piece) in *Paramecium tetraurelia* (14, 15), caudatum nonreversal (CNR)<sup>1</sup> in *Paramecium caudatum* (32, 34), and tetrahymena nonreversal in *Tetrahymena pyriformis* (33). Extensive electrophysiological studies show that these mutants fail in generating the calcium-action potential (18, 33, 34). Upon an abrupt depo-

larization with a voltage clamp, the  $Ca^{++}$  inward current in the wild type, which peaks within several ms, is greatly reduced or absent in pawns (24, 25, 28). Complementation in  $F_1$  and phenotypic segregation in  $F_2$  have led to the conclusion that the more than 200 lines of pawns (representing more than 100 independent mutations) belong to *pwA*, *pwB*, and *pwC* (4, 15, 16), and the seven lines of CNR belong to *cnrA*, *cnrB*, and *cnrC* (32).

Diffusible substances in the cytoplasm of the wild type can restore excitability in the mutants. This can be achieved naturally through the cytoplasmic bridge during conjugation (1) or artificially through microinjection (7–9, 11). Transferring the cytoplasm of a live wild-type donor or various fractions of a wild-type homogenate to a mutant recipient effects the “cure.” Curing occurs hours after injection, lasts 3–4 d, and corresponds to the recovery of  $Ca^{++}$  current upon step depolarization under voltage clamp (7, 8). Such cytoplasmic “transfusion” among many pawns shows that cytoplasmic complementation strictly parallels the genetic complementation so that microinjection can be used to classify the pawns (8).

The question of whether any pawns are genetically equiv-

<sup>1</sup>Abbreviations used in this paper. CNR, caudatum nonreversal.

alent to CNR cannot be answered by the standard complementation test since the interspecific conjugation is not fertile, although it can be artificially initiated (10). In this study, we tested the complementarity of these mutants by transferring their cytoplasm across the species boundary through injection. Much to our surprise, all pawns complemented all CNR and vice versa, indicating that they are likely to be mutants of different genes. There appeared to be at least six genes in all paramecia that controlled the calcium-channel function.

## MATERIALS AND METHODS

**Stocks and Cultures:** We used stock 51s (wild type), d4-500 (*pwA*), d4-95 (*pwB*), and d4-580 (*pwC*) of *P. tetraurelia*, and stock G3 (wild type), 16A712 (*cnrA*), 16B802 (*cnrB*), and 16D341 (*cnrC*) of *P. caudatum* (4, 5, 31, 32). Paramecia were cultured in Cerophyl medium (Cerophyl Laboratories, Inc., Kansas City, MO) buffered with sodium phosphates and bacterized with *Enterobacter aerogenes* (30). Cells were grown at 23°C except in the case of *pwC*, a heat-sensitive mutant, whose phenotype was best expressed after they were grown at 35°C (2, 3, 26).

**Microinjection and Behavioral Test:** Koizumi's method of microinjecting paramecia (13) was modified as described by Haga *et al.* (8): Approximately 15–20 pl of cytoplasm was injected into a *P. tetraurelia* and 45–50 pl into a *P. caudatum*, the larger paramecium. These volumes, ~12% of the total volume of the recipient, were the largest possible for single injections into live cells. The injected cells were incubated in a resting solution and were periodically withdrawn and transferred to a test solution with a micropipette. The test solution was the Dryl's solution enriched with 20 mM KCl. The duration of the backward swimming induced by the transfer was a quantitative measurement of the excitability of the paramecium (8).

**Dark-field Photography and Electrophysiology:** A polaroid camera mounted over a glass plate illuminated at a low angle with a Schott KL-1500 light source (Schott Optical Glassware, Inc., Duryea, PA) in a dark-room was used to register the behavior of the injected cells in a test solution. This has been described previously (2). Methods for capturing, rinsing, immobilizing, penetrating, and recording paramecia have been established (20). The voltage-clamp method was also standard (25). Cells were bathed in 1 mM K<sup>+</sup>, 1 mM Ca<sup>++</sup>, 10 mM tetraethylammonium<sup>+</sup>, 1 mM HEPES, and 10<sup>-5</sup> M EDTA, pH 7.3, during recording. All chemicals are reagent grade; cations were from chloride salts.

## RESULTS

### Interspecific Restorations of Membrane Excitability

We have previously shown that the ability of a pawn or a CNR mutant to swim backward can be restored by microinjection of cytoplasm from a wild type or a mutant of a different complementation group of the same species (7, 8, 11). We show here that transfer of cytoplasm between *P. tetraurelia* and *P. caudatum* also effects such a "curing" of the mutants. A mutant receiving the foreign cytoplasm, like those receiving potent cytoplasm of the same species, regained its ability to swim backward and have the avoiding reactions when properly stimulated. One standard method for showing the avoiding reactions was to transfer the cells into Ba<sup>++</sup>-containing solutions. Because of the generation of the all-or-none Ba-Ca action potentials, the excitable cells performed the "Ba-dance" with almost regular rapid backward and forward steps (17, 20). All inexcitable mutants performed this dance hours after receiving the cytoplasm from the other species. Fig. 1 shows that *cnrC* cells of *P. caudatum* injected with the cytoplasm of *pwC* of *P. tetraurelia* danced in the barium-solution, indicating restored calcium channel activity, whereas control cells (*cnrC* injected with *cnrC* cytoplasm) did not dance, showing no restoration.

### Electric Properties of the Mutant Membrane before and after Cytoplasmic Transfusion

The avoiding reaction resulted from an action potential




FIGURE 1 Behavior of the inexcitable mutant *cnrC* of *P. caudatum* 6–8 h after receiving ~50 pl of cytoplasm from (A) sister *cnrC* cells or from (B) the inexcitable mutant, *pwC*, of *P. tetraurelia*. Several cells were collected from the rest solution and transferred to a solution of 4 mM Ba<sup>++</sup>, 1 mM Ca, and 1 mM Tris, pH 7.2, and their reactions to this barium solution were registered by dark-field photography with 3-s exposure. The gentle helices of A indicate the usual forward swimming of the mutants that remained incapable of responding to the barium solution. The clustered barbs of B were due to repeated jerks of *cnrC* mutants after receiving the *pwC* cytoplasm. The jerks (avoiding reactions) were the consequence of the Ba-Ca action potentials.

and was therefore an indirect indicator of the excitability of the membrane. The membrane excitability can be directly measured with electrodes since paramecium is large enough for convenient intracellular recording (5, 19, 20, 27). The best quantitative analysis of membrane excitability employs a voltage clamp to control the voltage and examine the voltage-dependent ion currents across the membrane. Voltage clamp experiments with wild-type *Paramecium* show that a step depolarization from the resting level (–40 mV, inside negative) by several tens of millivolts (e.g., to –15 mV) triggers a transient inward current carried by Ca<sup>++</sup> that peaks in several milliseconds and subsides within another 5 ms (5, 19, 24). The peak inward current was ~7 nA in wild-type *P. caudatum* previously incubated in the resting buffer. Identical experiments with the pawn (24, 28) or the CNR mutants gave no inward current (Fig. 2A). We showed earlier that pawn mutants regain the calcium current hours after a transfusion from a wild type or from a mutant of a different complementation group of the same species (8). We report here that the same intraspecific restoration of calcium current was true of the CNR (Fig. 2, B–D). We also found that such a restoration of the calcium current could be achieved by transferring cytoplasm of a different species. For example, as shown in Fig. 2, E–G, *cnrC* of *P. caudatum* regained the calcium current after receiving cytoplasm from *pwA*, *pwB*, or *pwC* of *P. tetraurelia*. The restored currents could be as large as those in the wild type, i.e., ~7 nA. The electrophysiological findings show that the restoration of the avoiding reaction as demonstrated in Fig. 1 was not due to some trivial addition of ions to the recipient but reflected a profound change in the membrane excitability of these mutants.

### Complementation Matrix

A quantitative measurement of a paramecium's membrane excitability is easily obtained by transferring it from a resting buffer to a test solution containing a high concentration of K<sup>+</sup> and timing the duration of the first continuous backward swimming induced by this transfer (8). This duration is proportional to the inward calcium current under voltage clamp




FIGURE 2. Membrane currents ( $I_m$ ) induced by step depolarization ( $V_m$ ) in different *P. caudatum* cells. All currents were recorded from *cnrC* mutants except the top traces of A. The voltage steps are given at the top left and the two current traces from each cell are superposed in the figures. A step from  $-44$  mV to  $-30$  mV or to  $-15$  mV induced a small or a larger current in the excitable cells (5, 24, 25). The peak current induced by the larger voltage step was  $\sim 6$  nA (range: 4–9 nA) in the uninjected wild type, G3, previously kept in the rest solution for 6–8 h (A, top traces). The same voltage steps did not induce the inward current in the inexcitable mutant *cnrC* similarly treated (A, bottom traces). *cnrC* mutants became excitable and generated the inward current 6–8 h after they were injected with  $\sim 50$  pl of cytoplasm from (B) *cnrA*, a *P. caudatum* mutant of a

different complementation group; (C) *cnrB*, a *P. caudatum* mutant of yet another group; and (E) *pWA*, (F) *pWB*, and (G) *pWC*, mutants of three complementation groups in *P. tetraurelia*. The restored calcium currents were similar to that of an untreated wild-type paramecium. Injection of the cytoplasm from sister *cnrC* donors to *cnrC* recipients did not restore the calcium current in the recipients (D).




FIGURE 3. Time course of the restoration of the ability to swim backward after microinjection of the cytoplasm. The recipients were injected at time 0 (arrows), periodically withdrawn from the rest solution, and transferred into Dryl's solution with 20 mM  $K^+$ . The

(Y. Satow and N. Haga, unpublished results) and therefore provides an estimate of the amplitude of the calcium current restored by microinjection. Wild-type *P. tetraurelia* swam backward for  $\sim 45$  s and wild-type *P. caudatum* swam backward for  $\sim 60$  s in this test. Table 1 summarizes a complete study of the restoration of excitability, thus measured, with all eight cytoplasmic donors (the two wild types and the six mutants) and all six mutant recipients. Uninjected mutants or those injected with cytoplasm of the sister cells of the same clone showed no backward swimming (0 s for *pWA*, *pWB*, *cnrA*, and *cnrB*) or barely measurable backing (2–3 s for *pWC* and *cnrC*). In contrast, all pawns and *cnrC* showed complete excitability after being injected with other cytoplasm (30–50 s for *pWA*, *pWB*, and *pWC*; 70–75 s for *cnrC*). With the standard transfusion of 12% recipient cell volume, *cnrB* was cured to a lesser extent (10–20 s), whereas *cnrA* was not cured at all.

duration of backward swimming induced by this transfer, which is an estimate of the membrane excitability, was clocked and plotted over time. A shows that the cytoplasm of wild type, stock G3 of *P. caudatum*, injected into *cnrC*, a mutant of the same species, restored the ability of the mutant to swim backward (O, mean  $\pm$  SD,  $n = 6$ ), whereas the cytoplasm from sister *cnrC* donors did not restore (Δ,  $n = 6$ ). B shows that the cytoplasm of a wild type, stock 51 of *P. tetraurelia*, also restored the ability of *cnrC*, the *P. caudatum* mutant, to swim backward. The time courses of *cnrC* curing by the cytoplasms of the two species were very similar. C shows that cytoplasm of G3, twice injected into *cnrA*, a different mutant of *P. caudatum*, also restored the ability of this mutant to swim backward (O,  $n = 10$ ), whereas the cytoplasm of sister *cnrA* cells did not restore the ability (Δ,  $n = 6$ ). Note that the restored activity in *cnrA* was smaller and lasted shorter than in *cnrC* (see text).

However, a second injection, 1 h after the first to allow for recuperation, can partially restore the ability to swim backward even in *cnrA* (see Fig. 3C).

### Time Courses of Curing

Curing the inexcitable mutants with foreign cytoplasm took time. Upon periodic testing, we showed previously that pawns, for example, first regain detectable backward swimming by ~2 h after injection. Complete restoration occurs by 6–8 h. Full excitability is maintained for 1 d after which it gradually subsides until it is barely measurable after 3–4 d (8). The time course of the rise and fall of excitability after injection in *cnrC* was similar to those of the pawns (Fig. 3, A and B). The curing time course, like the extent of curing, was characteristic of the recipients and not the donors, even though the donors may be of two different species, e.g., the time course of curing of *cnrC* with G3 cytoplasm (Fig. 3A) was the same as that with 51s cytoplasm (Fig. 3B).

The course of curing of *cnrB* was similar to that of pawns and *cnrC*, although the extent of curing was smaller (data not shown). The course of curing of *cnrA*, however, was different. After two injections spaced 1 h apart, the excitability, as measured by the K<sup>+</sup>-induced backward swimming, gradually rose in the first 6 h as in other mutants. However, the excitability was not maintained for days. The bulk of the restored excitability was lost by 10 h after the second injection (Fig. 3C).

### DISCUSSION

Since the inexcitable mutants could be temporarily cured by transfusions of heterologous cytoplasm, these cytoplasm contained the gene products missing or defective in the mutants. Our previous study (8) shows that mutual curing among the pawn mutants through microinjection separates them into groups exactly as defined by genetic complementation. Thus, as long as a donor has the normal product of a particular gene, it is capable of curing a recipient that has a mutation in that gene. The normal gene product can be provided by the wild type or by mutants defective in different genes. The effectiveness and efficiency of curing were clearly functions

of the recipients (compare columns in Table 1 and Fig. 3, A and B with 3C) and not the donors (compare entries in each column in Table 1 and Fig. 3A and 3B). Because we found that cytoplasm from a different species could cure the mutants to the same extent and with the same time course, the normal gene products from the two species of *Paramecium* appeared to be functionally equivalent.

The curing was not due to the transfer of nuclear fragments because the nuclei were anchored and we were careful to place the micropipette away from the nuclei. In theory, the gene products transferred can be from either RNA (23) or protein (7, 8). In our experiments, curing of the pawns and *cnrC* could take place in the absence of translation. This, and other experiments with the fractions active in curing, demonstrated that these molecules are proteinaceous. Fractionations of the proteins that effect the cure in the mutants are in progress.

The finding that all pawns complement all CNR is very interesting. The simplest interpretation of these results is that these mutants defined six different complementation groups and therefore six different functional genes. This interpretation rests heavily on observing pawns, where extensive studies show that no curing of a recipient (cytoplasmic complementation) ever results from injection of a cytoplasm from a donor mutant previously defined by breeding analysis to be of the same complementation group. In other words, intragenic complementation has never been observed in pawns by either the genetic or the cytoplasmic tests. Formally, in the absence of intragenic complementation, a new complementing mutation defines a new gene. Since the number of alleles in each *cnr* group remained small, the interpretation that there were at least six genes involved in the excitability phenotype is not entirely secure. The possibility of intragenic complementation among different alleles of each of the *cnr* has not been completely ruled out. Nonetheless, two alleles of *cnrB* did not complement according to our microinjection tests and two alleles of *cnrA* also did not complement, even after double injections (N. Haga, unpublished results). Although the above formal reservation remains, there were other indications that the CNR harbored mutations in genes different from those of pawns. *cnrA* was most difficult to cure and the curing only lasted a short time. *cnrB* was significantly

TABLE I  
Curing Matrix Showing the Duration of Backward Swimming in the K<sup>+</sup>-test Solution after Injection\*

Wild type	<i>P. tetraurelia</i> <sup>‡</sup>			<i>P. caudatum</i> <sup>‡</sup>		
	<i>pwA</i>	<i>pwB</i>	<i>pwC</i>	<i>cnrA</i> <sup>‡</sup>	<i>cnrB</i>	<i>cnrC</i>
<i>P. tetraurelia</i> <sup>‡</sup>	34.4 ± 12.8 (8)	41.0 ± 5.2 (7)	50.8 ± 5.8 (6)	12.0 ± 1.4 (3)	12.2 ± 2.8 (6)	71.0 ± 5.5 (6)
<i>pwA</i>	0 ± 0 (47)	51.5 ± 18.3 (8)	41.5 ± 18.3 (8)	19.0 ± 1.4 (3)	15.0 ± 2.0 (6)	70.3 ± 9.4 (6)
<i>pwB</i>	39.0 ± 12.7 (12)	0 ± 0 (17)	36.0 ± 6.6 (6)	29.0 ± 2.6 (3)	17.7 ± 3.4 (6)	76.2 ± 12.2 (6)
<i>pwC</i>	35.0 ± 12.4 (6)	35.7 ± 12.0 (6)	3.0 ± 3.6 (11)	26.0 ± 6.1 (3)	11.7 ± 3.4 (6)	71.7 ± 11.1 (6)
<i>P. caudatum</i> <sup>‡</sup>	34.0 ± 3.7 (6)	47.3 ± 5.4 (6)	51.0 ± 4.4 (6)	14.7 ± 8.7 (10)	8.7 ± 3.4 (6)	76.1 ± 6.4 (6)
<i>cnrA</i>	37.2 ± 5.2 (6)	37.0 ± 2.4 (6)	50.7 ± 5.4 (7)	0 ± 0 (15)	15.8 ± 4.6 (6)	70.8 ± 8.3 (6)
<i>cnrB</i>	50.3 ± 11.7 (6)	37.0 ± 4.7 (6)	38.7 ± 4.1 (6)	61.5 ± 19.1 (2)	0 ± 0 (6)	69.2 ± 7.6 (6)
<i>cnrC</i>	36.8 ± 6.6 (6)	47.5 ± 8.9 (6)	47.8 ± 7.7 (6)	52.5 ± 3.5 (2)	19.2 ± 8.8 (6)	2.7 ± 2.3 (6)
No injection	0 ± 0 (10)	0 ± 0 (10)	2.4 ± 2.7 (10)	0 ± 0 (10)	0 ± 0 (10)	1.8 ± 1.8 (10)

\* Values are in seconds, mean ± SD (n); uninjected wild-type *P. tetraurelia* (51s): 45 ± 3.8 (10); uninjected wild-type *P. caudatum* (G3): 60.2 ± 4.8 (10). Cells were injected with ~12% of their volume and tested ~8 h after injection.

<sup>‡</sup> *P. tetraurelia* and *P. caudatum* acting as recipients.

<sup>‡</sup> Double injections were spaced 1 h apart. See fig. 3C and text.

<sup>‡</sup> *P. tetraurelia* and *P. caudatum* acting as donors.

more difficult to cure than the pawns (Table 1, Fig. 3), which may be due to certain inherent properties of *cnrB* that made it less receptive to the curing factor. Finally, *cnrC* is clearly a mutant of a different sort since the factor that cured it has now been traced to a postmicrosomal soluble fraction and is therefore most likely a soluble protein, whereas the factors that cured the three pawns have been traced to a microsomal pellet and are probably proteins bound to internal membranes (7, 9).

That *pwA*, *pwB*, *pwC*, *cnrA*, *cnrB*, and *cnrC* are likely to be six different genes in both species is an unexpected result. Over 100 independent mutational events leading to mutations in *pwA* and *pwB* have been analyzed (4, 16). Although relatively rare, more than five independent mutations of *pwC* have been observed. It is therefore surprising that we have not yet encountered mutations in the counterparts of *cnrA*, *cnrB*, and *cnrC* in *P. tetraurelia*. Although not as many mutants have been analyzed, it is equally surprising that the counterparts of *pwA* and *pwB* mutants have not been found in *P. caudatum* since they appear to be abundant, at least in the other species. One possible explanation may be that the methods employed to select pawns (4, 15, 29) and CNR (32) are different in actual execution, although based on the same principle. If pawns and CNR are different mutants, there is hope of finding all six kinds of mutants in both species simply by using different mutagens and/or different mutant-screening methods. Progress is being made toward identifying the factors in the cytoplasm that effect the cure and comparing the factors isolated from the two species. It would be interesting to find out whether these proteins act directly or indirectly (e.g., catalyze a reaction) to restore the calcium-channel activity.

We thank Drs. Michael Forte, Robert Hinrichsen, Rajeev Ramathan, and Joan Kung for critical reading of this manuscript.

This work was supported in part by National Science Foundation grant BNS-7918554 and PHS GM-22714-07 to C. Kung, and grant 57540357 from the Ministry of Education, Science and Culture, Japan, to M. Takahashi.

Received for publication 17 January 1983, and in revised form 7 April 1983.

## REFERENCES

- Berger, J. D. 1976. Gene expression and phenotypic change in *Paramecium tetraurelia* exconjugants. *Genet. Res.* 27:123-134.
- Chang, S. Y., and C. Kung. 1973. Temperature-sensitive pawns: conditional behavioral mutants of *Paramecium aurelia*. *Science (Wash. DC)*. 180:1197-1199.
- Chang, S. Y., and C. Kung. 1973. Genetic analysis of heat-sensitive pawn mutants of *Paramecium aurelia*. *Genetics*. 75:49-59.
- Chang, S. Y., J. Van Houten, L. Robles, S. Lui, and C. Kung. 1974. An extensive behavioral and genetic analysis of the Pawn mutants of *Paramecium aurelia*. *Genet. Res.* 23:165-173.
- Eckert, R., and P. Brehm. 1979. Ionic mechanisms of excitation in *Paramecium*. *Annu. Rev. Biophys. Bioeng.* 8:353-383.
- Eckert, R., Y. Naitoh, and H. Macheimer. 1976. Calcium in the bioelectric and motor functions of *Paramecium*. In *Calcium in Biological Systems*. C. J. Duncan, editor. Cambridge University Press, New York. 233-255.
- Haga, N., M. A. Forte, Y. Saimi, and C. Kung. 1981. Characterization and partial purification of a factor controlling the generation of action potential in *Paramecium*. In *Society for Neuroscience 11th Annual Meeting Abstracts*. Society for Neuroscience, Bethesda, Maryland. 15a. (Abstr.).
- Haga, N., M. Forte, Y. Saimi, and C. Kung. 1982. Microinjection of cytoplasm as a test of complementation in *Paramecium*. *J. Cell Biol.* 92:559-564.
- Haga, N., and K. Hiwatashi. 1982. A soluble gene product controlling membrane excitability in *Paramecium*. *Cell Biology International Report* 6:295-300.
- Hiwatashi, K. 1969. *Paramecium*. In *Fertilization II*. C. B. Metz and A. Monroy, editors. Academic Press, Inc., New York. 255-293.
- Hiwatashi, K., N. Haga, and M. Takahashi. 1980. Restoration of membrane excitability in a behavioral mutant of *Paramecium caudatum* during conjugation and by microinjection of wild-type cytoplasm. *J. Cell Biol.* 84:476-480.
- Jennings, H. S. 1906. *Behavior of Lower Organisms*. Indiana University Press, Bloomington, Indiana.
- Koizumi, S. 1974. Microinjection and transfer of cytoplasm in *Paramecium*. *Exp. Cell Res.* 88:74-78.
- Kung, C. 1971. Genic mutations with altered system of excitation in *Paramecium aurelia*. I. Phenotypes of the behavioural mutants. *Z. vegl. Physiol.* 71:142-164.
- Kung, C. 1971. Genic mutations with altered system of excitation in *Paramecium aurelia*. II. Mutagenesis, screening and genetic analysis of the mutants. *Genetics*. 69:29-45.
- Kung, C. 1979. The biology and genetics of paramecium behavior. In *Topics in Neurogenetics*. X. O. Breakfield, editor. Elsevier/North Holland Biomedical Press, New York. 1-26.
- Kung, C., S. Y. Chang, Y. Satow, J. Van Houten, and H. Hansma. 1975. Genetic dissection of behavior in *Paramecium*. *Science (Wash. DC)*. 188:898-904.
- Kung, C., and R. Eckert. 1972. Genetic modifications of electric properties in an excitable membrane. *Proc. Natl. Acad. Sci. USA*. 69:93-97.
- Kung, C., and Y. Saimi. 1982. The physiological basis of taxes in *Paramecium*. *Annu. Rev. Physiol.* 44:519-534.
- Naitoh, Y., and R. Eckert. 1972. Electrophysiology of ciliate protozoa. *Exp. Physiol. Biochem.* 5:17-31.
- Naitoh, Y., and H. Kaneko. 1972. Reactivated triton-extracted models of *Paramecium*: modification of ciliary movement by calcium ions. *Science (Wash. DC)*. 176:523-524.
- Nanney, D. L. 1980. *Experimental Ciliatology, An Introduction to Genetics and Developmental Analysis in Ciliates*. Wiley Interscience Div., John Wiley & Sons, Inc., New York.
- Nock, A. H., M. Kretschmar, H. J. Lipps, and J. E. Schultz. 1981. Restoration of membrane excitability by microinjection of cytoplasmic wild-type RNA into *Paramecium tetraurelia* Pawn C mutants. *FEBS (Fed. Eur. Biochem. Soc.) Lett.*
- Oertel, D., S. J. Schein, and C. Kung. 1977. Separation of membrane currents using a *Paramecium* mutant. *Nature (Lond.)*. 268:120-124.
- Saimi, Y., and C. Kung. 1982. Are ions involved in the gating of Ca-channels? *Science (Wash. DC)*. 218:153-156.
- Satow, Y., S. Y. Chang, and C. Kung. 1974. Membrane excitability: made temperature dependent by mutations. *Proc. Natl. Acad. Sci. USA*. 71:2703-2706.
- Satow, Y., and C. Kung. 1979. Voltage-sensitive Ca-channels and the transient inward current in *Paramecium tetraurelia*. *J. Exp. Biol.* 78:149-161.
- Satow, Y., and C. Kung. 1980. Membrane currents of pawn mutants of the *pwA* group in *Paramecium tetraurelia*. *J. Exp. Biol.* 84:57-71.
- Schein, S. J. 1976. Nonbehavioral selection for pawns, mutants of *Paramecium aurelia* with decreased excitability. *Genetics*. 84:453-468.
- Sonneborn, T. M. 1970. Methods in *Paramecium* research. In *Methods of Cell Physiology*. D. M. Prescott, editor. Academic Press, Inc., New York. 4:243-339.
- Sonneborn, T. M. 1974. *Paramecium aurelia*. In *Handbook of Genetics*. R. C. King, editor. Plenum Press, New York. 2:469-594.
- Takahashi, M. 1979. Behavioral mutants in *Paramecium caudatum*. *Genetics*. 91:393-408.
- Takahashi, M., H. Onimaru, and Y. Naitoh. 1980. A mutant of *Tetrahymena* with non-excitable membrane. *Proc. Jpn. Acad.* 56:585-590.
- Takahashi, M., and Y. Naitoh. 1978. Behavioral mutants of *Paramecium caudatum* with defective membrane electrogenesis. *Nature (Lond.)*. 271:656-658.