

Sequence of a Highly Divergent β Tubulin Gene Reveals Regional Heterogeneity in the β Tubulin Polypeptide

KEVIN F. SULLIVAN and DON W. CLEVELAND

Department of Biological Chemistry, The Johns Hopkins University School of Medicine, Baltimore, MD 21205

ABSTRACT The nucleotide sequence of a chicken genomic DNA segment containing the chicken $\beta 4$ tubulin gene has been determined. The predicted amino acid sequence of $\beta 4$ is surprisingly divergent from that of the chicken $\beta 2$ gene that encodes the dominant neural β tubulin. $\beta 4$ differs from $\beta 2$ at 36 residue positions and encodes a polypeptide that is four amino acids longer, yielding a divergence of 8.9% between the two β tubulin isotypes. While many of the amino acid substitutions are conservative, several involve significant alteration in the physiochemical properties of the residue. Furthermore, the amino acid substitution positions are not randomly located within the primary sequence but are distinctly clustered: major divergence occurs in the carboxy-terminal region beyond residue 430 and within the second protein coding exon segments of the genes. In addition, large regions of absolute sequence conservation are also present. Certain sequences within the heterogeneous regions are conserved in other species, indicating that these regions are under positive evolutionary selection pressure and are therefore probably essential for some aspect of β -tubulin function. These findings strongly suggest that regional amino acid sequence heterogeneity may play an important role in the establishment of functionally differentiated β tubulin polypeptides.

Microtubules, composed principally of dimeric subunits of 1 α and 1 β tubulin polypeptide, are filamentous polymers that are employed in a number of diverse functions in eucaryotic cells. For example, microtubules are the major components of mitotic and meiotic spindles and of cilia and flagella. Furthermore, they participate in various types of intracellular transport and, in conjunction with intermediate filaments and actin filaments, establish the structure and functional properties of the eucaryotic cytoskeleton. Yet in spite of this functional diversity, the ultrastructure of microtubules is, with a few notable exceptions (6, 10), identical in all cells. Moreover, the tubulin polypeptides themselves have been conserved throughout evolution to such an extent that tubulins from all sources investigated have been shown to co-polymerize efficiently *in vitro* (see reference 18 for review).

Despite these apparently conserved features, several lines of evidence suggest that the microtubule systems of different cells or within single cells are distinguishable on the basis of physical, biochemical, and pharmacological properties. For example, Benke and Forer (2) originally reported evidence for four classes of microtubules in individual cells which differed in their drug sensitivities or thermal stability. Similarly, Brinkley and Cartwright (5) have demonstrated that the interpolar and kinetichore microtubules of mammalian spindles differ in their response to drug- and cold-induced depolymerization.

Additional compelling evidence has come from the work of Thompson et al. (31) and Cumming et al. (9) who have independently demonstrated, using monoclonal antibodies to α tubulin determinants, that α tubulin related epitopes are apparently segregated into distinguishable subsets of microtubules within single cells or subsets of cells. Thus, it seems clear that although microtubules appear morphologically identical, they represent a heterogeneous class of structures.

One obvious potential level of control of microtubule properties is at the level of the tubulin dimers which constitute the major structural subunit. This hypothesis would require that multiple, functionally distinct tubulin subunits be present within cells. Immunological and biochemical support for this hypothesis has emerged from a number of investigators who have reported heterogeneous pools of α and β tubulins in a number of tissue and cell types (e.g., references 11, 12, 22, 24, 25, 30) and within single cells (13). More definitively, protein sequence analysis of porcine brain tubulins has demonstrated the presence of two β tubulins (19) and four α tubulins (27) that differ subtly in primary sequence. More recently, sequence analysis of two different human β tubulin genes has demonstrated the presence of at least two β tubulin isotypes that differ significantly in sequence at the carboxy terminus of the polypeptide (15).

We previously demonstrated that the chicken contains four

β tubulin genes that possess strong homology to a cloned β tubulin cDNA, each of which has been isolated by molecular cloning and designated $\beta 1$ – $\beta 4$ (20). Recent experiments (Havercroft, J. C., D. B. Murphy, and D. W. Cleveland, unpublished results) indicate the presence in the chicken genome of 3–5 additional divergent β tubulin sequences with only weak nucleic acid homology to other cloned β tubulin sequences. Overall, this suggests that the chicken genome possesses not more than eight or nine functional β tubulin genes. With the goal of determining the number and identity of tubulin polypeptides that constitute a complete repertoire of tubulins necessary for the development and differentiation of a vertebrate organism, we have begun nucleotide sequence analysis of the β tubulin gene family in the chicken. We report here the sequence of the chicken $\beta 4$ gene. We find that this gene encodes a polypeptide that is strikingly divergent from any previously reported vertebrate β tubulin. Comparison of these data with other currently available β tubulin sequences demonstrates the presence of regions of heterogeneity embedded within an otherwise highly conserved β tubulin polypeptide.

MATERIALS AND METHODS

DNA Sequence Analysis: DNA sequence analysis was carried out by the chain terminator method of Sanger et al. (28) using fragments of the cloned β G4 gene which had been subcloned into the Messing M13 vectors (23), mp8 or mp9. Random fragments of the plasmid β G4 were generated by digestion of the plasmid DNA with DNase I (Worthington Biochemical Corp, Freehold, NJ) in the presence of Mn^{2+} , as described by Anderson (1). Briefly, 100 μ g of supercoiled plasmid β G4 DNA (19) was digested with 5 μ g DNase I in the presence of 1 mM $MnCl_2$ at room temperature and 25 μ g samples were removed after 12, 15, 18, and 21 min of reaction and stopped by addition of 10 mM EDTA. Samples were pooled and, after deproteinization, ends were repaired by fill in with DNA polymerase I, Klenow fragment (Bethesda Research Laboratories, Gaithersburg, MD). The DNA was resolved on a polyacrylamide gel and fragments in the range from 600–2,000 bp were collected by electroelution. Recovered fragments were then ligated with M13 mp8, which had been cut with *HincII* and treated with calf intestinal phosphatase, and transformed into JM101. Phage plaques were then screened by the method of Benton and Davis (3) using the 5.3 kb *XhoI-KpnI* fragment of β G4 (see Fig. 1A) which had been labeled with ^{32}P using the method of Shank et al. (29). DNA was then prepared from positive plaques for sequencing. Some fragments were also obtained by directed cloning of restriction fragments into the appropriate double cut M13 vectors.

S1 analysis: S1 nuclease protection experiments were performed by a modification of the method of Berk and Sharp (4). To probe for the 3' end of $\beta 4$ transcripts, the 2.5 kb Bam HI fragment of β G4 was isolated by agarose gel electrophoresis and labeled at 3' ends by incubation with DNA polymerase I Klenow fragment in the presence of α - ^{32}P -dATP (400 Ci/mmol, Amersham Corp., Arlington Heights, IL) and 1 mM each dGTP, dCTP, and dTTP for 15 min at 30°C. Messenger RNA (mRNA) was prepared from whole 7-d chicken embryos using the guanidinium thiocyanate procedure of Chirgwin et al. (7). Probe (0.01–0.4 μ g) was co-precipitated with 0.25 to 1.0 μ g of mRNA plus 400 μ g of carrier yeast tRNA and then redissolved in 20 μ L of a solution containing 80% formamide, 40 mM PIPES, pH 6.4, 400 mM NaCl, and 1 mM EDTA. Samples were incubated at 85°C for 10 min to denature the probe and then at 60°C for 3 h to allow hybridization to occur. Hybridization mixtures were then diluted with 200 μ L of S1 reaction buffer containing 500 U of S1 nuclease (Bethesda Research Laboratories) and reactions were incubated at 22°C for 60 min. Samples were then prepared for electrophoresis and resolved on 6% polyacrylamide sequencing type gels and results were visualized by autoradiography.

RESULTS

Of the four chicken β tubulin genes that we have cloned, we have previously determined the coding region sequence for the chicken $\beta 2$ gene, the genomic progenitor of the full length cDNA clone pT2 (8, 32). This gene encodes the dominant neuronal β tubulin of the chicken (16, 20). To begin to analyze the diversity of β tubulin polypeptide sequences encoded within this organism we have now determined the sequence

of the $\beta 4$ gene. This gene was elected for analysis because RNA and DNA blot hybridization experiments had shown that it was divergent from the $\beta 2$ sequence (20). Plasmid β G4 containing the $\beta 4$ gene subcloned into the *SphI* site of pBR322 (as described previously [20]) is illustrated in Fig. 1A. The 5.3 kb *XhoI-KpnI* fragment of β G4 contains the entirety of the sequences homologous to gene $\beta 2$. This fragment was subjected to shotgun DNA sequence analysis using M13/dideoxy sequencing techniques as described in Materials and Methods. The sequences of 71 subcloned fragments, representing a total of 12584 bases, were used to assemble the final sequence. Overall, we have determined the sequence of 5358 contiguous bases spanning the entire *XhoI-KpnI* fragment. Greater than 90% of the sequence was read from both strands or from multiple overlapping clones; the actual sequencing strategy is shown in Fig. 1B.

The structure of the $\beta 4$ gene as determined by comparison with the known coding region sequence from $\beta 2$ is diagrammed in Fig. 1C. The organization is identical to that of the human β tubulin gene M40 (14) and consists of four blocks of protein coding exon sequences interrupted by three intervening sequences placed between codons 19 and 20, and within codons 56 and 93, respectively. Thus, it appears that the coding structure of β tubulin genes has been conserved throughout vertebrate evolution. A putative RNA polymerase II promoter element, TATAAA, is located 100 bases 5' to the initiator ATG codon and a consensus polyadenylation signal, AATAAA, is located 165 nucleotides 3' of the TGA translation termination codon. No additional polyadenylation signals were found in the 2.3 kb of 3' flanking DNA whose sequence was determined.

Amino Acid Sequence of the $\beta 4$ Polypeptide

From the nucleotide sequence we have deduced the predicted protein sequence of the polypeptide encoded by the $\beta 4$




FIGURE 1 Organization, sequencing strategy, and structure of the chicken $\beta 4$ tubulin gene. (A) The organization of the recombinant plasmid β G4 (20) which contains a 10-kb fragment of chicken genomic DNA cloned into the *Sph I* site of pBR322. The 5.3 kb *Xho I-Kpn I* fragment, which contains the entire $\beta 4$ coding region and 2.5 kb of 3' flanking sequences was chosen for DNA sequence analysis. (B) Schematic drawing of the fragments used to assemble the $\beta 4$ sequence. (C) The structure of the $\beta 4$ gene. Filled boxes denote protein coding exon sequences while noncoding sequences are shown by the thin line. The positions of putative transcription control sequences, the TATAAA promoter, and AATAAA polyadenylation site are noted. The direction of transcription is from left to right.

gene. The relevant DNA sequence and corresponding translation are shown in Fig. 2, along with the previously determined coding sequence of the dominant chicken neural β tubulin gene, $\beta 2$. It is readily apparent that the $\beta 4$ gene encodes a polypeptide that is surprisingly divergent compared with the dominant neural β tubulin. There are a total of 36 amino acid substitutions between these two proteins. In addition, the predicted $\beta 4$ polypeptide is four amino acids longer than the prototypic $\beta 2$ polypeptide. Most striking is the presence of a basic lysine residue that terminates the highly acidic carboxy-terminus of the protein. Thus, the overall divergence between these two proteins is 8.9%: this is to be compared with 1 and 5% divergence noted between the $\beta 2$ sequence and the sequence of a porcine brain (19) or human (15) β tubulin, respectively.

Comparison of the $\beta 4$ and $\beta 2$ Polypeptide Sequences

Detailed comparison of the amino acid differences between the $\beta 2$ and $\beta 4$ polypeptides shows generally conservative amino acid substitutions. Several substitutions, however, involve significant changes in the chemical properties or size of

the side chain. Three substitutions, at residues 37, 53, and 83, involve exchange of aliphatic or amide-containing residues for histidine, resulting in a net gain of two histidine residues in $\beta 4$. One marked change is $\beta 2$ Ser 275 to $\beta 4$ Arg 275, a change that involves not only the acquisition of an additional fixed positive charge in $\beta 4$, but also a significant alteration in the size of the side chain. Curiously, there is a complementary exchange of serine and cysteine residues in positions 124 and 239, substitutions likely to significantly alter the chemical properties of these residues. However, since these changes involve an exchange of serine and cysteine residues, it is tempting to speculate that these residues may be close to each other in the folded polypeptide and that the exchange preserves a functional requirement for the cysteine residue.

The major region of heterogeneity between $\beta 2$ and $\beta 4$ is localized to the extreme carboxy terminus of the proteins, including an additional four-amino acid extension for the $\beta 4$ polypeptide. Most of the substitutions in this region involve exchange or acquisition of acidic residues in $\beta 4$, preserving the overall net charge of this region. Two striking substitutions are found, however, replacing the peptide pair Phe436-Glu437 in $\beta 2$ with Met-Tyr in $\beta 4$. In addition, the four residue extension sequence, Gln-Gly-Ala-Lys, which in addition to




FIGURE 2 Comparison of the coding region of $\beta 4$ with the coding region of $\beta 2$ (pT2), the dominant chicken neural β tubulin. The coding sequence of $\beta 4$ is shown with the corresponding predicted translation to protein sequence above it. The three intron sequences, inserted between codons 19 and 20 and between bases 1 and 2 of codons 56 and 93, respectively, are not shown. The sequence is aligned with the nucleotide sequence of the full length β tubulin cDNA clone pT2 (32) which is derived from the chicken $\beta 2$ gene; identical nucleotides are denoted by dashes while nucleotide differences are explicitly shown. Amino acid residues of $\beta 2$ which differ from $\beta 4$ are noted beneath the $\beta 2$ nucleotide sequence.

the terminal lysine is comprised of small neutral amino acids, represents a significant departure from the highly acidic carboxy-terminal sequence that precedes it.

Regional Heterogeneity within β Tubulin Polypeptides

The amino acid substitutions between the $\beta 2$ and $\beta 4$ polypeptides do not appear to be homogeneously distributed, but rather are clustered within the primary sequence, giving rise to regions of high sequence heterogeneity as well as regions of high conservation. This perhaps best illustrated in Fig. 3 in which we have presented the parallel comparison of all currently available vertebrate β tubulin sequences. Immediately

apparent in this comparison is the prominent sequence divergence among these various β tubulins in the carboxy terminal region beyond amino acid 431 (see also discussion above). In particular, this region accounts for 33% of the divergence between $\beta 4$ and $\beta 2$ and also represents a major variable region between two human β tubulin genes (15). Moreover, a second variable region in which substantial divergence is also obvious is found between residues 33 and 57. This 24-amino acid portion of the polypeptide possesses seven substitution positions between $\beta 4$ and $\beta 2$. Inspection of Fig. 3 shows that this region is also a site of significant microheterogeneity between porcine brain β tubulins (19). Although most of the substitutions in this region are conservative, two substitutions involve exchange of neutral residues for histidine (as detailed above).





FIGURE 3 Comparison of the chicken $\beta 4$ tubulin sequence with the sequences of other vertebrate β tubulins. The sequence of the predicted chicken $\beta 4$ tubulin polypeptides is shown (using the one letter amino acid code) and has been aligned with the predicted protein sequences of the chicken $\beta 2$ tubulin gene (32), two human β tubulin sequences, D $\beta 1$ and 5 β (obtained by DNA sequencing of a fetal human brain cDNA clone and a partial sequence of a functional human β tubulin gene, respectively [reference 15]), and two variants, A and B, of porcine brain β tubulin (19). With the exception of residue position 275, porcine variants A and B differ only within the first 83 residues (the first line of the figure) and are therefore shown as a single sequence on subsequent lines. In each of the other sequences, only positions that differ from the chicken $\beta 4$ sequences are noted. The known sequence for human 5 β begins at residue 431. Dashed boxes enclose the regions of highest sequence divergence as discussed in the text. The diagram below shows the linear distribution of amino acid substitution positions between chicken $\beta 2$ and $\beta 4$ tubulins (vertical dashes) shown within the primary sequence of β tubulin (horizontal line).

In addition to regions of heterogeneity, regions of extreme sequence conservation are also strikingly evident in the comparison presented in Fig. 3. For example, the amino terminal portion up to residue 32 is invariant among the known vertebrate sequences, as is the carboxy terminal region between residues 366–431.

A summary of these observations is presented in the line drawing at the bottom of Fig. 3 which shows the linear distribution of amino acid substitutions between the chicken $\beta 4$ and $\beta 2$ tubulin isotypes. Overall, the data suggest that the β tubulin polypeptide is organized so as to incorporate regional heterogeneity within a highly conserved framework.

$\beta 4$ Gene Is Expressed during Chicken Development

The $\beta 4$ gene thus encodes a polypeptide with surprisingly divergent primary sequence. But is this predicted β tubulin subunit actually expressed and used in cellular microtubule systems? Blot analysis of mRNA transcripts of the chicken β tubulin genes has previously demonstrated that the $\beta 4$ genomic sequence hybridizes strongly and preferentially to a 3,500–3,700 base mRNA present at high abundance in most tissues and cell types tested (16, 20), thus suggesting that the $\beta 4$ gene product is a ubiquitous and perhaps constitutive β tubulin subunit. However, this initial blot data has proven to be misleading. Contrary to our initial expectations, more recent work has shown that neither the immediate 5' or 3' flanking regions of the $\beta 4$ gene are present on the 3,500–3,700 base β tubulin mRNA (16). In addition, the present sequence data demonstrate the presence of a consensus polyadenylation signal, AATAAA, 165 bp 3' to the termination codon, and a putative promoter element, TATAAA, located 100 bases 5' to the ATG translation initiation codon. Assuming both that transcription begins 30 bases downstream of the TATAAA sequence (reviewed, for example, in reference 26) and the presence of a 200 nucleotide 3' poly-A tract beginning 20 bases downstream of the polyadenylation signal, these data predict an mRNA species of ~1,800 bases, markedly different from the previously detected 3,500–3,700 base mRNAs. No additional polyadenylation sites are present in the 2.3 kb of downstream flanking sequence.

To determine whether the $\beta 4$ gene is a functional and expressed gene, we performed the following S1 nuclease protection experiment. A 2.3-kb fragment that begins in the coding region within codon 344 and extends through 317 bp of the C-terminal coding region and terminates in the flanking DNA ~2 kb 3' to the translation termination codon, was prepared from p β G4 by Bam H1 digestion. This fragment was labeled with 32 P at 3' ends as described in Materials and Methods and hybridized to poly-A+ mRNA from 7-d chick embryos. After hybridization, the mixture was digested with S1 nuclease to remove unhybridized single stranded regions of DNA, and the labeled fragments protected from digestion by hybridization to RNA were analyzed by electrophoresis and autoradiography as shown in Fig. 4. Clearly, the predominant mRNA species, which is homologous to the $\beta 4$ gene probe, protects a fragment ~510 bases in length. This is precisely the length of protected DNA expected if the polyadenylation signal sequence 165 bases downstream of the termination codon is used. Furthermore, there is no evidence for the presence of a larger RNA transcript that derives from this $\beta 4$ gene.




FIGURE 4 S1 nuclease protection analysis of $\beta 4$ gene transcripts. An S1 nuclease protection experiment was performed as described in Materials and Methods using the 2.5 kb Bam H1 fragment of p β G4 which spans the 3' end of the gene. This fragment is diagrammed at the top of the figure and extends 317 bp 5' of the TGA translation termination codon to 1921 bp 3' of the termination codon. The position of the final A of the polyadenylation signal, 171 bp 3' of the termination codon, is also noted. The autoradiogram is oriented with the bottom of the gel at left; the experimental lanes shown represent a titration of probe DNA from 12.5 to 400 ng in a series of twofold increments. Probe alone is noted in the next to the last lane and pBR322/Hpa II size markers are shown in the bottom lane. The positions of the two major bands are noted with respect to the sequence; the numbering refers to the size of the fragment with respect to the translation termination codon.

This experiment unambiguously demonstrates that the $\beta 4$ gene is transcribed into a polyadenylated mRNA. However, this mRNA represents a minor component of the β tubulin mRNA transcripts in most cells and tissues as we have not to date detected significant quantities of the $\beta 4$ transcript by RNA blot analysis using $\beta 4$ specific 5' or 3' untranslated region probes (16). Curiously, however, the experiment shown in Fig. 4 demonstrates the presence of a second, shorter protected fragment, corresponding to extensive nucleic acid homology with an mRNA that extends only up to amino acid residue 430 (~55 bases 5' to the stop codon). This protected fragment must arise from hybridization to an mRNA that is transcribed from a gene extremely homologous to, but not identical to, the cloned $\beta 4$ gene. From the known sequences (32; Sullivan and Cleveland, unpublished results), RNAs transcribed from the chicken $\beta 1$, $\beta 2$, and $\beta 3$ genes are unlikely to generate this protected fragment since for each of these the major region of sequence divergence from $\beta 4$ lies near residue 351 rather than near position 430. Therefore, we suggest that the mRNA which gives rise to the shorter protected probe is probably the 3,500–3,700 base transcript detected with overwhelming preference by the $\beta 4$ gene in blotting experiments and that this mRNA derives from a $\beta 4$ -like gene that we have not yet cloned.

In any case, irrespective of the surprisingly complex relationship between the chicken $\beta 4$ and $\beta 4$ -like tubulin gene, it is clear that the cloned $\beta 4$ gene does represent an authentic, functional β tubulin gene that is transcribed into a stable mRNA encoding a highly divergent β tubulin subunit.

DISCUSSION

Previous studies of tubulin polypeptide structure have stressed the conservation of β tubulins throughout evolution. For example, the partial sequence data of Luduena and Woodward (21) initially demonstrated that the amino termini of sea urchin flagellar and chicken brain tubulins were identical. More definitively, subsequent comparison of the entire se-

quence of a chicken brain cDNA for β tubulin (32) and porcine brain β tubulin (19) showed a conservation of 99% in the primary sequences of these two polypeptides. Although Krauhs et al. (19) identified sequence microheterogeneity between two porcine β tubulins, this heterogeneity was restricted to six residue positions, suggesting that the differences between different β tubulins were quite subtle. Together, these previous studies indicated that β tubulin was among the most highly conserved proteins known.

It has now become clear that this conclusion is both correct and misleading. Not only has the sequence of a human β tubulin (D β 1) demonstrated a much lower degree of conservation (15), we have now demonstrated a level of isotypic variation of β tubulins within a single species of 40 of 449 residues (8.9%). This level of divergence is greater than that seen in any interspecies comparison of available metazoan β tubulin sequences. However, certain isotypic forms of β tubulin, for example the neuronal polypeptides represented by the β 2 sequence and the porcine variant A, have remained highly conserved during evolution. Thus, while the individual isotypes that constitute the β tubulin family have diversified within species during evolution, individual isotypic forms of β tubulin have been highly conserved between species.

Collectively, it appears certain from the data currently available that the expression of different β tubulin isotypes is a general property of metazoan species. But are the isotypic differences which distinguish different tubulins within a single species functionally meaningful? Kempf, Raff, and co-workers (17) have presented an elegant genetic and morphological analysis of a series of mutants in a testes specific β tubulin variant and have demonstrated that this unique protein is the major β tubulin subunit of postmitotic spermatids and that it apparently functions in all postmitotic microtubule systems of the spermatid. Documentation of this multifunctional behavior presents a compelling argument against a unique testes specific function for this β subunit. This has led these workers to suggest instead that multiple tubulin genes may be functionally equivalent, representing duplicated genes that have evolved into different genetic regulatory systems, thereby enabling the cell to coordinate tubulin synthesis with specialized programs of differentiation.

However, the sum of the presently available sequence data (Fig. 3) offers equally compelling molecular evidence for the alternative hypothesis of different, functionally specialized β tubulin gene products. Many of the differences that distinguish the two chicken β tubulin isotypes are distinctly clustered between regions of the polypeptide that are remarkably conserved among other known vertebrate β tubulins. Of course, it may be argued that variable regions simply represent nonessential portions of the molecule that are not subject to strong selection at the level of protein sequence. This hypothesis seems unlikely, however, since it is clear from inspection of the interspecies comparisons in Fig. 3 that certain sequences within the clusters of divergence have been highly conserved during evolution. For example, the chicken β 2 sequence is virtually identical to porcine brain β variant A, even in the regions of high substitution frequency. It seems likely therefore that the conserved sequence exemplified by β 2 represents the predominant neural β tubulin isotype found in each of these organisms. Furthermore, several of the amino acid substitutions that characterize the amino-terminal variable cluster of β 4 are also observed in the porcine brain variant β (19). Since this area is entirely encoded on exons 2 and 3 of

the β tubulin genes, it is plausible that it constitutes a variable domain within the β tubulin polypeptide.

Since the specific sequence arrangements found in the heterogeneous regions cannot be essential for the highly conserved functions of β tubulin (polymerizability, etc.), the observed conservation of sequences in tubulins from different species argues strongly that the heterogeneous regions have been maintained by evolutionary selection for specific functional properties. An alternative, but perhaps less likely, explanation is that the variable sequences have not been conserved, but have undergone convergent evolution to similar sequences in response to similar structural requirements that have arisen during evolution. Both explanations, however, imply that specific sequence arrangements in the variable regions are the result of positive selective pressure. We cautiously suggest that the conservation of variable region sequences in β tubulin polypeptides forms the foundation of a convincing demonstration of a functional role for the variable domains, a suggestion that can be directly tested by further sequence analysis of β tubulins from different organisms.

Many thanks to Dr. Jane Havercroft for generously providing us with the mRNA for the S1 experiments, to Dr. Himan Sternlicht whose enthusiasm helped to get this project started, and to Mr. Floyd Benko for his excellent technical support.

This work has been supported by grants from the National Institutes of Health and March of Dimes to D. W. Cleveland, who is the recipient of a National Institutes of Health Research Career Development Award. K. F. Sullivan is the recipient of a National Institutes of Health Postdoctoral Fellowship.

Received for publication 2 July 1984, and in revised form 16 August 1984.

REFERENCES

- Anderson, S. 1981. Shotgun DNA sequencing using cloned DNase I generated fragments. *Nucl. Acids Res.* 13:3015-3027.
- Benke, O., and A. Forer. 1967. Evidence for four classes of microtubules in individual cells. *J. Cell Sci.* 2:169-192.
- Benton, W. D., and R. W. Davis. 1977. Screening gt recombinant clones by hybridization to single plaques in situ. *Science (Wash. DC)*. 196:180-183.
- Berk, A., and P. Sharp. 1978. Spliced early mRNAs of SV40. *Proc. Natl. Acad. Sci. USA* 75:1274-1278.
- Brinkley, B. R., and J. Cartwright. 1975. Cold-labile and cold-stable microtubules in the mitotic spindle of mammalian cells. *Ann. NY Acad. Sci.* 253:428-439.
- Chalfie, M., and J. N. Thompson. 1982. Structural and functional diversity in the neuronal microtubules of *Caenorhabditis elegans*. *J. Cell Biol.* 93:15-23.
- Chirgwin, J. M., A. E. Przybyla, R. J. MacDonald, and W. J. Rutter. 1979. Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry*. 24:5294-5299.
- Cleveland, D. W., M. A. Lopata, R. J. MacDonald, N. J. Cowan, W. J. Rutter and M. W. Kirschner. 1980. Number and evolutionary conservation of α and β tubulin and cytoplasmic β and γ actin genes using specific cloned cDNA probes. *Cell*. 20:95-105.
- Cumming, R., R. D. Burgoyne, and N. A. Lytton. 1984. Immunocytochemical demonstration of α tubulin modification during axonal maturation in the cerebellar cortex. *J. Cell Biol.* 98:347-351.
- Fujiwara, K., and L. G. Tilney. 1975. Structural analysis of the microtubule and its polymorphic forms. *Ann. NY Acad. Sci.* 253:27-50.
- George, H. J., L. Misra, D. J. Field, and J. C. Lee. 1981. Polymorphism of brain tubulin. *Biochemistry*. 20:2402-2409.
- Gozes, I., and U. Z. Littauer. 1978. Tubulin microheterogeneity increases with rat brain maturation. *Nature (Lond.)*. 276:411-413.
- Gozes, I., and K. J. Sweadner. 1981. Multiple tubulin forms are expressed by a single neurone. *Nature (Lond.)*. 294:477-480.
- Gwo-Shu Lee, M., S. A. Lewis, C. D. Wilde, and N. J. Cowan. 1983. Evolutionary history of a multigene family: an expressed human β tubulin gene and three processed pseudogenes. *Cell*. 33:477-487.
- Hall, J. L., L. Dudley, P. R. Dobner, S. A. Lewis, and N. J. Cowan. 1983. Identification of two human β tubulin isotypes. *Mol. Cell Biol.* 3:854-862.
- Havercroft, J. C., and D. W. Cleveland. 1984. Programmed expression of β tubulin genes during development and differentiation of the chicken. *J. Cell Biol.* In press.
- Kempf, K., T. C. Kaufman, R. A. Raff, and E. C. Raff. 1982. The testis specific β tubulin subunit in *Drosophila melanogaster* has multiple functions in spermatogenesis. *Cell*. 31:655-670.
- Kirschner, M. W. 1978. Microtubule nucleation and assembly. *Int. Rev. Cytol.* 54:1-71.
- Krauhs, E., M. Little, T. Kempf, R. Hofer-Warbinek, W. Ade, and H. Ponstingl. 1981. Complete amino acid sequence of β tubulin from porcine brain. *Proc. Natl. Acad. Sci. USA* 78:4156-4160.

20. Lopata, M. A., J. C. Havercroft, L. T. Chow, and D. W. Cleveland. 1983. Four unique genes required for β tubulin expression in vertebrates. *Cell*. 32:713-724.
21. Luduena, R. F., and D. O. Woodward. 1973. Isolation and partial characterization of α and β tubulin from outer doublets of sea urchin sperm and microtubules of chick embryo brain. *Proc. Natl. Acad. Sci. USA* 70:3594-3598.
22. Marotta, C. A., J. L. Harris, and J. M. Gilbert. 1978. Characterization of multiple forms of brain tubulin subunits. *J. Neurochem.* 30:1431-1440.
23. Messing, J., and J. Vieira. 1982. A new pair of M13 vectors for selecting either stand of double-digested restriction fragments. *Gene*. 19:269-276.
24. Murphy, D. B., and K. T. Wallis. 1983. Brain and erythrocyte microtubules from chicken contain different β tubulin polypeptides. *J. Biol. Chem.* 258:7870-7875.
25. Nelles, L. P., and J. R. Bamberg. 1979. Comparative peptide mapping and isoelectric focusing of isolated subunits from chick embryo brain tubulin. *J. Neurochem.* 32:477-489.
26. Nevins, J. 1983. The pathway of eukaryotic mRNA formation. *Ann. Rev. Biochem.* 52:441-466.
27. Ponstingl, H., E. Kraus, M. Little, and T. Kempf. 1981. Complete amino acid sequence for α tubulin from porcine brain. *Proc. Natl. Acad. Sci. USA*. 78:2757-2761.
28. Sanger, F., S. Nicken, and A. R. Coulson. 1977. DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA*. 74:5463-5467.
29. Shank, P. R., S. H. Hughes, H.-J. Kung, J. E. Majors, N. Quintrell, R. V. Guntaka, J. M. Bishop, and H. E. Varmus. 1978. Mapping unintegrated avian sarcoma virus DNA: termini of linear DNA bear 300 nucleotides present once or twice in two species of circular DNA. *Cell*. 15:1383-1395.
30. Sullivan, K. F., and L. Wilson. 1984. Developmental and biochemical analysis of chick brain tubulin heterogeneity. *J. Neurochem.* 42:1363-1371.
31. Thompson, W. C., D. J. Asai, and D. H. Carney. 1984. Heterogeneity among microtubules of the cytoplasmic microtubule complex detected by a monoclonal antibody to α tubulin. *J. Cell Biol.* 98:1017-1025.
32. Valenzuela, P., M. Quiroga, J. Zaldivar, W. J. Rutter, M. W. Kirschner, and D. W. Cleveland. 1981. Nucleotide and corresponding amino acid sequences encoded by α and β tubulin mRNAs. *Nature (Lond.)*. 289:650-655.