

CYTOTOXIC IMMUNE CELLS WITH SPECIFICITY FOR DEFINED SOLUBLE ANTIGENS

IV. ANTIBODY AS MEDIATOR OF SPECIFIC CYTOTOXICITY*

By VOLKER SCHIRRMACHER,† BENT RUBIN,§ HUGH PROSS,|| AND HANS WIGZELL

(From the Department of Tumorbiology, Karolinska Institute, Stockholm, Sweden)

(Received for publication 21 August 1973)

In an attempt to study model systems for allograft and tumor immunity, a number of in vitro cell-mediated cytotoxic systems have been investigated. Depending on the cytotoxic system studied, different types of cells, e.g. T lymphocytes, B lymphocytes, or macrophages, have been found to be involved in specific target cell destruction (see reference 1). Little is known about the parameters which determine the type of effector cell(s) that will function in a given experimental situation. Also, the mechanisms that lead to target cell destruction in the various systems and their possible in vivo interactions are poorly understood. In view of this complex situation, it seems desirable to start with the analysis of highly reproducible and well-defined systems.

In recent studies, we have investigated an in vitro cytotoxic system with specificity for defined soluble antigens. It has previously been found in an experimental autoimmune situation that red cells coated with soluble protein antigen can function as target cells for antigen specific cytolytic in vitro reactions (2). The present system consists of spleen cells from mice that have been immunized against conventional soluble antigens, such as heterologous proteins or hapten-carrier conjugates, and of ⁵¹Cr-labeled target cells (chicken red blood cells, CRBC) to which the respective antigens had been covalently attached (3). Cell-fractionation studies (4, 5) have revealed that the lytic effector cell in this system is neither an antigen-specific T cell (as tested by helper function) nor an antigen-specific B cell (as tested by antibody-forming cell precursor function) and that it has distinct adherence properties (see reference 6). Greenberg et al. (7), working with antibody-coated CRBC and non-

* This work was conducted under Contract no. NIH-NOI-CB-33 859, National Institutes of Health, United States Public Health Service. It was further supported by the Swedish Cancer Society, the Karolinska Institute, the Deutsche Forschungsgemeinschaft, the Statens Laegevidenskabelige Forskningsrad, Denmark, and the Medical Research Council of Canada.

† Present address: The London Hospital Medical College, Tissue Immunology, University of London, London E1 2AD, England.

§ Present address: Immunobiology Laboratory, Statens Serum Institute, DK-2300 Copenhagen, Denmark.

|| Fellow of the Medical Research Council of Canada.

Abbreviations used in this paper: Ag-CRBC, antigen-coated CRBC; BGG, bovine γ -globulin; CRBC, chicken red blood cells; DB, dinitrophenylated bovine serum albumin; DM, dinitrophenylated mouse serum albumin; DNP, dinitrophenyl; HSA, human serum albumin; OA, ovalbumin; PBS, phosphate-buffered saline, pH 7.4; and SRBC, sheep red blood cells.

immune mouse spleen cells, demonstrated that in their system the effector cells were membrane-Ig negative as well as θ -antigen negative. These effector cells could be separated from other cells and were characterized as monocyte-like cells.¹

The present paper reports investigations on the mechanism that leads to specific lysis of the antigen-coated target cells by immune spleen cells. Little is known about what confers specificity on monocyte- or macrophage-mediated cytotoxicity. Among the different factors that have been suggested are: specific macrophage-arming factor (SMAF, 8, 9), which might be a T cell product (9, 10), and cytophilic antibody (11, 12). Evidence will be presented that specific cytotoxicity in our system is mediated through specific antibody, which is produced by the immune cells, and which interacts with the target cell antigens through its combining sites. The interaction of antibody with the cytotoxic effector cells will be analyzed in the following report.²

Materials and Methods

The materials and methods used in the cytotoxic system have been described in detail previously (3-5) and therefore only a brief description will be given.

Mice and Immunization.—2-3-mo old CBA mice were used throughout the experiments. They were immunized subcutaneously with 0.1 ml antigen solution (1 mg/ml) emulsified in an equal volume of complete Freund's adjuvant (CFA).

Preparation of Antigen-Coated and ⁵¹Cr-Labeled Target Cells.—0.1 ml of a 50% suspension of washed fresh CRBC was added to 3 ml of phosphate-buffered saline (PBS), pH 7.4, containing 35 mg albumin antigen. 0.35 ml of a solution of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide-HCl (E CDI, 200 mg/ml) was added and the cells were incubated for 1 h at room temperature. They were then washed four times in PBS and tested for the presence of membrane-bound antigen using specific antisera and the passive hemagglutination microtest. For labeling with ⁵¹Cr, about 10⁷ Ag-CRBC in 0.2 ml PBS were incubated with 150 μ Ci Na₂⁵¹CrO₄ (1 mCi/ml) for 3-4 h at room temperature. The labeled cells were washed twice with culture medium (Medium F13, containing 10% heat-inactivated fetal calf serum, penicillin, and streptomycin) and adjusted to a concentration of 10⁵ cells/ml.

Test for Cell-Mediated Cytotoxicity.—The cytotoxic test was carried out in Wassermann plastic test tubes (11 \times 55 mm), each of which contained 5 \times 10⁴ ⁵¹Cr-labeled Ag-CRBC and varying amounts of immune or nonimmune spleen cells (see the corresponding ratio for each experiment) in a volume of 1 ml culture medium. Depending on the experiment, antibody or inhibitors were added to the incubation mixtures. Their final concentrations are indicated in each experiment. After incubation for about 18 h at 37°C in a 5% CO₂ atmosphere, the tubes were centrifuged and 0.5-ml aliquots of the supernatants removed. The radioactivity of the pellet and residual supernatant and the radioactivity of the 0.5-ml supernatant sample were measured in a well-type scintillation counter. After subtraction of the background, a release index (R.I.) was calculated:

$$\text{R.I.} = \frac{\text{radioactivity of the supernatant}}{\text{total radioactivity}} \times 100.$$

¹ A. H. Greenberg, L. Shen, and I. M. Roitt. 1973. Characterization of the antibody-dependent cytotoxic cell: a non-phagocytic monocyte? *Clin. Exp. Immunol.* **15**:251.

² V. Schirmmayer, B. Rubin, and H. Pross. 1973. Cytotoxic immune cells with specificity for defined soluble antigens. V. Analysis of the interaction of antibody with the cytotoxic effector cells in immune or nonimmune spleen cells. Manuscript submitted for publication.

The results are expressed as R.I. minus spontaneous release of target cells alone. Each value given is the mean of the figures obtained for three test tubes. The intragroup variance for a given experiment has been computed from the R.I. of all the test tubes. Use has been made of this variance to calculate the difference that would be significant ($P = 0.05$) in a student's t test between two experimental groups in the experiment.

Preparation and Use of Anti-Ig or Control Columns.—The preparation of glass bead columns coated with a polyvalent rabbit antimouse immunoglobulin serum (anti-Ig column) or with normal rabbit serum (control column) has been described (13). Sera or supernatants were absorbed on the columns by passing them through at room temperature and at a flow rate of about 1 ml/min. The eluates were concentrated to the original volume by negative pressure dialysis.

Separation of 19S and 7S Antibodies.—19S and 7S antibodies were separated by sucrose gradient centrifugation as described previously (14).

RESULTS

Absence of Carrier Specificity.—The secondary anti-DNP response to dinitrophenylated mouse serum albumin (DM) in mice shows the phenomenon of carrier specificity (15), while the anti-DNP antibodies induced are hapten specific (16).³ In order to evaluate the specificity characteristics of the cytotoxic reaction of DM-immune cells the role of the carrier molecule was investigated. The results in Table I show that DM-immune cells display the same degree of cytotoxicity to CRBC coated with dinitrophenylated bovine serum albumin (DB) as to CRBC-DM, while they are not cytotoxic to CRBC-OA. Since there is no cross-reactivity between the carrier determinants on DM and DB (16), it is concluded that the cytotoxic reaction of DM-immune cells is not carrier specific.

Inhibition of Cytotoxicity by Soluble Antigen or Hapten.—We further investigated the specificity of the cytotoxic system by inhibition experiments using as inhibitors free soluble antigen or hapten. Table II shows two such experiments. DM- and OA-immune cells were incubated in the presence or absence of either DNP-lysine (exp. I) or OA (exp. II) together with CRBC-DM, CRBC-DB, or CRBC-OA as target cells. Specific and complete inhibition was observed in both instances, and the existence of hapten-specific cytotoxicity is clearly demonstrated (exp. I).

The data in Table II show differences in the amount of OA or DNP-lysine that inhibited the respective cytotoxic system. On a molar basis, about a hundred times more DNP-lysine was needed to get the same degree of inhibition as with OA. When the serum antibodies of the OA- and DM-immune animals that were donating the immune cytotoxic cells were analyzed by passive hemagglutination inhibition, similar differences in the location of the inhibition curves were observed (Fig. 1). With increasing time after immunization both cytotoxicity inhibition curves and antibody inhibition curves shifted to lower antigen or hapten concentrations. From these data it is apparent that the cyto-

³ See Discussion section concerning the immune response against new antigenic determinants.

TABLE I
Absence of Carrier Specificity of DM-Immune Cell Cytotoxicity

Target cell	S.R.*	Cytotoxicity† of immune cells‡					
		Anti-DM			Anti-OA		
		150:1	75:1	25:1	150:1	75:1	25:1
CRBC-DM	7	35	26	20	5	9	7
CRBC-DB	8	37	28	23	9	9	10
CRBC-OA	7	6	5	7	37	30	23

* Spontaneous ⁵¹Cr release (S.R.) from target cells incubated alone.

† Expressed as release index minus S.R.

‡ The immune spleen cells (3 wk) were incubated with the target cells at the indicated ratios; the smallest difference that would be significant ($P = 0.05$) in a t test between two figures was 4.6.

TABLE II
Inhibition of Immune Cell Cytotoxicity by Free Hapten or Antigen

Exp.	Target cell	S.R.*	Cytotoxicity† of immune cells‡							Inhibitor	
			Anti-DM			Anti-OA					
I		F13	-8	-6	-4	F13	-8	-6	-4	DNP-lysine	
	CRBC-DM	5	42	37	16	8	4	—	3		3
	CRBC-DB	8	39	39	21	9	3	—	—		—
	CRBC-OA	6	6	5	2	3	49	—	40	42	
II		F13	-9	-8	-7	F13	-9	-8	-7	OA	
	CRBC-DM	4	40	—	—	42	9	—	—		7
	CRBC-OA	5	7	—	—	5	44	32	16		7

* Spontaneous ⁵¹Cr release (S.R.) of target cells incubated alone.

† Expressed as release index minus S.R.

‡ Immune spleen cells, 4 wk (exp. I) or 2 wk (exp. II) after immunization with 100 μ g DM or OA; the ratio of immune to target cells was 150:1. The smallest difference that would be significant ($P = 0.05$) in a t test between two figures was 3.5 (exp. I) and 5.5 (exp. II). —, not done.

|| The inhibitors were present during the whole incubation period (18 h). The \log_{10} molar inhibitor concentration in the final incubation mixture is indicated in the respective horizontal line. F13 means medium without inhibitor.

toxicity inhibition curves reflect the avidity or affinity (in the case of OA and DNP-lysine respectively) of the serum antibodies in the same animals.

Induction of Cytotoxicity in Nonimmune Spleen Cells by Target Cell-Bound Serum Antibodies.—Fig. 2 illustrates an experiment in which specific cytotoxicity was observed (a) with immune spleen cells, and (b) with immune




FIG. 1. Inhibition of antibody activity (A) and of immune cell cytotoxicity (B) by free soluble antigen or hapten. A Antibody activity in the sera from animals immunized 2 weeks (open symbols) or 6 weeks (closed symbols) before with 100 μ g OA (triangles) or DM (circles) was determined by passive hemagglutination inhibition using SRBC-OA or CRBC-DM as indicator cells. The log₂ titers for the pooled sera from three animals per group were: anti-OA (2 wk) 6.5; anti-OA (6 wk) 11.5; anti-DM (2 wk) 5.5; and anti-DM (6 wk) 5.5. Passive hemagglutination inhibition was carried out by titrating the sera in PBS containing OA or DNP-lysine respectively as inhibitors. B cytotoxicity of the immune spleen cells from the same animals was determined using ⁵¹Cr-labeled CRBC-OA or CRBC-DM as target cells. The ratio of immune to target cells was 150:1. The specific cytotoxicity in the absence of OA or DNP-lysine as inhibitors was: anti-OA (2 wk) 28%; anti-OA (6 wk) 33%; anti-DM (2 wk) 57%; and anti-DM (6 wk) 49%. The smallest difference that would be significant ($P = 0.05$) in a t test between two points was 2.6.




FIG. 2. Cytotoxicity of either OA-immune spleen cells (7 wk, circles) or of anti-OA antiserum from the same animals in presence of nonimmune spleen cells (= normal cells, triangles). The target cells were CRBC-OA (open symbols) and CRBC-HSA (solid symbols). The ratio of nonimmune spleen cells to target cells was 100:1. No specific cytotoxicity was observed with anti-OA antiserum at the indicated concentrations in the absence of nonimmune spleen cells. The smallest difference that would be significant ($P = 0.05$) in a t test between two points was 4.9.

serum from the same animals in the presence of nonimmune spleen cells. Cytotoxicity increased with increasing ratios of immune to target cells⁴ as well as with increasing amounts of antiserum (starting at a dilution of 10^{-7}) at a constant ratio of nonimmune cells to target cells (100:1). While minute amounts of antiserum (10^{-4} – 10^{-5} diluted) induced optimal cytotoxicity, higher concentrations ($\geq 10^{-3}$) usually had a suppressive effect.

In order to characterize the nature of the cytotoxicity inducing factor in the immune serum, the antiserum was absorbed on an anti-Ig or on a control column. The results (Table III) show that anti-OA antiserum absorbed on an

⁴ In earlier experiments (3) we observed a linear increase of cytotoxicity with increasing ratios of immune to target cells only up to a ratio of about 50:1. It was now possible to show that the suppression or relative reduction of cytotoxicity at higher ratios was due to the centrifugation of the tubes before the incubation period. This step, which increased the sensitivity of the system at low ratios, was omitted in the experiments reported in this paper.

TABLE III
Cytotoxicity of Nonimmune Spleen Cells in the Presence of Antiserum Absorbed on anti-Ig or Control Columns

Target cell	S.R.*	Cytotoxicity† of nonimmune spleen cells in presence of anti-OA antiserum‡ which was					
		Unfractionated		Absorbed on an anti-Ig column		Absorbed on a control column	
		10^{-4}	10^{-6}	10^{-4}	10^{-6}	10^{-4}	10^{-6}
CRBC-OA	10	57	39	14	1	54	16
CRBC-HSA	10	0	0	1	1	3	0

* Spontaneous ^{51}Cr release (S.R.) from target cells incubated alone.

† Expressed as release index minus S.R.

‡ Antiserum (inactivated 30 min at 56°C) from mice that had been immunized 6 mo before with $100\ \mu\text{g}$ OA in Freund's complete adjuvant. Its \log_2 passive hemagglutination titer with CRBC-OA was 9, after anti-Ig column absorption 1.5 and after control column absorption 8. The final dilution of the antiserum in the test is indicated. The ratio of spleen cells to target cells was 150:1. The smallest difference that would be significant ($P = 0.05$) in a t test between two figures was 4.1.

anti-Ig column was by a factor of about one hundred less active in inducing specific cytotoxicity in comparison with the same serum absorbed on a control column. These results indicate the immunoglobulin nature of the cytotoxicity inducing factor in the immune serum.

Induction of Cytotoxicity in Nonimmune Spleen Cells by Supernatants from Immune Cell Cultures.—The fact that serum antibodies bound to Ag-CRBC target cells induced cytotoxicity in nonimmune spleen cells does not necessarily mean that the present immune cell cytotoxic system functions according to the same principle, i.e. through the involvement of specific antibodies. In order to elucidate this point an attempt was made to directly demonstrate specific cytotoxicity inducing antibody activity in the supernatants of our immune cell cultures. Fig. 3 shows the positive outcome of such an experiment. DB-immune spleen cells were incubated in vitro under the conditions of the cytotoxic test, but in the absence of target cells, and the supernatants from fourteen such cultures were pooled. DNP specific cytotoxicity could be conferred on non-immune spleen cells by these supernatants, as well as by their γ -globulin fractions, provided, of course, that DNP-coated target cells were added to the system to demonstrate the effect (Fig. 3A and Fig. 3B). Supernatants absorbed on an anti-Ig column were inactive (Fig. 3C). It is thus shown that the immune spleen cells produce antibody in vitro in quantities sufficient to induce cytotoxicity in nonimmune spleen cells. Since we have also shown that the immune cell cytotoxic system has the specificity and affinity characteristics of serum antibodies (see above), we would conclude that cytotoxicity in our immune cell system is in fact mediated through specific antibodies.

Activity of Target Cell-Bound 19S and 7S Antibodies in the Induction of Cyto-




FIG. 3. Cytotoxicity of supernatants from DB-immune cell cultures in presence of non-immune spleen cells and CRBC-DM (white columns) or uncoated CRBC (black columns) as target cells. The ratio of nonimmune spleen cells to target cells was 100:1. 4×10^7 DB-immune cells (4 wk) were incubated in 1 ml culture medium for 18 h at 37°C in a 5% CO₂ atmosphere. The supernatants of fourteen such cultures were pooled. The γ -globulin fraction was prepared from a sample (7 ml) by two successive precipitations with half saturated ammonium sulfate, with extensive dialysis against saline in between and afterwards. Another sample of the supernatant was absorbed on a glass bead column coated with rabbit anti-mouse Ig antiserum.

toxicity.—The next step in the analysis of the Ag-CRBC lytic system was to investigate whether the ability of antibodies to mediate specific cytotoxicity was restricted to a certain class (19S/7S) of antibodies. Fig. 4 shows the results of several experiments. 19S and 7S anti-DNP antisera which had the same passive hemagglutination titer were titrated in the presence of nonimmune spleen cells and CRBC-DM (white columns) or CRBC-OA (black columns) as target cells. While 19S antibodies from whole sera or from a 19S sucrose gradient fraction were virtually inactive in the cytotoxic test (Fig. 4 A), 7S antibodies from whole sera or from a 7S sucrose gradient fraction were very efficient in inducing cytotoxicity (Fig. 4 B).

DISCUSSION

Antibody-dependent cell-mediated cytolytic systems have been described in a number of instances, where antibody complexed target cells were incubated




FIG. 4. Cytotoxicity of nonimmune spleen cells in presence of 19S (A) or 7S (B) anti-DNP antibodies and CRBC-DM (white columns) or CRBC-OA (black columns) as target cells. The ratio of spleen cells to target cells was 100:1. The DNP-specific hemagglutination titers of the different antibody fractions, indicated on the right part, were tested with DNP-HSA-SRBC. While the 19S antibody titers were sensitive to reduction with 0.2 M 2-mercaptoethanol, the 7S titers were reduction resistant. 19S antisera were obtained from mice 9 days (first column) or 13 days (second and third column) after a single intraperitoneal injection of 100 μ g DNP₆₀-BGG. The third column shows the activity of a 19S sucrose gradient fraction. 7S antisera were obtained from mice 6 wk (first column) or 6 mo (second column) after immunization with 100 μ g DNP₁₂-BGG in Freund's complete adjuvant. The second column shows the activity of a 7S sucrose gradient fraction. All sera were inactivated 30 min at 56°C.

in vitro with nonimmune lymphoid cells. Mostly xenogeneic systems have been studied (17-19) but allogeneic (20, 21) and syngeneic (autoimmune, reference 22, and tumor immune reference 23) systems have been described as well. The cytolytic effector cells in these systems are believed not to belong to the T lymphocytes (see references 24 and 25). It has not yet been demonstrated if immune cell cytolytic systems can function according to the same principle, i.e. through a cooperation between an antibody-producing system and non-immune effector cells. Synergistic and antagonistic effects of antibody on in vitro immune cell cytotoxicity have been described (26-28). The only safe conclusion that has been reached so far is that there exist in the immune system at least two alternative cell-mediated cytolytic pathways, one involving T lymphocytes (13, 29), the other non-T lymphoid or other cells (3, 31, 34). The exact nature of the latter effector cells is still under discussion. Non-T cell-mediated cytotoxicity has been attributed to any type of non-T cell-bearing membrane-bound F_c receptors: purified lymphocytes (18, 35), B cells (19, 25, 31, 36, 37); macrophages (11, 38-40); monocytes (41); polymorphonuclear cells (42); and also to a new type of cell (43-45). Some of the studies that involved the use of anti-Ig sera (31, 36) for characterizing the effector cells might have

been misleading, because inhibition of cytotoxicity might have reflected a blocking of F_c receptors via anti-Ig-Ig complexes rather than a direct effect on the effector cell surface structures (see references 4, 7, and 24).² Since the F_c receptors on lymphocytes and monocytes differ in their specificities for IgG subclasses (46, 47) aggregated myeloma proteins of different subclasses have been used in inhibition experiments in order to clarify whether the effector cells belong to the lymphocyte or to the monocyte lineage of cells. Unfortunately, different results have been obtained in different systems. The cytotoxic effector cells in the human peripheral blood showed subclass specificity characteristics of lymphocytes (45, 48), while in mouse spleen cells the cytotoxic effector cells shared the subclass specificities with the monocytes.¹ It therefore seems that depending on the cytotoxic system studied, different types of non-T cells may become activated by antibody-coated target cells to function as cytotoxic effector cells.

The present immune cell cytotoxicity system has certain advantages as compared with others: (a) it is generally applicable to the study of cytotoxic responses using any kind of soluble antigen (3),⁵ (b) the target cell membrane is relatively stable (3), reducing the risk of production of inhibitory complexes; (c) the exact control of specificity is made possible by using the same type of target cell coated with different antigens; and (d) it is also possible to do fine specificity studies and even affinity or avidity analysis through inhibition by defined soluble antigens.

In the present study, three lines of evidence were obtained for the involvement of specific antibodies in the immune cell cytotoxicity system: (a) DM-immune cell cytotoxicity has been shown to be hapten specific (Tables I and II) and thus to follow the specificity characteristics of anti-DNP antibodies rather than of cellular immune responses (see reference 50). No specificity was found for the new antigenic determinants (NAD's, reference 16) introduced into the autologous protein by the hapten-coupling reaction. This is due to the fact that NAD's were not expressed on the CRBC-DM target cells, probably due to the carbodiimide-coupling reaction (V. Schirmacher and B. Rubin, unpublished observations). (b) The cytotoxicity inhibition curves obtained in the presence of either free hapten (DNP-lysine in case of DM-immune cells) or soluble antigen (OA in case of OA-immune cells) reflected the affinity or avidity respectively of the serum antibodies from the same animals. Cytotoxicity inhibition curves and antibody inhibition curves were at a similar position in both the DNP and the OA system, and both shifted with increasing time after immunization towards lower inhibitor concentrations (Fig. 1). (c) Supernatants from in vitro immune cell cultures contained an antigen-specific factor, which, like serum antibodies (Fig. 2 and Table III), induced cytotoxicity in nonimmune spleen cells when specific target cells were added. This supernatant ac-

⁵ The system has already been successfully applied for studies on autoimmune thyroiditis (2, 49).

tivity was recovered in the γ -globulin fraction, and was absorbed on columns coated with polyvalent antimouse immunoglobulin serum (Fig. 3).

Thus, specificity is conferred on the effector cells in the immune cytolytic system by antibodies, which are produced during the incubation period of the test. We have no evidence for antibodies being brought into the system from the beginning, e.g. in form of cytophilic antibodies that were attached to cell surfaces (12), but we do know that cytotoxicity in the immune cell system starts only after a lag period of about 3 h, while in presence of antibodies it would start immediately.² Other evidence for the dependency of the immune cell cytotoxic system on active antibody synthesis *in vitro* comes from studies of MacLennan and Harding (51) who showed that their immune cell cytotoxic system was inhibited in presence of puromycin while the nonimmune cell system was not affected.

The ability of target cell-bound antibody to induce cytotoxicity in nonimmune spleen cells was restricted to the 7S antibody class (Fig. 4). Similar results have been obtained in other antibody-dependent cell-mediated cytotoxic systems (52, 53). It is difficult to explain why 19S antibodies were inactive in this cytolytic test, especially considering the fact that macrophages do possess a receptor for 19S antibodies (54).

SUMMARY

Spleen cells from mice immunized against ovalbumin (OA) or dinitrophenylated mouse serum albumin (DM) were found to be specifically cytotoxic *in vitro* towards target cells (chicken red blood cells) coated with these antigens. Inhibition of specific cytotoxicity was observed when free soluble antigen was added to the incubation mixtures. DM-immune cell cytotoxicity could be specifically and completely inhibited by DNP-lysine and was thus shown to be hapten specific. Complete and specific inhibition was also observed for OA-immune cell cytotoxicity using OA as inhibitor, but compared with the inhibition curves obtained with DNP-lysine, the OA cytotoxicity inhibition curves were shifted by a factor of about one hundred towards lower molar inhibitor concentrations. Very similar results were observed when the serum antibodies of DM- and OA-immune animals were analyzed by passive hemagglutination inhibition. With increasing time after immunization, both cytotoxicity inhibition curves and agglutination inhibition curves, shifted to lower antigen or hapten concentrations.

Specific cytotoxicity against antigen-coated target cells was induced in nonimmune spleen cells (*a*) by serum from immune animals, and (*b*) by supernatants from *in vitro* immune cell cultures. In both instances, the factor which induced antigen-specific cytotoxic activity could be absorbed on anti-mouse Ig columns, thus demonstrating its immunoglobulin nature. The ability of target cell bound antibodies to induce cytotoxicity in nonimmune spleen cells was restricted to the 7S antibody class.

We thank Prof. P. Perlmann and his staff for the continuous supply of fresh chicken blood.

REFERENCES

1. Cerottini, J.-C., and K. T. Brunner. 1973. Cell-mediated cytotoxicity, allograft rejection and tumor immunity. *Adv. Immunol.* In press.
2. Ringertz, B., J. Wassermann, Th. Packalen, and P. Perlmann. 1971. Cellular and humoral immune responses in experimental autoimmune thyroiditis. *Int. Arch. Allergy Appl. Immunol.* **40**:917.
3. Schirmacher, V., and P. Golstein. 1973. Cytotoxic immune cells with specificity for defined soluble antigens. I. Assay with antigen-coated target cells. *Cell. Immunol.* In press.
4. Golstein, P., V. Schirmacher, B. Rubin, and H. Wigzell. 1973. Cytotoxic immune cells with specificity for defined soluble antigens. II. Chasing the killing cells. *Cell. Immunol.* In press.
5. Schirmacher, V., B. Rubin, P. Golstein, and H. Wigzell. 1973. Cytotoxic immune cells with specificity for defined soluble antigens. III. Separation from helper cells and from antibody forming cell precursors. *Transplant. Proc.* In press.
6. Dennert, G., and E. Lennox. 1972. Cell interactions in humoral and cell-mediated immunity. *Nat. New Biol.* **238**:114.
7. Greenberg, A. H., L. Hudson, L. Shen, and I. M. Roitt. 1973. Antibody-dependent cell-mediated cytotoxicity due to a "null" lymphoid cell. *Nat. New Biol.* **242**:111.
8. Evans, R., and P. Alexander. 1971. Rendering macrophages specifically cytotoxic by a factor released from immune lymphoid cells. *Transplantation.* **12**:227.
9. Lohmann-Matthes, M.-L., F. G. Ziegler, and H. Fischer. 1973. Macrophage cytotoxicity factor. A product of in vitro sensitized thymus-dependent cells. *Eur. J. Immunol.* **3**:56.
10. Evans, R., C. K. Grant, H. Cox, K. Steele, and P. Alexander. 1972. Thymus-derived lymphocytes produce an immunologically specific macrophage arming factor. *J. Exp. Med.* **136**:1318.
11. Granger, G. A., and R. S. Weiser. 1966. Homograft target cells: contact destruction in vitro by immune macrophages. *Science (Wash. D.C.)*. **151**:97.
12. Temple, A., G. Loewi, P. Davies, and A. Howard. 1973. Cytotoxicity of immune guinea-pig cells. II. The mechanism of macrophage cytotoxicity. *Immunology* **24**:655.
13. Golstein, P., H. Wigzell, H. Blomgren, and E. A. J. Svedmyr. 1972. Cells mediating specific in vitro cytotoxicity. II. Probable autonomy of thymus-processed lymphocytes (T cells) for the killing of allogeneic target cells. *J. Exp. Med.* **135**:890.
14. Schirmacher, V., and K. Rajewsky. 1970. Determination of antibody class in a system of cooperating antigenic determinants. *J. Exp. Med.* **132**:1019.
15. Rubin, B., V. Schirmacher, and H. Wigzell. 1973. The immune response against hapten-autologous protein conjugates in the mouse. II. Carrier specificity in the secondary anti-hapten response and evidence for the existence of specific helper cells. *Scand. J. Immunol.* **2**:189.
16. Rubin, B. 1973. The immune response against hapten-autologous protein conjugates in the mouse. I. Specificity of antibodies produced during the primary response against dinitrophenylated mouse serum albumin. *Eur. J. Immunol.* **3**:26.

17. MacLennan, I. C. M., and G. Loewi. 1968. The effect of specific antibody to target cells on their specific and non-specific interactions with lymphocytes. *Nature (Lond.)*. **219**:1069.
18. Perlmann, P., and H. Perlmann. 1970. Contactual lysis of antibody-coated chicken erythrocytes by purified lymphocytes. *Cell. Immunol.* **1**:300.
19. Möller, G., and S.-E. Svehag. 1972. Specificity of lymphocyte-mediated cytotoxicity induced by in vitro antibody-coated target cells. *Cell. Immunol.* **4**:1.
20. Hasek, M., J. Hökl, J. Bubenik, and I. Karakoz. 1970. Cell to cell injury mediated through alloantigens. *Folia Biol.* **16**:173.
21. Wunderlich, J. R., E. B. Rosenberg, and J. M. Conolly. 1971. Human lymphocyte-dependent cytotoxic antibody and mechanism of target cell destruction. *In* Progress in Immunology. B. Amos, editor. Academic Press, New York. **1**:473.
22. Wassermann, J., Th. Packalen, and P. Perlmann. 1971. Antibody-induced lysis of thyroglobulin-coated chicken erythrocytes by lymphocytes of non-immune guinea pigs. *Int. Arch. Allergy Appl. Immunol.* **41**:910.
23. Pollack, S., G. Heppner, R. J. Brawn, and K. Nelson. 1972. Specific killing of tumor cells in vitro in the presence of normal lymphoid cells and sera from hosts immune to the tumor antigens. *Int. J. Cancer* **9**:316.
24. Perlmann, P., H. Perlmann, and H. Wigzell. 1972. Lymphocyte mediated cytotoxicity in vitro. Induction and inhibition by humoral antibody and nature of effector cells. *Transplant. Rev.* **13**:91.
25. MacLennan, I. C. M. 1972. Antibody in the induction and inhibition of lymphocyte cytotoxicity. *Transplant. Rev.* **13**:67.
26. Möller, E. 1965. Antagonistic effects of humoral isoantibodies on the in vitro cytotoxicity of immune lymphoid cells. *J. Exp. Med.* **122**:11.
27. Skurzak, H. M., E. Klein, T. O. Yoshida, and E. W. Lamon. 1972. Synergistic or antagonistic effect of different antibody concentrations on in vitro lymphocyte cytotoxicity in the moloney sarcoma virus system. *J. Exp. Med.* **135**:997.
28. Kiessling, R., and E. Klein. 1973. Cytotoxic potential of mouse spleen cells on H-2 antibody-treated target cells. *J. Exp. Med.* **137**:527.
29. Cerottini, J.-C., A. A. Nordin, and K. T. Brunner. 1970. Specific in vitro cytotoxicity of thymus-derived lymphocytes sensitized to alloantigens. *Nature (Lond.)*. **228**:1308.
30. Harding, B., D. J. Pudifin, F. Gotch, and I. C. M. MacLennan. 1971. Cytotoxic lymphocytes from rats depleted of thymus-processed cells. *Nat. New Biol.* **232**:80.
31. VanBoxel, J. A., J. D. Stobo, W. E. Paul, and I. Green. 1972. Antibody-dependent lymphoid cell-mediated cytotoxicity: No requirement for thymus-derived lymphocytes. *Science (Wash. D.C.)*. **175**:194.
32. Britton, S., H. Perlmann, and P. Perlmann. 1973. In vitro studies of antibody formation and cell mediated cytotoxicity against heterologous erythrocytes in mice. *Cell. Immunol.* In press.
33. Lamon, E. W., H. M. Skurzak, E. Klein, and H. Wigzell. 1972. In vitro cytotoxicity by a nonthymus-processed lymphocyte population with specificity for a virally determined tumor cell surface antigen. *J. Exp. Med.* **136**:1072.
34. Chia, E., and H. Festenstein. 1973. Specific cytostatic effect of lymph node cells from normal and T cell deficient mice on syngeneic tumor target cells in vitro

- and its specific abrogation by syngeneic ascites fluid from tumor-bearing mice. *Eur. J. Immunol.* **3**:483.
35. Gelfand, E. W., K. Resch, and M. Prester. 1972. Antibody-mediated target cell lysis by non-immune cells. Characterization of the antibody and effector cell populations. *Eur. J. Immunol.* **2**:419.
 36. Henney, C. S., J. Clayburgh, G. A. Cole, and R. A. Prendergast. 1972. B-lymphocyte mediated cytotoxicity: a complement independent phenomenon. *Immunol. Commun.* **1**:93.
 37. Lamon, E. W., H. Wigzell, E. Klein, B. Andersson, and H. M. Skurzak. 1973. The lymphocyte response to primary moloney sarcoma virus tumors in Balb/c mice: definition of the active subpopulations at different times after infection. *J. Exp. Med.* **137**:1472.
 38. Bennett, B. 1965. Specific suppression of tumor growth by isolated peritoneal macrophages from immunized mice. *J. Immunol.* **95**:656.
 39. Evans, R., and P. Alexander. 1970. Cooperation of immune lymphoid cells with macrophages in tumor immunity. *Nature (Lond.)*. **228**:620.
 40. Lohmann-Matthes, M.-L., H. Schipper, and H. Fischer. 1972. Macrophage-mediated cytotoxicity against allogeneic target cells in vitro. *Eur. J. Immunol.* **2**:45.
 41. Holm, G., and S. Hammarström. 1973. Haemolytic activity of human blood monocytes. Lysis of human erythrocytes treated with anti-A serum. *Clin. Exp. Immunol.* **13**:29.
 42. MacDonald, H. R. 1972. Ph.D. Thesis. University of Toronto.
 43. Allison, A. C. 1972. Immunity and Immunopathology in virus infections. *Ann. Inst. Pasteur* **123**:585.
 44. Wislöf, F., and S. S. Fröland. 1973. Antibody-dependent lymphocyte-mediated cytotoxicity in man: No requirement for lymphocytes with membrane-bound immunoglobulin. *Scand. J. Immunol.* **2**:151.
 45. Perlmann, P., H. Wigzell, P. Golstein, E. W. Lamon, Å. Larsson, C. O'Toole, H. Perlmann, and E. A. J. Svedmyr. 1973. Cell-mediated cytotoxicity in vitro. Analysis of active lymphocyte subpopulations in different experimental systems. *In Immunopathology Symposium. Advances in Biosciences*, Pergamon Press Inc., Elmsford, N.Y. II: In press.
 46. Basten, A., N. L. Warner, and T. Mandel. 1972. A receptor for antibody on B lymphocytes. II. Immunochemical and electron microscopic characteristics. *J. Exp. Med.* **135**:627.
 47. Huber, H., S. D. Douglas, J. Nusbacher, S. Kochwa, and R. E. Rosenfield. 1971. IgG subclass specificity of human monocyte receptor sites. *Nature (Lond.)* **229**:491.
 48. MacLennan, I. C. M., A. Howard, F. M. Gotch, and P. G. Quie. 1973. Effector activating determinants on IgG. I. The distribution and factors influencing the display of complement neutrophil and cytotoxic B cell determinants on IgG subclasses. *Immunology*. In press.
 49. Calder, E. A., W. J. Penhale, E. W. Barnes, and W. J. Irvine. 1973. Cytotoxic lymphocytes in Hashimoto Thyroiditis. An in vitro assay system using ⁵¹Cr-labelled chicken erythrocytes coated with thyroglobulin. *Clin. Exp. Immunol.* **14**:19.

50. Paul, W. E. 1970. Functional specificity of antigen-binding receptors of lymphocytes. *Transplant. Rev.* **5**:130.
51. MacLennan, I. C. M., and B. Harding. 1970. The role of immunoglobulins in lymphocyte-mediated cell damage in vitro. II. The mechanism of target cell damage by lymphoid cells from immunized rats. *Immunology* **18**:405.
52. MacLennan, I. C. M., C. Loewi, and B. Harding. 1970. The role of immunoglobulins in lymphocyte-mediated cell damage in vitro. I. Comparison of the effects of target cell specific antibody and normal serum factors on cellular damage by immune and non-immune lymphocytes. *Immunology*. **18**:397.
53. Carlsson, H. E., S. Hammarström, M. Lanteli, and P. Perlmann. 1971. Antibody-induced destruction of colon antigen-coated chicken erythrocytes by normal human lymphocytes. *Eur. J. Immunol.* **1**:281.
54. Hoy, W. E., and D. S. Nelson. 1969. *Aust. J. Exp. Biol. Med. Sci.* **47**:525.