

In the article entitled "Streptococcal M6 protein expressed in *Escherichia coli*: localization, purification, and comparison with streptococcal-derived M protein" by Fischetti et al. (April 1984, 159:1083), editorial and printing errors in Tables II-IV caused the terms "LysM6" and "ColiM6" to appear incorrectly. Tables II-IV, with appropriate corrections, appear below:

TABLE II
Comparison of the Amino Acid Composition of LysM6 and ColiM6 Proteins

| Amino acid | Residues/100 | |
|------------|--------------|---------|
| | LysM6* | ColiM6‡ |
| GLU | 20.3 | 20.1 |
| LYS | 17.4 | 16.9 |
| ASP | 13.9 | 14.0 |
| LEU | 12.3 | 11.3 |
| ALA | 10.4 | 9.9 |
| THR | 6.7 | 7.6 |
| GLY | 3.4 | 3.9 |
| ILE | 3.0 | 2.8 |
| ARG | 4.2 | 3.9 |
| SER | 3.4 | 3.2 |
| VAL | 3.0 | 3.3 |
| PRO | 0.8 | 1.4 |
| MET | 0.3 | 0.6 |
| TYR | 0.2 | 0.2 |
| PHE | 0.1 | 0.3 |
| HIS | —§ | — |
| CYS | — | — |
| TRP | — | — |

* Average of two determinations.

‡ Average of three determinations from two different lots.

§ No detectable amounts.

(Correction continued)