

CLONAL DELETION AND CLONAL ANERGY IN
THE THYMUS INDUCED BY CELLULAR ELEMENTS
WITH DIFFERENT RADIATION SENSITIVITIES

BY JOSEPH L. ROBERTS, SUSAN O. SHARROW, AND ALFRED SINGER

*From the Experimental Immunology Branch, National Cancer Institute,
National Institutes of Health, Bethesda, Maryland 20892*

Self-tolerance is an essential characteristic of the T cell repertoire, and results initially from negative selection processes that developing T cells undergo during differentiation in the thymus. Potentially autoreactive thymocytes expressing anti-self TCR specificities encounter self antigens in the thymus, and the signals resulting from engagement of their TCRs are thought to abort their differentiation, either by signaling the cells to die (clonal deletion) or by signaling the cells to become anergic (clonal inactivation). Using anti-V β mAbs specific for TCRs reactive against various self antigens, it has been possible to determine the fate of potentially autoreactive T cells as they differentiate in the thymus. Thus, it was found that TCR^{hi} V β 17a⁺ thymocytes are deleted in IE⁺ mice (1), and V β 6⁺ thymocytes are deleted in Mls^a mice (2). In the present study, we show that clonal deletion is not the only mechanism by which Mls^a- and IE-specific tolerance can be induced in the thymus, as clonal inactivation of developing V β 6⁺ and V β 17a⁺ T cells can also be induced, but the alternative tolerance mechanisms are mediated by cellular elements with different sensitivities to γ irradiation.

Material and Methods

Experimental Animals. Radiation bone marrow chimeras are designated as bone marrow donor \rightarrow irradiated recipient, and were constructed by injecting 1.5×10^7 T-depleted bone marrow cells into 950-rad γ -irradiated recipients. Chimeras were examined no earlier than 5 wk after irradiation and bone marrow reconstitution. At that time, no cells of host origin were detectable in the thymi of these chimeras, but spleen cell populations did contain <5% radiation-resistant host cells that were essentially all T cells, so that purified populations of spleen T cells were contaminated with, on average, 33.6% host cells.

Flow Cytometry (FCM). Cells were stained as indicated and samples were analyzed on a modified dual laser (488 nm, 590 nm) FACS II (Becton Dickinson Immunocytometry Systems, Mountain View, CA). Fluorescence data were collected using three-decade logarithmic amplification on viable cells as determined by forward light scatter intensity and propidium iodide exclusion.

T Cell Populations. Purified spleen T cells for FCM were obtained by passage over either nylon wool columns or anti-Ig plates. For functional studies, purified SJL (Ly5.1⁺) spleen

J. L. Roberts is a Ph.D. candidate in the Department of Microbiology and Immunology, Duke University, Durham, NC 27710.

Address correspondence to J. L. Roberts, Experimental Immunology Branch, National Cancer Institute, Building 10, Room 4B-17, National Institutes of Health, Bethesda, MD 20892.

T cells were obtained from chimeric animals by coating spleen cells with anti-Ly5.2 mAb, incubating them on anti-Ig plates, and collecting the nonadherent cells. The resultant cell populations were >80% Thy-1.2⁺ and >99.7% Ly5.1⁺.

Proliferation Assays. Triplicate cultures of responder T cells were stimulated with either mitomycin C-treated cells or anti-TCR mAb as indicated, and then pulsed with 1 μ Ci [³H]thymidine for 8–12 h before harvesting.

Results and Discussion

As previously reported, thymus and spleen T cell populations in SJL mice contain significant numbers of TCR^{hi} V β 6⁺ and V β 17a⁺ T cells (1), whereas T cell populations from SJL \times CBA/J mice expressing Mls^a and IE^k determinants do not (Fig. 1, Table I). To track the fate of developing V β 6⁺ and V β 17a⁺ thymocytes encountering Mls^a and IE^k determinants on either radiation-resistant or radiation-sensitive cellular elements, we constructed experimental animals by injecting SJL bone marrow stem cells into irradiated hosts of various genotypes (Fig. 1, Table I). Because only SJL cells express the Ly5.1 allelic marker, we were able to focus exclusively on SJL T cells in each experimental animal. TCR^{hi} V β 6⁺ and V β 17a⁺ SJL T cells were present in both the thymus and spleen of SJL \rightarrow B10 chimeras in which neither Mls^a nor IE determinants were expressed. Differences in frequency of V β 17a⁺ SJL T cells maturing in a normal SJL thymus vs. the chimeric B10 thymus (Table I) are consistent with the role of K^s as a positive selecting element for V β 17a⁺ T cells (3). In contrast, few if any TCR^{hi} V β 6⁺ and V β 17a⁺ SJL T cells were present in SJL + B6 \times CBA/J \rightarrow B10 experimental animals that were identical to SJL \rightarrow B10 animals, except for the injection of additional unirradiated B6 \times CBA/J (Mls^aIE^k) bone marrow cells, demonstrating that expression of Mls^a and IE^k de-

TABLE I
Development of V β 6⁺ and V β 17a⁺ T Cells
in Experimental Mice

Experimental animal	T cell source	Percent TCR ^{hi} SJL T cells expressing*:	
		V β 6	V β 17a
SJL	Thymus	9.6 \pm 0.3	8.4 \pm 0.4
	Spleen	11.5 \pm 0.2	10.5 \pm 0.2
SJL \times CBA/J	Thymus	0.9 \pm 0.05	0.4 \pm 0.1
	Spleen	0.2 \pm 0.03	0.8 \pm 0.6
SJL \rightarrow B10	Thymus	13.5 \pm 0.3	5.2 \pm 0.2
	Spleen	13.1 \pm 1.3	5.7 \pm 0.2
SJL \rightarrow B6 \times CBA/J	Thymus	11.6 \pm 0.8	6.3 \pm 0.5
	Spleen	4.5 \pm 1.0	3.2 \pm 0.6
SJL + B6 \times CBA/J \rightarrow B10	Thymus	1.8 \pm 0.2	1.5 \pm 0.2
	Spleen	1.0 \pm 0.2	1.0 \pm 0.3

* Two-color immunofluorescence staining with anti-Ly5.1 mAb in red and anti-TCR (CD3, V β 6, and V β 17a) was used to identify T cells of donor SJL origin. Values are V β ^{hi} SJL T cells as a percentage of CD3^{hi} SJL cells \pm SE for no fewer than four animals. T cells were defined as TCR^{hi} as described in Fig. 1.




FIGURE 1. Expression of CD3, V β 6, and V β 17a on thymocytes of SJL origin. One-color histograms of CD3, V β 6, and V β 17a expression on Ly5.1⁺ (SJL) thymocytes from the indicated experimental animals were obtained by two-color FCM after staining with anti-Ly5.1 vs. anti-CD3, anti-V β 6 (7), and anti-V β 17a (1). Fluorescence data were collected on 5–10 \times 10⁴ viable cells. Anti-TCR fluorescence data were plotted in solid lines as fluorescence intensity vs. cell number. Dashed lines represent background staining with FITC-conjugated negative control antibodies. The vertical dashed lines define the demarcation we used for defining TCR^{hi} T cells.

TABLE II
Developmental Phenotype of SJL T Cells from Experimental Mice

Thymocytes	TCR	CD4/CD8 and Qa2 phenotypes of TCR ^{hi} thymocytes*					
		4 ⁺ 8 ⁻	4 ⁻ 8 ⁺	4 ⁺ 8 ⁺	4 ⁻ 8 ⁻	Qa2 ⁺	Qa2 ⁻
	%						
SJL	CD3 (100) [†]	65.2	20.5	12.0	2.1		
SJL → B6 × CBA/J	CD3 (100)	73.8	14.3	8.5	2.7		
SJL	V β 6 (11.0)	49.9	29.9	18.0	2.0	22.8	77.2
SJL → B6 × CBA/J	V β 6 (11.8)	73.0	14.6	10.6	1.6	34.5	65.4
SJL	V β 17a (10.1)	71.7	14.6	10.2	3.3		
SJL → B6 × CBA/J	V β 17a (7.3)	79.4	8.6	9.8	2.0		

* Three-color FCM was used to assess CD4/CD8 expression on V β 6⁺ and V β 17a⁺ SJL thymocytes. Data were collected list mode and software gated to select V β 6^{hi} or V β 17a^{hi} cells. Two-color FCM was used to assess V β 6⁺ SJL thymocytes for Qa2 expression. Thymocytes were stained for V β 6 and counterstained for Qa2 with 695H1-1-2 mAb. Values represent the percentage of V β 6^{hi} thymocytes that are Qa2⁺ or Qa2⁻.

[†] Numbers in parentheses indicate percentage of V β 6^{hi} SJL thymocytes as a percentage of CD3^{hi} SJL thymocytes.

terminants on unirradiated bone marrow-derived cells (e.g., dendritic cells) is able to delete developing V β 6⁺ and V β 17a⁺ T cells (1). We next examined T cell populations from SJL → B6 × CBA/J animals in which Mls^a and IE^k determinants were expressed only on radiation-resistant host elements. Surprisingly, V β 6⁺ and V β 17a⁺ SJL T cells were present in SJL → B6 × CBA/J animals, indicating that radiation-resistant cellular elements such as thymic epithelium fail to delete developing V β 6⁺ and V β 17a⁺ thymocytes. Table I summarizes the frequencies of V β 6⁺ and V β 17a⁺ T cells observed in the thymi and spleens of all the experimental mice tested.

The failure of radiation-resistant B6 × CBA/J host cells to delete V β 6⁺ and V β 17a⁺ SJL T cells in SJL → B6 × CBA/J animals might have resulted in a failure

of the irradiated host to induce Mls^a- and IE^k-specific tolerance. To assess this possibility, purified Ly5.1⁺ (SJL) T cell populations from experimental animals were assessed for their proliferative responses against stimulator cells expressing third-party (DBA/2), IE^k (B10.BR), or Mls^aIE^k (CBA/J) alloantigens. It can be seen in Table III that SJL T cell populations from SJL → B6 × CBA/J as well as SJL + B6 × CBA/J → B10 animals were functionally tolerant to both Mls^a and IE^k, even though Vβ6⁺ and Vβ17a⁺ T cells were present in the former mice. To understand how Vβ6⁺ and Vβ17a⁺ T cells present in SJL → B6 × CBA/J mice could fail to react against Mls^a- and IE^k-bearing stimulator cells, we attempted to stimulate them by crosslinking their TCRs directly with anti-Vβ6 and anti-Vβ17a mAbs (Table III). To maximize responses, the cultures contained syngeneic accessory cells as well as exogenous T cell growth factors (Table III). In contrast to Mls^a- and IE^k-responsive Vβ6⁺ and Vβ17a⁺ SJL T cells from normal mice, tolerant Vβ6⁺ and Vβ17a⁺ SJL

TABLE III
Proliferative Responses of T cells from Experimental Mice

Experimental animal	T cell source*	Strain of stimulator cells			
		SJL (H-2 ^s , Mls ^c)	DBA/2 (H-2 ^d , Mls ^a)	B10.BR (H-2 ^k , Mls ^b)	CBA/J (H-2 ^k , Mls ^a)
<i>cpm × 10⁻³</i>					
SJL → B10	Thymus	0 ± 0.1	61.9 ± 1.4	16.5 ± 2.2	29.8 ± 0.9
	Spleen	0.2 ± 0.1	32.5 ± 0.8	2.0 ± 0.1	76.5 ± 7.2
SJL → B6 × CBA/J	Thymus	0 ± 1.5	18.2 ± 1.2	0 ± 1.3	0 ± 1.0
	Spleen	1.7 ± 0.2	18.0 ± 1.7	0.5 ± 0.1	4.0 ± 0.5
SJL + B6 × CBA/J → B10	Thymus	0 ± 0.2	23.4 ± 1.3	0 ± 0.2	1.4 ± 0.4
	Spleen	0.8 ± 0.2	11.3 ± 1.5	0.1 ± 0.2	1.9 ± 0.5
SJL	Thymus	2.5 ± 1.3	41.6 ± 2.6	24.5 ± 0.8	27.2 ± 1.6
	Spleen	1.1 ± 0.1	39.9 ± 2.4	8.6 ± 2.1	86.3 ± 1.4
B10.BR	Thymus	7.8 ± 0.6	31.1 ± 2.5	0 ± 0.1	28.9 ± 3.1
	Spleen	14.5 ± 1.7	54.9 ± 2.6	0.3 ± 0.2	83.3 ± 8.8
Specificity of stimulating mAb [†]					
SJL → B6 × CBA/J	Thymus	Vβ6: 1.8 ± 0.5	Vβ17a: 0.4 ± 0.5	α/β: 45.0 ± 5.2	CD3: 189.5 ± 33.1
	Spleen	0.1 ± 0.1	0 ± 1.0		87.4 ± 2.6
SJL	Thymus	14.9 ± 1.1	16.5 ± 1.6		142.2 ± 3.5
	Spleen	30.9 ± 2.7	34.7 ± 2.6	22.1 ± 1.9	49.0 ± 3.1
CBA/J	Thymus	0.5 ± 3.3	0 ± 2.2		371.0 ± 22.4
	Spleen	0 ± 2.1	0 ± 1.4	15.5 ± 2.3	289.1 ± 6.8

* Responder thymocytes (10⁶) and spleen T cells (5 × 10⁴) from individual animals were cultured with 5 × 10⁵ stimulator cells. Values are the mean ± SE of cultures containing stimulator cells minus the mean ± SE of cultures without stimulator cells. Results are representative of three experiments.

† 5 × 10⁵ responder thymocytes or 10⁵ responder Ly5.1⁺ spleen T cells were cultured with 25% culture supernatant from RR4-7 anti-Vβ6 (7), KJ23a anti-Vβ17a (1), H57-597 anti-TCR-α/β (8), or 145-2C11 anti-CD3. Mitomycin C-treated syngeneic spleen cells were added as FCR⁺ accessory cells. Cultures also were supplemented with exogenous T cell growth factors in the form of 25% Con A supernatant. Values represent the mean ± SE of cultures containing stimulating mAb minus the mean ± SE of cultures without stimulating mAb. Results are representative of three experiments.

T cells from SJL \rightarrow B6 \times CBA/J mice failed to proliferate in response to direct TCR engagement by either anti-V β 6 or anti-V β 17a mAbs, indicating that the undeleted T cells were anergic (Table III). Indeed, clonal anergy of undeleted but tolerant T cells developing in SJL \rightarrow B6 \times CBA/J mice could explain why it is only in these mice that the relative frequency of V β 6⁺ and V β 17a⁺ T cells is significantly lower among spleen T cells than thymic T cells (Table I), since anergic T cells would fail to clonally expand in the periphery in response to environmental antigens with a resultant decrease in their relative frequency.

We next assessed how far anergic V β 6⁺ and V β 17a⁺ thymocytes could differentiate in SJL \rightarrow B6 \times CBA/J thymi by phenotyping them for CD4/CD8 expression using three-color FCM. We found that the distribution among various CD4/CD8 thymus subpopulations of anergic SJL \rightarrow B6 \times CBA/J thymocytes resembled that of normal SJL thymocytes, with normal numbers of phenotypically mature CD4⁺8⁻ and CD4⁻8⁺ cells (Table II). Nevertheless, because Qa2 has been reported to be expressed only on functionally competent single-positive thymocytes and peripheral T cells (4), we thought that anergic V β 6⁺ thymocytes from SJL \rightarrow B6 \times CBA/J mice might have failed to differentiate into Qa2⁺ cells, but such was not the case (Table II). Thus, there was no identifiable differentiation step that the anergic thymocytes had failed to undergo, even though they apparently could not proliferate in response to TCR engagement. While the role of TCR signaling in driving either thymocyte maturation or proliferation is uncertain, it is likely to be involved in selective events occurring in the thymus. Consequently, we wanted to determine if the anergic thymocytes were at all responsive to TCR-mediated signals. Indeed, as has been observed in anergic T cell clones (5), we observed that anergic V β 6⁺ thymocytes do respond to TCR crosslinking by increasing their expression of IL-2-Rs (Fig. 2).

From the present study, it is clear that TCR engagement on developing thymocytes does not necessarily lead to clonal deletion. In fact, clonal deletion seems to require TCR engagement by immature T cells of self antigens on a specialized subpopulation of radiation-sensitive bone marrow-derived cells (probably dendritic cells), whereas TCR engagement of self antigens on other cells, possibly including thymic epithelium, induces clonal anergy. These results should help clarify conflicting reports in which T cell tolerance induced during development variably led to clonal deletion (1, 2, 6). Thus, TCR engagement without a competent second signal from bone




FIGURE 2. Increased IL-2-R expression on V β 6⁺ thymocytes from SJL \rightarrow B6 \times CBA/J after anti-TCR stimulation. Thymocytes were cultured for 16 h without exogenous growth factors at 37°C in plates coated with 5 μ g/ml purified anti-CD3 (a), 50 μ g/ml anti- α/β (H57-597) (b), or no mAb (c). Cells

were assessed by two-color FCM by staining for V β 6 and counterstaining for IL-2-R with 7D4 mAb. Software gating was used to select V β 6⁺ cells for analysis of IL-2-R expression. Because TCR stimulation causes downmodulation of cell surface TCR expression such that stimulated TCR^{hi} cells appear TCR^{lo}, the gates included both V β 6^{hi} and V β 6^{lo} cells. Single-color histograms depict the level of IL-2-R expression on stimulated V β 6⁺ SJL thymocytes from SJL \rightarrow B6 \times CBA/J mice.

marrow-derived APC may lead to clonal anergy in developing T cells as it does in mature T cell populations (5). The signals inducing clonal deletion in immature thymocytes, as well as the signals driving the differentiation of anergic thymocytes into phenotypic maturity, remain to be identified.

Summary

The present study demonstrates that immune tolerance can be achieved in the thymus both by clonal deletion and by clonal inactivation, but that the two tolerant states are induced by cellular elements with different radiation sensitivities. TCR engagement of self antigens on bone marrow-derived, radiation-sensitive (presumably dendritic) cells induces clonal deletion of developing thymocytes, whereas TCR engagement of self antigens on radiation-resistant cellular elements, such as thymic epithelium, induces clonal anergy. The nondeleted, anergic thymocytes can express IL-2-Rs but are unable to proliferate in response to either specific antigen or anti-TCR antibodies, and do develop into phenotypically mature cells that emigrate out of the thymus and into the periphery.

Received for publication 18 October 1989 and in revised form 18 December 1989.

References

1. Marrack, P., D. Lo, R. Brinster, R. Palmiter, L. Burkly, R. H. Flavell, and J. Kappler. 1988. The effect of thymus environment on T cell development and tolerance. *Cell*. 53:627.
2. MacDonald, H. R., R. Schneider, R. K. Lees, R. C. Howe, H. Acha-Orbea, H. Festenstein, R. M. Zinkernagel, and H. Hengartner. 1988. T-cell receptor V β use predicts reactivity and tolerance to Mls^a-encoded antigens. *Nature (Lond.)*. 332:40.
3. Zuniga-Pflucker, J. C., D. L. Longo, and A. M. Kruisbeek. 1989. Positive selection of CD4⁻CD8⁺ T cells in the thymus of normal mice. *Nature (Lond.)*. 338:76.
4. Rabinowitz, R., S. O. Sharrow, S. Chatterjee-Das, M. J. Rogers, and D. H. Sachs. 1986. Qa alloantigen expression on functional T lymphocytes from spleen and thymus. *Immunogenetics*. 24:391.
5. Jenkins, M. K., D. M. Pardoll, J. Mizuguchi, T. M. Chused, and R. H. Schwartz. 1987. Molecular events in the induction of a nonresponsive state in interleukin 2-producing helper T-lymphocyte clones. *Proc. Natl. Acad. Sci. USA*. 84:5409.
6. Lo, D., L. C. Burkly, R. A. Flavell, R. D. Palmiter, and R. L. Brinster. 1989. Tolerance in transgenic mice expressing class II major histocompatibility complex on pancreatic acinar cells. *J. Exp. Med.* 170:87.
7. Kanagawa, O., E. Palmer, and J. Bill. 1989. The T cell receptor V β 6 domain imparts reactivity to the Mls-1^a antigen. *Cell. Immunol.* 119:412.
8. Kubo, R. T., W. Born, J. W. Kappler, P. Marrack, and M. Pigeon. 1989. Characterization of a monoclonal antibody which detects all murine $\alpha\beta$ T cell receptors. *J. Immunol.* 142:2736.