

# The Mouse Vitronectin Receptor Is a T Cell Activation Antigen

By Kevin Moulder,\* Kevan Roberts,† Ethan M. Shevach,‡ and John E. Coligan\*

From the \*Biological Resources Branch and †Laboratory of Immunology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20892

## Summary

In this report, we demonstrate that the T cell activation antigen, recognized by monoclonal antibody H9.2B8, is the murine homologue of the vitronectin receptor (VNR) and, thereby, we provide initial evidence that VNR is expressed on lymphoid cells. VNR is expressed on a variety of T cell lines, tumors, and Con A-activated splenocytes, but not resting T cells, and is capable of binding to the extracellular matrix proteins fibronectin, fibrinogen, and vitronectin, via the tripeptide sequence RGD. There was no evidence of novel  $\beta$  chains pairing with the VNR  $\alpha$  chain, as has been demonstrated in some human cells. In view of recent studies demonstrating that this molecule functions as an accessory molecule in T cell activation, the VNR may play an important role in mouse T cell functions.

**I**ntegrins comprise a complex family of cell adhesion molecules involved in cell-cell and cell-extracellular matrix protein (ECMP)<sup>1</sup> interactions (1). At least three subfamilies have been defined on the basis of the association of any of a number of  $\alpha$  chains with a common  $\beta$  chain (2). We have previously characterized two mAbs (H9.2B8 and 8.18E12) that identified a novel integrin expressed on murine  $\gamma/\delta$  and  $\alpha/\beta$  T cell lines and hybridomas as well as on T cells activated for prolonged periods of time (7–21 d) by mitogens or alloantigens. This integrin mediated adhesion to fibronectin, vitronectin, and fibrinogen, and the two mAbs in combination, as well as the tetrapeptide RGDS alone, could inhibit the binding of cell lines to these ECMP (3). In the present report, we make use of a number of antisera to the vitronectin receptor (VNR) and demonstrate that this novel integrin is the murine homologue of the human VNR, thereby demonstrating for the first time that the VNR is present on mouse T lymphocytes and that the expression of this  $\beta 3$  integrin can be upregulated by activation.

## Materials and Methods

**Cell Lines.** Dendritic epidermal T cell (DETC) line T195 and its corresponding hybridoma (T195/BW) derived from fusion to BW5147 have been previously described (4–6).

**Antibodies.** mAb H9.2B8 is a hamster mAb obtained after immunizations with mouse DETC lines (3). Rabbit anti-human VNR

serum was purchased from Telios Pharmaceuticals, Inc. (La Jolla, CA). Rabbit antiserum to human GpIIIa (anti- $\beta 3$  chain) was a gift from Dr. R. McKeever (Oklahoma University, Norman, OK) and has previously been shown to react with mouse  $\beta 3$  integrin chains (7). Clone M18/2.a.8, secreting rat IgG reactive with mouse  $\beta 2$  integrin chains (8), was purchased from the American Type Culture Collection (Rockville, MD). Rat mAb to mouse LFA-1 (from clone M17/5.2) was purchased from Boehringer Mannheim Biochemicals (Indianapolis, IN). mAb H1.2F3, which is reactive with a T cell activation antigen, has been described previously (9).

**Antipeptide Sera.** Rabbit anti-peptide sera against the COOH-terminal sequences of human  $\beta 1$  (anti- $\beta 1$ ) (NPKYEGK) (10) and VNR  $\alpha$  chain (anti-VNR $\alpha$ ) (CEQLQPHENGEGNSET) (11) were prepared as previously described (12).

**Con A Stimulation of Splenocytes.** Con A-activated spleen cells from 8-wk-old female C3H/HeN mice were prepared as previously described (3).

**Radioiodination and Immunoprecipitation Analysis.** Cells were surface labeled with <sup>125</sup>I using the previously described lactoperoxidase method (13). Immunoprecipitations and immunodepletion experiments were performed as previously described (3). Precipitates were analyzed on SDS-PAGE (7% acrylamide) slab gels in the presence of 5% 2-ME.

**Two-dimensional Electrophoresis.** Nonequilibrium pH gradient electrophoresis (NEPHGE) was performed according to O'Farrell et al. (14) under nonreducing conditions using pH 4–6.5 ampholines (Pharmacia Fine Chemicals, Piscataway, NJ) in the first dimension and SDS-PAGE (7% acrylamide) in the second dimension.

**Ligand Binding.** Rat fibronectin (Calbiochem-Behring Corp., La Jolla, CA), rat fibrinogen (Sigma Chemical Co., St. Louis, MO), human vitronectin (Calbiochem-Behring Corp.), and rat collagen I (Sigma Chemical Co.) were coupled to CNBr-activated Sepharose 4B (Pharmacia Fine Chemicals). Peptide GRGDSPC (Telios Pharmaceuticals, Inc.) was coupled to thiopropyl-activated Sepharose

<sup>1</sup> Abbreviations used in this paper: DETC, dendritic epidermal T cells; ECMP, extracellular matrix proteins; NRS, normal rabbit serum; VNR, vitronectin receptor.

4B (Pharmacia Fine Chemicals). Magnetic laminin agarose was a gift from Advanced Magnetics (Cambridge, MA). Precipitation experiments were performed by incubating 250  $\mu$ l packed beads with 100  $\mu$ l cell lysate for 60 min room temperature. The beads were washed in 0.5% NP-40 buffer (3) and integrins eluted with RGDS peptide (2 mg/ml) in 0.5% NP-40 buffer for 60 min. Immunoprecipitations using anti-VNR $\alpha$  sera were performed with the eluate as previously described (3).

**Adhesion Assay.** 96-well plates (Costar, Cambridge, MA) were coated with vitronectin (2.0  $\mu$ g/ml) by incubating wells overnight at 37°C and then washing three times with binding buffer (PBS supplemented with 2.5 mg/ml BSA [Sigma Chemical Co.]). Remaining sites were blocked by adding 200  $\mu$ l binding buffer and incubating for 2 h at 37°C.


Resting T cells and Con A-activated lymphoblasts (10<sup>7</sup> cells/ml) were labeled with 300  $\mu$ Ci <sup>51</sup>Cr (Amersham Corp., Arlington Heights, IL) for 1 h at 37°C. The cells were then washed with HBSS and resuspended in 10 ml RPMI supplemented with 10% FCS for 30 min. After two washes, the cells were resuspended (5  $\times$  10<sup>5</sup>/ml) in binding buffer. The cell suspension (100  $\mu$ l) was added to plates precoated with human vitronectin (Telios Pharmaceuticals, Inc.). Adhesion was evaluated in the presence of RGDS or RGE8 (Peninsula Laboratories, Inc., Belmont, CA) or mAbs by adding these agents directly to the wells (100  $\mu$ l). After a 2-h incubation at 37°C, the wells were washed three times with binding buffer. The level of adhesion was subsequently evaluated by the addition of 200  $\mu$ l of 0.5% Triton X-100 and counting the lysate in a gamma counter. The percent cell adhesion is expressed as a percentage of the total counts per minute added to the well. Background binding represents the percent adhesion to plates coated with albumin alone.

## Results and Discussion


**mAb H9.2B8 Recognizes Mouse VNR.** We have previously shown with DETC cells that mAbs H9.2B8 and 8.18E12 recognize the  $\alpha$  chain of an integrin-like heterodimer of 95

kD ( $\beta$  chain) and 140 kD ( $\alpha$  chain) under nonreducing conditions, and that, upon reduction, the  $\alpha$  chain is cleaved into 116- and 24-kD components (3). To analyze the relationship of this heterodimer to other integrins, we performed a series of sequential immunoprecipitations. Cell lysates (T195 or T195/BW) were precleared of material reactive with antisera to  $\beta$ 3 or VNR $\alpha$  chains or, as a negative control, normal rabbit serum (NRS). These precleared lysates were then immunoprecipitated with various integrin-reactive antisera followed by analysis on SDS-PAGE under reducing conditions (Fig. 1). After preclearing material reactive with NRS, anti- $\beta$ 1 serum precipitated a heterodimer with an  $\alpha$  chain of 140 kD (intense band) and a  $\beta$  chain of 135 kD (faint band) (Fig. 1, lane 2). Antibody to integrin  $\beta$ 2 chains precipitated heterodimers of 98 kD ( $\beta$  chain) and 180 kD ( $\alpha$  chain) (Fig. 1, lane 3) that were identical in size to the chains precipitated by a mAb reactive with the  $\alpha$  chain of LFA-1 (Fig. 1, lane 4). Three antisera to the VNR were used. Anti- $\beta$ 3 and anti-VNR $\alpha$  chain sera, as well as antiserum to purified human VNR, all precipitated a heterodimer with a  $\beta$  chain of 98 kD and an  $\alpha$  chain of 116 kD (Fig. 1, lanes 5, 6, and 8). The 24-kD component of the  $\alpha$  chain can be visualized on 12% acrylamide gels (not shown). The molecular sizes of the component VNR chains were similar to those of the integrin recognized by mAb H9.2B8 (Fig. 1, lane 7).

To determine if mAb H9.2B8 recognizes VNR, T195/BW cell lysates were cleared of material reactive with antisera to  $\beta$ 3 chains or VNR $\alpha$  chains by repeated immunoprecipitations. Subsequent precipitations revealed that both anti- $\beta$ 3 and anti-VNR $\alpha$  sera cleared the integrin recognized by mAb H9.2B8 (Fig. 1, B and C), thus proving that mAb H9.2B8 recognizes the VNR. Other immunodepletion experiments demonstrated that mAb H9.2B8 precleared the material recognized by anti-VNR sera (data not shown). The process of



**Figure 1.** mAb H9.2B8 recognizes the vitronectin receptor. Lysates of <sup>125</sup>I cell surface-labeled T195/BW were precleared by serial immunoprecipitations with antiserum to the  $\beta$ 3 chain or the VNR $\alpha$  chain, or as a negative control, NRS. Precleared cell lysates were immunoprecipitated with NRS, anti- $\beta$ 1 chain peptide serum, mAb M18/2.a.8 to the  $\beta$ 2 chain, mAb M17/5.2 to LFA-1  $\alpha$  chain, anti- $\beta$ 3 chain serum, anti-VNR $\alpha$  serum, anti-human VNR serum, and mAb H9.2B8. Precipitates were analyzed by SDS-PAGE with 7% acrylamide under reducing conditions. The position of prestained molecular mass markers is indicated on the left. The bands below the 98-kD molecular mass marker in lanes 2, 3, 4, 14, and 20 are background bands as evidenced by their presence in lane 1. Identical results were obtained with the parent T195 DETC cell line.



**Figure 2.** VNR is expressed by activated T cells. Spleen cells from C3H/HeN mice were grown in the presence of Con A (3  $\mu\text{g}/\text{ml}$ ) for 4 d and IL-2 (20 U/ml) for a further 10 d. These, and fresh spleen cells as nonactivated controls, were surface labeled with  $^{125}\text{I}$  and cell lysates immunoprecipitated with NRS, mAb M18/2.a.8 to  $\beta 2$  chain, anti- $\beta 3$  chain serum, anti-VNR $\alpha$  serum, and anti-human VNR serum. Immunoprecipitates were analyzed by SDS-PAGE with 7% acrylamide under reducing conditions. The position of prestained molecular mass markers is indicated on the right. The cells that proliferated in response to Con A were found by FACS analysis to be >95% Thy-1 $^{+}$ .

immunodepletion did not interfere with subsequent immunoprecipitations, as  $\beta 1$  integrins (not shown) and LFA-1 (Fig. 1, lanes 14 and 20) could still be precipitated from precleared lysates.

**Con A-activated Splenocytes Express VNR.** We have previously shown by FACS $^{\circledR}$  analysis that the  $\alpha$  chain recognized by mAb H9.2B8 (here shown to be VNR $\alpha$ ) is present on a population of activated splenocytes (3). In view of the fact that VNR has not previously been reported to be expressed by mouse lymphoid cells, we examined the structure of the receptor expressed by Con A-activated T cells. As shown in Fig. 2, no material could be immunoprecipitated from lysates of resting lymphocytes with the anti-VNR $\alpha$  peptide serum, while the typical LFA-1 heterodimer was seen with the anti- $\beta 2$  precipitate (A). In contrast, when similar studies were performed on lysates of T cells after 14 d of activation with Con A, all three anti-VNR sera immunoprecipitated a 116- and 98-kD dimer (Fig. 2 B). To determine if the VNR expressed by the activated splenocytes was functional, quantitative cell attachment studies were performed. About 30% of splenic T cells activated in vitro with Con A and IL-2 adhered to wells coated with vitronectin, and this binding could be completely inhibited by mAb H9.2B8 and by the RGDS peptide, but not by the RGEs peptide or the control mAb H1.2F3 (Table I). No binding of resting T cells to vitronectin-coated plates could be detected.


Previous studies with human VNR have shown the VNR  $\alpha$  chain may associate with  $\beta$  chains other than  $\beta 3$ , namely  $\beta 1$  (15, 16),  $\beta x$  (now named  $\beta 5$ ) (17),  $\beta s$  (18), and  $\beta 3b$  (19). Some of these novel  $\beta$  chains are not easily distinguished by their mobilities in one-dimensional SDS-PAGE, but can be clearly distinguished from  $\beta 3$  chains by pI (17, 19). To examine whether some of the murine lymphoid VNR  $\alpha$  chains might be associated with chains other than  $\beta 3$ , we compared immunoprecipitations from T195/BW and Con A-activated splenocytes by NEPHGE analysis (Fig. 3). It can be seen that both T195/BW and Con A-activated splenocytes apparently express only a single heterodimer when examined after precipitation with anti- $\beta 3$  or anti-VNR $\alpha$  sera, indicating that the VNR expressed by Con A-activated splenocytes is composed solely of the VNR $\alpha$ / $\beta 3$  heterodimer.

**VNR Expression by Activated T Cells Bind Multiple ECMP.** Integrins have been shown to bind various ECMP (20), and the key amino acid sequence in many of these binding specificities has been shown to be the RGD motif (1). To determine the specificity of murine VNR, lysates from cells were reacted with ECMP- or RGD-containing peptide coupled to

**Table 1.** The Binding of Activated Murine T Cells to Vitronectin Is Inhibited by RGDS and mAb H9.2B8

|                     | Media | Inhibitor ( $\mu\text{g}/\text{ml}$ ) |     |      |      |        |     |        |      |
|---------------------|-------|---------------------------------------|-----|------|------|--------|-----|--------|------|
|                     |       | RGDS                                  |     | RGEs |      | H9.2B8 |     | H1.2F3 |      |
|                     |       | 100                                   | 10  | 100  | 10   | 50     | 5   | 50     | 5    |
| Day 10 lymphoblasts | 32.1  | 4.5                                   | 7.4 | 29.0 | 26.8 | 7.6    | 7.3 | 23.5   | 23.5 |
| Uncultured          | 7.9   | 4.7                                   | 5.9 | 5.5  | 4.8  | 5.5    | 4.5 | 5.9    | 4.7  |


The results are expressed as the percent cell adhesion. The background binding for the lymphoblasts was 7.3% and for the uncultured cells was 3.8%.



**Figure 3.** NEPHGE analysis of VNR. T195/BW and Con A-activated splenocytes were surface labeled with  $^{125}\text{I}$  and cell lysates immunoprecipitated with anti- $\beta 3$  serum or anti-VNR $\alpha$  serum. Immunoprecipitates were first separated by NEPHGE using pH 4–6.5 ampholines in the first dimension, then SDS-PAGE with 7% acrylamide under nonreducing conditions in the second dimension. The position of prestained molecular mass markers is indicated on the right.

Sephacose beads, eluted with RGDS peptide, and immunoprecipitated with anti-VNR $\alpha$  sera. The results of such analyses shown in Fig. 4 indicate that VNR from Con A-stimulated splenocytes bound to RGD, fibronectin, fibrinogen, and vitronectin. Binding to all ligands was blocked by the presence of RGDS peptide (data not shown). A similar binding pattern was seen with VNR expressed by T195/BW (Fig. 4).

Our previous studies (3), together with the results of this report of the mouse VNR, demonstrate that it is similar to the human VNR (21). Both integrins comprise an  $\alpha$  chain, which is cleaved upon reduction, and a  $\beta$  chain, which has an apparently larger molecular mass upon reduction than when nonreduced, indicative of intrachain disulfide bonds. The mouse VNR differs from the human VNR in that it is capable of binding to fibronectin in addition to fibrinogen and vitronectin. Binding to all three ECMP is mediated by the VNR $\alpha/\beta 3$  heterodimer, and we have not been able to demonstrate any other chains associated with VNR $\alpha$  or  $\beta 3$  that may contribute to this unique binding specificity. In the case of human cells, adherence to fibronectin is associated with VNR $\alpha$  paired with the  $\beta 1$  chain (15, 16) and the  $\beta x$  chain



**Figure 4.** Binding of VNR to extracellular matrix proteins. T195/BW and Con A-activated splenocytes were cell surface labeled with  $^{125}\text{I}$  and cell lysates precipitated with fibrinogen, fibronectin, vitronectin, laminin, or collagen bound to Sepharose 4B. Integrins were eluted with 2 mg/ml RGDS peptide and immunoprecipitated with NRS or anti-VNR $\alpha$  serum. Precipitates were analyzed at 7% SDS-PAGE under reducing conditions. The faint bands of 90, 130, and 170 kDa were only seen in precipitates of material eluted from extracellular matrix protein-coupled beads. Their identities are currently under investigation. The position of prestained molecular mass markers is indicated on the right.

(17, 22); however, the VNR $\alpha/\beta 1$  heterodimer does not mediate adherence to vitronectin, and the VNR $\alpha/\beta x$  heterodimer does not bind fibrinogen. One possible explanation for the difference is that the studies in man have characterized the VNR on nonlymphoid cells, whereas our studies in the mouse have characterized a VNR found on lymphoid cells.

Previous studies have shown that activation is required for integrin expression and/or function on T lymphocytes. Integrins that are expressed constitutively are only able to bind their ligand after T cell activation (20, 23–25), whereas integrins not normally expressed by resting T cells, such as VLA-1 and VLA-2, are induced after T cell activation. In this study, we have shown that only activated murine T cells express the VNR. These results differ from those of Klingemann and Dedhar (26), who demonstrated VNR expression on resting human T cells. Although a species difference in the regulation of expression of the VNR on lymphoid cells may exist, these latter studies were performed with polyclonal antisera to the human VNR and have not, as yet, been confirmed with mAbs.

The function of VNR on activated T lymphocytes is still unclear; however, it seems likely that VNR functions as a costimulatory molecule (27), as has been recently described for several other integrins (LFA-1, VLA-4, -5, -6) (23–25).

We thank Drs. Scott Wadsworth, Gary Kikuchi, and Thomas McConnell, and Mr. Jeffrey Marine for critical review of the manuscript. We appreciate the expert typing assistance of Ms. Brenda Rae Marshall.

Address correspondence to John E. Coligan, Building 4, Room 413, NIAID, National Institutes of Health, Bethesda, MD 20892.

Received for publication 25 June 1990 and in revised form 18 October 1990.

## References

1. Ruoslahti, E., and M.D. Pierschbacher. 1987. New perspectives in cell adhesion: RGD and integrins. *Science (Wash. DC)*. 238:491.
2. Hynes, R.O. 1987. Integrins: a family of cell surface receptors. *Cell*. 48:549.
3. Maxfield, S.R., K. Moulder, F. Koning, A. Elbe, G. Stingl, J.E. Coligan, E.M. Shevach, and W.M. Yokoyama. 1989. Murine T cells express a cell surface receptor for multiple extracellular matrix proteins: identification and characterization with monoclonal antibodies. *J. Exp. Med.* 169:2173.
4. Stingl, G., F. Koning, H. Yamada, W.M. Yokoyama, E. Tschachler, J.A. Bluestone, G. Steiner, L.E. Samelson, A.M. Lew, J.E. Coligan, and E.M. Shevach. 1987. Thy-1<sup>+</sup> dendritic epidermal cells express T3 and T cell receptor  $\gamma$  chain. *Proc. Natl. Acad. Sci. USA*. 84:4586.
5. Koning, F., G. Stingl, W.M. Yokoyama, H. Yamada, W.L. Maloy, E. Tschachler, E.M. Shevach, and J.E. Coligan. 1987. Identification of a T3-associated  $\gamma\delta$  T cell receptor on Thy-1<sup>+</sup> dendritic epidermal cell lines. *Science (Wash. DC)*. 236:834.
6. Yokoyama, W.M., F. Koning, G. Stingl, J.A. Bluestone, J.E. Coligan, and E.M. Shevach. 1987. Production of a T cell hybridoma that expresses the T cell receptor  $\gamma/\delta$  heterodimer. *J. Exp. Med.* 165:1725.
7. Ignatz, R.A., J. Heino, and J. Massague. 1989. Regulation of cell adhesion receptors by transforming growth factor  $\beta$ . *J. Biol. Chem.* 264:389.
8. Sanchez-Madrid, F., P. Simon, S. Thompson, and T.A. Springer. 1983. Mapping of antigenic and functional epitopes on the  $\alpha$  and  $\beta$  subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. *J. Exp. Med.* 158:586.
9. Yokoyama, W.M., F. Koning, P.J. Kehn, G.M.B. Peteira, G. Stingl, J.E. Coligan, and E.M. Shevach. 1988. Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation. *J. Immunol.* 141:369.
10. Argraves, W.S., S. Suzuki, H. Arai, K. Thompson, M.D. Pierschbacher, and E. Ruoslahti. 1987. Amino acid sequence of the human fibronectin receptor. *J. Cell Biol.* 105:1183.
11. Suzuki, S., W.S. Argraves, H. Arai, L.R. Languino, M.D. Pierschbacher, and E. Ruoslahti. 1987. Amino acid sequence of the vitronectin receptor  $\alpha$  subunit and comparative expression of adhesion receptor mRNAs. *J. Biol. Chem.* 262:14080.
12. Maloy, W.L., A.M. Lew, R.W. Anderson, and J.E. Coligan. 1988. H-2K molecules have two different C-termini, one of which is K-region specific. *Mol. Immunol.* 25:453.
13. Lew, A.M., D.H. Margulies, W.L. Maloy, E.P. Lillehoj, J. McCluskey, and J.E. Coligan. 1986. Alternative protein products with different carboxyl termini from a single class I gene, H-2K<sup>b</sup>. *Proc. Natl. Acad. Sci. USA*. 83:6084.
14. O'Farrell, P.Z., H.M. Goodman, and P.M. O'Farrell. 1977. High resolution two-dimensional electrophoresis of basic as well as acidic proteins. *Cell*. 12:1133.
15. Vogel, B.E., G. Tarone, F.G. Giancotti, J. Gailit, and E. Ruoslahti. 1990. A novel fibronectin receptor with an unexpected subunit composition ( $\alpha\beta 1$ ). *J. Biol. Chem.* 265:5934.
16. Bodary, S.C., and J.W. McLean. 1990. The integrin  $\beta 1$  subunit associates with the vitronectin receptor  $\alpha$  subunit to form a novel vitronectin in a human embryonic kidney cell line. *J. Biol. Chem.* 265:5938.
17. Cheresch, D.A., J.W. Smith, H.M. Cooper, and V. Quaranta. 1989. A novel vitronectin receptor integrin ( $\alpha\beta x$ ) is responsible for distinct adhesive properties of carcinoma cells. *Cell*. 57:59.
18. Freed, E., J. Gailit, P. van der Geer, E. Ruoslahti, and T. Hunter. 1989. A novel integrin  $\beta$  subunit is associated with the vitronectin receptor  $\alpha$  subunit ( $\alpha$ ) in a human osteosarcoma cell line and is a substrate for protein kinase C. *EMBO (Eur. Mol. Biol. Organ.) J.* 8:2955.
19. Krissansen, G.W., M.J. Elliott, C.M. Lucas, F.C. Stomski, M.C. Berndt, D.A. Cheresch, A.F. Lopez, and G.F. Burns. 1990. Identification of a novel integrin  $\beta$  subunit expressed on cultured monocytes (macrophages). *J. Biol. Chem.* 265:823.
20. Hemler, M.E. 1990. VLA proteins in the integrin family: structures, functions and their role on leukocytes. *Annu. Rev. Immunol.* 8:365.
21. Pytela, R., M.D. Pierschbacher, and E. Ruoslahti. 1985. A 125/115 kDa cell surface receptor specific for vitronectin interacts with the arginine-glycine-aspartic acid adhesion sequence from fibronectin. *Proc. Natl. Acad. Sci. USA*. 82:5766.
22. Smith, J.W., D.J. Vestal, S.V. Irwin, T.A. Burke, and D.A. Cheresch. 1990. Purification and functional characterization of integrin  $\alpha\beta 5$ . *J. Biol. Chem.* 265:11008.
23. Dustin, M.L., and T.A. Springer. 1989. T-cell receptor cross-linking transiently stimulates adhesiveness through LFA-1. *Nature (Lond.)*. 341:619.
24. Matsuyama, T., A. Yamada, J. Kay, K.M. Yamada, S.K. Akiyama, S.F. Schlossman, and C. Morimoto. 1989. Activation of CD4 cells by fibronectin receptor and anti-CD3 antibody: a synergistic effect mediated by the VLA-5 fibronectin complex. *J. Exp. Med.* 170:1133.
25. Shimizu, Y., G.A. van Seventer, K.J. Horgan, and S. Shaw. 1990. Regulated expression and binding of three VLA ( $\beta 1$ ) integrin receptors on T-cells. *Nature (Lond.)*. 345:250.
26. Klingemann, H.G., and S. Dedhar. 1989. Distribution of integrins on human peripheral blood mononuclear cells. *Blood*. 74:1348.
27. Roberts, K., W.M. Yokoyama, P.J. Kehn, and E.M. Shevach. 1991. The vitronectin receptor serves as an accessory molecule for the activation of a subset of  $\gamma/\delta$  T cells. *J. Exp. Med.* 173:231.