

γ/δ T Cell-deficient Mice Have Impaired Mucosal Immunoglobulin A Responses

By Kohtaro Fujihashi,* Jerry R. McGhee,** Mi-Na Kweon,* Max D. Cooper,^{‡§||} Susumu Tonegawa,** Ichiro Takahashi,^{‡‡} Takachika Hiroi,^{§§} Jiri Mestecky,^{‡§} and Hiroshi Kiyono^{**‡§§}

From The Immunobiology Vaccine Center, Mucosal Immunization Research Group, Departments of *Oral Biology, [‡]Microbiology, [§]Medicine, and ^{||}Pediatrics; [¶]Howard Hughes Medical Institute, the University of Alabama at Birmingham, Medical Center, Birmingham, Alabama 35294; **Center for Cancer Research, Department of Biology, Howard Hughes Medical Institute and Massachusetts Institute of Technology, Cambridge, Massachusetts 02139; ^{‡‡}Department of Oral Microbiology, Faculty of Dentistry; and ^{§§}Department of Mucosal Immunology, Research Institute for Microbial Diseases, Osaka University, Suita, Osaka 565, Japan

Summary

Mucosal tissues of mice are enriched in T cells that express the γ/δ T cell receptor. Since the function of these cells remains unclear, we have compared mucosal immune responses in γ/δ T cell receptor-deficient (TCR $\delta^{-/-}$) mice versus control mice of the same genetic background. The frequency of intestinal immunoglobulin (Ig) A plasma cells as well as IgA levels in serum, bile, saliva, and fecal samples were markedly reduced in TCR $\delta^{-/-}$ mice. The TCR $\delta^{-/-}$ mice produced much lower levels of IgA antibodies when immunized orally with a vaccine of tetanus toxoid plus cholera toxin as adjuvant. Conversely, the antigen-specific IgM and IgG antibody responses were comparable to orally immunized control mice. Direct assessment of the cells forming antibodies against the tetanus toxoid and cholera toxin antigens indicated that significantly lower numbers of IgA antibody-producing cells were present in the intestinal lamina propria and Peyer's patches of TCR $\delta^{-/-}$ mice compared with the orally immunized control mice. The selective reduction of IgA responses to ingested antigens in the absence of γ/δ T cells suggests a specialized role for γ/δ T cells in mucosal immunity.

The mucosal immune system is considered as a separate functional entity quite independent of the systemic immune compartment, since it possesses unique anatomical features and is composed of specialized subsets of lymphoid cells. For example, the γ/δ T cells are prevalent among the intestinal intraepithelial lymphocyte (IEL) population (1–5). The underlying lamina propria of the small intestine also contains a higher frequency of γ/δ T cells than the systemic lymphoid tissues (6). Most of the resident γ/δ T cells in intestinal epithelium express cell surface CD8 molecules composed of $\alpha\alpha$ homodimeric chains (7, 8), and their V γ gene repertoire differs from that of the systemic γ/δ T cells. Thus the intestinal, uterine, and skin γ/δ T cells express the V γ 7, V γ 6, and V γ 5 genes, respectively, whereas systemic γ/δ T cells predominantly express the V γ 4 gene (9, 10).

Studies of TCR-deficient mice suggest an important role for γ/δ T cells in immune responses to intracellular bacteria and parasites (11–13). The γ/δ T cells appear to be required for control of mycobacterial infection (11) and contribute to immunity after *Plasmodium yoelii* vaccination, since γ/δ TCR-deficient (TCR $\delta^{-/-}$) mice do not respond

normally to these intracellular microorganisms (12). The γ/δ T cells also play an accessory role in the late stages of protective immune responses to *Mycobacterium bovis* Bacillus Calmette-Guerin (13). These observations clearly implicate γ/δ T cells in microbial immunity, but the precise function of γ/δ T cells in specific immune responses remains unclear. Our previous studies suggested that γ/δ T cells are important for the maintenance of mucosal IgA responses in the presence of systemic unresponsiveness or tolerance induced by oral immunization (14, 15).

To examine the role of γ/δ T cells in the induction and regulation of mucosal immunity, we compared the immune responses in TCR $\delta^{-/-}$ and control mice to oral immunization with tetanus toxoid (TT) and cholera toxin (CT) as mucosal adjuvants. This immunization regimen induces intestinal secretory IgA (S-IgA) responses as well as circulating IgG and IgA antibodies to both antigens in normal mice (16, 17), thus allowing examination of the role of γ/δ T cells in both mucosal and systemic responses. In this communication, we report that in contrast to their normal counterparts, TCR $\delta^{-/-}$ mice display significant alterations in IgA responses induced by oral immunization.





Figure 1. Effect of TCR- δ gene disruption on immunoglobulin production (A) Antibody-producing cells of IgM (\square), IgG (\square), and IgA (\blacksquare) isotypes were examined in gut-associated tissues and the spleen by an ELISPOT assay (B) Immunofluorescence staining of small intestine tissue sections from TCR $\delta^{-/-}$ and control TCR $\delta^{+/+}$ mice. IgA-containing cells were visualized by incubation with biotinylated goat anti-mouse α F(ab')₂ followed by avidin-AMCA. Results represent the values (mean \pm SEM) for 12 mice in each experimental group.

Materials and Methods

γ/δ T Cell-deficient Mice. The TCR $\delta^{-/-}$ mice were constructed by introducing germline mutations in the TCR- δ chain gene on a (129/Ola \times C57Bl/6) (H-2^b) background (18). The δ chain-deficient and (129/O1a \times C57Bl/6)_F₂ normal mice (TCR $\delta^{+/+}$) were barrier maintained in Trexler isolators and have remained pathogen antibody negative. At 5–6 wk of age, the mice were removed from the colony isolator unit, housed in microisolator cages in horizontal laminar flow cabinets, and provided sterile food and water ad lib. The mice were between 7 and 10 wk of age at the beginning of the experiment.

Immunization. Mice were immunized orally on days 0, 7, and

14 with 0.25 ml PBS containing a mixture of vaccine-grade TT (250 μ g/mouse; kindly provided by Dr. Patricia J. Freda Pietrobon [Connaught Laboratories Inc., Swiftwater, PA]) and CT (10 μ g/mouse; List Biologic Laboratories, Inc., Campbell, CA) (16, 17).

Antibody Assays. Antibody titers in serum and fecal extracts were determined by ELISA (16, 17). Assay plates (Falcon Microtest; Becton Dickinson & Co., Oxnard, CA) were coated with an optimal concentration of TT (100 μ l of 5 μ g/ml TT, equivalent to 0.8 flocculation units/ml) or recombinant CT-B (100 μ l of 5 μ g/ml of CT-B; List Biological Laboratories Inc.) in PBS. End

point titers were expressed as the last dilution yielding an optical density at 414 nm (OD₄₁₄) of >0.1 U above negative control values after a 15-min incubation. In some experiments, the plates were coated with goat anti-mouse Ig (Southern Biotechnology Associates, Birmingham, AL) at 2 µg/ml (100 µl/well). Total IgM, IgG, and IgA levels in serum saliva, bile, and fecal extracts were estimated by comparison with serial dilutions of mouse IgM, IgG, and IgA standards (Southern Biotechnology Associates).

Enumeration of Antibody-producing Cells. The spleen was removed aseptically, and single-cell suspensions were prepared as described (14, 15). Peyer's patches carefully excised from the intestinal wall were dissociated using the neutral protease enzyme Dispase® (Boehringer Mannheim Corp., Indianapolis, IN) in Joklik-modified medium (Life Technologies, Inc., Gaithersburg, MD) to obtain single-cell preparations (16, 17). Mononuclear cells in the lamina propria were isolated after removal of PP from the small intestine using a combination of enzymatic dissociation and discontinuous Percoll gradients (Pharmacia, Uppsala, Sweden) (6, 19). Mononuclear cells in the interface between the 40 and 75% layers were removed, washed, and resuspended in RPMI 1640 containing 10% FCS. An enzyme-linked immunospot assay was used to detect cells producing IgM, IgG, and IgA antibodies (14, 16). 96-well nitrocellulose plates (Millititer HA; Millipore Corp., Bedford, MA) were coated with goat anti-mouse Ig at 2 µg/ml (100 µl/well) to detect total IgM, IgG, and IgA antibody-forming cells (AFC), 5 µg/ml of CT-B (100 µl/well) for anti-CT-B-specific AFCs, or 5 µg/ml of TT (100 µl/well) for anti-TT-specific AFC (16, 19).

Immunohistology of the Small Intestine. Samples of mouse jejunum and ileum were obtained from TCRδ^{-/-} and control mice for conventional histology and staining of antibody-containing cells (19). 1-cm portions of intestine were opened longitudinally, mounted on thin cards, and washed in three changes of PBS at 4°C over a period of 2 h to remove tissue-associated Ig. The tissue was then fixed in 5% glacial acetic acid in 95% ethanol at -20°C for 24 h before paraffin embedding. 4-µm-thick serial tissue sections were mounted on glass slides, and Ig-containing cells were visualized by three-color staining with FITC-labeled goat (Fab')₂ anti-mouse µ (Southern Biotechnology Associates), RITC-labeled goat F(ab')₂ anti-mouse γ (Southern Biotechnology Associates), and biotinylated F(ab')₂ anti-mouse α (Southern Biotechnology Associates) followed by avidin-7-amino-4-methylcoumarin-3-acetic acid (AMCA; Jackson ImmunoResearch Laboratories, Inc., West Grove, PA [19]).

Statistics. Mean numbers of AFC in the different tissues were compared among experimental and control groups by Wilcoxon signed-rank test. Mean percentages of Ig-containing cells per villi were calculated from multiple determinations of experimental and control groups and compared by Wilcoxon signed-rank test.

Results

Deficiency of IgA-producing Cells in TCRδ^{-/-} Mice. The vast majority of γ/δ T cells are located in the epithelium of the small intestine of normal mice (1-3), suggesting that absence of this T cell subset in this mucosal effector site could influence immunological homeostasis. Since previous studies have not assessed γ/δ T cells for their participation in regulation of humoral and mucosal immune responses, we initially examined the effects of TCR-δ gene disruption on the numbers of IgM-, IgG-, and IgA-producing cells in systemic and mucosal tissues, and levels of IgM, IgG, and

IgA present in serum, saliva, bile, and fecal extracts. When the frequency of Ig-producing cells was compared between spleens of nonimmunized TCRδ^{-/-} mice and control mice of the same (129 × B6)F₂ background (TCRδ^{+/+}), comparable numbers of IgM- or IgG-producing cells were seen. In contrast, the numbers of IgA-secreting cells in the intestinal lamina propria and Peyer's patches of TCRδ^{-/-} mice were significantly lower than in control TCRδ^{+/+} mice ($P < 0.02$, Fig. 1 A).

The frequency of IgA-containing cells was also evaluated in tissue sections of jejunum and ileum by immunohistological analysis. Enumeration of the IgA-producing cells in the lamina propria of the small intestine indicated a reduction of IgA plasma cells in TCRδ^{-/-} mice. The numbers of intestinal IgA plasma cells in normal and mutant mice were 460 ± 63 and 116 ± 22/10 fields, respectively ($P < 0.002$, Fig. 1 B). The reduction in IgA-producing cells in TCRδ^{-/-} mice was confirmed by an assessment of antibody levels in serum, saliva, bile, and fecal extracts using an isotype-specific ELISA. The IgA levels were reduced by ~80% in fecal extracts obtained from TCRδ^{-/-} mice compared with fecal IgA levels in control TCRδ^{+/+} mice ($P < 0.001$, Table 1). Serum IgA levels were also reduced in TCRδ^{-/-} mice ($P < 0.003$), whereas IgM and IgG levels were normal (Fig. 2, Table 1). Further, IgA levels in saliva and bile of TCRδ^{-/-} mice were significantly lower than controls ($P < 0.005$, Table 1).

Reduction of TT-specific IgA Responses in Orally Immunized TCRδ^{-/-} Mice. The antibody response to TT was examined in TCRδ^{-/-} mice that were immunized orally with a vaccine containing TT and CT to examine the potential involvement of γ/δ T cells in antigen-specific IgA responses. When TT-specific serum antibody responses were compared after three oral doses of the combined vaccine, the TCRδ^{-/-} mice and their normal littermates (TCRδ^{+/+}) produced almost identical levels of IgG antibodies to both proteins, whereas lower serum IgA responses were seen in TCRδ^{-/-} mice ($P < 0.003$, Table 2). When TT-specific antibodies were assessed in fecal samples, lower IgA responses were noted in the TCRδ^{-/-} mice ($P < 0.005$, Table 2). To evaluate these findings at a cellular level, we examined the frequency of antigen-specific AFC in different tissues of mice immunized orally with the combined vaccine by using the ELISPOT assay. A reduction in the numbers of TT-specific IgA AFC was noted in both Peyer's patches and intestinal lamina propria in the TCRδ^{-/-} mice when compared with control mice ($P < 0.01$, Fig. 3 A).

Characterization of Antibody Response to Cholera Toxin B Subunit in TCRδ^{-/-} Mice. It is possible that hyporesponsiveness of TCRδ^{-/-} mice to combined TT was due to unresponsiveness to the adjuvant effect of CT. Since the CT molecule possesses strong immunogenicity in addition to mucosal adjuvant activity for IgA responses, we could also measure IgA responses to the CT antigen. Serum and fecal extracts as well as mononuclear cells from various tissues of both TCRδ^{-/-} mice and controls were obtained 1 wk after the last oral dose of the combined vaccine for analysis of CT-B-specific responses by ELISA and ELISPOT assays.

Table 1. Low Levels of Total Serum and Secretory IgA Antibodies in TCR $\delta^{-/-}$ Mice

Mouse strains	Fecal	Saliva	Bile	Serum
	$\mu\text{g/ml}$	ng/ml	$\mu\text{g/ml}$	$\mu\text{g/ml}$
TCR $\delta^{-/-}$	2 \pm 1	244 \pm 19	270 \pm 30	25 \pm 3
TCR $\delta^{+/+}$	11 \pm 2	753 \pm 100	849 \pm 155	125 \pm 10

Values represent the mean endpoint titer \pm SEM for nine mice in each experimental group.

When CT-B-specific serum responses were compared between TCR $\delta^{-/-}$ and normal mice, lower CT-B-specific IgA titers were detected in the former ($P < 0.008$, Table 2). When CT-B-specific antibody titers were examined in fecal extracts, a significant decrease in the antigen-specific IgA levels was observed in TCR $\delta^{-/-}$ mice ($P < 0.01$, Table 2). Reductions in CT-B-specific IgA-producing cells in both intestinal lamina propria and Peyer's patches were also observed in orally immunized TCR $\delta^{-/-}$ mice ($P < 0.03$, Fig. 3 B).

Discussion

These studies suggest that γ/δ T cells exert an unexpected role in mucosal immunity. Decreased levels of total serum and secretory IgA as well as the reduction of IgA responses to both tetanus toxoid and cholera toxin antigens in TCR $\delta^{-/-}$ mice suggests that γ/δ T cells can regulate the IgA response. In contrast, we found that IgM and IgG antibody responses were unimpaired in TCR $\delta^{-/-}$ mice when compared with normal controls.

CD4 $^{+}$ α/β T cells have been shown to be essential for IgA B cell responses (for review see reference 20), whereas the results of this study suggest that γ/δ T cells also may regulate the mucosal IgA immune response. The helper T

cells that preferentially produce IL-4, IL-5, IL-6, and IL-10, the Th2 subset of CD4 $^{+}$ cells, promote IgA responses (16, 21–23). Our earlier studies indicated that oral immunization of normal mice with the combined TT and CT vaccine induced Th2-type cells in the IgA mucosal response (16), and TCR- $\beta^{-/-}$ mice are essentially devoid of mucosal IgA-producing cells (Fujihashi, K., M.-N. Kweon, M. Marinaro, K. Imaoka, R.J. Jackson, J.R. McGhee, and H. Kiyono, manuscript in preparation). Function-loss mutations in the CD40L gene in humans and mice result in the failure to generate T cell-dependent secondary humoral immune responses, including antibodies of IgG and IgA isotype (24, 25). A severe impairment of IgA responses is also observed in anti-CD4-treated and athymic mice, which also have reduced germinal center development and few IgA-producing cells in the intestinal lamina propria (19, 26). The interaction between CD4 $^{+}$ Th2 cells and B cells that are induced to undergo isotype switching normally occurs in germinal centers (20), where the interaction between CD40L on the activated CD4 $^{+}$ α/β T cells and the CD40 molecule on B cells also initiates a signal needed for the isotype switch process (for review see reference 27). A special subset of the CD4 $^{+}$ α/β T cells is thus essential for the induction of systemic IgA responses.

Our present results suggest that γ/δ T cells may serve an important regulatory role for mucosal IgA responses in that TCR $\delta^{-/-}$ mice possess normal numbers of functional α/β T cells but display markedly decreased IgA responses to oral immunization. The previous studies also suggested that γ/δ T cells may play an important role in the maintenance of mucosal IgA responses in the presence of oral tolerance (14, 15). One possible explanation for this unexpected finding is that γ/δ T cells positively influence α/β Th2 cells that regulate IgA immune responses in mucosal tissues. In this scenario, a triad interaction between mucosal γ/δ T cells, α/β Th2-type cells, and IgA B cell precursors is involved in the induction of maximal IgA responses to oral antigens. Experiments in mice (11, 13, 28) and in chickens (29, 30) in-




Figure 2. IgM, IgG, and IgA levels in serum and fecal extracts of TCR $\delta^{-/-}$ and TCR $\delta^{+/+}$ mice. These levels were determined by an ELISA using mouse IgM, IgG, and IgA standards. Values are mean \pm SEM for nine mice in each experimental group.

Table 2. Selective Impairment of IgA Responses to Oral Immunization with TT and CT Antigens in TCR $\delta^{-/-}$ Mice

Antigen	Mouse strains	Antibody titer (Log ₂)			
		Fecal		Serum	
		IgA	IgA	IgG	IgM
TT	TCR $\delta^{-/-}$	5.0 ± 0.3	8.0 ± 0.3	16.6 ± 0.3	11.6 ± 0.3
	TCR $\delta^{+/+}$	7.5 ± 0.5	10.8 ± 0.4	16.8 ± 0.7	11.4 ± 0.4
CT-B	TCR $\delta^{-/-}$	5.7 ± 0.5	10.6 ± 0.4	16.2 ± 0.2	11.0 ± 0.2
	TCR $\delta^{+/+}$	8.3 ± 0.3	12.5 ± 0.3	17.0 ± 0.5	10.7 ± 0.4

Values represent the mean end point titer ± SEM for nine mice in each experimental group. The antibody end point titration was performed by ELISA as described in Materials and Methods.

deed suggest that bidirectional interactions between α/β and γ/δ T cells may have functionally important consequences. Alternatively, the γ/δ T cells could have a direct effect either on the differentiation of sIgA⁺ B cells or in the induction of the mucosal IgA switch process. A recent study indicates that activated γ/δ T cells can express

CD40L and thereby induce an IgE B cell response (31). It thus seems possible that mucosal CD40L⁺ γ/δ T cells are capable of direct interaction with CD40⁺ B cells to induce IgA responses. The observation of significant IgA production in CD40^{-/-} and CD40L^{-/-} mice (25, 32, 33) also raises the possibility of an alternative set of interaction molecules that could allow γ/δ (or α/β) T cells in the mucosal compartment to induce IgA responses to ingested antigens.

Our previous studies and those of others indicate that γ/δ IEL can produce an array of Th1- and Th2-type cytokines, as well as TNF- α and TGF- β (34, 35). Specifically, the γ/δ IELs can secrete IL-5 and IL-6, which are key cytokines for inducing sIgA⁺ B cells to differentiate into IgA plasma cells (34, 35). Moreover, γ/δ T cells in other mucosal effector tissues, such as the salivary glands, are committed to produce IL-5 and IL-6 (36), and the frequency of IgA-producing cells is reduced in the salivary glands of TCR $\delta^{-/-}$ mice (data not shown). IgA-producing cells are also greatly reduced in intestinal tissues of IL-6^{-/-} mice (37). The impaired IgA responses in TCR $\delta^{-/-}$ mice could therefore reflect the absence of mucosal γ/δ T cells that produce IL-5 and IL-6.

Yet another possible explanation for our findings is that the lack of γ/δ T cells in intestinal epithelium negatively influences epithelial cell production of TGF- β and IL-6,




Figure 3. Comparison of (A) TT- and (B) CT-B-specific antibody responses in TCR $\delta^{-/-}$ and control mice immunized orally with a combined vaccine containing TT and CT. Mononuclear cells isolated from spleen, Peyer's patch, and lamina propria of both TCR $\delta^{-/-}$ and control mice were subjected to the TT- and CT-B-specific ELISPOT assay. Antigen-specific IgM (□), IgG (▤) and IgA (■) AFC were enumerated. Results represent the values (mean ± SEM) for nine mice in each experimental group.

which serve as IgA isotype switching and differentiation factors, respectively (38, 39). T cell-derived cytokines, including IFN- γ , TNF- α , and IL-4, can influence epithelial cell functions (40, 41). All of these cytokines can be produced by γ/δ IEL (34, 35), and γ/δ T cells may influence epithelial cell growth and function. In fact, reduction in both the numbers of intestinal epithelial cells and their level of MHC class II expression have been observed in TCR- δ gene-disrupted mice (42). Further, intraepithelial γ/δ T cells have been shown to modulate growth of epithelial cells via the production of keratinocyte growth factor (43). The absence of γ/δ T cells in the intestinal epithelium could therefore compromise TGF- β and IL-6 production

by epithelial cells, which, in turn, may result in diminished IgA responses.

In conclusion, γ/δ T cells may influence IgA B cell responses to ingested antigens via their interactions with other T cells and mucosal epithelial cells. Mucosal γ/δ T cells may thus regulate production of key cytokines for IgA B cell development by CD4⁺ α/β T cells (e.g., IL-5, IL-6, and IL-10) and epithelial cells (e.g., TGF- β and IL-6). The interactions between γ/δ T cells, α/β T cells, and epithelial cells in the induction and regulation of mucosal immune responses could thus represent a fertile field for future investigation.

We thank Annette M. Pitts, Thaddeus V. Bamberg, and Steve H. Yoon for technical assistance, and Sheila Shaw for the preparation of this manuscript.

This work was supported by U.S. Public Health Service grants AI-35932, DE-09837, AI-35544, AI-30366, DE-04217, DK-44240, AI-18958, AI-39816, and contract AI-15128 as well as grants from Ministry of Education, Science Sports and Cultures, Ministry of Health and Welfare, and Asahi Chemical Industry Co. Ltd. in Japan. M.D. Cooper and S. Tonegawa are Howard Hughes Medical Institute investigators.

Address correspondence to Hiroshi Kiyono, Departments of Oral Biology and Microbiology, BBRB Room 761, Immunobiology Vaccine Center, University of Alabama at Birmingham, Medical Center, Birmingham, AL 35294-2170.

Received for publication 27 November 1995.

References

1. Goodman, T., and L. Lefrançois. 1988. Expression of the $\gamma\delta$ T cell receptor on intestinal CD8⁺ intraepithelial lymphocytes. *Nature (Lond.)*. 333:855–858.
2. Bonneville, M., C.A. Janeway, Jr., K. Ito, W. Haser, I. Ishida, N. Nakanishi, and S. Tonegawa. 1988. Intestinal intraepithelial lymphocytes are a distinct set of $\gamma\delta$ T cells. *Nature (Lond.)*. 336:479–481.
3. Kyes, S., E. Carew, S.R. Carding, C.A. Janeway, Jr., and A.C. Hayday. 1989. Diversity in T-cell receptor γ gene usage in intestinal epithelium. *Proc. Natl. Acad. Sci. USA*. 86:5227–5531.
4. Bucy, R.P., C.H. Chen, J. Cihak, U. Losch, and M.D. Cooper. 1988. Avian T cells expressing $\gamma\delta$ receptors localize in the splenic sinusoids and the intestinal epithelium. *J. Immunol.* 141:2200–2205.
5. Bucy, R.P., C.H. Chen, and M.D. Cooper. 1989. Tissue localization and CD8 accessory molecule expression of T $\gamma\delta$ cells in humans. *J. Immunol.* 142:3045–3049.
6. Aicher, W.K., K. Fujihashi, M. Yamamoto, H. Kiyono, A.M. Pitts, and J. R. McGhee. 1992. Effects of the *lpr/lpr* mutation on T and B cell populations in the lamina propria of the small intestine, a mucosal effector site. *Int. Immunol.* 4: 959–968.
7. Guy-Grand, D., N. Cerf-Bensussan, B. Malissen, M. Malassis-Seris, C. Briottet, and P. Vassalli. 1991. Two gut intraepithelial CD8⁺ lymphocyte populations with different T cell receptors: a role for the gut epithelium in T cell differentiation. *J. Exp. Med.* 173:471–481.
8. Lefrançois, L. 1991. Phenotypic complexity of intraepithelial lymphocytes of the small intestine. *J. Immunol.* 147:1746–1751.
9. Takagaki, Y., A. DeCloux, M. Bonneville, and S. Tonegawa. 1989. Diversity of $\gamma\delta$ T-cell receptors on murine intestinal intraepithelial lymphocytes. *Nature (Lond.)*. 339:712–714.
10. Itohara, S., A.G. Farr, J.J. Lafaille, M. Bonneville, Y. Takagaki, W. Haas, and S. Tonegawa. 1990. Homing of $\gamma\delta$ thymocyte subset with homogeneous T-cell receptors to mucosal epithelia. *Nature (Lond.)*. 343:754–757.
11. Mombaerts, P., J. Arnoldi, F. Russ, S. Tonegawa, and S.H.E. Kaufmann. 1993. Different roles of $\alpha\beta$ and $\gamma\delta$ T cells in immunity against an intracellular bacterial pathogen. *Nature (Lond.)*. 365:53–56.
12. Tsuji, M., P. Mombaerts, L. Lefrançois, R.S. Nussenzweig, F. Zavala, and S. Tonegawa. 1994. $\gamma\delta$ T cells contribute to immunity against the liver stages of malaria in $\alpha\beta$ T-cell-deficient mice. *Proc. Natl. Acad. Sci. USA*. 91:345–349.
13. Ladel, C.H., J. Hess, S. Daugelat, P. Mombaerts, S. Tonegawa, and S.H.E. Kaufmann. 1995. Contribution of α/β and γ/δ T lymphocytes to immunity against *Mycobacterium bovis* Baccillus Calmette Guerin: studies with T cell receptor-deficient mutant mice. *Eur. J. Immunol.* 25:838–846.
14. Fujihashi, K., T. Taguchi, J.R. McGhee, J.H. Eldridge, M.G. Bruce, D.R. Green, B. Singh, and H. Kiyono. 1990. Regulatory function for murine intraepithelial lymphocytes: two subsets of CD3⁺, T cell receptor-1⁺ intraepithelial lymphocyte T cells abrogate oral tolerance. *J. Immunol.* 145:2010–2019.
15. Fujihashi, K., T. Taguchi, W.A. Aicher, J.R. McGhee, J.A. Bluestone, J.H. Eldridge, and H. Kiyono. 1992. Immunoregulatory functions for murine intraepithelial lymphocytes: γ/δ

- T cell receptor-positive (TCR⁺) T cells abrogate oral tolerance, while α/β TCR⁺ T cells provide B cell help. *J. Exp. Med.* 175:695–707.
16. Xu-Amano, J., H. Kiyono, R.J. Jackson, H.F. Staats, K. Fujihashi, P.D. Burrows, C.O. Elson, S. Pillai, and J.R. McGhee. 1993. Helper T cell subset for immunoglobulin A responses: oral immunization with tetanus toxoid and cholera toxin as adjuvant selectively induces Th2 cells in mucosa-associated tissues. *J. Exp. Med.* 178:1309–1320.
 17. Jackson, R.J., K. Fujihashi, J. Xu-Amano, H. Kiyono, C.O. Elson, and J.R. McGhee. 1993. Optimizing oral vaccines: induction of systemic and mucosal B-cell and antibody responses to tetanus toxoid by use of cholera toxin as an adjuvant. *Infect. Immun.* 61:4272–4279.
 18. Itohara, S., P. Mombaerts, J. Lafaille, J. Iacomini, A. Nelson, A. Farr, and S. Tonegawa. 1993. T cell receptor δ gene mutant mice: independent generation of α/β T cells and programmed rearrangements of γ/δ TCR gene. *Cell.* 72:337–348.
 19. Mega, J., M.G. Bruce, K.W. Beagley, J.R. McGhee, T. Taguchi, A.M. Pitts, M.L. McGhee, R.P. Bucy, J.H. Eldridge, J. Mestecky, and H. Kiyono. 1991. Regulation of mucosal responses by CD4⁺ T lymphocytes: effects of anti-L3T4 treatment on the gastrointestinal immune system. *Int. Immunol.* 3:793–805.
 20. McGhee, J.R., J. Mestecky, C.O. Elson, and H. Kiyono. 1989. Regulation of IgA synthesis and immune response by T cells and interleukins. *J. Clin. Immunol.* 9:175–199.
 21. Beagley, K.W., J.H. Eldridge, H. Kiyono, M.P. Everson, W.J. Koopman, T. Honjo, and J.R. McGhee. 1988. Recombinant murine IL-5 induces high rate IgA synthesis in cycling IgA-positive Peyer's patch B cells. *J. Immunol.* 141:2035–2042.
 22. Beagley, K.W., J.H. Eldridge, F. Lee, H. Kiyono, M.P. Everson, W.J. Koopman, T. Hirano, T. Kishimoto, and J.R. McGhee. 1989. Interleukins and IgA synthesis. Human and murine interleukin 6 induce high rate Igs secretion in IgA-committed B cells. *J. Exp. Med.* 169:2133–2148.
 23. Defrance, T., B. Vanbervliet, F. Briere, I. Durand, F. Rousset, and J. Banchereau. 1992. Interleukin 10 and transforming growth factor β cooperate to induce anti-CD40-activated naive human B cells to secrete immunoglobulin A. *J. Exp. Med.* 175:671–682.
 24. Aruffo, A., M. Farrington, D. Hollenbaugh, X. Li, A. Milatovich, S. Nonoyama, J. Bajorath, L.S. Grosmaire, R. Stenkamp, M. Neubauer, et al. 1993. The CD40 ligand, gp39, is defective in activated T cells from patients with X-linked hyper-IgM syndrome. *Cell.* 72:291–300.
 25. Renshaw, B.R., W.C. Fanslow III, R.J. Armitage, K.A. Campbell, D. Liggitt, B. Wright, B.L. Davison, and C.R. Maliszewski. 1994. Humoral immune responses in CD40 ligand-deficient mice. *J. Exp. Med.* 180:1889–1900.
 26. Guy-Grand, D., C. Griscelli, and P. Vassalli. 1975. Peyer's patches, gut IgA plasma cells and thymic function: study in nude mice bearing thymic grafts. *J. Immunol.* 115:361–364.
 27. Aversa, G., J. Punnonen, J.M. Carballido, B.G. Cocks, and J.E. de Vries. 1994. CD40 ligand-CD40 interaction in Ig isotype switching in mature and immature human B cells. *Semin. Immunol.* 6:295–301.
 28. Kaufmann, S.H., C. Blum, and S. Yamamoto. 1993. Crosstalk between α/β T cells and γ/δ T cells in vivo: activation of α/β T-cell responses after γ/δ T-cell modulation with the monoclonal antibody GL3. *Proc. Natl. Acad. Sci. USA.* 90:9620–9624.
 29. Arstila, T.P., P. Toivanen, and O. Lassila. 1993. Helper activity of CD4⁺ α/β T cells is required for the avian γ/δ T cell response. *Eur. J. Immunol.* 23:2034–2037.
 30. Kasahara, Y., C.H. Chen, and M.D. Cooper. 1993. Growth requirement for avian γ/δ T cells include exogenous cytokines, receptor ligation and *in vivo* priming. *Eur. J. Immunol.* 23:2230–2236.
 31. Horner, A.A., H. Jabara, N. Ramesh, and R.S. Geha. 1995. γ/δ T lymphocytes express CD40 ligand and induce isotype switching in B lymphocytes. *J. Exp. Med.* 181:1239–1244.
 32. Kawabe, T., T. Naka, K. Yoshida, T. Tanaka, H. Fujiwara, S. Suematsu, N. Yoshida, T. Kishimoto, and H. Kikutani. 1994. The immune responses in CD40-deficient mice: impaired immunoglobulin class switching and germinal center formation. *Immunity.* 1:167–178.
 33. Castigli, E., F.W. Alt, L. Davidson, A. Bottaro, E. Mizoguchi, A.K. Bhan, and R.S. Geha. 1994. CD40-deficient mice generated by recombination-activating gene-2-deficient blastocyst complementation. *Proc. Natl. Acad. Sci. USA.* 91:12135–12139.
 34. Taguchi, T., W.K. Aicher, K. Fujihashi, M. Yamamoto, J.R. McGhee, J.A. Bluestone, and H. Kiyono. 1991. Novel function for intestinal intraepithelial lymphocytes: murine CD3⁺, γ/δ TCR⁺ T cells produce IFN- γ and IL-5. *J. Immunol.* 147:3736–3744.
 35. Barrett, T.A., T.F. Gajewski, D. Danielpour, E.B. Chang, K.W. Beagley, and J.A. Bluestone. 1992. Differential function of intestinal intraepithelial lymphocyte subsets. *J. Immunol.* 149:1124–1130.
 36. Hiroi, T., K. Fujihashi, J.R. McGhee, and H. Kiyono. 1995. Polarized Th2 cytokine expression by both mucosal γ/δ and α/β T cells. *Eur. J. Immunol.* 25:2743–2751.
 37. Ramsay, A.J., A.J. Husband, I.A. Ramshaw, S. Bao, K.I. Matthaei, G. Koehler, and M. Kopf. 1994. The role of interleukin-6 in mucosal IgA antibody responses in vivo. *Science (Wash. DC).* 264:561–563.
 38. Anzano, M.A., D. Riemann, W. Prichett, D.F. Bowen-Pope, and R. Greig. 1989. Growth factor production by human colon carcinoma cell line. *Cancer Res.* 49:2898–2904.
 39. McGee, D.W., K.W. Beagley, W.K. Aicher, and J.R. McGhee. 1993. Transforming growth factor β and IL-1 β act in synergy to enhance IL-6 secretion by the intestinal epithelial cell line, IEC-6. *J. Immunol.* 151:970–978.
 40. Kvale, D., D. Lovhaug, L.M. Sollid, and P. Brandtzaeg. 1988. Tumor necrosis factor- α up-regulates expression of secretory component, the epithelial receptor for polymeric Ig. *J. Immunol.* 140:3086–3089.
 41. Phillips, J.O., M.P. Everson, Z. Moldoveanu, C. Lue, and J. Mestecky. 1990. Synergistic effect of IL-4 and IFN- γ on the expression of polymeric Ig receptor (secretory component) and IgA binding by human epithelial cells. *J. Immunol.* 145:1740–1744.
 42. Komano, H., Y. Fujiura, M. Kawaguchi, S. Matsumoto, Y. Hashimoto, S. Oana, P. Mombaerts, S. Tonegawa, H. Yamamoto, S. Itohara, et al. 1995. Homeostatic regulation of intestinal epithelia by intraepithelial γ/δ T cells. *Proc. Natl. Acad. Sci. USA.* 92:6147–6151.
 43. Boismenu, R., and W.L. Havran. 1994. Modulation of epithelial cell growth by intraepithelial γ/δ T cells. *Science (Wash. DC).* 266:1253–1255.