

## **Dendritic Cells Are Recruited into the Airway Epithelium during the Inflammatory Response to a Broad Spectrum of Stimuli**

By Andrew S. McWilliam, Sylvia Napoli, Amanda M. Marsh, Francis L. Pemper, Delia J. Nelson, Carolyn L. Pimm, Philip A. Stumbles, Timothy N.C. Wells, and Patrick G. Holt

---

*From the Division of Cell Biology, TVW Telethon Institute for Child Health Research, West Perth, Western Australia 6872*

### **Summary**

A key rate-limiting step in the adaptive immune response at peripheral challenge sites is the transmission of antigen signals to T cells in regional lymph nodes. Recent evidence suggests that specialized dendritic cells (DC) fulfill this surveillance function in the resting state, but their relatively slow turnover in most peripheral tissues brings into question their effectiveness in signaling the arrival of highly pathogenic sources of antigen which require immediate mobilization of the full range of host defenses for maintenance of homeostasis. However, the present report demonstrates that recruitment of a wave of DC into the respiratory tract mucosa is a universal feature of the acute cellular response to local challenge with bacterial, viral, and soluble protein antigens. Consistent with this finding, we also demonstrate that freshly isolated respiratory mucosal DC respond *in vitro* to a variety of CC chemokines as well as complementary cleavage products and N-formyl-methionyl-leucine-phenylalanine. This suggests that rapid amplification of specific antigen surveillance at peripheral challenge sites is an integral feature of the innate immune response at mucosal surfaces, and serves as an “early warning system” to alert the adaptive immune system to incoming pathogens.

The host response to invading pathogens is classically viewed as a two-tiered system, comprising a series of innate (inflammatory) and acquired (adaptive) immune mechanisms which operate over distinct time scales (1). Thus, the first line of defense is provided by the rapid recruitment of phagocytic granulocytes (usually neutrophils) into sites of tissue injury in response to locally produced chemotactic factors. This is followed up to 48 h later by a second wave of mononuclear cells containing large numbers of macrophages which are effective in both the uptake of persisting antigen and its subsequent presentation to the T cell system. In this classical scheme, the adaptive immune system serves as an optional backup to innate host defenses, and is called upon only in situations where significant amounts of antigen persist at the challenge site beyond the time frame of the acute inflammatory response.

While this scheme appears inherently economical, it could equally be viewed as flawed in at least one respect; viz, the delayed recruitment of adaptive immune mechanisms into the host response provides a temporal window for the establishment and spread of incoming pathogens. This danger would appear to be greatest in the case of pathogens not previously encountered by the host, i.e., for which neither antibody nor T-effector memory cells are available. Under

such circumstances, host survival may ultimately be determined by the rapidity with which naive T cells are primed against antigens displayed by the pathogen at the challenge site, a process which occurs initially in local draining lymph nodes.

Recent evidence suggest that dendritic cells (DC) function as first-line sentinels in immune surveillance of peripheral tissues (2), including mucosal surfaces such as those in the lung and airways (3, 4). These DC migrate into peripheral tissues from a circulating monocytelike precursor pool, and differentiate locally to a stage in which they are specialized for acquisition and processing of antigen, but remain unable to effectively present the antigen locally to T cells (2). This latter function, in particular a unique capacity for potent activation of naive T cells, is acquired after their migration to regional lymph nodes (2).

Thus, individual DC present a “snapshot” of the antigens encountered during their transient sojourn through their respective peripheral tissue sites, presumably including those antigens derived from incoming pathogens.

The effectiveness of such a surveillance system in the context of infectious disease is presumably a direct function of the DC traffic (i.e., cell number/unit time) between peripheral tissue sites and their respective regional lymph

nodes. In relatively quiescent tissues such as skin and muscle, mean DC transit times are estimated to normally be in the order of weeks (5–7). However, DC turnover is considerably more rapid at the main mucosal surfaces in direct contact with the outside environment (viz, the gastrointestinal and respiratory tracts) where resident populations are renewed every 3–4 d (7, 8).

## Materials and Methods

**Experimental Animals.** Specified pathogen-free adult PVG rats were supplied by the Animal Resource Centre (Murdoch, Western Australia). All animal experimentation was carried out with the prior approval of the Institute for Child Health Animal Ethics and Experimentation Committee which complies with conditions set down by the Australian National Health and Medical Research Council.

**Bacterial Models.** (a) *Moraxella catarrhalis* was grown in Mueller Hinton broth, washed extensively with saline, and suspended at  $\sim 10^9$  CFU/ml. The suspension was heated at 60°C for 1 h and passed through a 26 gauge needle several times to break up any bacterial clumps. Rats were exposed by aerosol to the suspension for 1 h using a Tri-R inhalation exposure apparatus (Tri-R Instruments, New York). (b) *Bordetella pertussis* (Welcome strain 28; provided by Dr. P. Novotny, Kent, England) was grown on Charcoal agar, washed in sterile PBS, and  $\sim 10^9$  organisms in 50  $\mu$ l were deposited directly onto the tracheal surface of adult rats by intratracheal intubation.


**Viral Model.** Sendai virus was provided by Dr. Jane Allan (University of Western Australia, Perth, Western Australia) and

grown for 3 d in the allantoic cavity of eggs. Allantoic fluid was stored at  $-70^\circ\text{C}$  until used. Adult rats were inoculated intranasally with  $10^3$  HAU in 50  $\mu$ l of virus containing allantoic fluid. Control animals were similarly inoculated with virus-free allantoic fluid. Infection of airway epithelium was confirmed by staining with mAb WS16 against the nucleoprotein antigen of Sendai virus (provided by Dr. A. Portner, St. Jude Children's Research Hospital, Memphis, TN). There was no influx of DC or T cells into the airway epithelium of control animals.

**Antigen Sensitization.** For antigen sensitization experiments, animals were intraperitoneally primed with 100  $\mu$ g of ovalbumin (OVA; Sigma Chemical Co., St. Louis, MO) in 0.5 ml PBS containing 10 mg aluminium hydroxide (Wyeth Amphojel). 14 d later, the animals were challenged with a 30 min aerosol of 1% (wt/vol) OVA in PBS.

**Immunohistochemical Analysis.** Tracheas were removed and immediately fixed in cold ethanol for 30 min. The tissue was then rehydrated in PBS, embedded in 100% OCT, and frozen in liquid nitrogen-cooled iso-pentane. Tangential sections 8–10  $\mu$ m were cut on a cryostat and immunostained as detailed in reference 4. Eosinophils were identified by cytochemical staining (4). Sections were counterstained with hematoxylin, dehydrated, and mounted. Primary antibodies used were Ox6 (rat MHC class II; 9), Ox12 (rat kappa light chains on B cells; 9), R73 (rat TcR $\alpha\beta$ ; 10), ED2 (rat macrophage; 11), and RP3 (rat neutrophils; 12). DC identification criteria were pleiomorphic morphology, together with positive staining with Ox6 and negative staining for Ox12, R73, and ED2, as detailed previously (4, 13).

**DC Chemotaxis.** The assay system was based upon that described in 14, except that 6.5 mm Costar Transwells (Costar Corp., Cambridge, MA) with polycarbonate membranes (3- $\mu$ m pore



**Figure 1.** Quantitative cellular responses within tracheal epithelium during inflammation. Data shown are from representative experiments ( $\geq 5$  for each model); each point is a mean value derived from four to six animals. (A) Inhalation of aerosolized heat-killed *M. catarrhalis* produced an acute inflammatory reaction in which Ox6<sup>+</sup> DC and RP3<sup>+</sup> neutrophils were the only inflammatory cell type detected in the epithelium. Peak numbers were detected at 1 d after aerosol. (B) Live *B. pertussis* organisms inoculated intratracheally resulted in an influx of Ox6<sup>+</sup> DC and RP3<sup>+</sup> neutrophils with maximum cell numbers occurring at day 1. The T cell influx began at day 2 and peaked at day 4. (C) Intranasal Sendai virus initiated a cellular response which commenced with an Ox6<sup>+</sup> DC influx shortly after the earliest detection of intraepithelial Sendai virus nucleoprotein immunostaining at day 2 (C, arrow) and shortly before changes could be detected in other cell types. Maximum DC numbers were found at day 5, and coincided with maximal neutrophil and T cell numbers. (D) After intraperitoneal priming with ovalbumin and Al(OH)<sub>3</sub>, animals were challenged 14 d later with an aerosol of ovalbumin and cellular changes within the epithelium measured. In this instance Ox6<sup>+</sup> DC, eosinophil, and T cell numbers peaked at day 1 after challenge.

Downloaded from <http://jci.org/> by guest on 18 September 2021



lenged airway tissue to levels that are two- to threefold those of resting tissues.

We have additionally screened a broad panel of chemotactic agents for capacity to attract respiratory mucosal DC in vitro (Fig. 2). Consistent with findings in Fig. 1 indicating rapid in vivo recruitment of DC in response to bacterial stimuli, complement cleavage products, and fMLP exhibit potent chemoattractant activity in vitro. Additionally, MCP-1, -4, RANTES, and eotaxin, all of which are members of the CC chemokine family, exhibited significant chemoattractant activity, whereas the CXC chemokines IL-8, GRO/KC, and GRO/ $\beta$ , were inactive. These findings are comparable to those recently reported using putative DC cultured from human blood in the presence of GM-CSF and IL-4, which also responded to a broad spectrum of chemotactic stimuli (14); however, the blood-derived DC were unresponsive

to MCP-1, which may reflect subtle species differences in receptor expression, or possibly variations related to the degree of maturation of respective DC populations at the time of assay.

These findings indicate that current perceptions of the scope of innate host defense responses need to be broadened to include the rapid activation of an in-built "early warning system" at sites of inflammation to facilitate the most rapid possible flow of information on the presence of previously unencountered antigens to the adaptive immune system. This mechanism fulfills an essential prediction of the model proposed recently (1, 15) for the evolution of the immune system, viz, that DC may be evolution's answer to the problem posed by organisms that evade primitive (innate) first-line defense systems.

---

This work was supported by Glaxo Wellcome, the Raine Foundation of Western Australia, and the National Health and Medical Research Council of Australia.

Address correspondence to Andrew S. McWilliam or Patrick G. Holt, Division of Cell Biology, TVW Telethon Institute for Child Health Research, PO Box 855, West Perth, Western Australia 6872. T.N.C. Well's present address is Geneva Biomedical Research Institute, Glaxo Wellcome Research & Development, Geneva.

Received for publication 10 June 1996 and in revised form 30 September 1996.

## References

1. Janeway, C.A. 1992. The immune response evolved to discriminate infectious nonself from noninfectious self. *Immunol. Today*. 13:11–16.
2. Steinman, R.M. 1991. The dendritic cell system and its role in immunogenicity. *Annu. Rev. Immunol.* 9:271–296.
3. Holt, P.G., M.A. Schon-Hegrad, and J. Oliver. 1988. MHC class II antigen-bearing dendritic cells in pulmonary tissues of the rat: regulation of antigen presentation activity by endogenous macrophage populations. *J. Exp. Med.* 167:262–274.
4. Schon-Hegrad, M.A., J. Oliver, P.G. McMenamin, and P.G. Holt. 1991. Studies on the density, distribution, and surface phenotype of intraepithelial class II MHC antigen (Ia)-bearing dendritic cells (DC) in the conducting airways. *J. Exp. Med.* 173:1345–1356.
5. Hart, D.N.J., and J.W. Fabre. 1981. Demonstration and characterization of Ia-positive dendritic cells in the interstitial connective tissues of rat heart and other tissues, but not brain. *J. Exp. Med.* 154:347–361.
6. Chen, H.-D., C. Ma, J.-T. Yuan, Y.-K. Wang, and W.K. Silvers. 1986. Occurrence of donor Langerhans cells in mouse and rat chimeras and their replacement in skin grafts. *J. Invest. Dermatol.* 86:630–633.
7. Fossum, S. 1989. Dendritic leukocytes: features of their *in vivo* physiology. *Res. Immunol.* 140:883–891.
8. Holt, P.G., S. Haining, D.J. Nelson, and J.D. Sedgwick. 1994. Origin and steady-state turnover of class II MHC-bearing dendritic cells in the epithelium of the conducting airways. *J. Immunol.* 153:256–261.
9. Mason, D.W., R.P. Arthur, M.J. Dallman., J.R. Green., G.P. Spickett, and M.L. Thomas. 1983. Functions of rat T-lymphocyte subsets isolated by means of monoclonal antibodies. *Immunol. Rev.* 74:57–82.
10. Hünig, T., H.J. Wallny, J.K. Hartley, A. Lawetzky, and G. Tiefenthaler. 1989. A monoclonal antibody to a constant determinant of the rat T cell antigen receptor that induces T cell activation. Differential reactivity with subsets of immature and mature T lymphocytes. *J. Exp. Med.* 169:73–86.
11. Damoiseaux, J.G., E.A. Dopp, J.J. Neeffjes, R.H. Beelen, and C.D. Dijkstra. 1989. Heterogeneity of macrophages in the rat evidenced by variability in determinants: two new anti-rat macrophage antibodies against a heterodimer of 160 and 95 kd (CD11/CD18). *J. Leukocyte Biol.* 46:556–564.
12. Sekiya, S., S. Gotoh, T. Yamashita, T. Watanabe, S. Saitoh, and F. Sendo. 1989. Selective depletion of rat neutrophils by *in vivo* administration of a monoclonal antibody. *J. Leukocyte Biol.* 46:96–102.
13. McWilliam, A.S., D. Nelson, J.A. Thomas, and P.G. Holt. 1994. Rapid dendritic cell recruitment is a hallmark of the acute inflammatory response at mucosal surfaces. *J. Exp. Med.* 179:1331–1336.
14. Sozzani, S., F. Sallusto, W. Luini, D. Zhou, L. Piemonti, P. Allavena, J. Van Damme, S. Valitutti, A. Lanzavecchia, and A. Mantovani. 1995. Migration of dendritic cells in response to formyl peptides, C5a, and a distinct set of chemokines. *J. Immunol.* 155:3292–3295.
15. Ibrahim, M.A.A., B.M. Chain, and D.R. Katz. 1995. The injured cell: the role of the dendritic cell system as a sentinel receptor pathway. *Immunol. Today*. 16:181–186.