

# A Novel Antioxidant Gene from *Mycobacterium tuberculosis*

By Sabine Ehrt,\* Michael U. Shiloh,† Jia Ruan,‡ Michael Choi,\*  
Stuart Gunzburg,\* Carl Nathan,† Qiao-wen Xie,‡ and Lee W. Riley\*

From the \*Division of International Medicine and Infectious Disease and †Division of Hematology-Oncology, Department of Medicine, Cornell University Medical College, New York 10021

## Summary

Among the major antimicrobial products of macrophages are reactive intermediates of the oxidation of nitrogen (RNI) and the reduction of oxygen (ROI). Selection of recombinants in acidified nitrite led to the cloning of a novel gene, *noxR1*, from a pathogenic clinical isolate of *Mycobacterium tuberculosis*. Expression of *noxR1* conferred upon *Escherichia coli* and *Mycobacterium smegmatis* enhanced ability to resist RNI and ROI, whether the bacteria were exposed to exogenous compounds in medium or to endogenous products in macrophages. These studies provide the first identification of an RNI resistance mechanism in mycobacteria, point to a new mechanism for resistance to ROI, and raise the possibility that inhibition of the *noxR1* pathway might enhance the ability of macrophages to control tuberculosis.

Each year, *Mycobacterium tuberculosis* kills nearly three million among the one-third of the world's population who are infected (1), making it the most deadly as well as one of the most successful bacterial pathogens of the human species. Tuberculosis arises in a small proportion of infected individuals in whose macrophages the bacteria replicate extensively, when, for example, malnutrition (2) or HIV (3) impede the cell-mediated immune response that normally leads to the activation of macrophages. However, in some cases, no host immune impairment is evident, and the question arises whether some disease-causing strains of *M. tuberculosis* have means to resist the toxic molecules produced by activated macrophages. Identification of a gene in a pathogen conferring resistance to a putative antimicrobial product of the host could be considered evidence that the host product exerts evolutionary pressure on the pathogen. Moreover, such genes may be considered virulence factors, and inhibition of their action might enhance host defense.

Two of the major antimicrobial mechanisms of activated macrophages depend on the synthesis of inorganic radical gasses by immunologically regulated flavocytochrome complexes that use NADPH to reduce molecular oxygen. When oxygen is the sole cosubstrate, the product is superoxide ( $O_2^-$ ; reference 4) when L-arginine is an additional cosubstrate, the product is nitric oxide (NO; reference 5).<sup>1</sup>

<sup>1</sup>Abbreviations used in this paper: ASN, acidified sodium nitrite; GSNO, S-nitrosoglutathione; LB, Luria-Bertani; NADPH, nicotinamide adenine dinucleotide phosphate (reduced); NO, nitric oxide; NOS2, nitric oxide synthase type 2; ORF, open reading frame; pBS, pBluescript; RNI, reactive nitrogen intermediates; ROI, reactive oxygen intermediates; RT, reverse transcriptase.

C. Nathan and L.W. Riley contributed equally to this study.

These radicals react with themselves, oxygen, transition metals, halides, sulfhydryls, and each other to produce a series of broadly cytotoxic products termed reactive oxygen intermediates (ROI) and reactive nitrogen intermediates (RNI), as well as at least one compound with features of both, peroxyxynitrite ( $OONO^-$ ; references 6 and 7).

*M. tuberculosis* resists ROI by a diversity of mechanisms. Phenolic glycolipids (8) and cyclopropanated mycolic acids (9) protect the cell wall, while catalase, alkylhydroperoxide reductase (10), and superoxide dismutase (11, 12) guard the cytosol. Moreover, *M. tuberculosis* may enter macrophages via complement receptors (13, 14), a pathway that fails to stimulate generation of ROI in some populations of macrophages (15). The ability of *M. tuberculosis* to mount such a broad defense against ROI implies that other products of the activated macrophage may be more important for tuberculosistasis. Indeed, activated murine macrophages selectively deficient in production of ROI were nonetheless mycobactericidal (16).

In contrast, abundant evidence establishes the importance of RNI in the control of mycobacteria, at least in the mouse. *M. tuberculosis* proliferates exuberantly in mice rendered selectively deficient in nitric oxide synthase type 2 (NOS2 or iNOS; reference 17). The organism also grows rapidly in mice made deficient in components of the cell-mediated immune response that normally leads to the induction of NOS2 (for review see reference 17), as well as in mice dosed with organochemicals (for review see references 2 and 17) or glucocorticoids (17) that inhibit the action or expression of NOS2. NOS2 was recently shown to be expressed by alveolar macrophages collected from the lungs of patients with tuberculosis (18) as well as pulmonary fibrosis (19).

However, in contrast to the situation with ROI, no spe-

cific mechanisms have been identified by which mycobacteria resist RNI. There was variability in the degree to which RNI inhibited several mycobacterial strains in vitro (20, 21), and the more resistant strains of *M. tuberculosis* were more virulent in guinea pigs (21). *M. tuberculosis* strain CB3.3, a drug-susceptible clinical isolate, caused >10% of the tuberculosis cases reported in New York City between 1992 and 1993 (22) and was the most RNI resistant of those tested (23). Reasoning that RNI-resistant strains may express RNI resistance genes, we used a library from *M. tuberculosis* CB3.3 to clone a gene that does not resemble previously recognized antioxidants, but which protects transformed enteric and mycobacteria from both RNI and ROI.

## Materials and Methods

**Bacterial Strains and Growth Conditions.** The following mycobacterial strains were used in this study: *M. smegmatis* mc<sup>2</sup>155 (24), *M. tuberculosis* CB3.3 (22), *M. tuberculosis* H37Ra (American Type Culture Collection/[ATCC, Rockville, MD] No. 25177), *M. tuberculosis* H37Rv (ATTC No. 25618), *M. bovis* (ATTC No. 19210), *M. africanum* (ATTC No. 25420), *M. goodii* (ATTC No. 14470), *M. fortuitum* (ATTC No. 6841), *M. avium* (ATTC No. 25291), *M. intracellulare* (ATTC No. 13950), *M. kansasii* (ATTC No. 12478). Mycobacterial strains were grown in Middlebrook 7H9 broth (Difco Labs. Inc., Detroit, MI) supplemented with 2% glycerol, 0.05% Tween 80, and ADC supplement (Difco Labs. Inc.) or plated on 7H11 agar (Difco Labs. Inc.). Luria-Bertani (LB) broth or LB agar were used for *Escherichia coli* strains XL1-Blue (Stratagene Inc., La Jolla, CA), HB101 (ATTC), DH5 $\alpha$  (GIBCO/BRL, Gaithersburg, MD), M15 (QIAGEN Inc., Chatsworth, CA), GC4468 and DJ109 (a *soxRS* mutant, provided by T. Nunoshiba and B. Demple, Harvard School of Public Health, Boston, MA; reference 25), and JTG100 and its *oxyR*-deficient derivative, JTG101 (26), also the gift of B. Demple. Ampicillin and hygromycin B (Sigma Chemical Co., St. Louis, MO) were used at 100  $\mu$ g/ml and 200  $\mu$ g/ml, respectively, to grow *E. coli*. Hygromycin B was used at 50  $\mu$ g/ml to grow mycobacteria.

**Plasmids.** pBluescript (pBS; Stratagene Inc.) served as a cloning vector and pQE31 (QIAGEN Inc.) as an expression vector in *E. coli*. The shuttle vector pOLYG (a gift from Peadar O'Gaora and D.B. Young, Department of Medical Microbiology, St. Mary's Hospital Medical School, London, UK) is a derivative of p16R (27). pSMT3 is derived from pOLYG and contains the hsp60 promoter for overexpression of genes in mycobacteria (also a gift from P. O'Gaora).

**Cloning of *M. tuberculosis* DNA Fragment Associated with Resistance against RNI.** Chromosomal DNA was isolated from *M. tuberculosis* CB3.3 as previously described (28) and digested with EcoRI and BamHI. A genomic library was constructed by ligation of the DNA fragments into the *E. coli* vector pBS. *E. coli* XL1-Blue was initially tested for growth in LB at various pH and NaNO<sub>2</sub> concentrations. At pH 6.0 and 10 mM NaNO<sub>2</sub>, its growth was completely suppressed. Therefore, the genomic library was electroporated into strain XL1-Blue and the recombinants were screened for growth in acidified sodium nitrite (ASN; LB at pH 6.0 containing 10 mM NaNO<sub>2</sub>).

**Construction of pNO14.1 and its Open Reading Frame Mutant.** A HindIII-SmaI fragment of pNO14 was cloned into the Hind-

III and HincII sites of pBS. The resulting plasmid, pNO14.1, still contains the complete open reading frame (ORF)1. A point mutation was introduced at codon 12 of ORF1 (nucleotide 36 G  $\rightarrow$  T), which created a stop codon (TGA) at that position. This mutation does not affect the amino acid sequence of the putative protein encoded by ORF2. The mutation was introduced by PCR mutagenesis. The final construct, pNO14.1-mut1, was confirmed by sequencing.

**Analysis of the Resistant Phenotype.** The relative resistance to chemically generated RNI was tested by inoculating a 1:100 dilution of the overnight or log-phase culture of the strain to be tested into 3 ml of LB, pH 5.3, or 7H9, pH 5.3, containing NaNO<sub>2</sub> (Sigma Chemical Co.) at various concentrations. Resistance to paraquat (Sigma Chemical Co.), nalidixic acid (Sigma Chemical Co.), NaOCl (Aldrich Chemical Company, Inc., Milwaukee, WI), and ethanol was measured in LB, pH 7.0. Cultures were incubated at 37°C for the indicated periods. The chosen conditions resulted in at least a 10-fold reduction in CFU of the control *E. coli* HB101. The number of viable bacteria was determined by plating on LB agar. Alternatively, where indicated, a microplate assay was used that detects the ability of surviving bacteria to reduce a formulation of resazurin termed AlamarBlue<sup>®</sup> (Sensititre/Alamar; AccuMed International Companies, Westlake, OH) to a fluorescent product (29).

**Northern Blot Analysis and Reverse Transcriptase PCR.** RNA was isolated from 20 ml logarithmically growing mycobacterial cultures according to the FastPrep FP120 bead beater apparatus (Bio-101, La Jolla, CA) protocol. The integrity of the RNA preparation was verified by the presence of two sharp rRNA bands, 1,500 and 3,100 nucleotides in length.

Two oligonucleotides, RNA-1 (5'-gacgcgctgatcgccgatcgcgcgctggtggtcg-3') and RNA-2 (5'-cggcaacgccggtgacaacgcgcgggcatcctcgccc-3') were labeled with dioxigenin using the oligonucleotide tailing kit (Boehringer Mannheim Biochemicals, Indianapolis, IN) to serve as probes for the transcripts of the two open reading frames, ORF1 and ORF2, respectively. Northern blots were carried out according to the Genius<sup>™</sup> System User's Guide For Membrane Hybridization (Boehringer Mannheim Biochemicals). Reverse transcriptase (RT)-PCR was performed using the Perkin Elmer Gene Amp RNA PCR Kit (Perkin Elmer Corp., Norwalk, CT). In brief, 0.2–1  $\mu$ g total RNA was transcribed by Moloney MuLV RT into cDNA with either random hexamer primers or a specific primer for the ORF1 RNA at 42°C for 15 min. cDNA specific primers were added (Ia: 5'-ctaccgcgcgaggtagcagctgacc-3'; Ib: 5'-cggcaacgccggtgacaacgcgcgggcatcctcgccc-3'; IIa: 5'-gggatggcggtgggtgctgctcg-3'; IIb: 5'-gacgcgctgatcgccgatcagcgcgcatggtggtcg-3'), and the reaction was carried out with AmpliTaq (Perkin-Elmer, Norwalk, CT) DNA polymerase in a volume of 100  $\mu$ l. The combined annealing and extending reaction was done at 60°C for 30 s.

**Protein Expression.** *noxR1* was cloned behind an inducible T5 promoter into the expression vector pQE-31 (QIAGEN Inc.). M15 (pREP4) pQE-31-ORF1 were grown in LB containing 100 mg/l ampicillin and 25 mg/l kanamycin to an OD<sub>580</sub> of 1.0 and induced with 1.5 mM IPTG (isopropyl- $\beta$ -thiogalactopyranoside). After 4 h, bacteria were harvested and a sample of lysate was subjected to SDS-PAGE and Coomassie blue staining to check for overexpression of recombinant protein. Protein containing an NH<sub>2</sub>-terminal histidine tag was purified on Ni-NTA resin columns (QIAGEN Inc.) and analyzed by SDS-PAGE. The NH<sub>2</sub>-terminal sequence of the purified protein was established for 19 residues, sufficient to read beyond the tag and 7 residues into NoxR1 proper. The purified protein was injected in female New Zealand White rabbits (4 injections of 100  $\mu$ g NoxR1, at 4 wk

intervals). The resulting antiserum was used for immunoblot analysis of bacterial lysates or purified protein by standard procedures. Affinity-purified antibody was prepared as described in Results.

**Studies in Macrophages.** An assay modified from that previously described (30) was used to determine the survival of *M. smegmatis* strains inside macrophages. Wild-type (iNOS<sup>+/+</sup>) and iNOS-deficient mice (iNOS<sup>-/-</sup>; C57BL/6x129/SvEv) (31) and wild-type and *phox-91*-deficient mice (C57BL6/J) (32) were injected intraperitoneally with 1.0 ml of sterile, freshly prepared 5 mM sodium periodate (Sigma Chemical Co.) in PBS 4 d before harvest. The mice were killed by cervical dislocation or CO<sub>2</sub> inhalation. Peritoneal cells were harvested by lavage with 10 ml ice-cold sterile RPMI (Sigma Chemical Co.), pH 7.2. Cells were collected by centrifugation for 10 min at 250 *g* at 4°C and resuspended in RPMI medium supplemented with 10% heat-inactivated fetal bovine serum (HyClone, Logan, UT), 1% glutamine (complete medium), and 10 μg/ml gentamicin. Viable cells were counted on a hemocytometer in the presence of trypan blue and the proportion of macrophages determined by differential count of Diff-Quik stained Cytospin (Shandon, Inc., Pittsburgh, PA) preparations. Peritoneal cells (4 × 10<sup>5</sup>, ~50% macrophages) were plated in 96-well tissue-culture plates (Corning Medical and Scientific, Medfield, MA) at 100 μl per well. In some experiments, recombinant mouse IFN-γ (Genentech, South San Francisco, CA) was added at 50 U/ml. The plates were incubated at 37°C in 5% CO<sub>2</sub>, and 12–24 h later the adherent monolayers were washed twice with sterile PBS to remove gentamicin containing medium; complete removal required that the plates be emptied with a hard flick in each wash. Fresh complete medium ± mIFN-γ was added, and 24–48 h after the initial plating the cells were washed again with sterile PBS, and reconstituted with complete medium before infection.

Freshly electroporated *M. smegmatis* were grown to mid-log phase. Bacteria were opsonized in 10% fresh mouse serum for 30 min at 37°C, and 10 μl of the opsonized bacteria (~2 × 10<sup>5</sup>) were added to each well. The plates were centrifuged for 5 min at 250 *g* to synchronize the infection, and were then incubated at 37°C for 30 min to allow phagocytosis. The wells were washed three times with sterile PBS to remove free bacteria. Complete medium (100 μl) containing 10 μg/ml gentamicin was added to each well, and the plates were incubated at 37°C. Gentamicin was added to prevent extracellular replication of mycobacteria that may not have been internalized or may have escaped from dying macrophages. Samples were taken at the indicated time points from individual wells in triplicate, as follows. The medium was removed and 50-μl aliquots were saved for nitrite determination. The cell monolayer was washed twice with PBS and lysed with 100 μl 0.1% sodium deoxycholate in PBS. Appropriate dilutions of the lysates were plated onto LB plates containing 50 μg/ml hygromycin B for CFU determinations.


To monitor macrophage production of NO, we measured nitrite in the culture supernatants as an accumulating oxidation product. The Griess reaction was performed as previously described (33). *M. smegmatis* itself produced no detectable nitrite under the conditions of these experiments, as evidenced in cultures with iNOS<sup>-/-</sup> macrophages (see Fig. 7 B, inset).

Macrophage production of hydrogen peroxide was assessed by the horseradish peroxidase-catalyzed oxidation of fluorescent scopoletin to a nonfluorescent product, using a microplate format (34).


Decreased production of NO<sub>2</sub><sup>-</sup> or hydrogen peroxide and diminished bactericidal activity could not be attributed to differential loss of macrophages from the monolayers, as monitored by measurements of adherent cell protein in the same cultures (34).

## Results

**Cloning of an *M. tuberculosis* Gene Associated with Resistance to RNI.** *E. coli* XL1-Blue was electroporated with a genomic library of *M. tuberculosis* CB3.3 and exposed for 24 h to ASN (6 mM NaNO<sub>2</sub>, pH 6.0). Protonation of NaNO<sub>2</sub> generates HNO<sub>2</sub>, whose dismutation provides NO and nitrate, and, through reaction of NO with oxygen, other RNI (35, 36). The main products of these reactions are probably dinitrogen tri- and tetra-oxides (N<sub>2</sub>O<sub>3</sub> and N<sub>2</sub>O<sub>4</sub>) as well as *S*-nitrosothiols, which have profound bacteriostatic effects (37, 38, 39). ASN plays a physiologic role in the microbicidal system of the stomach (40) and the combination of RNI and low pH also mimics aspects of the intraphagolysosomal milieu of the macrophage (5, 41). The chosen conditions killed *E. coli* XL1-Blue efficiently (>7



**Figure 1.** Plasmid pNO14 confers enhanced survival on *E. coli* in ASN. (A) Survival of *E. coli* HB101 pBS (open squares) and HB101 pNO14 (solid squares) in ASN medium. Overnight cultures were diluted 100-fold into LB, pH 5.3, containing 10 mM NaNO<sub>2</sub>, and incubated at 37°C. CFUs were determined at indicated times by plating serial dilutions on LB agar containing ampicillin. In this and subsequent panels, results are means ± SE for triplicates; some error bars fall within the symbols. (B) Growth of *M. smegmatis* pOLYG (open symbols) and *M. smegmatis* pOLYG-NO14 (solid symbols) in 7H9 medium containing 30 mM sodium nitrite at pH 7.4 (triangles) or 30 mM sodium nitrate at pH 5.3 (squares). CFU were determined by plating on LB agar containing 50 μg/ml hygromycin, either at onset of culture or after 12 h. (C and D) Survival of *M. smegmatis* pOLYG (open squares) and *M. smegmatis* pOLYG-NO14 (solid squares) in ASN medium. Bacteria were grown to an optical density of A<sub>600</sub> = 4.5 (stationary phase cultures, C) or to A<sub>600</sub> = 1.5 (logarithmic phase cultures, D) and diluted 100-fold in 7H9 medium, pH 5.3, containing 30 mM NaNO<sub>2</sub> for stationary phase cultures and 20 mM NaNO<sub>2</sub> for logarithmic phase cultures. At indicated times CFUs were determined by plating appropriate dilutions on LB agar containing 50 μg/ml hygromycin.



**Figure 2.** Molecular analysis of the cloned DNA. (A) Presence of *noxR1* in different species of mycobacteria. Chromosomal DNA (1  $\mu$ g) of each strain was digested to completion with *EcoRI* and analyzed by Southern blotting with a digoxigenin-labeled probe corresponding to the 0.68-kb fragment from pNO14. Lane 1,  $M_r$  markers; lane 2, *M. tuberculosis* CB3.3; lane 3, *M. tuberculosis* H37Ra; lane 4, *M. tuberculosis* H37Rv; lane 5, *M. bovis*; lane 6, *M. africanum*; lane 7, *M. goodii*; lane 8, *M. fortuitum*; lane 9, *M. smegmatis*; lane 10, *M. avium*; lane 11, *M. intracellulare*; lane 12, *M. kansasii*. (B) Map of the inserts in pNO14 and pNO16. (C) Nucleotide sequence of the NO14 fragment and amino acid sequence of the putative protein encoded by ORF1 (*noxR1*). These sequence data are available from the EMBL/GenBank/DBJ under accession number Y08323. (D) Hydrophilicity plot of putative protein (NoxR1) encoded by ORF1.

log<sub>10</sub> in 24 h). Surviving transformants were detected with a frequency of  $<10^{-6}$ . One recombinant plasmid, called pNO14, consistently conferred upon XL1-Blue and four other *E. coli* hosts (DH5 $\alpha$ , HB101, GC4468, and JTG100) as well as *Salmonella typhimurium* LT2 and 14028 (data not shown) an enhanced ability to resist ASN. By 12 h, transformation of HB101 with pNO14 afforded a 50-fold increase in survival above transformation of the same *E. coli* host with the vector alone (Fig. 1 A).

Means are not yet available for the facile genetic manipulation of *M. tuberculosis*. As a substitute mycobacterial species in which to analyze the function of pNO14, we chose *M. smegmatis*; the organism is fast-growing and easily transformable, and, as shown below, lacks a chromosomal copy of the gene carried by pNO14. The 680-bp mycobacterial DNA fragment of pNO14 was cloned into the shuttle plasmid pOLYG (27) and electroporated into *M. smegmatis* mc<sup>2</sup>155 (24). Neither the pOLYG-NO14- nor the vector-transformed strain was killed by sodium nitrate at pH 5.3, nor by sodium nitrite at pH 7.4 (Fig. 1 B). These findings excluded a mycobactericidal effect of acid plus nitrate (the nonreactive dismutation product of HNO<sub>2</sub>), and of nitrite under conditions that disfavor the formation of NO. In contrast, when recombinant and control strains were exposed to nitrite at pH 5.3, both strains succumbed. Compared to its vector-transformed counterpart, *M. smegmatis* transformed with pOLYG-NO14 was increasingly less susceptible after 6 h, attaining a 100-fold relative advantage by 15 h (Fig. 1 C). Logarithmically growing *M. smegmatis* (Fig. 1 D) were more susceptible to ASN than those at stationary growth phase (Fig. 1 C). However, the protective effect of

pOLYG-NO14 was manifest in both phases. Thus, a gene product encoded by NO14 conferred relative resistance to RNI upon both *E. coli* and *M. smegmatis*.

**Expression Analyses.** The cloned 680-bp fragment hybridized to genomes of the members of the *M. tuberculosis* complex. None of the other mycobacterial strains tested hybridized with this probe (Fig. 2 A). Thus, among mycobacteria, the cloned DNA appears to be specific for the members of the *M. tuberculosis* complex associated with human and rodent disease.

Plasmid pNO14's 680-bp *EcoRI*/*BamHI* fragment from *M. tuberculosis* included two overlapping ORFs oriented in opposite directions (Fig. 2 B). ORF1 encodes a putative protein of 152 amino acids (Fig. 2 C) in an unusual pattern of alternating hydrophilic and hydrophobic segments that are shorter than typical membrane-spanning domains (Fig. 2 D). The putative protein was predicted to have an  $M_r$  of 15,514 daltons and a pI of 10.6, the basicity chiefly reflecting the relative abundance of Arg residues (9.2 mole percent). Despite the presence of four cysteines, no structural motifs were recognized, nor were there homologous sequences in the database. ORF2 did not contain a stop codon. Therefore, we isolated by colony hybridization a plasmid termed pNO16, which contained the full length ORF2, consisting of 1074 bases, from a library of *M. tuberculosis* CB3.3 (Fig. 2 B). The putative protein encoded in ORF2 is 40% identical to XP55, a *Streptomyces lividans* secretory product of unknown function (42).

To find out which ORF was responsible for the observed phenotype, we determined which was transcribed in recombinant *M. smegmatis*. In Northern blots with an oli-




Methods). Wild-type *noxR1* encoded by pNO14.1, but not its ORF-1 mutant pNO14.1-mut1, protected the bacteria from ASN (Fig. 3 C). Thus, we have been able to detect, purify, and sequence NoxR1 protein only when a NoxR1 fusion was overexpressed in association with toxicity; we have not been able to detect NoxR1 protein under conditions where lower levels of expression were presumed and a phenotype was conferred. Nonetheless, translation of the *noxR1* transcript appears to be required to confer resistance to ASN.

*noxR1* also Confers Resistance to ROI and H<sup>+</sup>. To more fully explore the phenotype afforded by expression of *noxR1*, we made use of a fluorescent, dye-reduction microplate assay whose results corresponded almost perfectly (correlation coefficients  $r^2 > 0.96$ ) to the results of the more laborious colony-forming agar-plate assay after exposing bacteria to RNI in vitro or to the intraphagosomal milieu of macrophages (29). Not only was *E. coli* HB101 rendered far more resistant to ASN (2.5 mM NaNO<sub>2</sub>, pH 5) by expression of *noxR1* (data not shown), but the bacteria also better resisted *S*-nitrosoglutathione (GSNO; 2 mM, pH 5.0), a physiologic and bacteriostatic (37, 43, 44) source of several RNI, including ammonia (45; Fig. 4, A and C). By 6 h, the survival advantage was close to 100-fold. GSNO was more bactericidal at pH 5.0 (Fig. 4, A and C) than at pH 7.0 (Fig. 4 E), but *noxR1* conferred protection under both conditions.


Unexpectedly, *noxR1* also conferred resistance to H<sub>2</sub>O<sub>2</sub> (survival advantage, >100-fold) by 6 h, and in this case H<sup>+</sup> also appeared to show a small synergistic effect on the killing (pH 5.0, Fig. 4, B and D; pH 7.0, Fig. 4 F). Furthermore, *noxR1* protected *E. coli* from the synergistic cytotoxicity afforded by GSNO plus H<sub>2</sub>O<sub>2</sub> at concentrations of each agent that were harmless singly (Fig. 4 G), and from the cytotoxicity afforded by the combination of three species likely to be simultaneously present in some phagosomes: GSNO, H<sub>2</sub>O<sub>2</sub>, and H<sup>+</sup> (Fig. 4 H).

The results to this point did not allow us to distinguish whether *noxR1* might protect relatively specifically against RNI and ROI, or might instead confer a general survival or repair function effective against virtually any threat to bacterial viability. The distinction can be hard to draw, since so many forms of insult, such as heat shock and ultraviolet irradiation, lead directly or indirectly to formation of ROI. Nevertheless, we subjected *noxR1*- and vector-transformed *E. coli* to several other stresses: elevated concentrations of H<sup>+</sup>, sodium chloride, detergent, ethanol, nalidixic acid, paraquat, hypochlorous acid, and heat. *noxR1* did confer relative resistance to acid at pH 4.5, the lowest level reported in phagosomes (46) as well as at pH 4.0 (Fig. 5 A). At pH 5 and above, no growth inhibition was detectable (Fig. 1 B and data not shown). In addition, *noxR1* protected *E. coli* against killing by hypochlorous acid (10 mM). In contrast, *noxR1* afforded no protection against the growth-inhibiting effects of high salt, SDS, heat, ethanol, nalidixic acid, or paraquat (Fig. 5, B–E).

*Independence of noxR1 Effects from the oxyR and soxRS Regulons.* *oxyR* and *soxRS* are multigenic regulons (25),




**Figure 4.** Survival advantage conferred by *noxR1* on *E. coli* cultured in GSNO and/or hydrogen peroxide. (A and B) Time course of survival of *E. coli* HB101 transformed with pBS (open squares) or pNO14 (solid squares) and cultured in LB at pH 5.0 with (A) 2 mM GSNO or (B) 0.5 mM H<sub>2</sub>O<sub>2</sub>, as determined by the dye reduction microplate assay. (C–F) Concentration-response curves for GSNO (C and E) and H<sub>2</sub>O<sub>2</sub> (D and F) after 6 h incubation at a pH of 5 (C and D) or pH 7 (E and F). (G and H) Resistance to the combination of GSNO and H<sub>2</sub>O<sub>2</sub>. (Black bars) *E. coli* HB101 transformed with pBS. (Gray bars) *E. coli* HB101 transformed with pNO14. (G) pH 7.0 with the following treatments: I, none; II, H<sub>2</sub>O<sub>2</sub> (0.1 mM); III, GSNO (5 mM); IV, H<sub>2</sub>O<sub>2</sub> (0.05 mM) + GSNO (5 mM). (H) pH 5.0 with the following treatments: I, none; II, H<sub>2</sub>O<sub>2</sub> (0.1 mM); III, GSNO (1 mM); IV, H<sub>2</sub>O<sub>2</sub> (0.05 mM) + GSNO (1 mM). Data in panels A–F are means ± SE for triplicates; some error bars fall within the symbols. Data in panels G and H are means of duplicates.



**Figure 5.** Effect of transformation with *noxR1* on resistance of *E. coli* to various stress conditions. Survival of *E. coli* HB101 transformed with pBS (black bars) or pNO14 (gray bars) after 3 or 6 h incubation in LB medium with the following treatments: (A) LB at pH 4.5 and pH 4.0 for 3 h; (B) addition of 0.5 M and 1.0 M NaCl for 3 h; (C) addition of 2 and 5% SDS for 3 h; (D) addition of 10% ethanol or 200 µg/ml nalidixic acid for 6 h or incubation at 50°C for 3 h; (E) addition of 100 mM paraquat for 6 h or 10 mM NaOCl for 3 h. Control bars reflect growth at pH 7.0. Survival was quantified by plating appropriate dilutions on LB agar or by a dye reduction microplate assay (see Materials and Methods). Results are means ± SE for triplicates.

each of which is activated by and confers resistance to both ROI and RNI in *E. coli* (47–49). In *M. tuberculosis*, *oxyR* is disrupted and *soxRS* is undescribed (9, 10, 50, 51), whereas in *E. coli* neither regulon contains any sequence homologous to *noxR1*. Nonetheless, we wished to test whether *noxR1* from *M. tuberculosis* might function in recombinant *E. coli* through activation of *oxyR* or *soxRS*. The genes controlled by these factors number at least 19, including those encoding superoxide dismutase, NADPH:ferredoxin oxidoreductase, fumarase, DNA repair endonuclease IV, catalase, alkylhydroperoxide reductase, and glutathione reductase. Accordingly, we transformed *E. coli* deficient in either of these two regulons and their corresponding wild-type strains with pNO14 or pBS vector alone. In all four hosts, *noxR1* conferred resistance to ASN (Fig. 6). The degree of protection depended on the host cell type, but not on its expression of *oxyR* or *soxRS*. Thus, genes dependent




**Figure 6.** *noxR1*-mediated resistance to ASN is independent of *oxyR* and *soxRS*. Survival of the following *E. coli* hosts transformed with pBS (open squares) or pNO14 (solid squares) in ASN medium: (A) JTG100 (*oxyR* wild type); (B) JTG101 (*oxyR* deficient); (C) GC4468 (*soxRS* wild type); (D) DJ901 (*soxRS* deficient). Overnight cultures were diluted 100-fold into LB pH 5.0 containing 10 mM NaNO<sub>2</sub> and incubated at 37°C. Resistance was quantified by a dye reduction microplate assay (see Materials and Methods). Results are means ± SE for triplicates; some error bars fall within the symbols.

upon *oxyR* or *soxRS* are dispensable for the function of *noxR1*, unless *noxR1* can substitute for *oxyR* or *soxRS* to induce their expression.

**Effect of *noxR1* on Survival of *M. smegmatis* within Macrophages.** The observation that *noxR1* protects *M. smegmatis* from the antibacterial effects of RNI and ROI in vitro prompted us to explore the effect of this gene on the survival of *M. smegmatis* inside activated macrophages, where the full complement of antibacterial mechanisms is undefined.

Macrophages were collected from the peritoneal cavity of mice 4 d after intraperitoneal injection of sodium periodate (52). Periodate, a lymphocyte mitogen, stimulates cytokine production (53). Macrophages from periodate-injected mice have a respiratory burst capacity typical of macrophages activated by infection of the host with mycobacteria (54) or macrophages treated in vitro with cytokines (55). Periodate-elicited macrophages respond to further inductive signals, such as in vitro incubation with IFN-γ, by producing NO (33). Both properties were confirmed in wild-type macrophages in this experiment; in addition, *M. smegmatis* alone was sufficient to trigger NO production in periodate-elicited macrophages without exposure to exogenous cytokines in vitro (Fig. 7, insets).

To vary the genotype of the macrophages along with that of the mycobacteria, wild-type mice of C57BL/6x129/SvEv background were matched with NOS2-deficient mice



**Figure 7.** Survival of *noxR1*-transformed *M. smegmatis* in wild-type and genetically altered macrophages. Periodate-elicited peritoneal macrophages from (A) wild-type C57BL/6x129/SvEv, (B) *iNOS*<sup>-/-</sup> C57BL/6x129/SvEv, (C) wild-type C57BL/6, or (D) *phox91*<sup>-/-</sup> C57BL/6 mice were pretreated or not with mouse IFN- $\gamma$  and infected with *M. smegmatis* transformed with either pOLYG (black bars) or pOLYG-NO14 (gray bars). At indicated times, macrophages were lysed and surviving bacterial CFU were determined. Values are means  $\pm$  SE for triplicate macrophage cultures as a percentage of the starting CFU, defined as the numbers of CFUs recoverable from the cells after the 30 min uptake period; the latter averaged  $2 \times 10^5$ /well, or  $\sim 1$ /macrophage. (Insets) Nitrite accumulation (nmol/well) in the same cultures of macrophages infected with *M. smegmatis*-pOLYG (squares) or pOLYG-NO14 (circles). Values are means  $\pm$  SE; error bars fall within the symbols. One of six similar experiments (three with IFN- $\gamma$  and three without).

on the same background (31). Similarly, wild-type C57BL/6 mice were compared with *Phox91*-deficient (respiratory burst, oxidase-null) mice backcrossed to C57BL/6. As expected, H<sub>2</sub>O<sub>2</sub> production was preserved and NO production was abolished in macrophages from NOS2-deficient mice, while the reverse was the case in macrophages from *Phox91*-deficient mice; neither class of knock-out macrophages overproduced the opposite product (Shiloh, M.U., unpublished data). In all settings described below, results were similar with or without addition of exogenous IFN- $\gamma$  to the already activated macrophages (data not shown), and have been pooled.

On average, activated macrophages from wild-type C57BL/6x129/SvEv mice killed 80% of ingested pOLYG-transformed (control) *M. smegmatis* by 5.6 h and 97% by 24 h (Fig. 7 A; interpolation from data in Table 1). When *M. smegmatis* was transformed with *noxR1*, it took the wild-type macrophages twofold longer to kill 80% of them. Moreover, at each time point tested (6, 12, and 24 h), approximately twofold more bacteria survived (Fig. 7 A; Table 1).

Macrophages that were genetically incapable of expressing NOS2 (31) were no less efficient at killing *M. smegmatis* than wild-type macrophages of the same genetic background. 80% of the bacteria were killed by 6 h and 96% were killed by 24 h. (Fig. 7 B; Table 1). This indicated that

RNI do not play a major role in the control of *M. smegmatis* in vitro, in contrast to the situation with *M. tuberculosis* (see references 16 and 17, and the references cited therein) and *M. leprae* (56). Nonetheless, expression of *noxR1* protected the bacteria, delaying the time of 80% killing by a factor of 2.7-fold and resulting in 1.6–1.7-fold more surviving organisms at each time point tested (Table 1). This indicated that the protective action of *noxR1* is not directed exclusively against RNI, consistent with the evidence in vitro showing that *noxR1* also protects against ROI.

Wild-type C57BL/6 macrophages killed wild-type (pOLYG-transformed) *M. smegmatis* more slowly (80% killing by 12.5 h) than did wild-type C57BL/6x129/SvEv macrophages (80% killing by 5.6 h). Nonetheless, expression of *noxR1* in *M. smegmatis* delayed the 80% killing time by a factor of 1.75-fold and resulted in 1.6–2.5-fold more surviving organisms at each time point tested (Fig. 7 C; Table 1).

C57BL/6 macrophages deficient in *Phox91* were strikingly impaired in killing wild-type *M. smegmatis*; within the 24 h period of observation, 80% killing was not often attained (Fig. 7 D; Table 1). This indicated that ROI play a prominent, albeit not an exclusive, role in killing *M. smegmatis*, in contrast to the situation with *M. tuberculosis*, where a minimal role for ROI was evident (16). In ROI-deficient macrophages, *M. smegmatis* that expressed *noxR1* survived 1.4–1.9-fold better than vector-transformed *M. smegmatis* at



**Table 1.** Percentage of Survival of Vector- or *noxR1*-transformed *M. smegmatis* after the Indicated Time Periods Compared to their CFU at Time 0 in Mouse Macrophages

Mouse genotype	Number of Exp.	Time	Surviving vector-transformed <i>M. smegmatis</i> * (± SE)	Surviving <i>noxR1</i> -transformed <i>M. smegmatis</i> † (± SE)	<i>P</i> §	Fold protection   (± SE)
		<i>h</i>				
C57BL/6x	6	6	14.2 (1.9)	28.3 (3.7)	<.005	2.03 (0.21)
129Sv/Ev		12	9.7 (1.5)	18.3 (2.5)	<.01	1.97 (0.21)
		24	2.8 (0.4)	5.2 (0.3)	<.0001	2.06 (0.29)
iNOS <sup>-/-</sup>	6	6	19.0 (3.6)	29.9 (5.3)	>.05	1.56 (0.08)
		12	14.0 (2.6)	24.9 (4.5)	<.05	1.56 (0.14)
		24	4.1 (0.7)	7.0 (1.0)	<.05	1.71 (0.20)
C57BL/6	6	6	32.1 (3.1)	49.8 (3.8)	<.001	1.61 (0.16)
		12	21.0 (2.8)	38.9 (4.1)	<.001	2.06 (0.14)
		24	6.7 (1.2)	15.8 (2.8)	<.01	2.49 (0.33)
Phox91 <sup>-/-</sup>	6	6	84.2 (9.1)	110.0 (8.3)	<.05	1.42 (0.18)
		12	61.9 (4.7)	96.4 (4.2)	<.0001	1.67 (0.20)
		24	22.0 (3.3)	39.1 (3.7)	<.005	1.93 (0.27)

\*‡Means ± SE for the number of independent experiments indicated, each in triplicate. The number of CFUs per well at time 0 (see Materials and Methods) was defined as 100%, and averaged  $2 \times 10^5$ , or  $\sim 1$ /macrophage.

\* *M. smegmatis* transformed with pOLYG vector alone.

† *M. smegmatis* transformed with pOLYG-NO14.

§ Two-tailed Student's *t* test.

|| The percentage of surviving *M. smegmatis* pOLYG-NO14 divided by the percentage of surviving *M. smegmatis* pOLYG at each time point was calculated for each experiment and averaged.

each time point tested (Fig. 7 D; Table 1). These findings suggested that ROI and another product represent redundant killing mechanisms for *M. smegmatis*, the former more effectively than the latter; in the absence of ROI, less potent killing by RNI, H<sup>+</sup>, or another product is manifest, against which *noxR1* affords protection.

## Discussion

To our knowledge, this is the first study of macrophage-pathogen interactions in which both the macrophages and the pathogens have been genetically modified, such that the host cells do or do not express specific cytotoxic mechanisms, and the bacteria do or do not express a presumptive resistance pathway directed against those mechanisms. By this analysis, *noxR1*, a novel gene from *M. tuberculosis*, confers partial resistance to three of the major antimicrobial products of macrophages, the cells ultimately responsible for controlling tuberculosis. However, the greater resistance conferred on *noxR1*-transformed *M. smegmatis* in vitro than in macrophages strongly suggests that there are macrophage antimycobacterial products other than RNI, ROI, and H<sup>+</sup>, and that *noxR1* protects against some physiologically relevant stresses but not others. Likewise, the relative selectivity of *noxR1*'s protective effects in vitro and

the restriction of the *noxR1* gene to a subset of pathogenic mycobacterial species argue that *noxR1* is neither a general stress resistance gene nor a housekeeping gene.

Cloned from a prevalent clinical isolate of *M. tuberculosis*, *noxR1* was only identified in the genome of members of the *M. tuberculosis* complex. *noxR1* was absent from the chromosome of mycobacteria considered nonpathogenic or opportunistically pathogenic for humans, including *M. smegmatis*, *M. avium*, and *M. intracellulare*. We do not know if a *noxR1*-like gene is present in any other organisms, nor whether *noxR1* is transcribed naturally by any mycobacteria besides the *M. tuberculosis* strain we tested. It remains to be determined whether the natural gene may be regulated by environmental conditions, including the stresses against which it confers resistance. It would be of particular relevance to know how much NoxR1 is expressed by *M. tuberculosis* that resides in phagolysosomes.

A major mystery is *noxR1*'s mechanism of action. With no homologies or motifs recognized at nucleotide or amino acid levels, the sequence afforded few clues. Because so little NoxR1 appears to be expressed, it is unlikely that its four cysteine residues merely serve to titrate ROI or RNI, as homocysteine is thought to do in *Salmonella typhimurium* (44), or as metallothionein may do when overexpressed in hepatocytes (57). In *E. coli*, the DNA-binding protein en-

coded by *dps* protects DNA from oxidative damage (58, 59), and *noxR1* might work in a similar manner. Its effectiveness in a heterologous mycobacterium whose genome lacks *noxR1* may argue against a role as a transcription factor, and its effectiveness in *oxyR*- and *soxRS*-deficient *E. coli* argues against *noxR1* activating those two regulons in particular. The *oxyR* homolog of *M. tuberculosis* contains numerous deletions and frameshifts and is nonfunctional (9, 60). Perhaps NoxR1 compensates for the loss of OxyR in *M. tuberculosis*, similar to the suggested role of AhpC (10, 50).

If NoxR1 is an enzyme, the novelty of its sequence suggests that it may work differently than those already known to affect RNI. The latter serve to alter the proportions of various RNI in a mixture. Thus, in vitro, mammalian thioredoxin reductase can catalyze the NADPH-dependent reduction of *S*-nitrosoglutathione to glutathione and an NO-like species (61), whereas superoxide dismutase favors the accumulation of NO at the expense of its conversion to peroxynitrite. At present, the yield of recombinant NoxR1 has been compromised by its apparent autotoxicity upon

overexpression, and this has precluded biochemical assays of hypothesized actions.

The physiologic role of NoxR1 cannot be established until it is possible to inactivate *noxR1* selectively in *M. tuberculosis* and compare the growth of the organism in the mammalian host with the growth of isogenic *M. tuberculosis* to which *noxR1* has been restored. Until then, the possibility remains that the actions of *noxR1* observed in transformed bacteria are artefacts of over or heterologous expression. Weighing against this concern is that *noxR1* conferred a protective phenotype only when expressed at low levels, and did so in diverse species and strains.

There is an urgent need for new antitubercular drugs. Almost all currently used antibacterials manifest their antibacterial activity in pure culture, and new candidates are screened on that basis. Such screens could miss compounds that inhibit a pathway which is dispensable for bacterial growth in vitro but which confers a survival advantage on the pathogen within the infected host. In this sense, inhibitors of NoxR1 may warrant investigation as possible prototypes of a new class of anti-infectious agents.

---

We thank M. Dinauer for *phox91*-deficient breeders; J. MacMicking for preliminary experiments with recombinant *E. coli* in macrophages; A. Nunoshiba, B. Demple, P. O'Gaora, and D.B. Young for bacterial strains; F. Fang for helpful comments; D. Schnappinger for fruitful discussions; Genentech, Inc. (South San Francisco, CA) for recombinant mouse IFN- $\gamma$ ; and Stuart Shuman for access to equipment.

This work was funded by National Institutes of Health (NIH) grant HL-51967 to L.W. Riley and C. Nathan; by a German Cancer Research Center fellowship to S. Ehrt; by a Richard Lounsbery Foundation fellowship to J. Ruan; and by NIH Medical Scientist Training Program grant GM-07739(MS) to M. Shiloh. We also acknowledge financial support from the Auxiliary of the Society of the New York Hospital for computer equipment.

Address correspondence to Dr. Lee W. Riley at his current address, Division of Public Health Biology and Epidemiology, School of Public Health, 140 Warren Hall, UC Berkeley, Berkeley, CA 94720. Phone: 510-642-9200; FAX: 510-643-8236; E-mail: lriley@uclink4.berkeley.edu S. Ehrt's current address is Division of Public Health Biology and Epidemiology, School of Public Health, UC Berkeley, Berkeley, CA 94720.

Received for publication 19 September 1997.

## References

1. Raviglione, M.C., D.E. Snider, Jr., and A. Kochi. 1995. Global epidemiology of tuberculosis. Morbidity and mortality of a worldwide epidemic. *JAMA (J. Am. Med. Assoc.)*. 273:220-226.
2. Chan, J., Y. Tian, K.E. Tanaka, M.S. Tsang, K. Yu, P. Salgame, D. Carroll, Y. Kress, R. Teitelbaum, and B.R. Bloom. 1996. Effects of protein malnutrition on tuberculosis in mice. *Proc. Natl. Acad. Sci. USA*. 93:14857-14861.
3. Daley, C.L., P.M. Small, G.F. Schecter, G.K. Schoolnik, R.A. McAdam, W.R. Jacobs, Jr., and P.C. Hopewell. 1992. An outbreak of tuberculosis with accelerated progression among persons infected with the human immunodeficiency virus. An analysis using restriction-fragment-length polymorphisms. *N. Engl. J. Med.* 326:231-235.
4. Nathan, C.F. 1983. Mechanisms of macrophage antimicrobial activity. *Trans. R. Soc. Trop. Med. Hyg.* 77:620-630.
5. Nathan, C.F., and J.B. Hibbs, Jr. 1991. Role of nitric oxide synthesis in macrophage antimicrobial activity. *Curr. Opin. Immunol.* 3:65-70.
6. Butler, A.R., F.W. Flitney, and D.L. Williams. 1995. NO, nitrosonium ions, nitroxide ions, nitrosothiols and iron-nitrosyls in biology: a chemist's perspective. *Trends Pharmacol. Sci.* 16: 18-22.
7. De Groote, M.A., and F.C. Fang. 1995. NO inhibitions: antimicrobial properties of nitric oxide. *Clin. Infect. Dis.* 21: 162-165.
8. Neill, M.A., and S.J. Klebanoff. 1988. The effect of phenolic glycolipid-1 from *Mycobacterium leprae* on the antimicrobial activity of human macrophages. *J. Exp. Med.* 167:30-42.
9. Sherman, D.R., P.J. Sabo, M.J. Hickey, T.M. Arain, G.G. Mahairas, Y. Yuan, C.E. Barry III, and C.K. Stover. 1995. Disparate responses to oxidative stress in saprophytic and pathogenic mycobacteria. *Proc. Natl. Acad. Sci. USA*. 92: 6625-6629.

10. Sherman, D.R., K. Mdluli, M.J. Hickey, T.M. Arain, S.L. Morris, C.E. Barry III, and C.K. Stover. 1996. Compensatory *ahpC* gene expression in isoniazid-resistant *Mycobacterium tuberculosis*. *Science*. 272:1641–1643.
11. Dumarey, C.H., V. Labrousse, N. Rastogi, B.B. Vargaftig, and M. Bachelet. 1994. Selective *Mycobacterium avium*-induced production of nitric oxide by human monocyte-derived macrophages. *J. Leukocyte Biol.* 56:36–40.
12. Zhang, Y., M.J. Garcia, R. Lathigra, B. Allen, C. Moreno, J.D. van Embden, and D. Young. 1992. Alterations in the superoxide dismutase gene of an isoniazid-resistant strain of *Mycobacterium tuberculosis*. *Infect. Immun.* 60:2160–2165.
13. Chan, J., and S.H.E. Kaufmann. 1994. Immune mechanisms of protection. In *Tuberculosis: Pathogenesis, Protection and Control*. B.R. Bloom, editor. American Society for Microbiology, Washington. 389–415.
14. Schlesinger, L.S., C.G. Bellinger-Kawahara, N.R. Payne, and M.A. Horwitz. 1990. Phagocytosis of *Mycobacterium tuberculosis* is mediated by human monocyte complement receptors and complement component C3. *J. Immunol.* 144:2771–2780.
15. Wright, S.D., and S.C. Silverstein. 1983. Receptors for C3b and C3bi promote phagocytosis but not the release of toxic oxygen from human phagocytes. *J. Exp. Med.* 158:2016–2023.
16. Chan, J., Y. Xing, R.S. Magliozzo, and B.R. Bloom. 1992. Killing of virulent *Mycobacterium tuberculosis* by reactive nitrogen intermediates produced by activated murine macrophages. *J. Exp. Med.* 175:1111–1122.
17. MacMicking, J.D., R.J. North, R. LaCourse, J.S. Mudgett, S.K. Shah, and C.F. Nathan. 1997. Identification of *NOS2* as a protective locus against tuberculosis. *Proc. Natl. Acad. Sci. USA.* 94:5243–5248.
18. Nicholson, S., M. da G. Bonecini-Almeida, J.R. Lapa e Silva, C. Nathan, Q.W. Xie, R. Mumford, J.R. Weidner, J. Calaycay, J. Geng, N. Boechat, et al. 1996. Inducible nitric oxide synthase in pulmonary alveolar macrophages from patients with tuberculosis. *J. Exp. Med.* 183:2293–2302.
19. Nozaki, Y., Y. Hasegawa, S. Ichiyama, I. Nakashima, and K. Shimokata. 1997. Mechanism of nitric oxide-dependent killing of *Mycobacterium bovis* BCG in human alveolar macrophages. *Infect. Immun.* 65:3644–3647.
20. Doi, T., M. Ando, T. Akaike, M. Suga, K. Sato, and H. Maeda. 1993. Resistance to nitric oxide in *Mycobacterium avium* complex and its implication in pathogenesis. *Infect. Immun.* 61:1980–1989.
21. O'Brien, L., J. Carmichael, D.B. Lowrie, and P.W. Andrew. 1994. Strains of *Mycobacterium tuberculosis* differ in susceptibility to reactive nitrogen intermediates in vitro. *Infect. Immun.* 62:5187–5190.
22. Friedman, C.R., M.Y. Stoeckle, B.N. Kreiswirth, W.D. Johnson, Jr., S.M. Manoach, J. Berger, K. Sathianathan, A. Hafner, and L.W. Riley. 1995. Transmission of multidrug-resistant tuberculosis in a large urban setting. *Am. J. Respir. Crit. Care Med.* 152:355–359.
23. Friedman, C., G.C. Quinn, B.N. Kreiswirth, D.C. Perlman, N. Salomon, N. Schluger, M. Lutfey, J. Berger, N. Poltoratskaia, and L.W. Riley. 1997. Widespread dissemination of a single drug-susceptible strain of *Mycobacterium tuberculosis*. *J. Infect. Dis.* 176:478–484.
24. Snapper, S.B., R.E. Melton, S. Mustafa, T. Kieser, and W.R. Jacobs, Jr. 1990. Isolation and characterization of efficient plasmid transformation mutants of *Mycobacterium smegmatis*. *Mol. Microbiol.* 4:1911–1919.
25. Greenberg, J.T., P. Monach, J.H. Chou, P.D. Josephy, and B. Demple. 1990. Positive control of a global antioxidant defense regulon activated by superoxide-generating agents in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA.* 87:6181–6185.
26. Greenberg, J.T., and B. Demple. 1988. Overproduction of peroxide-scavenging enzymes in *Escherichia coli* suppresses spontaneous mutagenesis and sensitivity to redox-cycling agents in oxyR-mutants. *EMBO (Eur. Mol. Biol. Organ.) J.* 7: 2611–2617.
27. Garbe, T.R., J. Barathi, S. Barnini, Y. Zhang, C. Abou-Zeid, D. Tang, R. Mukherjee, and D.B. Young. 1994. Transformation of mycobacterial species using hygromycin resistance as selectable marker. *Microbiology (Reading)*. 140:133–138.
28. van Soolingen, D., P.W. Hermans, P.E. de Haas, D.R. Soll, and J.D. van Embden. 1991. Occurrence and stability of insertion sequences in *Mycobacterium tuberculosis* complex strains: evaluation of an insertion sequence-dependent DNA polymorphism as a tool in the epidemiology of tuberculosis. *J. Clin. Microbiol.* 29:2578–2586.
29. Shiloh, M.U., J. Ruan, and C. Nathan. 1997. Evaluation of bacterial numbers and phagocyte function with a fluorescence-based microplate assay. *Infect. Immun.* 65:3193–3197.
30. Buchmeier, N., and F. Heffron. 1989. Intracellular survival of wild-type *Salmonella typhimurium* and macrophage-sensitive mutants in diverse populations of macrophages. *Infect. Immun.* 57:1–7.
31. MacMicking, J.D., C. Nathan, G. Hom, N. Chartrain, D.S. Fletcher, M. Trumbauer, K. Stevens, Q.W. Xie, K. Sokol, N. Hutchinson, et al. 1995. Altered responses to bacterial infection and endotoxic shock in mice lacking inducible nitric oxide synthase. *Cell.* 81:641–650 (see erratum *Cell.* 1995. 81: 1171).
32. Pollock, J.D., D.A. Williams, M.A. Gifford, L.L. Li, X. Du, J. Fisherman, S.H. Orkin, C.M. Doerschuk, and M.C. Dinauer. 1995. Mouse model of X-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide production. *Nat. Genet.* 9:202–209.
33. Ding, A.H., C.F. Nathan, and D.J. Stuehr. 1988. Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages: comparison of activating cytokines and evidence for independent production. *J. Immunol.* 141:2407–2412.
34. De la Harpe, J., and C.F. Nathan. 1985. A semi-automated micro-assay for H<sub>2</sub>O<sub>2</sub> release by human blood monocytes and mouse peritoneal macrophages. *J. Immunol. Methods.* 78:323–336.
35. Stuehr, D.J., and C.F. Nathan. 1989. Nitric oxide. A macrophage product responsible for cytostasis and respiratory inhibition in tumor target cells. *J. Exp. Med.* 169:1543–1555.
36. Taylor, T.W.J., E.W. Wignall, and J.F. Cowley. 1927. The decomposition of nitrous acid in aqueous solutions. *J. Chem. Soc. (Lond.)*. 11:1923.
37. Morris, S.L., and J.N. Hansen. 1981. Inhibition of *Bacillus cereus* spore outgrowth by covalent modification of a sulfhydryl group by nitrosothiol and iodoacetate. *J. Bacteriol.* 148: 465–471.
38. Stamler, J.S. 1995. S-nitrosothiols and bioregulatory actions of nitrogen oxides through reactions with thiol groups. *Curr. Topics Microbiol. Immunol.* 196:19–36.
39. Stamler, J.S., D.J. Singel, and J. Loscalzo. 1992. Biochemistry of nitric oxide and its redox-activated forms. *Science.* 258: 1898–1902.
40. Dykhuizen, R.S., R. Frazer, C. Duncan, C.C. Smith, M.

- Golden, N. Benjamin, and C. Leifert. 1996. Antimicrobial effect of acidified nitrite on gut pathogens: importance of dietary nitrate in host defense. *Antimicrob. Agents Chemother.* 40: 1422–1425.
41. Sturgill-Koszycki, S., P.H. Schlesinger, P. Chakraborty, P.L. Haddix, H.L. Collins, A.K. Fok, R.D. Allen, S.L. Gluck, J. Heuser, and D.G. Russell. 1994. Lack of acidification in *Mycobacterium* phagosomes produced by exclusion of the vesicular proton-ATPase. *Science.* 263:678–681.
  42. Burnett, W.V., J. Henner, and T. Eckhardt. 1987. The nucleotide sequence of the gene coding for XP55, a major secreted protein from *Streptomyces lividans*. *Nucleic Acids Res.* 15: 3926.
  43. De Groote, M.A., D. Granger, Y. Xu, G. Campbell, R. Prince, and F.C. Fang. 1995. Genetic and redox determinants of nitric oxide cytotoxicity in a *Salmonella typhimurium* model. *Proc. Natl. Acad. Sci. USA.* 92:6399–6403.
  44. De Groote, M.A., T. Testerman, Y. Xu, G. Stauffer, and F.C. Fang. 1996. Homocysteine antagonism of nitric oxide-related cytotoxicity in *Salmonella typhimurium*. *Science.* 272:414–417.
  45. Lewis, R.S., S. Tamir, S.R. Tannenbaum, and W.M. Deen. 1995. Kinetic analysis of the fate of nitric oxide synthesized by macrophages in vitro. *J. Biol. Chem.* 270:29350–29355.
  46. Ohkuma, S., and B. Poole. 1978. Fluorescence probe measurement of the intralysosomal pH in living cells and the perturbation of pH by various agents. *Proc. Natl. Acad. Sci. USA.* 75:3327–3331.
  47. Hausladen, A., C.T. Privalle, T. Keng, J. Deangelo, and J.S. Stamler. 1996. Nitrosative stress: activation of the transcription factor OxyR. *Cell.* 86:719–729.
  48. Nunoshiba, T., T. DeRojas-Walker, S.R. Tannenbaum, and B. Demple. 1995. Roles of nitric oxide in inducible resistance of *Escherichia coli* to activated macrophages. *Infect. Immun.* 63:794–798.
  49. Nunoshiba, T., T. DeRojas-Walker, J.S. Wishnok, S.R. Tannenbaum, and B. Demple. 1993. Activation by nitric oxide of an oxidative-stress response that defends *Escherichia coli* against activated macrophages. *Proc. Natl. Acad. Sci. USA.* 90: 9993–9997.
  50. Dhandayuthapani, S., Y. Zhang, M.H. Mudd, and V. Deretic. 1996. Oxidative stress response and its role in sensitivity to isoniazid in mycobacteria: characterization and inducibility of *ahpC* by peroxides in *Mycobacterium smegmatis* and lack of expression in *M. aurum* and *M. tuberculosis*. *J. Bacteriol.* 178: 3641–3649.
  51. Zhang, Y., S. Dhandayuthapani, and V. Deretic. 1996. Molecular basis for the exquisite sensitivity of *Mycobacterium tuberculosis* to isoniazid. *Proc. Natl. Acad. Sci. USA.* 93:13212–13216.
  52. Weinberg, J.B., and J.B. Hibbs, Jr. 1979. In vitro modulation of macrophage tumoricidal activity: partial characterization of a macrophage-activating factor(s) in supernatants of NaIO<sub>4</sub>-treated cells. *J. Reticuloendothel. Soc.* 26:283–293.
  53. Novogrodsky, A. 1974. Selective activation of mouse T and B lymphocytes by periodate, galactose oxidase and soybean agglutinin. *Eur. J. Immunol.* 4:646–648.
  54. Nathan, C.F., and R.K. Root. 1977. Hydrogen peroxide release from mouse peritoneal macrophages: dependence on sequential activation and triggering. *J. Exp. Med.* 146:1648–1662.
  55. Nathan, C.F., H.W. Murray, M.E. Wiebe, and B.Y. Rubin. 1983. Identification of interferon- $\gamma$  as the lymphokine that activates human macrophage oxidative metabolism and antimicrobial activity. *J. Exp. Med.* 158:670–689.
  56. Adams, L.B., S.G. Franzblau, Z. Vavrin, J.B.J. Hibbs, and J.L. Krahenbuhl. 1991. L-Arginine-dependent macrophage effector functions inhibit metabolic activity of *Mycobacterium leprae*. *J. Immunol.* 147:1642–1646.
  57. Schwarz, M.A., J.S. Lazo, J.C. Yalowich, W.P. Allen, M. Whitmore, H.A. Bergonia, E. Tzeng, T.R. Billiar, P.D. Robbins, J.R.J. Lancaster, and B.R. Pitt. 1995. Metallothionein protects against the cytotoxic and DNA-damaging effects of nitric oxide. *Proc. Natl. Acad. Sci. USA.* 92:4452–4456.
  58. Almiron, M., A.J. Link, D. Furlong, and R. Kolter. 1992. A novel DNA-binding protein with regulatory and protective roles in starved *Escherichia coli*. *Genes Dev.* 6:2646–2654.
  59. Altuvia, S., M. Almiron, G. Huisman, R. Kolter, and G. Storz. 1994. The *dps* promoter is activated by OxyR during growth and by IHF and sigma S in stationary phase. *Mol. Microbiol.* 13:265–272.
  60. Deretic, V., W. Philipp, S. Dhandayuthapani, M.H. Mudd, R. Curcic, T. Garbe, B. Heym, L.E. Via, and S.T. Cole. 1995. *Mycobacterium tuberculosis* is a natural mutant with an inactivated oxidative-stress regulatory gene: implications for sensitivity to isoniazid. *Mol. Microbiol.* 17:889–900.
  61. Nikitovic, D., and A. Holmgren. 1996. S-nitrosoglutathione is cleaved by thioredoxin system with liberation of glutathione and redox regulating nitric oxide. *J. Biol. Chem.* 271:19180–19185.