

The IKK β Subunit of I κ B Kinase (IKK) is Essential for Nuclear Factor κ B Activation and Prevention of Apoptosis

By Zhi-Wei Li,* Wenming Chu,* Yinling Hu,* Mireille Delhase,*
Tom Deerinck,[§] Mark Ellisman,[§] Randall Johnson,[‡] and Michael Karin*

From the *Department of Pharmacology, Laboratory of Gene Regulation and Signal Transduction,
[‡]Department of Biology, and [§]Department of Neuroscience, University of California, San Diego,
La Jolla, California 92093-0636

Summary

The I κ B kinase (IKK) complex is composed of three subunits, IKK α , IKK β , and IKK γ (NEMO). While IKK α and IKK β are highly similar catalytic subunits, both capable of I κ B phosphorylation in vitro, IKK γ is a regulatory subunit. Previous biochemical and genetic analyses have indicated that despite their similar structures and in vitro kinase activities, IKK α and IKK β have distinct functions. Surprisingly, disruption of the *Ikk α* locus did not abolish activation of IKK by proinflammatory stimuli and resulted in only a small decrease in nuclear factor (NF)- κ B activation. Now we describe the pathophysiological consequence of disruption of the *Ikk β* locus. IKK β -deficient mice die at mid-gestation from uncontrolled liver apoptosis, a phenotype that is remarkably similar to that of mice deficient in both the RelA (p65) and NF- κ B1 (p50/p105) subunits of NF- κ B. Accordingly, IKK β -deficient cells are defective in activation of IKK and NF- κ B in response to either tumor necrosis factor α or interleukin 1. Thus IKK β , but not IKK α , plays the major role in IKK activation and induction of NF- κ B activity. In the absence of IKK β , IKK α is unresponsive to IKK activators.

Key words: inflammation • tumor necrosis factor α • interleukin 1 • knockout mice • signal transduction

The nuclear factor (NF)- κ B¹ transcription factor plays a key role in activation of inflammatory and innate immune responses (1, 2). In nonstimulated cells, NF- κ B dimers are kept as cytoplasmic latent complexes through binding of specific inhibitors, the I κ Bs, which mask their nuclear localization signal (NLS). Upon exposure to proinflammatory stimuli, such as bacterial LPS, TNF- α , or IL-1, the I κ Bs are rapidly phosphorylated at two conserved NH₂-terminal serines, a posttranslational modification that is rapidly followed by their polyubiquitination and proteasomal degradation (3–6). This results in unmasking of the NLS of NF- κ B dimers followed by their translocation to the nucleus, binding to specific DNA sites (κ B sites), and target gene activation. NF- κ B target genes include many of the cytokine and chemokine genes, as well as genes coding for adhesion molecules, cell surface receptors, and enzymes that produce secondary inflammatory mediators (7, 8).

The protein kinase that phosphorylates I κ Bs in response to proinflammatory stimuli has been identified biochemically and molecularly (9–11). Named IKK, this protein kinase is a complex composed of at least three subunits: IKK α , IKK β and IKK γ (for a review, see reference 12). IKK α and IKK β are highly similar protein kinases that act as the catalytic subunits of the complex (9, 11, 13, 14). In vitro, both IKK α and IKK β form homo- and heterodimers that can phosphorylate I κ B proteins at their NH₂-terminal regulatory serines (15). In mammalian cells, IKK α and IKK β form a stable heterodimer that is tightly associated with the IKK γ (NEMO) subunit (16, 17). As cell lines that fail to express IKK γ (NEMO) exhibit a major defect in I κ B degradation and NF- κ B activation in response to proinflammatory stimuli and double-stranded RNA, this regulatory subunit plays an essential function (at least in the examined cell lines) in IKK and NF- κ B activation (17). The physiological function of the two catalytic subunits has been less clear. Initially, overexpression of catalytically inactive forms of IKK α and IKK β that blocked IKK and NF- κ B activation suggested that both subunits play similar and possibly redundant roles in I κ B phosphorylation and NF- κ B activation (13, 14). This hypothesis was fostered by finding that in vitro IKK α and

¹Abbreviations used in this paper: EF, embryonic fibroblast; EMSA, electrophoretic mobility shift assay; ES, embryonic stem; H&E, hematoxylin and eosin; IKK, I κ B kinase; NF, nuclear factor; NIK, NF- κ B inducing kinase; TUNEL, terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling.

IKK β can directly phosphorylate I κ B α and I κ B β at the serines that trigger their degradation *in vivo* (15). However, it was also suggested that IKK α rather than IKK β is responsible for activation of the entire complex in response to certain stimuli, such as the NF- κ B inducing kinase, NIK (18). Recently, we found that in addition to an IKK γ subunit with an intact COOH terminus (16), IKK activation requires the phosphorylation of IKK β at two serines within its activation loop (19). Replacement of these serines, whose phosphorylation is stimulated by proinflammatory stimuli or NIK, with alanines abolishes IKK activation. Interestingly, although the entire activation loop is identical in sequence between IKK α and IKK β , replacement of the same two serines in IKK α with alanines has no effect on IKK activation (19). These results were further substantiated by gene targeting (knockout) experiments. Cells and tissues from mice that no longer express IKK α (*Ikk α ^{-/-}* mice) exhibit normal IKK activation in response to TNF, IL-1, or LPS (20). Although NF- κ B is fully inducible, for an unknown reason, IKK α -deficient fibroblasts exhibit approximately twofold reduction in both basal and induced NF- κ B binding activity (20). Thus, IKK α may somehow stimulate NF- κ B DNA binding despite not being required for I κ B phosphorylation and degradation in most cell types. The gene targeting experiments reveal that, although not involved in activation of IKK by proinflammatory stimuli, IKK α plays an instrumental role in morphogenesis (20). The most important function of IKK α appears to be in the control of keratinocyte differentiation and formation of the epidermis (20). It is not yet clear whether these morphogenetic functions of IKK α are exerted through localized NF- κ B activation in response to developmental cues.

To determine the physiological function(s) of IKK β , we have used gene targeting to create *Ikk β* knockout mice. We now show that the loss of IKK β results in embryonic lethality at mid-gestation due to extensive apoptosis of the developing liver. This phenotype is similar to that of mice

deficient in the RelA (p65) subunit of NF- κ B (21). It was recently shown that the lethality of *RelA^{-/-}* mice is completely suppressed by the loss of TNF- α (22). As NF- κ B is required for protection of cells from TNF- α -induced apoptosis (23–25), the apoptotic phenotype of *Ikk β ^{-/-}* mice strongly suggests that the absence of IKK β results in a severe defect in NF- κ B activation. Indeed, neither IKK nor NF- κ B can be activated by TNF- α or IL-1 in IKK β -deficient cells. Furthermore, we show that in the absence of IKK β , the IKK α subunit is not responsive to NIK even though it can still associate with the IKK γ subunit.

Materials and Methods

Generation of IKK β -deficient Mice. Using a 0.2-kb BstEII-Bsu36I restriction fragment from the 5' end of human IKK β cDNA as a probe, three murine IKK β genomic fragments were isolated from a 129/SvJ mouse genomic library (Stratagene, Inc.). One of the clones contained at least the first three coding exons and was used to construct the targeting vector IKK β KO. A 1.4-kb SacI restriction fragment harboring part of the second exon was used as the short homology arm, and the long arm was a 5.5-kb EcoRV-XhoI restriction fragment containing part of the third intron. The two arms were inserted into the XmnI and SmaI sites, respectively, of pGNA, which contains the G418 resistance gene (*Neo^r*) and *LacZ* (26). As a negative selection marker, a diphtheria toxin gene cassette (*DT*) was inserted into the KpnI site of pGNA. After cutting with PmeI, 20 μ g of the linearized targeting vector was electroporated into 10⁷ mouse embryonic stem (ES) cells (line GS from Genome Systems). After selection with G418 at 0.4 mg/ml, G418-resistant colonies were picked and screened by PCR. The genotype of the PCR-positive clones was confirmed by Southern blotting analysis. Homologous recombinants were karyotyped and analyzed for mycoplasma. Two homologous recombinant ES clones were injected into C57BL/6 blastocysts. Resulting male chimeras were crossed with C57BL/6 females, and germline transmission was scored by coat color. Heterozygous mice were identified by PCR and Southern analysis of




Figure 1. Generation of IKK β -deficient mice. (A) The mouse *Ikk β* locus and the targeting vector. Map of the *Ikk β* genomic fragment used for gene targeting is shown. The exons are indicated by solid black boxes, the introns are indicated by bold lines, and the selection markers, lengths of restriction fragments, restriction enzyme sites, the probes used for Southern analysis, and the location of primers used in PCR screening are also shown. RI, EcoRI; P, PstI; S, SacI; RV, EcoRV; Xh, XhoI. (B) Southern blot analysis of mouse genomic DNA. Mouse genomic DNA was digested with EcoRI and probed with probe A (1.2-kb HindIII-PstI fragment of *Ikk β*). After homologous recombination, the 9.7-kb EcoRI fragment of wild-type *Ikk β* is replaced by a 7.2-kb EcoRI fragment, as indicated in panel A. (C) Western blot analysis of mouse proteins using antibody H470 specific for IKK β . Location of the IKK β band is indicated. The lower band is nonspecific (ns). The same blot was also probed with antibodies to IKK α , IKK γ , p65(RelA), and p50(NF- κ B1). The genotypes are as indicated.




Figure 2. Appearance of an *Ikkβ*^{-/-} E13.5 embryo and a normal littermate. Wild-type (*Ikkβ*^{+/+}, WT) and mutant (*Ikkβ*^{-/-}, M) embryos were isolated at E13.5 and photographed. The genotypes of the embryos were later determined by PCR and Southern blot analysis.

mouse tail DNA. Embryos from intercrosses of heterozygous (*Ikkβ*^{+/-}) mice, as well as mouse embryonic fibroblasts (EFs), were genotyped by PCR and Southern analysis using DNA isolated from a piece of each embryo or a cell pellet, respectively.

PCR and Southern Blotting Analysis. PCR was performed in the presence of 10% DMSO with Taq DNA polymerase using a Perkin-Elmer 9600 thermocycler programmed for denaturation at 95°C for 5 min, amplification for 35 cycles (94°C for 30 s, 55°C for 30 s, 65°C for 2 min), and elongation at 72°C for 10 min. Primers used were: P1 (5'-AGTCCAACCTGGCAGCGA-ATA-3') located outside of the homology arm and P2 (5'-CAACATTAAATGTGAGCGAG-3') located within the *LacZ* gene. Southern blotting analysis was performed according to a standard

protocol (27) except that hybridization was performed in phosphate-SDS buffer (28).

Kinase Assay, Immunoprecipitation, Immunoblotting, and Electrophoretic Mobility Shift Assays. *Ikkβ*^{-/-}, *Ikkβ*^{+/-}, and *Ikkβ*^{+/+} ES and EF cells were treated with TNF-α or IL-1 at 20 ng/ml. Kinase assays and immunoprecipitations were performed as described (9). Immunoblotting was performed as described (14, 16). Electrophoretic mobility shift assays (EMSA) using the consensus κB and NF-1 sequences were performed as described (16, 29).

Histology, In Situ TUNEL Assay, and Transmission Electron Microscopy. Mouse embryos or embryo livers were fixed in 10% buffered formalin and embedded in paraffin. After routine processing, the sections (5-μm thick) were stained with hematoxylin and eosin (H&E) for histological analysis. In situ TUNEL (terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling) assay was done using the in situ cell death detection kit according to the manufacturer's instructions (Boehringer Mannheim). For electron microscopy, embryonic day 13 (E13) embryos were removed and the livers were dissected out and fixed for 1 h in 2% formaldehyde and 2% glutaraldehyde in 0.15 M sodium cacodylate buffer (pH 7.4) at 4°C. The remainder of the embryos were placed in PBS for subsequent PCR and Southern analysis. After washing in cacodylate buffer, the livers were postfixed in 1% osmium tetroxide in cacodylate buffer for an additional 1 h. After postfixation, the samples were rinsed in double distilled water, dehydrated in a graded ethanol series, and infiltrated and polymerized in Durcupan ACM resin (Electron Microscopy Sciences). Sections 80-nm thick were stained with Sato lead and examined at 80 keV with either a JEOL 100CX or 2000EX transmission electron microscope.

Results

Generation of *Ikkβ* Knockout Mice. To create a strain of IKKβ-deficient mice, we used gene targeting technology (30). Mouse genomic *Ikkβ* DNA was cloned from a 129 strain library and, after mapping and sequencing, was used




Figure 3. Analysis of wild-type (WT) and mutant (M) livers. E13.5 embryos were fixed and sectioned. Paraffin-embedded transverse sections at the area of the liver were subjected to H&E (top; original magnification: 400×) or TUNEL (bottom; original magnification: 600×) staining. The stained sections were photographed.

to construct the targeting vector (Fig. 1 A). To eliminate IKK β kinase activity, part of the second and the entire third coding exon that specifies an essential part of the kinase domain were replaced with a DNA fragment encoding β -galactosidase (*LacZ*) and neomycin resistance (*Neo^r*). Because the *Neo^r* gene contains transcription termination and polyadenylation signals, the COOH-terminal three quarters of IKK β including its protein interaction motifs are unlikely to be expressed from the targeted allele.

After selection and screening by Southern blotting, six ES cell clones with homologous integration of the targeting vector into the *Ikk β* locus were isolated, and two of them were used to generate chimeric mice. Chimeric mice derived from these clones transmitted the targeted *Ikk β* allele to their progeny (Fig. 1 B). Although *Ikk β ^{+/-}* male and female mice appeared normal and were fertile, upon intercrossing they did not give rise to live *Ikk β ^{-/-}* progeny.

Analysis of protein extracts of *Ikk β ^{+/+}*, *Ikk β ^{+/-}*, and *Ikk β ^{-/-}* cells revealed that, as expected, no IKK β protein was expressed from the targeted allele (Fig. 1 C). In addition, *Ikk β ^{+/-}* cells expressed approximately half the dose of IKK β present in wild-type cells. No compensatory increases in IKK α , IKK γ , p65(RelA), or p50(NF- κ B1) expression were observed.

Phenotype of *Ikk β ^{-/-}* Mice. Given the expected importance of IKK β for NF- κ B activation and the embryonic lethality of *RelA^{-/-}* mice (21), we suspected that the loss of IKK β would result in a similar phenotype. Therefore, we analyzed embryos from timed pregnancies of *Ikk β ^{+/-}* intercrosses. Although *Ikk β ^{-/-}* embryos isolated at E11.5 were alive and had perfectly normal appearance (data not shown), *Ikk β ^{-/-}* embryos isolated at E13.5 were no longer alive and were rather anemic in appearance (Fig. 2). Even external examination suggested that the liver of E13.5 *Ikk β ^{-/-}* embryos had degenerated. Notably, however, the limbs and head of *Ikk β ^{-/-}* embryos were normally developed, unlike those of *Ikk α ^{-/-}* E13.5 embryos (20). Histochemical examination of transverse sections of normal and mutant E13.5 mouse embryos stained with H&E revealed massive cell death in livers of *Ikk β ^{-/-}* embryos (Fig. 3 A). Essentially, no viable hepatocytes could be detected, and the numbers of dead cells with highly condensed and fragmented nuclei were markedly increased. However, hematopoietic precursors retained their normal appearance in *Ikk β ^{-/-}* livers. TUNEL staining revealed that the observed cell death is most likely due to apoptosis, whose rate was increased manifold (Fig. 3 B). Examination of E13 *Ikk β ^{-/-}* embryos revealed close to normal external appearance (data not shown), but electronmicroscopic examination of ultra-thin sections from their livers revealed massive numbers of dead hepatocytes with highly condensed nuclei characteristic of apoptotic cell death (Fig. 4). The livers of *Ikk β ^{+/+}* or *Ikk β ^{+/-}* littermates had perfectly normal appearance.

Defective NF- κ B Activation in *Ikk β ^{-/-}* Cells. We used two different approaches to determine the consequences of the loss of IKK β expression on IKK and NF- κ B activation. First, we prepared *Ikk β ^{-/-}* ES cell lines by subjecting *Ikk β ^{+/-}* ES cells to selection at higher G418 concentration.

One *Ikk β ^{-/-}* cell line was identified. As shown in Fig. 5 A, stimulation of these cells with either TNF- α or IL-1 did not result in IKK activation, whereas a normal activation response was observed in *Ikk β ^{+/-}* cells. Note, however, that *Ikk β ^{+/-}* cells had ~50% of the IKK activity of wild-type (*Ikk β ^{+/+}*) ES cells, consistent with the reduced amount of IKK β protein (data not shown). In addition to the defect in IKK activation, hardly any induction of NF- κ B DNA binding activity was observed in *Ikk β ^{-/-}* cells after stimulation with either IL-1 or TNF- α (Fig. 5 B). Even the basal level of NF- κ B DNA binding activity was considerably reduced in *Ikk β ^{-/-}* cells, despite no detectable changes in p65(RelA) or p50(NF- κ B1) abundance (data not shown). The second approach to evaluate the function of IKK β was to prepare cultures of EFs from E11.5 mouse embryos of all three genotypes. As shown in Fig. 6, essentially no induc-




Figure 4. Electron microscopic analysis of livers from E13 *Ikk β ^{+/+}*, *Ikk β ^{+/-}*, and *Ikk β ^{-/-}* embryos. Both E13 *Ikk β ^{+/+}* (A) and *Ikk β ^{+/-}* (B) livers exhibited normal morphology. The *Ikk β ^{-/-}* liver (C) exhibited varying degrees of apoptosis characterized by collapsed and condensed nuclei and general cellular degeneration. Bars = 5 μ m.




Figure 5. Defective IKK and NF- κ B activation in IKK β -deficient ES cells. (A) IKK activity. Lysates of TNF- α - or IL-1-treated *Ikk β ^{+/-}* and *Ikk β ^{-/-}* cells were prepared at the indicated time points (in min) after stimulation and immunoprecipitated with antibody M280 to IKK α . IKK activity (KA) was measured by an immunocomplex kinase assay using GST-I κ B α (1-54) as a substrate. The kinase assay products were separated by SDS-PAGE, transferred to nitrocellulose membrane, and autoradiographed. The membrane was reprobbed with antibody M280 (IB: IKK α) for loading control. (B) NF- κ B binding activity. Nuclear extracts of *Ikk β ^{+/-}* and *Ikk β ^{-/-}* cells stimulated with IL-1 or TNF- α for the indicated times (in min) were incubated with ³²P-labeled κ B oligonucleotide probe and subjected to EMSA. Binding to an NF-1 probe was used to control the quality and amount of nuclear protein extracts.

tion of IKK or NF- κ B activity could be detected in *Ikk β ^{-/-}* EF cells treated with either IL-1 or TNF- α . Interestingly, *Ikk β ^{+/-}* EF cells exhibited an ~50% reduction in IKK activity (consistent with the reduction in IKK β expression) but a much larger decrease in NF- κ B DNA binding activity.

IKK α Cannot Be Activated by NIK in the Absence of IKK β . The results described above indicate that IKK α , which is expressed in normal levels in *Ikk β ^{-/-}* cells, cannot be activated by either TNF- α or IL-1. To further examine this point, we cotransfected an HA epitope-tagged IKK α expression vector into *Ikk β ^{-/-}* ES cells in the absence or

presence of an NIK expression vector. NIK is the most potent IKK activator identified to date (31) and was suggested to be a direct IKK α kinase (18). Recently, however, we obtained results that suggested that NIK-induced IKK α phosphorylation is not direct and is likely to be dependent on IKK β (19). Consistent with this hypothesis, we found no increase in IKK activity towards I κ B α (1-54) substrate upon coexpression of HA-IKK α with NIK in *Ikk β ^{-/-}* cells (Fig. 7 A). Yet, when an IKK β expression vector was included in these transfections, NIK elicited a clear increase in IKK activity. As shown previously, NIK coexpression efficiently stimulates IKK α -associated IKK activity in IKK β -expressing cells (19).

One reason for the inability of IKK α to respond to proinflammatory stimuli or NIK in the absence of IKK β could be its inability to directly associate with IKK γ , the regulatory subunit of the IKK complex. Previous experiments indicate that IKK γ is essential for recruitment of upstream activators to IKK (16). In addition, using recombinant proteins, it was found that IKK β directly interacts with IKK γ much more efficiently than does IKK α (16, 17). Having available IKK β -deficient cells, we reexamined the ability of IKK α to interact with IKK γ . In contrast to the results obtained with recombinant proteins, very efficient coprecipitation of IKK α by anti-IKK γ antibodies was observed using lysates of *Ikk β ^{-/-}* cells as a starting material (Fig. 7 B). Therefore, the refractoriness of IKK α to IKK activators in IKK β -deficient cells is not due to its inability to associate with IKK γ .




Figure 6. Defective IKK and NF- κ B activation in IKK β -deficient EF cells. Second passage EFs from E11.5 *Ikk β ^{+/+}*, *Ikk β ^{+/-}*, and *Ikk β ^{-/-}* embryos were stimulated with TNF- α or IL-1. At the indicated times, whole cell extracts were prepared and used to measure (A) IKK activity (KA), and (B) NF- κ B DNA binding activity. IB, immunoblotting.

Discussion

The enzymatic activity of the IKK complex, composed of two catalytic subunits, IKK α and IKK β , and one regulatory subunit, IKK γ , is rapidly stimulated by proinflamma-




Figure 7. IKK α is refractory to activation in *Ikk β ^{-/-}* cells despite its association with IKK γ . (A) *Ikk β ^{-/-}* ES cells were transiently transfected by electroporation with an HA-IKK α expression vector alone or together with XpressNIK or HA-IKK β and XpressNIK expression vectors. 24 h after transfection, HA-IKK proteins were immunoprecipitated (IP) with anti-HA antibody and their associated IKK activity (KA) was

determined using GST-I κ B α (1-54) as a substrate. Protein expression levels were determined by immunoblotting (IB) with anti-HA. (B) Lysates of *Ikk β ^{+/+}*, *Ikk β ^{+/-}*, and *Ikk β ^{-/-}* cells were immunoprecipitated (IP) with either anti-IKK α or anti-IKK γ antibodies as indicated. The immunocomplexes were dissolved in SDS loading buffer and separated by SDS-PAGE. After transfer to an Immobilon membrane, the proteins were analyzed by immunoblotting (IB) with anti-IKK α antibody. A lysate of 3T3 cells was used as a control (Cont).

tory cytokines and LPS (for a review, see reference 12). Activated IKK phosphorylates the different I κ Bs at the two NH $_2$ -terminal serines that trigger their polyubiquitination and proteasome-mediated degradation. Once the I κ Bs are degraded, the freed NF- κ B dimers migrate to the nucleus and activate target gene transcription. Based on their similar primary structures (11, 13, 14) and substrate specificities (15), IKK α and IKK β were expected to play redundant and interchangeable roles in proinflammatory signaling to NF- κ B. Therefore, it was rather surprising that only IKK β was found to be involved in IKK activation. Alanine substitutions of the two serines in the activation loop of IKK β , whose phosphorylation is stimulated by either TNF- α treatment or NIK overexpression, prevented IKK activation. Yet, the same mutations introduced into the activation loop of the IKK α subunit had no effect on the response of IKK to TNF- α or NIK (19). These results were confirmed by the analysis of IKK α -deficient cells and tissues which revealed no defect in IKK activation and I κ B α degradation in response to TNF- α , IL-1, or LPS (20). However, it remained possible that the function of IKK α in I κ B phosphorylation in response to proinflammatory stimuli can be fully replaced by IKK β . The results described here indicate that IKK β and IKK α have different physiological functions and that IKK α cannot substitute for IKK β .

To determine the physiological function of IKK β , we generated *Ikk β ^{-/-}* knockout mice and cell lines. The loss of IKK β results in embryonic death at mid-gestation due to massive hepatocyte apoptosis. This phenotype is remarkably similar to that of *RelA* knockout mice (21), with one exception: while *Ikk β ^{-/-}* embryos die around E13, *RelA*^{-/-} embryos die around E15. The earlier death of *Ikk β ^{-/-}* embryos is likely to be due to a more extensive reduction in NF- κ B activity, as embryos that are deficient in both the p65 (RelA) and the p50 (NF- κ B1) subunits of NF- κ B die at E12.5, the same time as IKK β -deficient embryos, from massive hepatocyte apoptosis (32). Thus, IKK β and RelA are genetically proven to be components of the same pathway. Accordingly, cells that lack IKK β are completely defective in IKK and NF- κ B activation in response to either TNF- α or IL-1. Therefore, the IKK β subunit is absolutely

essential for mounting a response to proinflammatory stimuli. This function is not replaced by IKK α , whose expression is not diminished in the absence of IKK β . In addition, as indicated by the normal morphology of the head and limbs of E13.5 *Ikk β ^{-/-}* embryos, IKK α can carry out its developmental function (20) in the complete absence of IKK β . Interestingly, a 50% reduction in IKK β expression, as in *Ikk β ^{+/-}* cells, results in a similar decrease in IKK activity but a much more severe defect in NF- κ B activation. These results underscore the importance of the IKK β subunit and indicate that the NF- κ B activation response does not follow a simple linear relationship to the magnitude of IKK activation. It also appears from these results that a low level of NF- κ B activity may be sufficient for protecting the liver from TNF- α -induced apoptosis.

One possible cause for the inability of IKK α to substitute for IKK β was its relatively lower affinity to IKK γ , the regulatory subunit that is absolutely required for IKK activation (17). Using recombinant proteins, it was observed that IKK α does not form a stable complex with IKK γ in vitro, whereas IKK β readily associates with IKK γ (16, 17). However, immunoprecipitation experiments indicate that a similar amount of IKK α is precipitated by IKK γ antibodies from *Ikk β ^{-/-}* cells as from *Ikk β ^{+/+}* cells. Despite its ability to associate with IKK γ in the absence of IKK β , IKK α is refractory to upstream activators involved in proinflammatory signaling, including the most potent IKK activator identified so far, NIK, in IKK β -deficient cells. These results underscore the differences in regulation of IKK α and IKK β activities.

In summary, together with the previous analysis of IKK α -deficient mice, the analysis of IKK β -deficient mice, described here, indicates that the two catalytic subunits of the IKK complex, although similar in structure, have very different functions. Although IKK β is responsible both for activation of the entire complex in response to proinflammatory stimuli, through phosphorylation at its activation loop, and for activation of NF- κ B, through I κ B phosphorylation, IKK α is assigned the control of epidermal and skeletal morphogenesis. Although the stimuli that activate IKK β and the substrates that mediate its biological activity are known, the stimuli and the relevant substrates for IKK α remain to be identified.

We thank G. Hageman for the gift of mouse TNF- α , D.M. Rothwarf for advice and technical assistance, and B. Thompson for assistance with manuscript preparation.

Y. Hu was supported by a postdoctoral fellowship from the Arthritis Foundation. This work was supported by grants from the National Institutes of Health (AI43477, ES04151, AG05131, and RR04050) and the Department of Energy (DE-FG03-86ER60429). M. Karin is the Frank and Else Schilling-American Cancer Society Research Professor.

Address correspondence to Michael Karin, Laboratory of Gene Regulation and Signal Transduction, Department of Pharmacology, University of California, San Diego, 9500 Gilman Dr., La Jolla, CA 92093-0636. Phone: 619-534-1361; Fax: 619-534-8158; E-mail: karinoffice@ucsd.edu

Received for publication 7 April 1999.

References

1. Baldwin, A.S. 1996. The NF- κ B and I κ B proteins: new discoveries and insights. *Annu. Rev. Immunol.* 14:649–681.
2. May, M.J., and S. Ghosh. 1998. Signal transduction through NF- κ B. *Immunol. Today.* 19:80–88.
3. Brown, K., S. Gerstberger, L. Carlson, G. Franzoso, and U. Siebenlist. 1995. Control of I κ B α proteolysis by site-specific, signal-induced phosphorylation. *Science.* 267:1485–1491.
4. Chen, Z., J. Hagler, V.J. Palombella, F. Melandri, D. Scherer, D. Ballard, and T. Maniatis. 1995. Signal-induced site-specific phosphorylation targets I κ B to the ubiquitin-proteasome pathway. *Genes Dev.* 9:1586–1597.
5. DiDonato, J.A., F. Mercurio, C. Rosette, J. Wu-li, H. Suyang, S. Ghosh, and M. Karin. 1996. Mapping of the inducible I κ B phosphorylation sites that signal its ubiquitination and degradation. *Mol. Cell. Biol.* 16:1295–1304.
6. Scherer, D.C., J. Brockman, Z. Chen, T. Maniatis, and D. Ballard. 1995. Signal-induced degradation of I κ B α requires site-specific ubiquitination. *Proc. Natl. Acad. Sci. USA.* 92:11259–11263.
7. Baeuerle, P.A., and T. Henkel. 1994. Function and activation of NF- κ B in the immune system. *Annu. Rev. Immunol.* 12:141–179.
8. Barnes, P.J., and M. Karin. 1997. Nuclear factor- κ B: a pivotal transcription factor in chronic inflammatory diseases. *N. Engl. J. Med.* 336:1066–1071.
9. DiDonato, J.A., M. Hayakawa, D.M. Rothwarf, E. Zandi, and M. Karin. 1997. A cytokine-responsive I κ B kinase that activates the transcription factor NF- κ B. *Nature.* 388:548–554.
10. Régnier, C.H., H.Y. Song, X. Gao, D.V. Goeddel, Z. Cao, and M. Rothe. 1997. Identification and characterization of an I κ B kinase. *Cell.* 90:373–383.
11. Mercurio, F., H. Zhu, B.W. Murray, A. Shevchenko, B.L. Bennett, J. Li, D.B. Young, M. Barbosa, M. Mann, A. Manning, and A. Rao. 1997. IKK-1 and IKK-2: cytokine-activated I κ B kinases essential for NF- κ B activation. *Science.* 278:860–866.
12. Zandi, E., and M. Karin. 1999. Bridging the gap: the I κ B kinase complex, composition and regulation. *Mol. Cell. Biol.* In press.
13. Woronicz, J.D., X. Gao, Z. Cao, M. Rothe, and D.V. Goeddel. 1997. I κ B kinase- β : NF- κ B activation and complex formation with I κ B kinase- α and NIK. *Science.* 278:866–869.
14. Zandi, E., D.M. Rothwarf, M. Delhase, M. Hayakawa, and M. Karin. 1997. The I κ B kinase complex (IKK) contains two kinase subunits, IKK α and IKK β , necessary for I κ B phosphorylation and NF- κ B activation. *Cell.* 91:243–252.
15. Zandi, E., Y. Chen, and M. Karin. 1998. Direct phosphorylation of I κ B by IKK α and IKK β : discrimination between free and NF- κ B-bound substrate. *Science.* 281:1360–1363.
16. Rothwarf, D.M., E. Zandi, G. Natoli, and M. Karin. 1998. IKK γ is an essential regulatory subunit of the I κ B kinase complex. *Nature.* 395:297–300.
17. Yamaoka, S., G. Courtois, C. Bessia, S.T. Whiteside, R. Weil, F. Agou, H.E. Kirk, R.J. Kay, and A. Israel. 1998. Complementation cloning of NEMO, a component of the I κ B kinase complex essential for NF- κ B activation. *Cell.* 93:1231–1240.
18. Ling, L., Z. Cao, and D.V. Goeddel. 1998. NF- κ B-inducing kinase activates IKK-alpha by phosphorylation of Ser-176. *Proc. Natl. Acad. Sci. USA.* 95:2791–2797.
19. Delhase, M., M. Hayakawa, Y. Chen, and M. Karin. 1999. Positive and negative regulation of I κ B kinase activity through IKK β subunit phosphorylation. *Science.* 284:309–313.
20. Hu, Y., V. Baud, M. Delhase, P. Zhang, T. Deerinck, M. Ellisman, R. Johnson, and M. Karin. 1999. Abnormal morphogenesis but intact IKK activation in mice lacking the IKK α subunit of the I κ B kinase. *Science.* 284:316–320.
21. Beg, A.A., W.C. Sha, R.T. Bronson, S. Ghosh, and D. Baltimore. 1995. Embryonic lethality and liver degeneration in mice lacking the RelA component of NF- κ B. *Nature.* 376:167–169.
22. Doi, T.S., M.W. Marino, T. Takahashi, T. Yoshida, T. Sakakura, L.J. Old, and Y. Obata. 1999. Absence of tumor necrosis factor rescues RelA-deficient mice from embryonic lethality. *Proc. Natl. Acad. Sci. USA.* 96:2994–2999.
23. Beg, A.A., and D. Baltimore. 1996. An essential role for NF- κ B in preventing TNF- α induced cell death. *Science.* 274:782–784.
24. Liu, Z.-G., H. Hu, D.V. Goeddel, and M. Karin. 1996. Dissection of TNF receptor 1 effector functions: JNK activation is not linked to apoptosis, while NF- κ B activation prevents cell death. *Cell.* 87:565–576.
25. Wang, C.-Y., M.W. Mayo, and A.S. Baldwin, Jr. 1996. TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF- κ B. *Science.* 274:784–787.
26. Sabapathy, K., Y. Hu, T. Kallunki, M. Schreiber, J.-P. David, W. Jochum, E.F. Wagner, and M. Karin. 1999. JNK2 is required for efficient T-cell activation and apoptosis but not for normal lymphocyte development. *Curr. Biol.* 9:116–125.
27. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. Molecular Cloning, A Laboratory Manual. 2nd ed. Cold Spring Harbor Laboratory Press, Plainview, NY. 9.31–9.46.
28. Church, G.M., and W. Gilbert. 1984. Genomic sequencing. *Proc. Natl. Acad. Sci. USA.* 81:1991–1995.
29. Li, N., and M. Karin. 1998. Ionizing radiation and short wavelength UV activate NF- κ B through two distinct mechanisms. *Proc. Natl. Acad. Sci. USA.* 95:13012–13017.
30. Joyner, A.L. 1993. Gene Targeting: A Practical Approach. Practical Approach Series. IRL Press at Oxford University Press, New York. 234 pp.
31. Karin, M., and M. Delhase. 1998. JNK or IKK, AP-1 or NF- κ B, which are the targets for MEK kinase 1 action? *Proc. Natl. Acad. Sci. USA.* 95:9067–9069.
32. Horwitz, B.H., M.L. Scott, S.R. Cherry, R.T. Bronson, and D. Baltimore. 1997. Failure of lymphopoiesis after adoptive transfer of NF- κ B-deficient fetal liver cells. *Immunity.* 6:765–772.