

Körner et al. Vol. 191, No. 1, January 3, 2000. Pages 89–96.

The authors regret that Table I contained a typographical error. Below, the correct value is indicated by boldface type.

Table I. Proportional Changes in Lymphocyte Populations in the Blood of 12- and 18–22-wk-old B6.WT, B6.gld, B6.TNF^{−/−}, and B6.gld.TNF^{−/−} Mice

	Age of mice					
	12 wk			18–22 wk		
	T cells	B220 ⁺ DN T cells	B cells	T cells	B220 ⁺ DN T cells	B cells
B6.WT*	16 ± 2.6	0.7 ± 0.2	75 ± 1.3	ND	ND	ND
B6.TNF ^{−/−} ‡	ND	ND	ND	12.7 ± 4	0.8 ± 0.2	78.5 ± 6
B6.gld§	5 ± 2.2	2 ± 0.7	87 ± 1.7	16 ± 15	11.3 ± 7	63 ± 23
B6.gld.TNF ^{−/−}	6 ± 2.2	2.1 ± 2	88 ± 3.8	5.1 ± 2	4.1 ± 2.4¶	83.1 ± 5

Peripheral blood lymphocytes were stained with anti-TCR- α/β and anti-B220. Data are recorded as mean percent ± SD.

*n = 4.

‡n = 4.

§n = 5 (<12 wk of age) and 8 (18–22 wk of age).

||n = 5 (<12 wk of age) and 9 (18–22 wk of age).

¶The level of B220⁺ DN T cells in B6.gld.TNF^{−/−} mice is significantly different from that in B6.gld (P < 0.01) mice.