

The μ Switch Region Tandem Repeats Are Important, but Not Required, for Antibody Class Switch Recombination

By Thomas M. Luby,* Carol E. Schrader,[‡] Janet Stavnezer,[‡] and Erik Selsing*

From the *Department of Pathology, Program in Immunology, Sackler School of Biomedical Sciences, Tufts University School of Medicine, Boston, Massachusetts 02111; and the [‡]Department of Molecular Genetics and Microbiology, Program in Immunology and Virology, University of Massachusetts Medical School, Worcester, Massachusetts 01655

Abstract

Class switch DNA recombinations change the constant (C) region of the antibody heavy (H) chain expressed by a B cell and thereby change the antibody effector function. Unusual tandemly repeated sequence elements located upstream of H chain gene exons have long been thought to be important in the targeting and/or mechanism of the switch recombination process. We have deleted the entire switch tandem repeat element ($S\mu$) from the murine μ H chain gene. We find that the $S\mu$ tandem repeats are not required for class switching in the mouse immunoglobulin H-chain locus, although the efficiency of switching is clearly reduced. Our data demonstrate that sequences outside of the $S\mu$ tandem repeats must be capable of directing the class switch mechanism. The maintenance of the highly repeated $S\mu$ element during evolution appears to reflect selection for a highly efficient switching process rather than selection for a required sequence element.

Key words: gene rearrangement • B lymphocyte • heavy chain • class switching • immunoglobulin isotype

Introduction

Antibody class switching is a process allowing B cells expressing IgM to give rise to cells that produce IgG, IgA, or IgE. The different classes of Ig molecules are defined by the constant region of the H chain. Class switching allows B cells to change effector function against a foreign antigen without losing antigen receptor specificity and takes place via DNA recombination events that involve looping out and deletion of DNA sequences (for reviews, see references 1–3). Unlike V(D)J recombination, which is targeted by specific sequence elements (4), no clear consensus sites for class switch recombination are apparent. The switch recombination mechanism appears to involve stretches of tandemly repeated DNA sequences, called switch (S)¹ regions, which are located upstream of each Ig H chain constant region gene except δ .

The roles of S regions in class switching are not clear, but evidence suggests that the tandemly repeated elements are used as the sites for cleavage (5), as well as being involved

in the joining reaction (6–8). Studies of integrated or extrachromosomal switch substrates (9–16) have suggested that S regions may be sufficient for targeting switch recombination and that two S regions are required for switch recombination to take place. Finally, transcribed S region tandem repeats have been found to form RNA–DNA complexes, and these have been suggested to play a role in switch recombination (17–20).

In all species that have been analyzed, switch recombination sites in H chain genes are found within or near a tandemly repeated S region. Furthermore, all identified breakpoints from DNA circles excised during switching in normal mouse splenocytes have been found within the repeated S elements (21–23). However, when chromosomal breakpoints in a variety of cell types were analyzed (7, 24), it was found that some mapped outside the tandemly repeated elements, especially for the $S\mu$ element. Some breakpoints found in DNA circles that arise from an apparent class switching process in a lymphoma cell line are also found outside of the $S\mu$ region (24). These data suggest that the function of and requirement for the $S\mu$ tandem repeats is not clearly understood.

In the mouse, the $S\mu$ element consists of repeated (GAGCT)_nGGGGT sequences, where n varies from one to

Address correspondence to E. Selsing, Department of Pathology, Tufts University School of Medicine, Boston, MA 02111. Phone: 617-636-0467; Fax: 617-636-6903; E-mail: eselsing@opal.tufts.edu

¹Abbreviations used in this paper: Ach, acetylcholine; Ars, *p*-phenylarsenate; DC, digestion circularization; NHEJ, nonhomologous end joining; S, switch.

seven in different repeats, but has an average value of three. To directly test whether $S\mu$ is required for antibody class switching, we have removed all the $S\mu$ tandem repeats from the mouse *Igh* locus to determine the impact of this deletion on the switching process. We find that the $S\mu$ tandem repeats are not required for class switching in the mouse *Igh* locus, although the efficiency of switching is reduced. The maintenance of the highly repeated $S\mu$ element during evolution appears, therefore, to reflect selection for a highly efficient switching process rather than selection for a required sequence element.

Materials and Methods

Targeting of $S\mu$ in Embryonic Stem Cells. The targeting construct is shown in Fig. 1. The 5' homology region is a 1.4-kb segment upstream of $S\mu$ bounded by *EcoRI* and *HindIII* sites. The 3' homology region is a 4.6-kb segment downstream of $S\mu$ bounded by *HindIII* and *KpnI* sites. The *neo/loxP* cassette (a gift from Dr. F. Alt, Harvard Medical School, Boston, MA) was inserted in between the two homology regions. The targeting construct was used in standard gene targeting approaches to obtain chimeric mice that carried the targeted allele. Chimeric mice that transmitted the mutation were mated to Cre recombinase transgenic mice (a gift of Dr. J. Chen, Massachusetts Institute of Technology, Cambridge, MA) in order to remove the neomycin cassette. This provided the $\Delta S\mu$ allele that has one *loxP* site replacing $S\mu$.

Immunizations and Immunoassays. Mice were immunized with 100 $\mu\text{g/ml}$ *p*-phenylarsenate-KLH (Ars-KLH) in CFA (GIBCO BRL), and boosted two times with Ars-KLH in incomplete adjuvant. For hybridoma production, animals were boosted with a final injection of 100 $\mu\text{g/ml}$ Ars-KLH in PBS 3 d before fusion.

Total IgG titers were determined by ELISA on plates coated with 4 $\mu\text{g/ml}$ rat anti-mouse IgG (Zymed Laboratories) and blocked with PBS/3% BSA (Sigma-Aldrich). Samples were applied and detected with 0.5 $\mu\text{g/ml}$ biotin-labeled rat anti-mouse IgG (Zymed Laboratories) and 0.5 $\mu\text{g/ml}$ alkaline phosphatase-streptavidin (Boehringer). Plates were developed with 20 $\mu\text{g/ml}$ of *p*-nitrophenyl phosphate (Sigma-Aldrich) and analyzed on a Dynatech MR700 plate reader.

For quantitation of the different isotypes in the sera, the same basic protocol described for the total IgG ELISA was used with the following modifications: plates were coated with unlabeled goat anti-mouse IgG3, IgG1, IgG2b, and IgA (all from Southern Biotechnology Associates, Inc.). Detecting antibodies used were: goat anti-mouse IgG3, goat anti-mouse IgG1, goat anti-mouse IgG2b, and (for IgA) both goat anti-mouse κ and λ (all from Southern Biotechnology Associates, Inc.). Total IgM was determined by ELISA using the same basic protocol as above with goat anti-IgM as the coating reagent and rat anti-IgM as the detecting reagent.

Surface IgM staining was done on splenocytes from wild-type and $\Delta S\mu$ mice using FITC-goat anti-mouse IgM (Southern Biotechnology Associates, Inc.). Samples were collected on a FAC-Scan™ (Becton Dickinson) and analyzed using FloJo (Treestar) analysis software.

In Vitro Switching and Digestion Circularization PCR. T cell-depleted splenocytes were cultured at $2 \times 10^5/\text{ml}$ in 5-ml cultures with LPS alone to induce IgG3, or with cytokines added to induce IgG1 (IL-4, 800 U/ml), IgG2b (dextran sulfate, 30 $\mu\text{g}/$

ml), or IgG2a (IFN- γ , 10 U/ml). After 4 d, cells were acid treated (50 mM sodium acetate, pH 5.2, 85 mM sodium chloride, 5 mM potassium chloride, 1% FCS) for 2 min on ice to remove Ig bound to Fc receptors, and washed and stained for FACS® analyses with FITC-goat anti-mouse IgM and PE-goat F(ab')₂ anti-mouse IgG1, IgG3, IgG2a, or IgG2b (Southern Biotechnology Associates, Inc.).

Digestion circularization (DC)-PCR is described briefly here and in detail elsewhere (25, 26). Genomic DNAs from LPS and LPS plus IL-4-induced cultures were digested with *EcoRI* (2 $\mu\text{g}/100 \mu\text{l}$) overnight, ligated overnight at a concentration of 180 ng/ μl , and then dialyzed against distilled water. PCR primers are as published for acetylcholine receptor (AChR) and IgG1 (25) and IgG2b (10). Twofold dilutions of ligated DNA and a plasmid standard for AChR, P2AO (25), were analyzed to demonstrate that the amount of product depends on the amount of input DNA in a linear fashion. For quantitation, [α -³²P]dCTP




Figure 1. Generation of the $\Delta S\mu$ mice. (A) Diagram of $S\mu$ tandem repeats and map of the J_H - $C\mu$ intron. $E\mu$, intronic enhancer; $C\mu$, μ constant region exons; μmem , μ membrane exons; E, *EcoRI*; H, *HindIII*; B, *BamHI*; K, *KpnI*. In the expanded view of $S\mu$, each vertical line represents either a GAGCT or GGGGT sequence and the two *HindIII* sites represent the pair of sites flanking $S\mu$ in the diagram of the wild-type (WT) allele. (B) Southern blot analyses of knockout mice. Genomic DNAs from mice with the indicated phenotypes were digested and hybridized. The left blot contains *BamHI* digests and was hybridized with pJ11 (reference 52), a 1.8-kb *BamHI*-*EcoRI* fragment containing JH3 and JH4. Sizes of wild-type (wt) and $\Delta S\mu$ alleles are ~8 and 5.3 kb, respectively. The right blot contains *EcoRI* digests and was hybridized sequentially with, first, a 900-bp *KpnI*-*XbaI* probe containing the μ membrane exons and, second, pM2-20 (reference 52), a 1.8-kb probe containing ~1.4-kb of $S\mu$ tandem repeat sequences. Sizes of the wild-type and $\Delta S\mu$ alleles are ~12 and 8.9 kb, respectively. Positions of λ *HindIII* DNA fragments are indicated. μmemb , μ membrane exons.

was incorporated, and the amounts of $S\mu$ - $S\gamma 1$ and $S\mu$ - $S\gamma 2b$ PCR products were normalized to the amount of AchR product to control for the efficiency of digestion and circularization.

Filter Hybridization. Northern blots were prepared and analyzed as described previously (27) using a 0.7-kb EcoRI-HindIII probe containing the $I\mu$ exon. For Southern blots, genomic DNA was digested, electrophoresed, and transferred to nitrocellulose (Sartorius). The membranes were UV cross-linked and hybridized ($1\times$ SSC, $1\times$ Blotto, 0.1% SDS) with the indicated probes (48 h at 65°C). The blots were washed sequentially with $3\times$ SSC, 5 mM EDTA, 0.1% SDS, $1\times$ Blotto, and 50 $\mu\text{g}/\text{ml}$ salmon sperm DNA; $0.2\times$ SSC, 1.25 mM EDTA, and 0.1% SDS; and $0.1\times$ SSC, 1.25 mM EDTA, and 0.1% SDS (all at 65°C), and autoradiographed.

$S\mu/S\gamma 1$ Junction Analysis. PCR and sequencing of recombination junctions was generally as described previously (28). Nested μ primers were 5'-AAGTTGAGGATTCAGCCGA-ACTGGAGAGG-3' (MUSICD07 nos. 3434-3463) followed by 5'-GCTTGAGTAGTTCTAGTTTCC-3' (MUSICD07 nos. 3502-3522). The sequence of the nested set of $\gamma 1$ primers was 5'-CAATTAGCTCCTGCTTCTGTGG-3' (D78344 nos. 8464-8488) followed by 5'-TCTAATCTGCCCTGTTCCCTCTACAAC-3' (D78344 nos. 8365-8395). PCR products were directly sequenced or cloned and sequenced using a TA cloning kit (Invitrogen). Switch junctions in $\Delta S\mu$ and wild-type mice were analyzed for mutation frequency. Mutations in $\Delta S\mu$ junctions were identified by comparison with μ and $\gamma 1$ germline sequences that were determined by compiling sequences from several independent PCR clones. These germline sequences (derived from the 129 strain) were found to be identical to the corresponding published BALB/c sequences. Mutations in junction regions were confirmed from two independent PCR clones. The frequencies of mutation per basepair in the donor (5') and acceptor (3') S regions were determined by dividing the total number of mutations by the total number of basepairs in the sequences. Mutation frequencies in wild-type junctions were similarly calculated using published sequences (7, 8).

Results

Mutant Mice that Lack $S\mu$. The first step in the creation of the $S\mu$ knockout was defining which sequences to remove from the JH- $C\mu$ intron. The goal was to remove all

of the tandem repeats without deleting elements required for transcription of the locus. The upstream border of the $S\mu$ deletion was set at the second HindIII site downstream of $E\mu$ (MUSICD07 base no. 4995, sequence data available from Genbank/EMBL/DDBJ under accession no. J00440), leaving intact important elements such as the μ enhancer ($E\mu$), matrix attachment regions (29), and the $I\mu$ exon (Fig. 1). The downstream border was set at the HindIII site ~ 1.2 kb upstream of $C\mu$ (MUSICD09 base no. 1208, sequence data available from Genbank/EMBL/DDBJ under accession no. J00442, leaving a region of DNA indicated to be important in the transcription of Ig H chain transgenes (27, 30). This deletion removes all of the $S\mu$ tandem repeat region. As indicated in Fig. 1 A, some GAGCT or GGGGT sequences remain in the JH- $C\mu$ intron after the deletion of the $S\mu$ element. One GAGCT is left in the intron between $E\mu$ and the 5' border of the deletion, while 11 additional nontandemly spaced GAGCT sequences are located between the end of the deletion and $C\mu$. Also, a single GGGGT is ~ 150 bp upstream of the deletion. Gene targeting was used to produce mice carrying the $\Delta S\mu$ deletion, which was confirmed by Southern blotting (Fig. 1 B).

Serum IgG Antibodies in $\Delta S\mu$ Mice. Once the $\Delta S\mu$ deletion was bred to homozygosity, we bled the animals at various time points to determine whether the mice had detectable IgG titers. Sera from the mice were analyzed at weeks 6, 9, 12, and 15. The results show that $\Delta S\mu$ animals have the ability to produce IgG, although the levels appear to be reduced approximately twofold relative to wild-type mice (Fig. 2 A). We also analyzed the serum isotype profile of these mice at week 15 (Fig. 2 B). It appears that $\Delta S\mu$ mice have a slight reduction in IgG1 and IgG3, a relatively large reduction in IgG2b, and no decrease in IgA. When IgM titers were checked on a subset of these week 15 sera, an increase in the amount of serum IgM in the $\Delta S\mu$ mice was noted (Fig. 2 C). However, when surface IgM of splenocytes was examined by flow cytometry, both C57BL/6 and $\Delta S\mu$ mice had approximately equivalent numbers of IgM expressing B cells and the same levels of surface IgM




Figure 2. Analysis of IgM and IgG production in unimmunized $\Delta S\mu$ mice and their littermate controls. (A) Animals were bled and total IgG titers were determined for three wild-type, six heterozygous, and eight mutant mice at weeks 6, 9, 12, and 15. (B) Isotype profile of serum from wild-type, heterozygous, and mutant mice were analyzed at week 15. (C) Serum IgM titers at week 15 were determined on a subset of wild-type, heterozygous, and mutant mice. (D) Surface IgM staining of wild-type and mutant mice was examined by flow cytometry using a FITC-conjugated monoclonal antibody reactive with mouse IgM.




Figure 3. Reduced, but easily detectable IgG expression on B cells from $\Delta S\mu$ mice. IgG2b is more severely affected than IgG1 by the deletion of $S\mu$ tandem repeats. Splenic B cells from wild-type or $\Delta S\mu$ mice were cultured with either LPS plus IL-4 to induce IgG1 or with LPS plus dextran sulfate (DXS) to induce IgG2b. On day 4 the cells were stained for surface IgG and IgM and analyzed by flow cytometry. The percentage of cells in the upper gate (indicated in the top right corner of each panel) includes cells positive for IgG alone and for IgG and IgM together. The gates for IgG1 were set on wild-type cells treated with LPS alone, which does not induce switching to IgG1, and then were adjusted for the other isotypes to optimize separation of the populations. Gates for wild-type and $\Delta S\mu$ cells were always the same for a given isotype.

expression (Fig. 2 D). Northern blot analyses of unstimulated and LPS-stimulated splenocytes also showed identical levels of spliced $I\mu$ germline transcripts in $\Delta S\mu$ and wild-type mice (data not shown). These results suggest that the $S\mu$ deletion has little or no effect on early B cell differentiation or on the accessibility of the $I\mu$ - $C\mu$ locus.

Switching of $\Delta S\mu$ B Cells in Culture. To determine if the differences in serum isotype profiles of the $\Delta S\mu$ mice were due to an intrinsic difference in the B cells, we polyclonally stimulated splenic B cells from the mice in vitro for 4 d and measured the amount of switching to the IgG subclasses by surface staining of Ig isotypes and flow cytometry (representative data shown in Fig. 3). The results (summarized in Table I) extend the serological data and show that switching by the $\Delta S\mu$ mice is partially reduced for all isotypes analyzed: two- to threefold for IgG1, IgG3, and IgG2a, and fivefold for IgG2b.

To determine whether the class switching detected in the $\Delta S\mu$ mice is due to a DNA recombination event occurring within the $S\mu$ intron and to obtain a semiquantitative estimate for the level of switch recombination in these mice, we analyzed DNA from in vitro-activated B cells by DC-PCR (25, 26, 31). Using one primer upstream of $S\mu$ and one primer downstream of either $S\gamma 1$ or $S\gamma 2b$, this

Table I. FACS[®] Analysis of In Vitro Switching on Day 4 by Splenic B Cells from $\Delta S\mu$ and Wild-type Littermates

			Percent switched cells		Percent WT
			Exp. 1	Exp. 2	
IgG1	WT	LPS	1.9	0.6	
		LPS plus IL-4	24.7	41.1	
	$\Delta S\mu$	LPS	1.4	0.3	
		LPS plus IL-4	19.6	19.8	63 ± 15*
IgG3	WT	LPS	13.4	9.8	
		LPS plus DxSO4	10.7	11.6	
	$\Delta S\mu$	LPS	6.3	4.5	
		LPS plus DxSO4	5.2	4.1	44.2 ± 2*
IgG2b	WT	LPS	7.6	9.1	
		LPS plus DxSO4	23.9	19.7	
	$\Delta S\mu$	LPS	3.3	1.2	
		LPS plus DxSO4	5.7	3.3	24.3 ± 12*
IgG2a	WT	LPS plus IFN- γ	ND	11.0	
	$\Delta S\mu$	LPS plus IFN- γ	ND	3.6	33

Exp., experiment; ND, not determined. The percentages of cells expressing the switched isotypes are shown for two experiments (Exp. 1 and 2). For each isotype, average values for $\Delta S\mu$ switching as percentages of WT are also shown.

* ± SEM.

technique yields a single size band that can be quantified and is independent of the location of switch recombination junctions. An $S\mu/S\gamma 1$ product was clearly detectable in DNA from $\Delta S\mu$ B cells stimulated with LPS plus IL-4, but not with LPS alone (Fig. 4), and was reduced fourfold relative to wild-type in both of two independent PCRs. In the experiment shown, the reduction in IgG1 switching as assayed by cytometry was twofold (Table I, experiment 2). The $S\mu/S\gamma 2b$ PCR product from LPS-stimulated $\Delta S\mu$ B cells was reduced by an average of ninefold relative to wild-type in two independent PCR assays, compared with eightfold reduction assayed by cytometry. Thus, decreases in IgG expression observed by flow cytometry are concordant with the reduction in switch recombination in the $\Delta S\mu$ B cells as determined by DC-PCR.

Switch Recombination Sites in IgG-Producing $\Delta S\mu$ Hybridomas. To characterize the nature of switch recombination events in $\Delta S\mu$ mice, we produced IgG1-expressing hybridomas from animals immunized with Ars-KLH. DNA fragments containing recombination sites were PCR-amplified from hybridoma DNAs. Approximately 30% of the hybridomas yielded PCR fragments. Fragments from 10 hybridomas were sequenced to locate the switch joining




Figure 4. DC-PCR demonstrates switch recombination at the DNA level is reduced in B cells from $\Delta S\mu$ mice stimulated with LPS or LPS plus IL-4 (L+4). DNA template levels were adjusted by normalizing to the amount of AchR DC-PCR product and then twofold dilutions of adjusted levels of input or a plasmid standard were amplified for AchR, and for $S\mu/S\gamma 1$ and $S\mu/S\gamma 2b$ recombination, with the incorporation of [α - ^{32}P]dCTP. The fold reductions relative to wild-type (WT) from two independent PCR reactions were: IgG1, 3.6 and 3.65; IgG2b, 9.7 and 8.5.

sites. Sequenced fragments were compared with germline JH-C μ sequences and germline C $\gamma 1$ flanking sequences to locate the sites of recombination.

As shown in Fig. 5 A, the recombination sites in the $\Delta S\mu$ JH-C μ intron are found at a variety of locations; ~ 2 kb separate the most 5' and 3' sites. In wild-type mice, most switch sites in IgG1-producing cells occur within $S\mu$; for those sites that are outside of $S\mu$, most occur in the 5' flanking sequence with a much smaller percentage in the 3' flanking sequence. In the $\Delta S\mu$ mice, most μ switch sites in IgG1-producing cells are in the 3' flanking sequence with a smaller proportion in the 5' flanking sequence. Although our data set is limited, there does not appear to be any shared motif at the sites of recombination. Only one site is within a GAGCT sequence, but all sites are located < 250 bp from the nearest GAGCT.

The sites of switch recombination within the $S\gamma 1$ region of the mutant mice are shown in Fig. 5 B and these are compared with analogous sites in wild-type animals. The $\gamma 1$ recombination sites observed for $\Delta S\mu$ hybridomas are clearly clustered to the 3' end of $S\gamma 1$ relative to the $\gamma 1$ sites in wild-type mice. However, because the $\Delta S\mu$ sites were from PCR amplifications, whereas the published wild-type sites were generally obtained by cloning genomic DNA fragments, we believe that this clustering may be due to smaller fragments being more readily amplified by PCR. Shorter PCR products will mostly exhibit recombination sites located at the 3' end of the long (10 kb) $S\gamma 1$ element. On the other hand, there is no indication that the use of PCR has affected our analysis of the μ switch recombination sites in $\Delta S\mu$ JH-C μ intron. PCR effects would result in μ switch sites clustered at the 5' end of the JH-C μ intron. Instead, the actual observed μ switch sites (Fig. 5 A) are located toward the 3' end of this intron.

The DNA sequences surrounding the recombination sites in H chain genes that have undergone class switching frequently display single basepair mutations. These muta-




Figure 5. (A) $\Delta S\mu$ and wild-type (WT) recombination junctions in the JH-C μ intron. The wild-type junctions pictured are all from $\mu - \gamma 1$ recombinations and were derived from published sequences. Junctions nos. 1 and 2 are from reference 53, nos. 3 and 4 from reference 7, nos. 5-7 from reference 54, no. 8 from reference 21, nos. 9-19 from reference 23, nos. 20 and 21 from reference 55, no. 22 from reference 56, and no. 23 from reference 57. $\Delta S\mu$ recombination junctions were generated by PCR from IgG1-producing hybridomas. Sites were from hybridomas 16B3 (A), 18A4 (B), 8B5 (C), 8A2 (D), 18B6 (E), 18B2 (F), 9A6 (G), 8C4 (H), 4D5 (I), and 16B5 (J). Only the relevant restriction sites are shown on the map. E, EcoRI; H, HindIII; X, XbaI. Location of the PCR primer used to generate $\Delta S\mu$ junctions is represented by small arrowheads. (B) $\Delta S\mu$ and wild-type (WT) switch junctions upstream of C $\gamma 1$. The wild-type junctions are all $\mu - \gamma 1$ recombinations and are derived from (reference 55). The wild-type junctions shown here are a subset of the group shown in A because we could not accurately place all the breakpoints within the $\gamma 1$ tandem repeats. $\Delta S\mu$ junctions were generated by PCR from IgG1-producing hybridomas; the small arrowheads represents the location of the PCR primer used.

tions have been suggested to reflect a role for error-prone DNA synthesis in the switching mechanism (6, 32). Mutations are seen surrounding the switch sites in $\Delta S\mu$ mice and, within the μ sequences, the frequency of mutation (1.8%) is similar to that previously reported for wild-type mice (1.4%). However, within the $\gamma 1$ sequences flanking the switch junctions in $\Delta S\mu$ mice there appears to be an elevated level of mutation (5.5%) relative to the levels found in analogous flanking S regions in wild-type mice (1.3%). This elevated level is mainly due to a rather large number of mutations found in 3 of the 11 cloned recombination sites that have been analyzed (Fig. 6).

Hybridomas from $\Delta S\mu$ Mice Frequently Have Switch Recombination at Only One μ Allele. The panel of $\Delta S\mu$ IgG1-producing hybridomas was analyzed by Southern blot




Figure 6. Sequence analysis of $\Delta S\mu$ IgG1 hybridoma switch recombination junctions. Approximately 80 bp of each clone is shown, 40 bp upstream and 40 bp downstream of each breakpoint. The sequence on top is the corresponding germline μ sequence, and the sequence on the bottom is the germline $\gamma 1$ sequence. A small vertical line is drawn between bases if they are identical. The switch junctions are shown as either a large vertical line if there are not any shared base pairs, or as a box if there are shared base pairs. Junctions are assigned according to the following criteria: if there is a base change close to the potential breakpoint it must be followed by at least one base that is the same as germline; if there are two or more base changes they must be followed by at least the same number of germline bases. Base changes are marked with an asterisk, insertions have no corresponding germline sequence, and deletions have the corresponding germline base placed below the germline sequence.

hybridization to assess the recombination events that had occurred during class switching in immunized $\Delta S\mu$ B cells. The hybridomas in this panel were isolated as clones from soft agar but were not re-cloned after isolation; thus, a fraction of the hybridomas might represent multiple clones. This fraction appears to be small because only one of the hybridomas displays more than two B cell-derived Igh alleles when the allele derived from the X63.Ag8 tumor fusion parent is discounted.

Hybridoma DNAs were analyzed both with 5' $S\mu$ probes (Fig. 7, left) and $\gamma 1$ probes (Fig. 7 right). Consistent with the sequenced PCR clones described above, recombined DNA fragments that hybridize with both 5' $S\mu$ and $S\gamma 1$ probes are found in almost all of the analyzed hybridomas, including some of those that did not yield detectable

fragments in PCR assays. The inability to obtain PCR products from some hybridomas is likely due to a long distance between the μ and $\gamma 1$ primers used in the amplification. The large size of some recombined bands, together with the known size of the $\Delta S\mu$ JH-C μ intron, suggests that these bands reflect $S\gamma 1$ recombination sites located further 5' in the $S\gamma 1$ element.

The most notable finding from the Southern blot analyses of $\Delta S\mu$ IgG1-producing hybridomas is that most of the hybrids display germline JH-C μ alleles (indicated by the arrowheads in Fig. 7). This result indicates that only one of the two JH-C μ alleles in these B cells has undergone switch recombination. In contrast, three separate studies of wild-type mice showed that 80–100% of IgG hybridomas displayed switch recombinations at both μ alleles (33–35). Because the $\Delta S\mu$ hybridomas produce IgG1, it appears that the nonproductive chromosome is the one that has not undergone switching within the μ region. On the other hand, most of the $\Delta S\mu$ hybridomas have undergone recombination at both $S\gamma 1$ regions as indicated by the small number of germline $S\gamma 1$ bands among the hybridoma panel (Fig. 7). Therefore, the lack of switching in the $\Delta S\mu$ JH-C μ intron indicates a defect in the ability of the μ locus to participate in switch recombination rather than an overall defect in switch recombination of the nonproductive chromosome.

Discussion

S μ Is Not Required for Class Switch Recombination. The tandemly repeated S region sequences are the most unique sequence features found within the DNA regions known to be involved in class switch recombination. Since their discovery (36), the tandem repeats have been the focus of investigations into the targeting and mechanism of class switching. These repetitive sequences have been suggested to be the sites of DNA breakage for switch recombination (5, 7), and transfection experiments using switch constructs have suggested that S regions may be sufficient to direct switch recombination (9–16). However, our analyses of the $\Delta S\mu$ mice, in which all the tandem repeats in the JH-C μ intron have been deleted, demonstrate that the $S\mu$ element is not required for antibody class switching to occur.

Tandem Repeats Appear to be Important for Efficient Switching of Both Igh Alleles. In normal mice, >90% of hybridomas that exhibit switching on a productive H chain allele also exhibit switch recombination on the nonproductive allele, and these are frequently to the same isotype (34, 37, 38). There is no evidence to suggest that the switch mechanism can distinguish between the productive and nonproductive alleles in a B cell; therefore, this efficient recombination of both alleles has probably been evolutionarily selected to ensure that the productive allele undergoes class switching when the cell is appropriately stimulated.

In the $\Delta S\mu$ knockout mouse, however, a large percentage of the hybridomas that have undergone switching exhibit recombination only at the functional μ allele. In these cells, the nonfunctional μ allele does not appear to have




Figure 7. Southern blots of $\Delta S\mu$ IgG1 hybridomas. DNAs from 32 hybridomas were digested with EcoRI and hybridized. The blots on the left were hybridized with a 700-bp EcoRI-HindIII fragment located just 3' of the E μ enhancer (pJ14c; reference 53). On the right, the same blots were probed with a 10-kb EcoRI fragment containing the S γ 1 S region (p γ 1EH10.0; reference 58). The germline bands are marked with an arrowhead and the sizes are \sim 8.9 kb for the μ intron and 16 kb for the γ 1 intron. Positions of Hind III fragments of λ phage DNA are indicated for each gel.

undergone any recombination event. This result suggests that during immune responses in $\Delta S\mu$ mice some cells that have been stimulated to switch might recombine only a nonfunctional μ allele, and some may fail to recombine either μ allele. This notion would appear to be consistent

with the observed increased IgM production in the $\Delta S\mu$ mice. Thus, the efficiency of switching that is provided by the presence of the tandem repeats appears to increase the probability that B cells participating in an immune response undergo switching.




Figure 8. Analysis of the frequency of G nucleotides in the $\Delta S\mu$ J_H-C μ intron. The top three graphs wrap around to form a continuous sequence from J_H to the beginning of C μ in the $\Delta S\mu$ mouse. The bottom graph represents a portion of the sequence that has been removed in the $\Delta S\mu$ mouse; however, \sim 2.5 kb of tandem repeats is not included in this graph because the sequence is not known. The analysis was done by counting the number of G residues for nucleotide nos. 1–10 (numbers not shown), and plotting the value on the graph. This was repeated for nucleotide nos. 2–11, nos. 3–12, etc. Essentially, a peak in the graph indicates a stretch of DNA with numerous G residues. A random sequence would have 2.5 G residues per 10 nucleotides. An oval on the top line of the graph represents runs of G that are three nucleotides or greater. The positions of the recombination sites in the $\Delta S\mu$ IgG1 hybridomas are indicated by a horizontal line between the 7th and 8th line of the graph. Relevant restriction sites and features of the intron are indicated.

Potential Roles for S μ in Switch Recombination. Class switch recombination occurs by a type of nonhomologous end joining (NHEJ) as indicated by the involvement of DNA protein kinase and Ku in the process (39–41), and by the absence of any extensive homology surrounding the joining sites (7). Switch recombination can be divided into three major steps: targeting, initiation (cleavage), and resolution (rejoining; reference 42). The S μ tandem repeats could act at one or more of these steps. Targeting of isotype switching appears to be relatively intact in the Δ S μ mice. The expected isotypes are produced upon in vitro stimulation with cytokines, and the locations of the switch junctions in both the JH-C μ intron and in S γ 1 appear consistent with normal targeting of switch recombination. Thus, it seems more likely that S μ has a role in either switch initiation or resolution.

Our results provide some data suggesting that S μ tandem repeats could be involved during the resolution phase of switch recombination. Δ S μ mice and mismatch repair (MMR)-deficient mice exhibit switching defects that have similar isotype profiles (26, 28). It has been proposed that the MMR protein Msh2 is involved in end processing that occurs after formation of switch double strand breaks but before ligation (26, 28). Therefore, the similar defects in isotype expression could suggest that Msh2 and S μ both affect switch resolution. However, Msh2 and S μ do not appear to impinge on switch recombination in the identical manner because Δ S μ mice do not exhibit the focus of μ switch sites within GAGCT sequences that has been reported in Msh2 knockout animals (28). A defect in switch resolution might also result in increased mutations around switch sites; our limited data from Δ S μ switch junctions show potential increases in mutation frequency that would be consistent with this suggestion.

The S μ element could certainly also be important in the initiation of switch recombination. The observation that many Δ S μ B cells have Igh alleles that are germline in the JH-C μ intron but rearranged at γ 1 suggests that initiation is intact at S γ 1, whereas it is decreased at μ . One proposed role for S regions in the initiation of class switching is as the site of double strand breaks; this role is supported by the detection of strand breaks within S γ 3 in B cells undergoing switching (5). Although no site for break induction has yet been identified in S μ , it is likely that some sites are located within S μ and it is possible that the loss of these sites is responsible for the decrease in switching. Furthermore, NHEJ is important in the resolution step of switch recombination and is generally considered to be sequence independent. Therefore, the presence or absence of specific S μ sequences would seem less likely to affect the efficiency of NHEJ during resolution, suggesting that S μ might be more likely to be important for switch initiation. However, because little is known about the nature of the DNA ends generated during switch initiation, or about possible processing of these ends before DNA joining, it may well be that S μ sequences play a role at both the initiation and the resolution of switch breakpoints.

Locations of Δ S μ Switch Sites Suggest that Sequences 3' of S μ Might Be Important for Switch Recombination. Because S μ is not required for isotype switching, it appears that sequences located outside of S μ must be capable of directing the switch recombination mechanism. RNA transcription and splicing have been shown to be important in S region function (43–46), although they cannot be sufficient for specific targeting of the switch mechanism and must be coupled with some additional sequence or structural information. Perhaps targeting is focused by sequences located between E μ and S μ or between S μ and C μ ; both of these regions have not yet been tested by deletional mutagenesis.

S μ , as well as other S regions, is made up of tandemly repeated sequences that exhibit relatively high levels of G nucleotide residues and GGG or GGGG sequences on the nontranscribed DNA strand. Fig. 8 diagrams the location of G-rich segments and GGG sequences for both the wild-type S μ region and the Δ S μ JH-C μ intron. A segment of \sim 200–300 bp located 3' of the loxP site in the Δ S μ intron exhibits G richness, clusters of GGG sequences, and some isolated GAGCT sequences (Figs. 1 and 8). This 3' element does not exhibit the repetitive pattern of G-richness that is apparent in the 3,000-bp wild-type S μ , part of which is shown at the bottom of Fig. 8. This 200–300 bp 3' region has not been considered part of S μ because there are no tandem repeats present and the frequency of switch recombination sites in wild-type mice within this element is lower than within S μ . Indeed, the frequency of recombination sites within the 3' element in wild-type mice is lower than the frequencies within other JH-C μ regions that exhibit no GAGCT sequences (see wild-type breakpoints summarized in Fig. 5 A). The recombinational activity of this 3' region has never been tested in switch substrate experiments. Nevertheless, the 200–300-bp element is found centrally situated in the region where Δ S μ switch sites occur. Perhaps this region is important in the switch recombination process.

It has been hypothesized that S regions might function in class switching by forming a special secondary structure that starts the recombinational process (5, 47–50). The proposed mechanisms all involve single-stranded S region DNA. Transcription of tandemly repeated S regions has been shown to lead to stable RNA–DNA complexes and single-stranded S region sequences which could play an important role in class switch recombination (17–20).

It is not known what causes RNA–DNA complexes to form in transcribed S regions, but extensive polypurine or GGG stretches have been suggested to play a role. In Δ S μ mice, the JH-C μ intron exhibits considerably fewer of these types of sequence elements than do the S μ tandem repeats. However, it is possible that the complexes can still form in the Δ S μ JH-C μ intron, perhaps using the shorter stretches of GGG clusters remaining. The reduction in Δ S μ switching might be due to a decrease in the efficiency of formation or the stability of the complex in vivo.

RNA–DNA complexes are an intriguing mechanism for targeting switch recombination; however, there is no direct

evidence that demonstrates a role for these complexes in class switch recombination. In fact, switching in some species appears to involve sequences that are not G rich or even purine rich (50, 51). Although switching processes could differ in different species, it is also possible that RNA–DNA complexes are not actually needed for switching or that the sequence features that promote complex formation are not yet understood. The location of switch sites in $\Delta S\mu$ mice do, however, suggest that the 200–300-bp element in the $\Delta S\mu$ JH–C μ intron could contain the minimal sequences required for switching. The long S μ tandem repeat element found in a variety of animal species appears to be present for optimal efficiency of the switch recombination process.

We thank Naomi Rosenberg for helpful comments on the manuscript and Jørgen Selsing for help with sequence analyses.

Supported by National Institutes of Health grants AI23283 and AI42108 (J. Stavnezer), AI24465 and AI42569 (E. Selsing), and AI07077 (T.M. Luby).

Submitted: 11 August 2000

Accepted: 6 December 2000

References

- Gritzmacher, C.A. 1989. Molecular aspects of heavy chain class switching. *Crit. Rev. Immunol.* 9:173–200.
- Stavnezer, J. 1996. Antibody class switch. In *Advances in Immunology*. Vol. 61. Academic Press, Inc., San Diego, CA. 79–146.
- Stavnezer, J. 2000. Molecular processes that regulate class switching. *Curr. Top. Microbiol. Immunol.* 245:127–168.
- Sakano, H., K. Huppi, G. Heinrich, and S. Tonegawa. 1979. Sequences at the somatic recombination sites of immunoglobulin light chain genes. *Nature*. 280:288–294.
- Wuerffel, R.A., J. Du, R.J. Thompson, and A.L. Kenter. 1997. Ig Sy3 DNA-specific double strand breaks are induced in mitogen-activated B cells and are implicated in switch recombination. *J. Immunol.* 159:4139–4144.
- Dunnick, W., M. Wilson, and J. Stavnezer. 1989. Mutations, duplication and deletion of recombined switch regions suggest a role for DNA replication in the immunoglobulin heavy chain switch. *Mol. Cell. Biol.* 9:1850–1856.
- Dunnick, W., G.Z. Hertz, L. Scappino, and C. Gritzmacher. 1993. DNA sequences at immunoglobulin switch region recombination sites. *Nucleic Acids Res.* 21:365–372.
- Du, J., Z. Ye, A. Shanmugam, and A. Kenter. 1997. Analysis of immunoglobulin Sy3 recombination breakpoints by PCR: implications for the mechanism of isotype switching. *Nucleic Acids Res.* 25:3066–3073.
- Daniels, G.A., and M.R. Lieber. 1995. Strand specificity in the transcriptional targeting of recombination at immunoglobulin switch sequences. *Proc. Natl. Acad. Sci. USA.* 92:5625–5629.
- Ballantyne, J., D. Henry, J. Muller, F. Briere, C. Snapper, M. Kehry, and K. Marcu. 1998. Efficient Recombination of a switch substrate retrovector in CD40-activated B lymphocytes: implications for the control of CH gene switch recombination. *J. Immunol.* 161:1336–1347.
- Leung, H., and N. Maizels. 1992. Transcriptional regulatory elements stimulate recombination in extrachromosomal substrates carrying immunoglobulin switch-region sequences. *Proc. Natl. Acad. Sci. USA.* 89:4154–4158.
- Leung, H., and N. Maizels. 1994. Regulation and targeting of recombination in extrachromosomal substrates carrying immunoglobulin switch region sequences. *Mol. Cell. Biol.* 14:1450–1458.
- Kinoshita, K., J. Tashiro, S. Tomita, C.-G. Lee, and T. Honjo. 1998. Target specificity of immunoglobulin class switch recombination is not determined by the nucleotide sequences of S regions. *Immunity.* 9:849–858.
- Petry, K., G. Siebenkotten, R. Christine, K. Hein, and A. Radbruch. 1999. An extrachromosomal switch recombination substrate reveals kinetics and substrate requirements of switch recombination in primary murine B cells. *Int. Immunol.* 11:753–763.
- Stavnezer, J., S.P. Bradley, N. Rousseau, T. Pearson, A. Shanmugam, D.J. Waite, P.R. Rogers, and A.L. Kenter. 1999. Switch recombination in a transfected plasmid occurs preferentially in a B cell line that undergoes switch recombination of its chromosomal Ig heavy chain genes. *J. Immunol.* 163:2028–2040.
- Shanmugam, A., M.J. Shi, L. Yauch, J. Stavnezer, and A.L. Kenter. 2000. Evidence for class-specific factors in immunoglobulin isotype switching. *J. Exp. Med.* 191:1365–1380.
- Reaban, M.E., and J.A. Griffin. 1990. Induction of RNA-stabilized DNA conformers by transcription of an immunoglobulin switch region. *Nature*. 348:342–344.
- Reaban, M.E., J. Lebowitz, and J.A. Griffin. 1994. Transcription induces the formation of a stable RNA:DNA hybrid in the immunoglobulin alpha switch region. *J. Biol. Chem.* 269:21850–21857.
- Daniels, G.A., and M.R. Lieber. 1995. RNA:DNA complex formation upon transcription of immunoglobulin switch regions: implications for the mechanism and regulation of class switch recombination. *Nucleic Acids Res.* 23:5006–5011.
- Tian, M., and F.W. Alt. 2000. Transcription-induced cleavage of immunoglobulin switch regions by nucleotide excision repair nucleases in vitro. *J. Biol. Chem.* 275:24163–24172.
- Matsuoka, M., K. Yoshida, T. Maeda, S. Usuda, and H. Sakano. 1990. Switch circular DNA formed in cytokine-treated mouse splenocytes: evidence for intramolecular DNA deletion in immunoglobulin class switching. *Cell.* 62:135–142.
- von Schwedler, U., H.-M. Jack, and M. Wabl. 1990. Circular DNA is a product of immunoglobulin class switch rearrangement. *Nature*. 345:452–456.
- Iwasato, T., A. Shimizu, T. Honjo, and H. Yamagishi. 1990. Circular DNA is excised by immunoglobulin class switch recombination. *Cell.* 62:143–149.
- Lee, C., S. Kondo, and T. Honjo. 1998. Frequent but biased class switch recombination in the S μ flanking regions. *Curr. Biol.* 8:227–230.
- Chu, C.C., W.E. Paul, and E.E. Max. 1992. Quantitation of immunoglobulin μ - γ 1 heavy chain switch region recombination by a digestion-circularization polymerase chain reaction method. *Proc. Natl. Acad. Sci. USA.* 89:6978–6982.
- Schrader, C., W. Edelmann, R. Kucherlapati, and J. Stavnezer. 1999. Reduced isotype switching in splenic B cells from mice deficient in mismatch repair enzymes. *J. Exp. Med.* 190:323–330.
- Sigurdardottir, D., J. Sohn, J. Kass, and E. Selsing. 1995. Regulatory regions 3' of the immunoglobulin heavy chain

- intronic enhancer differentially affect expression of a heavy chain transgene in resting and activated B cells. *J. Immunol.* 154:2217–2225.
28. Ehrenstein, M.R., and M.S. Neuberger. 1999. Deficiency in Msh2 affects the efficiency and local sequence specificity of immunoglobulin class-switch recombination: parallels with somatic hypermutation. *EMBO (Eur. Mol. Biol. Organ.) J.* 18: 3484–3490.
 29. Cockerill, P.N., M.-H. Yuen, and W.T. Garrard. 1987. The enhancer of the immunoglobulin heavy chain locus is flanked by presumptive chromosomal loop anchorage elements. *J. Biol. Chem.* 262:5394–5397.
 30. Gram, H., G. Zenke, S. Geisse, B. Kleuser, and K. Bürki. 1992. High-level expression of a human immunoglobulin γ 1 transgene depends on switch region sequences. *Eur. J. Immunol.* 22:1185–1191.
 31. Ballantyne, J., D.L. Henry, and K.B. Marcu. 1997. Antibody class switch recombinase activity is B cell stage specific and functions stochastically in the absence of ‘targeted accessibility’ control [published erratum at 9:1773]. *Int Immunol* 9:963–974.
 32. Dunnick, W., and J. Stavnezer. 1990. Copy choice mechanism of immunoglobulin heavy-chain switch recombination. *Mol. Cell. Biol.* 10:397–400.
 33. Bottaro, A., F. Young, J. Chen, M. Serwe, F. Sablitzky, and F.W. Alt. 1998. Deletion of the IgH intronic enhancer and associated matrix-attachment regions decreases, but does not abolish, class switching at the mu locus. *Int. Immunol.* 10: 799–806.
 34. Hummel, M., J.K. Berry, and W. Dunnick. 1987. Switch region content of hybridomas: the two spleen cell Igh loci tend to rearrange to the same isotype. *J. Immunol.* 138:3539–3548.
 35. Winter, E., U. Krawinkel, and A. Radbruch. 1987. Directed Ig class switch recombination in activated murine B cells. *EMBO (Eur. Mol. Biol. Organ.) J.* 6:1663–1671.
 36. Davis, M.M., S.K. Kim, and L.E. Hood. 1980. DNA sequences mediating class switching in α -immunoglobulins. *Science.* 209:1360–1365.
 37. Radbruch, A., W. Muller, and K. Rajewsky. 1986. Class switch recombination is IgG1 specific on active and inactive IgH loci of IgG1-secreting B-cell blasts. *Proc. Natl. Acad. Sci. USA.* 83:3954–3957.
 38. Siebenkotten, G., C. Esser, M. Wabl, and A. Radbruch. 1992. The murine IgG1/IgE class switch program. *Eur. J. Immunol.* 22:1827–1834.
 39. Casellas, R., A. Nussenzweig, R. Wuerffel, R. Pelanda, A. Reichlin, H. Suh, X.F. Qin, E. Besmer, A. Kenter, K. Rajewsky, and M.C. Nussenzweig. 1998. Ku80 is required for immunoglobulin isotype switching. *EMBO (Eur. Mol. Biol. Organ.) J.* 17:2404–2411.
 40. Manis, J.P., Y. Gu, R. Lansford, E. Sonoda, R. Ferrini, L. Davidson, K. Rajewsky, and F.W. Alt. 1998. Ku70 is required for late B cell development and immunoglobulin heavy chain class switching. *J. Exp. Med.* 187:2081–2089.
 41. Rolink, A., F. Melchers, and J. Andersson. 1996. The SCID but not the RAG-2 gene product is required for S mu-S epsilon heavy chain class switching. *Immunity.* 5:319–330.
 42. Kinoshita, K., C.-G. Lee, J. Tashiro, M. Muramatsu, X.-C. Chen, K. Yoshikawa, and T. Honjo. 1999. Molecular mechanism of immunoglobulin class switch recombination. *Cold Spring Harbor Symp. Quant. Biol.* 64:217–226.
 43. Stavnezer-Nordgren, J., and S. Sirlin. 1986. Specificity of immunoglobulin heavy chain switch correlates with activity of germline heavy chain genes prior to switching. *EMBO (Eur. Mol. Biol. Organ.) J.* 5:95–102.
 44. Lorenz, M., S. Jung, and A. Radbruch. 1995. Switch transcripts in immunoglobulin class switching. *Science.* 267:1825–1828.
 45. Hein, K., G.O. Lorenz, G. Siebenkotten, K. Petry, R. Christine, and A. Radbruch. 1998. Processing of switch transcripts is required for targeting of antibody class switch recombination. *J. Exp. Med.* 188:2369–2374.
 46. Bottaro, A., R. Lansford, L. Xu, J. Zhang, P. Rothman, and F.W. Alt. 1994. S region transcription per se promotes basal IgE class switch recombination but additional factors regulate the efficiency of the process. *EMBO (Eur. Mol. Biol. Organ.) J.* 13:665–674.
 47. Barr, J., N.M. Pennell, and M.J. Shulman. 1996. Analysis of a hot spot for DNA insertion suggests a mechanism for Ig switch recombination. *J. Immunol.* 157:3430–3435.
 48. Dempsey, L.A., H. Sun, L.A. Hanakahi, and N. Maizels. 1999. G4 DNA binding by LR1 and its subunits, nucleolin and hnRNP D, A role for G-G pairing in immunoglobulin switch recombination. *J. Biol. Chem.* 274:1066–1071.
 49. Kinoshita, K., and T. Honjo. 2000. Unique and unprecedented recombination mechanisms in class switching. *Curr. Opin. Immunol.* 12:195–198.
 50. Mussmann, R., M. Courtet, J. Schwager, and L. Du Pasquier. 1997. Microsites for immunoglobulin switch recombination breakpoints from Xenopus to mammals. *Eur. J. Immunol.* 27:2610–2619.
 51. Kitao, H., H. Arakawa, K. Kuma, H. Yamagishi, N. Nakamura, S. Furusawa, H. Matsuda, M. Yasuda, S. Ekino, and A. Shimizu. 2000. Class switch recombination of the chicken IgH chain genes: implications for the primordial switch region repeats. *Int. Immunol.* 12:959–968.
 52. Marcu, K.B., J. Banerji, N.A. Penncavage, R. Lang, and N. Arnheim. 1980. 5′ flanking region of immunoglobulin heavy chain constant region genes displays length heterogeneity in germ lines of inbred mouse strains. *Cell.* 22:187–196.
 53. Katzenberg, D.R., and B.K. Birshtein. 1988. Sites of switch recombination in IgG2b- and IgG2a-producing hybridomas. *J. Immunol.* 140:3219–3227.
 54. Petrini, J., B. Shell, M. Hummel, and W. Dunnick. 1987. The immunoglobulin heavy chain switch: structural features of gamma 1 recombinant switch regions. *J. Immunol.* 138: 1940–1946.
 55. Iwasato, T., H. Arakawa, A. Shimizu, T. Honjo, and H. Yamagishi. 1992. Biased distribution of recombination sites within S regions upon immunoglobulin class switch recombination induced by transforming growth factor beta and lipopolysaccharide. *J. Exp. Med.* 175:1539–1546.
 56. Dunnick, W., T.H. Rabbitts, and C. Milstein. 1980. An immunoglobulin deletion mutant with implications for the heavy-chain switch and RNA splicing. *Nature.* 286:669–675.
 57. Yoshida, K., M. Matsuoka, S. Usuda, A. Mori, K. Ishizaka, and H. Sakano. 1990. Immunoglobulin switch circular DNA in the mouse infected with *Nippostrongylus brasiliensis*: evidence for successive class switching from mu to epsilon via gamma 1. *Proc. Natl. Acad. Sci. USA.* 87:7829–7833.
 58. Mowatt, M., and W. Dunnick. 1986. DNA sequence of murine γ 1 switch segment reveals novel structural elements. *J. Immunol.* 136:2674–2683.