

Takaki et al. Vol. 195, No. 2, January 21, 2002. Pages 151–160.

There was a typographical error in Table III. The third column said 3.6 ± 4.8^a , but it should be 33.6 ± 4.8^a . The corrected table appears below.**Table III.** B Cell Overproduction and Increased in $\text{Lin}^- \text{Sca-1}^+ \text{c-Kit}^+$ Hematopoietic Progenitors in $\text{lnk}^{-/-}$ Mice Is Partly Normalized by Reducing Expression of c-Kit

Genotype	Cell number	B220 ^{lo} IgM ⁻	B220 ^{lo} IgM ⁺	Li ⁻ Sca-1 ⁺ c-Kit ⁺		
		(10^7 cells/femur)	(%)	(%)	(%)	(% in Lin ⁻ cells)
<i>lnk</i> ^{+/+} <i>Kit</i> ^{+/+}	1.65 ± 0.16	19.5 ± 5.1	8.9 ± 3.6	($n = 12$)	0.45 ± 0.07	16.2 ± 2.2
<i>lnk</i> ^{+/+} <i>Kit</i> ^{W/+}	1.71 ± 0.19	18.2 ± 4.6	8.0 ± 3.1	($n = 10$)	0.38 ± 0.06	14.9 ± 1.7
<i>lnk</i> ^{-/-} <i>Kit</i> ^{+/+}	2.03 ± 0.37^a	33.6 ± 4.8^a	14.1 ± 2.1^a	($n = 11$)	0.70 ± 0.06^a	26.1 ± 2.1^a
<i>lnk</i> ^{-/-} <i>Kit</i> ^{W/+}	1.91 ± 0.28	28.9 ± 3.9^b	11.8 ± 1.7^b	($n = 8$)	0.56 ± 0.13	21.3 ± 1.4^b

Bone marrow cells obtained from 8–10-wk-old mice were analyzed, and mean cell number (\pm SD), mean percentage (\pm SD) of the indicated cell fractions and the number of mice analyzed are shown.

^a $P < 0.05$ compared to *lnk*^{+/+}*Kit*^{+/+}.

^b $P < 0.05$ compared to *lnk*^{-/-}*Kit*^{+/+}.