

Perini et al. Vol. 195, No. 7, April 1, 2002. Pages 907–918.

Due to a production error, the column heading of BETrka was missing. The corrected table appears below.

**Table I.** *Cell Death Induced by A $\beta$  Peptides*

	BENTR-free		BEp75		BETrka	BEp75Trka
	24 h	48 h	24 h	48 h	24 h	24 h
Controls	5.9 $\pm$ 2.6 <sup>a</sup> (18)	7.8 $\pm$ 2.6 (18)	8.2 $\pm$ 2.2 (62)	9.2 $\pm$ 2.4 (20)	5.3 $\pm$ 2.5 (3)	10.3 $\pm$ 2.1 (3)
A $\beta$ (25–35)	5.2 $\pm$ 2.6 (12)	6.2 $\pm$ 2.8 (12)	29.7 $\pm$ 4.5 <sup>b</sup> (65)	34.0 $\pm$ 4.8 <sup>b</sup> (14)	7.1 $\pm$ 3.1 (5)	28.5 $\pm$ 6.3 <sup>b</sup> (5)
A $\beta$ (1–40)	5.6 $\pm$ 2.3 (5)	5.5 $\pm$ 2.8 (4)	27.6 $\pm$ 2.3 <sup>b</sup> (5)	33.2 $\pm$ 6.2 <sup>b</sup> (4)		
A $\beta$ (1–42)	5.6 $\pm$ 1.6 (5)	7.3 $\pm$ 1.8 (4)	30.0 $\pm$ 2.5 <sup>b</sup> (3)	30.9 $\pm$ 2.2 <sup>b</sup> (4)		

Cells were treated with A $\beta$ (25–35) (20  $\mu$ M), A $\beta$ (1–40) (5.0  $\mu$ M), or A $\beta$ (1–42) (5.0  $\mu$ M) and cell death was assessed by epifluorescence microscopy after 24 and 48 h.

<sup>a</sup>The values express percentages of cell death and are means  $\pm$ SD of the experiments indicated within brackets.

<sup>b</sup> $P < 0.001$  with respect to the controls of the corresponding time point.