

A B Cell Receptor with Two Ig α Cytoplasmic Domains Supports Development of Mature But Anergic B Cells

Amy Reichlin,¹ Anna Gazumyan,¹ Hitoshi Nagaoka,³ Kathrin H. Kirsch,⁴ Manfred Kraus,⁵ Klaus Rajewsky,⁵ and Michel C. Nussenzweig^{1,2}

¹Laboratory of Molecular Immunology, ²Howard Hughes Medical Institute, The Rockefeller University, New York, NY 10021

³Department of Medical Chemistry and Molecular Biology, Graduate School of Medicine, Kyoto University, Yoshida Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan

⁴Department of Biochemistry, Boston University School of Medicine, Boston, MA 02118

⁵Center for Blood Research, Harvard Medical School, Boston, MA 02115

Abstract

B cell receptor (BCR) signaling is mediated through immunoglobulin (Ig) α and Ig β a membrane-bound heterodimer. Ig α and Ig β are redundant in their ability to support early B cell development, but their roles in mature B cells have not been defined. To examine the function of Ig α –Ig β in mature B cells in vivo we exchanged the cytoplasmic domain of Ig α for the cytoplasmic domain of Ig β by gene targeting (Ig β_c → α_c mice). Ig β_c → α_c B cells had lower levels of surface IgM and higher levels of BCR internalization than wild-type B cells. The mutant B cells were able to complete all stages of development and were long lived, but failed to differentiate into B1a cells. In addition, Ig β_c → α_c B cells showed decreased proliferative and Ca²⁺ responses to BCR stimulation in vitro, and were anergic to T-independent and -dependent antigens in vivo.

Key words: B cell receptor • immunoglobulin • signal transduction • anergy • B cell development

Introduction

Signaling by membrane Ig (mIg) regulates antigen-independent B cell development, antigen-dependent adaptive immune responses, and B cell survival (1, 2). Signals are transmitted from mIg to the cytoplasm through Ig α and Ig β , which form a disulfide-linked membrane-bound heterodimer that is noncovalently associated with mIg through polar residues in the plane of the cell membrane (3–7). Ig α and Ig β contain tyrosine residues that are imbedded in immunoreceptor tyrosine-based activation motifs (ITAMs) that are essential for B cell receptor (BCR) signaling (6–11). Upon BCR cross-linking these tyrosine residues recruit and serve as substrates for src and syk family kinases (12–21). Deletion of src kinases blk, fyn, and lyn results in a severe, but incomplete, block at the preB cell stage of development and the combination of syk and zap70 mutation abrogates B cell development (15, 22–24).

Although the cytoplasmic domains of both Ig α and Ig β contain essential ITAMs they are otherwise nonhomologous and they display different signaling properties in transfected cell lines (9, 16). Only Ig α has non-ITAM tyrosines that

bind SLP-65, which in turn recruit Grb2, Nck, Vav, Btk, and PLC- γ 2 (20, 25–27). These differences, and the observation that Ig α and Ig β cytoplasmic domains bind to distinct sets of kinases in cytoplasmic extracts, led to the suggestion that the two signal transducers have different functions in vivo (6, 7, 9, 11, 16, 28–30). However, studies in transgenic and gene-targeted mice showed that Ig α and Ig β have nearly equivalent function for all aspects of early B cell development (31–35). Either the Ig α or the Ig β cytoplasmic domain is sufficient to induce early B cell development in transgenic mice by a mechanism that requires ITAM tyrosine phosphorylation (32, 33). In addition, selective deletion of Ig α or Ig β cytoplasmic domains by gene targeting (Ig α Δ C and Ig β Δ C mice, respectively) results in a block in B cell development that is more severe in early B cells in Ig α Δ C than in Ig β Δ C mice (34, 35). Both Ig α Δ C and Ig β Δ C mice display a near absence of mature B cells in the periphery (34–36).

To determine whether the cytoplasmic domains of Ig α and Ig β have distinct signaling activities in mature B cells

Address correspondence to Michel C. Nussenzweig, 1230 York Ave., Box 220, New York, NY 10021. Phone: (212) 327-8067; Fax: (212) 327-8370; email: nussen@mail.rockefeller.edu

Abbreviations used in this paper: BCR, B cell receptor; BrdU, bromodeoxyuridine; CFDA,SE, 5-carboxyfluorescein diacetate succinimidyl ester; CGG, chicken gamma globulin; IP, immunoprecipitation; ITAM, immunoreceptor tyrosine activation motif; mIg, membrane Ig; NP, nitrophenol.

in vivo we replaced the cytoplasmic tail of Ig β with the cytoplasmic domain of the Ig α and produced a BCR that contains a homodimer of Ig α tails instead of the normal physiologic heterodimer of Ig α and Ig β . Here we report that all B cell subsets, with the exception of B1 cells, developed normally in Ig $\beta_c \rightarrow \alpha_c$ mice, but mature B cells had reduced cell surface BCR and were anergic.

Materials and Methods

Mice. Ig $\beta_c \rightarrow \alpha_c$ mice carry an Ig β gene in which the cytoplasmic tail of Ig β from positions 183–228 was replaced with the cytoplasmic tail of Ig α from positions 161–220 by gene targeting in 129/Sv embryonic stem cells. A cDNA containing the coding sequence for Ig α from positions 161–220 followed by a stop codon, TGA, was inserted after amino acid position 182 in the Ig β gene. A unique HindIII site was placed into the targeting vector as indicated (see Fig. 1 A). The long arm of the targeting construct was 16 kb, the short arm was 6 kb, a LoxP-flanked neomycin-resistance gene was used for positive selection, and a diphtheria toxin gene for negative selection (37). Homologous integrants were confirmed by Southern blotting after digestion with HindIII. The genomic fragment used as a probe for Southern blotting was generated by PCR with the primers 5' GGA TTC GAA TGG TGA ATG TTG G 3' and 5' AGG CTC TAGG CTC AGT GAA GGC AG 3'. Screening by PCR was performed using the primers, 5' AGC AGG AAG ATG GGC ACA ATG ATG AAG 3' and 5' TGG TAT CTA CTT CTG CAA GCA GAA ATG 3': a 250-bp product represented the mutant allele and a 186-bp product corresponded to the wild-type allele. The rate of homologous recombination was 1:35. Neomycin primers 5' ATG ATT GAA CAA GAT GGA TTG CAC 3' and 5' TCG TCC AGA TCA TCC TGA TCG AC 3' were used to confirm neomycin gene deletion. Heterozygous Ig $\beta_c \rightarrow \alpha_c$ mice were backcrossed to C57BL/6 for two generations after Cre deletion before intercrossing. All mice were maintained under specific pathogen-free conditions. All experiments shown were performed with homozygous mice.

B Cell Purification and Western Blotting. B cells were purified by negative selection from spleen using anti-CD43 magnetic microbeads (Miltenyi Biotec). For cross-linking experiments B cells were resuspended at 10^8 /ml in RPMI with 1% FBS and were cross-linked with 20 μ g/ml goat anti-mouse IgM, μ chain specific F(ab')₂ fragments (Jackson ImmunoResearch) for the indicated times at 37°C. Cells were lysed in TX-100 buffer (50 mM Tris, pH 7.5, 137 mM NaCl, 2 mM EDTA, 10% glycerol, 1% Triton X-100, and inhibitors) or 50 mM Tris/HCl, pH 7.7, 150 mM NaCl, 1% NP-40, 1 mM EDTA, 1 mM EGTA, 0.1% deoxycholic acid, 10% glycerol, 1 mM PMSF, 0.2 mM Na₃VO₄, and protease inhibitors) for Western blotting and in RIPA buffer (50 mM Tris/HCl, pH 7.7, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.1% deoxycholic acid, 1 mM PMSF, 0.2 mM Na₃VO₄, and protease inhibitors) for immunoprecipitation (IP). Lysates were incubated for 15 min on ice and cellular debris were sedimented at 14,000 rpm for 15 min at 4°C. Lysates were resolved on NuPAGE 4–12% Bis-Tris gels (Invitrogen) and immunoblotted on Immobilon polyvinylidene fluoride membranes (Millipore). For IP, cellular lysates were precleared with protein A-Sepharose (Amersham Biosciences), incubated for 2 h with primary antibodies, and captured with protein A-Sepharose. Visualization was with protein A horse radish peroxidase (Amersham Biosciences), donkey anti-rabbit IgG (Jackson ImmunoResearch

Laboratories) or goat and mouse IgG (Jackson ImmunoResearch Laboratories) and Western Plus chemiluminescent reagent (NEN Life Science Products). Antibodies used for IP and Western blotting were anti-Ig α and anti-Ig β (38), 4G10 (Upstate Biotechnology), anti-Syk (Santa Cruz Biotechnology, Inc.), and Phospho-Syk (Cell Signaling Technologies).

Flow Cytometry. Single cell suspensions from bone marrow, spleen, and peritoneal cavity were stained with FITC, PE, APC, and biotin-conjugated mAbs visualized with Streptavidin red 613 (GIBCO BRL) or streptavidin PerCp (Becton Dickinson). mAbs were anti-CD43, anti-IgM, anti-B220, anti-CD25, anti-IgD, anti-CD19, anti-CD5, anti-CD21, anti-HSA, and anti-CD23 (BD Biosciences). Data collection was performed on a FACSCalibur™ and analyzed using CellQuest™ software (Becton Dickinson).

In Vitro B Cell Cultures and Proliferation Assays. Purified spleen B cells were cultured in RPMI 1640 (GIBCO-BRL) supplemented with 10% FCS, 2 mM L-glutamine (GIBCO-BRL), 100 IU/ml penicillin/streptomycin (GIBCO-BRL), and 100 μ M 2-mercaptoethanol (Sigma-Aldrich). Cells were stimulated with CD40L at a concentration of 1:100 (CD8-gp39 fusion protein derived from insect cells; a gift from Dr. Randolph Noelle, Dartmouth Medical School, Hanover, NH; reference 39), or 25 μ g/ml LPS (Sigma-Aldrich), or 10 μ g/ml goat anti-mouse F(ab')₂ IgM (Jackson ImmunoResearch), or 5 μ g/ml RP105 (a gift from A. Tarahkovsky, The Rockefeller University), and 1:2,000 CpG (MWG Biotech).

For thymidine incorporation assays 3×10^5 cells were cultured in triplicate for 48 h in 96-well flat bottom plates and pulsed with 1 μ Ci/ml of [³H]thymidine for 6 h. For labeling with 5-carboxy-fluorescein diacetate succinimidyl ester (CFDA,SE; Molecular Probes) 10^7 cells were incubated at 37°C for 10 min in 1 ml RPMI 1640 supplemented with 5 μ M CFDA,SE. Cultures were stained with PE-CD19 (BD Biosciences) and analyzed on a FACSCalibur™ using CellQuest™ software.

BCR Internalization Assays. Single cell suspensions were equilibrated in ice water baths and stained with biotin-SP goat anti-mouse IgM Fab' fragment (μ specific; Jackson ImmunoResearch). An aliquot of cells was fixed with PBS containing 0.5% paraformaldehyde for the time 0 (T₀) cell surface expression value. The remaining cells were incubated at 37°C for the indicated times (T_n). Reactions were terminated with ice-cold PBS containing 0.5% paraformaldehyde and cells stained with PE-B220 (BD Biosciences) and Streptavidin red 670 (GIBCO BRL). Percent internalization was calculated by the formula % sIgM(T₀) – % sIgM(T_n)/% IgM(T₀) \times 100.

Bromodeoxyuridine (BrdU) Labeling. 8–12-wk-old Ig $\beta_c \rightarrow \alpha_c$ and age- and sex-matched control (B6/129F1) mice received BrdU (Sigma-Aldrich) in their drinking water at a concentration of 0.5 mg/ml. The number of B220⁺ bone marrow and spleen cells was not significantly altered during the 63-d time course. Staining was with anti-BrdU mAb using the BrdU flow kit™ (BD Biosciences) and data was collected using a FACSCalibur™ (Becton Dickinson). Fitted curves were drawn using Microsoft Excel software. Approximate half-life was equal to time point where 50% of cells stained positively for BrdU.

Ca²⁺ Flux. Spleen cell suspensions from C57BL/6 mice and Ig $\beta_c \rightarrow \alpha_c$ mice were adjusted to 5×10^6 /ml in PBS containing 1% FCS, 1 mM CaCl₂, 1 mM MgCl₂ (loading buffer), and incubated with 1.5 μ M Indo-1-AM (Molecular Probes) for 30 min at 37°C. Cells were stained with PE-anti-B220 and Fab' FITC goat anti-mouse IgM (Jackson ImmunoResearch Laboratories) to determine gating. Calcium flux was measured by fluorescence emission ratios of Indo-1-AM on a dual laser FACS Advantage™ (Bec-

ton Dickinson) at 395/510 nm on B220⁺IgM⁺ cells. A baseline reading was collected for 60 s before cross-linking the BCR with 30 $\mu\text{g/ml}$ goat anti-mouse IgM F(ab')₂ fragments (Southern Biotechnology Associates, Inc.).

Immunization. 6–8-wk-old Ig $\beta_c \rightarrow \alpha_c$ and B6/129F1 controls were immunized intraperitoneally with either 50 μg alum precipitated 4-hydroxy-3-nitrophenylacetyl coupled to chicken gamma globulin (NP-CGG; Biosearch Technologies) or 50 μg NP-Ficoll in PBS. Blood was collected from the tail vein of each mouse before immunization and then again at days 7, 14, 21, and 31 postimmunization for the mice immunized with NP-CGG and at days 7, 14, 21, and 28 post immunization for the NP-Ficoll group. NP-specific IgM and IgG levels were measured by ELISA using NP₁₆BSA as a capture and developed with goat anti-IgM or anti-IgG coupled to horse radish peroxidase (Southern Biotechnology Associates, Inc.). Immunoabsorbance was read at 415 nm and titers were calculated relative to control sera from un-immunized mice. Three mice were used in each group.


Results

Ig $\beta_c \rightarrow \alpha_c$ Mice. To determine whether Ig α and Ig β have unique or redundant functions in mature B cells we replaced the cytoplasmic tail of Ig β with the cytoplasmic tail of Ig α by gene targeting (Fig. 1 A, Ig $\beta_c \rightarrow \alpha_c$). Ig $\beta_c \rightarrow \alpha_c$ protein expression in B cells was confirmed by Western

blotting of purified B cell lysates using antibodies specific for the cytoplasmic domains of Ig α and Ig β (38, 40). Control lysates showed the presence of both Ig α and Ig β (Fig. 1 B). In contrast, there was no wild-type Ig β in the lysates from Ig $\beta_c \rightarrow \alpha_c$ B cells. Instead we found a new species reactive with the anti-Ig α antibody that had the predicted mobility of the Ig $\beta_c \rightarrow \alpha_c$ protein and no additional Ig β species (Fig. 1 B). We conclude that B cells from Ig $\beta_c \rightarrow \alpha_c$ mice do in fact express only the chimeric Ig β protein with the cytoplasmic tail of Ig α .

B Cell Development in Ig $\beta_c \rightarrow \alpha_c$ Mice. To examine the effects of the Ig α cytoplasmic tail on development of B lineage cells we compared B cells from Ig $\beta_c \rightarrow \alpha_c$ mice (BCR with two Ig α tails) with B cells from Ig $\beta\Delta\text{C}$ mice (BCR with a single Ig α tail; reference 35) and wild-type controls (Ig α -Ig β).

Like Ig $\beta\Delta\text{C}$, Ig $\beta_c \rightarrow \alpha_c$ mice showed an increase in IgM⁻B220^{low}CD43⁺ pro-B cells, normal levels of IgM⁻B220^{low}CD25⁺ preB cells and a decrease in the number of immature IgM⁺B220^{low}IgD^{-/low} B cells. In addition, immature Ig $\beta_c \rightarrow \alpha_c$ B cells displayed decreased surface IgM and IgD expression (Figs. 1 C and 2 A). Lower IgM levels were also found in Ig $\beta\Delta\text{C}$ B cells but these cells were severely impaired in B cell development (35). Ig $\beta_c \rightarrow \alpha_c$ differed from Ig $\beta\Delta\text{C}$ in that BCRs with two Ig α tails sup-






Figure 2. B cells in bone marrow and spleen in wild-type and $Ig\beta_c \rightarrow \alpha_c$ mice. Numbers show percentages of bone marrow or splenic lymphocytes. (A) Bone marrow. Pro B cells were $CD43^+B220^+$, pre BII cells were $CD25^+IgM^-$, immature B cells were IgM^+IgD^{lo} , and mature B cells were IgM^+IgD^{hi} . (B) Spleen. T1, T2, and mature B cells were defined as in Fig. 1. Marginal zone B cells were $CD21^+CD23^-$ and follicular B cells were $CD21^+CD23^+$. Filled diamonds represent wild-type and open squares represent $Ig\beta_c \rightarrow \alpha_c$ mice. Each symbol shows an individual mouse. Bars represent the means of all mice in a group.

ported development of mature recirculating IgM^+B220^{high} B cells whereas a single $Ig\alpha$ tail in $Ig\beta\Delta C$ mice did not (Figs. 1 C and Fig. 2 A; reference 35), but the number of mature recirculating IgM^+B220^{high} B cells in the bone marrow in $Ig\beta_c \rightarrow \alpha_c$ was somewhat lower than wild-type controls (12% Wt vs. 4% $Ig\beta_c \rightarrow \alpha_c$; Figs. 1 C and Fig. 2 A). The total number of cells in spleens of $Ig\beta_c \rightarrow \alpha_c$ mice was similar to wild type (Table I). However, we found a 30% overall decrease in $B220^+IgM^+$ B cells with a small relative increase in marginal zone B cells (Figs. 1 D and Fig. 2 B). In the peritoneal cavity $Ig\beta_c \rightarrow \alpha_c$ mice showed a sevenfold relative decrease in the number of B1a cells but normal number of B1b and B2 cells (Fig. 1 D). Heterozygous $Ig\beta_c \rightarrow \alpha_c$ mice were no different than wild-type controls in B cell development or any of the signaling assays discussed below. We conclude that BCRs with two $Ig\alpha$ tails support development of all B cell subtypes except B1a cells.

Decreased Levels of Cell Surface BCR. Mutation of the $Ig\alpha$ ITAM ($Ig\alpha^{FF/FF}$ mice) increased the level of cell surface

Table I. Total Numbers of Nucleated Cells in Bone Marrow and Spleens of 8–12-wk-old $Ig\beta_c \rightarrow \alpha_c$ Mice and Wild-type Controls

	Bone marrow (n = 9)	Spleen (n = 17)
Wild type	27.4×10^6 SD ± 6.80	71.5×10^6 SD ± 21.6
$Ig\beta_c \rightarrow \alpha_c$	24.2×10^6 SD ± 7.26	61.5×10^6 SD ± 19.9

n, number of mice tested.

BCR expression suggesting that phosphorylation of ITAM tyrosines in the cytoplasmic tail of $Ig\alpha$ might regulate cell surface BCR levels (41). Consistent with this idea B cells with two cytoplasmic $Ig\alpha$ chains showed decreased levels of cell surface BCR (Fig. 3 A). The decrease in surface BCR expression on $Ig\beta_c \rightarrow \alpha_c$ B cells was not due to a change in steady-state levels of $Ig\mu$ mRNA or in $Ig\mu$ protein synthesis as measured by RNase protection and [35 S]methionine metabolic labeling experiments (unpublished data). However, the decreased surface BCR level was associated with increased BCR internalization as measured with a biotinylated Fab' anti-IgM (Fig. 3 B). $Ig\beta_c \rightarrow \alpha_c$ B cells showed increased and more rapid receptor internalization than wild-type controls. We conclude that lower levels of cell surface BCR in $Ig\beta_c \rightarrow \alpha_c$ mice are associated with increased BCR internalization.

To determine whether decreased cell surface BCR expression in $Ig\beta_c \rightarrow \alpha_c$ mice is the result of BCR signaling we bred $Ig\beta_c \rightarrow \alpha_c$ and $Ig\alpha^{FF/FF}$ mice ($Ig\beta_c \rightarrow \alpha_c/Ig\alpha^{FF/FF}$ mice; reference 41). $Ig\beta_c \rightarrow \alpha_c/Ig\alpha^{FF/FF}$ B cells differ from $Ig\beta_c \rightarrow \alpha_c$ B cells in that only one of the two cytoplasmic $Ig\alpha$ tails in $Ig\beta_c \rightarrow \alpha_c/Ig\alpha^{FF/FF}$ B cells carries a functional ITAM. We found that B cells with two functional $Ig\alpha$ ITAMs ($Ig\beta_c \rightarrow \alpha_c$) had the lowest BCR levels, those with no functional $Ig\alpha$ ITAMs ($Ig\alpha^{FF/FF}$) had the highest levels and B cells with one functional $Ig\alpha$ ITAM ($Ig\beta_c \rightarrow \alpha_c/Ig\alpha^{FF/FF}$) were intermediate (Fig. 3 A). We conclude that an active feedback signaling mechanism mediated by tyrosine phosphorylation of the $Ig\alpha$ cytoplasmic domain is responsible for the low levels of BCR on $Ig\beta_c \rightarrow \alpha_c$ B cells.

Half-Life of $Ig\beta_c \rightarrow \alpha_c$ B Cells In Vivo. $Ig\beta_c \rightarrow \alpha_c$ mice displayed a small decrease in the numbers of mature B cells in spleen and a more significant decrease in immature B cells in the bone marrow (Fig. 2 B). B cell half-life is influenced by the rate of B cell influx and is increased in the absence of bone marrow B cell production (42). To determine the life span of $Ig\beta_c \rightarrow \alpha_c$ B cells we performed continuous labeling experiments using BrdU. In these experiments the half-life of cells in any compartment is the time taken to label 50% of the population. We found an increase in the half-life of $Ig\beta_c \rightarrow \alpha_c$ B cells in spleen (Fig. 4 A, 58.3 d vs. 23.4 d for the wild type). We conclude that $Ig\beta_c \rightarrow \alpha_c$ B cells have a significant defect in passing early B cell checkpoints in the bone marrow but once established in the periphery these cells have a longer life span than wild-type B cells.





Figure 3. Cell surface BCR expression. (A) IgM and IgD surface expression on B cells from bone marrow and spleen of wild-type, $Ig\beta_c \rightarrow \alpha_c$, $Ig\beta_c \rightarrow \alpha_c / Ig\alpha^{FF/FF}$, and $Ig\alpha^{FF/FF}$ mice. Splenic B cells were purified by negative selection before staining. (B) IgM internalization assessed by flow cytometry. Percent internalization was calculated by the formula $\% \text{ sIgM}(T_n) - \% \text{ sIgM}(T_0) / \% \text{ IgM}(T_0) \times 100$. The graph shows three separate experiments. Wild type (filled diamonds) and $Ig\beta_c \rightarrow \alpha_c$ (filled squares).

$Ig\beta_c \rightarrow \alpha_c$ B Cells Are Relatively Unresponsive to T Cell-dependent and -independent Antigens In Vivo. To determine whether $Ig\beta_c \rightarrow \alpha_c$ B cells respond to antigen in vivo we immunized mice with nitrophenol-28 coupled to NP-CGG, a T cell-dependent antigen or nitrophenol-59 coupled to ficoll (NP-Ficoll) a T cell-independent antigen.

$Ig\beta_c \rightarrow \alpha_c$ mice showed a three- to fivefold reduction in specific IgM and IgG responses to T-dependent antigens when compared with wild-type controls and they were unable to respond to NP-Ficoll (Fig. 5). We conclude that despite their longevity $Ig\beta_c \rightarrow \alpha_c$ B cells are relatively anergic to stimulation with antigen in vivo.




Figure 4. (A) B cell lifespan in bone marrow and spleen in wild-type and $Ig\beta_c \rightarrow \alpha_c$ mice measured by BrdU. Diamonds show wild-type B cells and circles show the $Ig\beta_c \rightarrow \alpha_c$ B cells. Immature B cells (bone marrow) and mature B cells (spleen) were defined as in previous figures. Half-life was the time when 50% of the cells were BrdU labeled. Immature B cells were as in previous figures. Spleen immature B cells were $IgM^+ IgD^{lo}$, mature B cells were $IgM^{lo} IgD^{hi}$, and T cells were $CD3^+$.




Figure 5. Antibody responses in $Ig\beta_c \rightarrow \alpha_c$ mice. Graphs show anti-NP IgM and IgG responses measured by ELISA on days 7, 14, 21, and 28 (for NP-CGG) and days 7, 14, 21, and 31 for (NP-Ficoll) after immunization with T-dependent (NP-CGG) or -independent (NP-Ficoll) antigens. The filled diamonds and the open squares represent serum titers in individual wild-type and $Ig\beta_c \rightarrow \alpha_c$ mice, respectively. The ordinate shows OD 415 nm relative to nonimmunized controls. The line shows the means for each group at a given time point. There were three mice in each group.

Proliferation Responses In Vitro. To determine whether the inability to respond to antigen reflects a B cell autonomous defect we stimulated $Ig\beta_c \rightarrow \alpha_c$ B cells with mitogens in vitro. Purified B cells were labeled with CFDA,SE and stimulated with anti-IgM, CD40L, RP105, CpG, or LPS (Fig. 6, A and E). $Ig\beta_c \rightarrow \alpha_c$ B cells responded to anti-IgM stimulation by up-regulating CD69 and CD86 expression (Fig. 6 B). However, $Ig\beta_c \rightarrow \alpha_c$ B cells failed to proliferate in response to BCR cross-linking. Only a small number of the B cells stimulated with anti-IgM diluted CFSE; thymidine incorporation was also severely decreased compared with the wild-type control (Fig. 6, A and C). To determine whether the lack of responsiveness to BCR stimulation was due to a signaling defect as opposed to a problem with receptor assembly we measured responses to anti-IgM by $Ig\beta_c \rightarrow \alpha_c / Ig\alpha^{FF/FF}$ B cells, which carry ITAM tyrosine mutations in one of the two $Ig\alpha$ tails. We found that mutating the ITAM tyrosines in $Ig\alpha$ partially rescued the unresponsive phenotype in $Ig\beta_c \rightarrow \alpha_c / Ig\alpha^{FF/FF}$ B cells (Fig. 6 A). Whereas $Ig\beta_c \rightarrow \alpha_c$ B cells fail to proliferate in response to anti-IgM treatment, $Ig\beta_c \rightarrow \alpha_c / Ig\alpha^{FF/FF}$ B cells divided, although the average number of divisions was somewhat lower than wild-type or $Ig\alpha^{FF/FF}$ B cells (Fig. 6 A). Thus, the lack of responsiveness found in $Ig\beta_c \rightarrow \alpha_c$ B cells is at least in part due to a process that requires $Ig\alpha$ ITAM tyrosine phosphorylation.

$Ig\beta_c \rightarrow \alpha_c$ showed normal levels of cell surface CD40L and RP105, but diminished responses to CD40L, RP105, CpG, and LPS stimulation (Fig. 6, C–E). There was less cell division, as measured by CFSE dye dilution, in response to CD40L and RP105, and fewer cells were induced to divide by LPS or CpG (Fig. 6 E). These results were confirmed by measuring [3 H]thymidine uptake in response to the same stimuli (Fig. 6 C). These differences

were not due to a change in the number of cells undergoing apoptosis (Fig. 6; and unpublished data). We conclude that $Ig\beta_c \rightarrow \alpha_c$ B cells show modestly decreased responses to a broad range of mitogenic stimuli in vitro.

To determine whether proximal receptor signaling is decreased in $Ig\beta_c \rightarrow \alpha_c$ B cells we measured Ca^{2+} flux responses to BCR cross-linking. We found that $Ig\beta_c \rightarrow \alpha_c$ B cells showed impaired Ca^{2+} responses compared with wild-type controls (Fig. 6 F). To further characterize the proximal BCR signaling in $Ig\beta_c \rightarrow \alpha_c$ mice we examined total phosphorylation, as well as $Ig\alpha$ and Syk phosphorylation after BCR cross-linking (12–18). We found an overall decrease in total phosphorylation consistent with the lower levels of Ca^{2+} signaling in $Ig\beta_c \rightarrow \alpha_c$ B cells (Fig. 7 A). In addition, $Ig\beta_c \rightarrow \alpha_c$ B cells showed little $Ig\alpha$ phosphorylation in response to BCR cross-linking (Fig. 7 B). Finally, phosphorylation of Syk was substantially reduced in $Ig\beta_c \rightarrow \alpha_c$ mice when compared with wild-type controls (Fig. 7 C). We conclude that decreased surface BCR levels on $Ig\beta_c \rightarrow \alpha_c$ B cells are associated with decreased intensity of proximal signaling.

Discussion

Our experiments extend previous work by examining the function of $Ig\alpha$ and $Ig\beta$ in mature B cells in vivo. We show that a BCR containing a signaling module composed of a homodimer of $Ig\alpha$ cytoplasmic tails can support the development of all currently defined subgroups of B cells with the exception of B1a cells and therefore $Ig\alpha$ and $Ig\beta$ are redundant for all stages of B cell development. In contrast, $Ig\alpha$ and $Ig\beta$ have distinct functions in regulating BCR surface expression and setting thresholds for mature B cell activation.





Figure 7. Protein tyrosine phosphorylation. Purified B cells were stimulated with 10 μ g/ml goat anti-mouse F(ab')₂ for the indicated times. (A) Tyrosine phosphorylation of total cellular proteins (5×10^6 cells/lane) detected by antiphosphotyrosine immunoblotting with the mAb 4G10. (B) Tyrosine phosphorylation in $Ig\alpha$ immunoprecipitates. $Ig\alpha$ immunoprecipitates were immunoblotted with mAb 4G10 to detect tyrosine phosphorylation. The lower panel shows the same blot stripped and reprobed with $Ig\alpha$ antibody. (C) Syk tyrosine phosphorylation. Anti-Syk immunoprecipitates were immunoblotted with mAb 4G10 to detect tyrosine phosphorylation. The lower panel shows the same blot stripped and reprobed with anti-Syk antibody.

Ig α and Ig β and B Cell Development. Early B cell development requires preBCR signaling and either $Ig\alpha$ or $Ig\beta$ cytoplasmic domains are sufficient to satisfy this requirement (32, 33). Signaling through BCR ITAM tyrosines was required for early B cell development but the BCR ITAMs were redundant and a single functional $Ig\beta$ ITAM was sufficient to reconstitute development (32, 33, 41, 43). Transgenic Fc chimeras fused to the cytoplasmic tails of $Ig\alpha$ or $Ig\beta$ differed from $Ig\mu$ chimeras in that they appeared to support mature B cell development, but the distribution of different B cell subpopulations was not examined and the Fc chimeras were not antigen receptors so B cell function could not be studied (32, 33).

Gene targeting confirmed the transgenic experiments and showed that BCRs lacking either the cytoplasmic domain of $Ig\alpha$ or $Ig\beta$ ($Ig\alpha\Delta C$ and $Ig\beta\Delta C$) were sufficient for B cells to progress to the immature stage of development but not the mature B cell stage (34, 35). $Ig\alpha\Delta C$ and $Ig\beta\Delta C$ differed in that $Ig\beta\Delta C$ progressed to the immature stage of development nearly normally whereas $Ig\alpha\Delta C$ B cells showed a partial block in the transition between preB and immature stage of development (34, 35). These findings

were reminiscent of the observation that thymocyte development is dependent on the number of T cell receptor ζ ITAMs (44–46). Single ITAM-containing BCRs in $Ig\alpha\Delta C$ and $Ig\beta\Delta C$ mice were simply not sufficient for complete B cell maturation (34, 35).

In vitro experiments with isolated $Ig\alpha$ and $Ig\beta$ cytoplasmic domains suggested that the two transducers bind to different but overlapping sets of kinases (16). However, transfection experiments with chimeric receptors and transgenic experiments failed to show qualitative differences in early B cell development or signaling stimulated by $Ig\alpha$ or $Ig\beta$ (6, 7, 9, 11, 28–35). Mature B cells are therefore unique in their requirement for both $Ig\alpha$ and $Ig\beta$ for normal levels of BCR expression and physiologic responses to BCR cross-linking.

As both $Ig\alpha$ and $Ig\beta$ have a single ITAM, the physiologic differences in mature B cell signaling could be due to the non-ITAM residues. Among the non-ITAM residues in $Ig\alpha$ implicated in signaling are two non-ITAM tyrosines, 176 and 204, that bind the adaptor molecule BLNK (20, 21, 26, 47, 48), which links the BCR with phospholipase $C\gamma$, Vav, Grb2, Syk, Btk, and HPK1 (20, 25–27). Mice that are BLNK deficient show a block in early B cell development, decreased numbers of peripheral B cells, and defective B cell activation (20, 21, 25, 27, 49–53). BLNK-deficient mice also show an accumulation of mature B cells whose phenotype is $IgM^{high}IgD^{low}$ and increased expression of surface preBCR (54, 55). These features would lend credence to the idea that BCRs with two $Ig\alpha$ cytoplasmic domains in $Ig\beta_c \rightarrow \alpha_c$ B cells may simply recruit abnormally high levels of BLNK and its signaling partners, however, we find lower levels of BLNK phosphorylation in $Ig\beta_c \rightarrow \alpha_c$ B cells (unpublished data) and therefore the two non-ITAM tyrosines in $Ig\alpha$ that recruit BLNK are not likely to be responsible for the difference in signaling by $Ig\alpha$ and $Ig\beta$ in mature B cells in vivo.

$Ig\beta_c \rightarrow \alpha_c$ mice show a small relative enlargement of the MZ compartment but a near absence of B1 cells. Several signaling molecules have been shown to be essential for normal MZ B cell development including CD19, *ltk*, *pyk2*, NF- κ B, *aiolos*, *DOCK2*, and *Lsc* (56–61). In addition, Kraus et al. showed that impaired $Ig\alpha$ signaling in a BCR with point mutations of the $Ig\alpha$ ITAM tyrosines results in decreased MZ B cells production (41). Together our data suggests that the size of the MZ compartment is directly related to the number of $Ig\alpha$ ITAM tyrosines.

Similarly B1 development is BCR dependent and requires positive selection by low affinity interaction with self-antigen (62–64). The reduced cell surface BCR and decreased signaling by $Ig\beta_c \rightarrow \alpha_c$ B cells may simply be insufficient for normal B1 selection. Alternatively, there may be a specific requirement for $Ig\beta$ signaling in the development of the B1 population but this seems unlikely as $Ig\beta$ is not sufficient to produce B1 cells in $Ig\alpha^{FF/FF}$ mice.

Decreased Surface BCR Expression in $Ig\beta_c \rightarrow \alpha_c$ Mice. Cell surface expression of mIg requires coexpression of either $Ig\alpha$ – $Ig\beta$ or $Ig\beta$ (6, 38, 65). This requirement is linked to the presence of polar residues in the transmembrane domain of mIg that interact with $Ig\alpha$ – $Ig\beta$ (6). Once on the

cell surface the BCR is constitutively recycled from the plasma membrane to endosomes where captured antigens are processed for presentation to cognate T cells. Signaling by the cytoplasmic domains of $Ig\alpha$ and $Ig\beta$ is essential for BCR internalization and targeting to MHC II-containing endosomes (47, 66). The idea that tyrosines in the cytoplasmic domains of $Ig\alpha$ and $Ig\beta$ may be critical for constitutive BCR internalization derived from studies of endocytosis of the transferrin and LDL receptors (67, 68). Bulky hydrophobic amino acids such as phenylalanine or tyrosine in the cytoplasmic domains of those receptors were required for constitutive endocytosis. Consistent with the role of such residues in endocytosis $Ig\alpha$ tyrosines were found to be essential for both signal transduction and antigen presentation in cells lines transfected with Fc– $Ig\alpha$ chimeras (69). In contrast $Ig\beta$ tyrosine residues were dispensable for internalization (70). In addition, experiments with PDGFR chimeras that contained both $Ig\alpha$ and $Ig\beta$ showed that both cytoplasmic domains were required to regulate ligand-induced internalization (47). The implication of these in vitro studies was that $Ig\alpha$ and $Ig\beta$ had distinct but complementary roles in mediating BCR internalization.

A role for $Ig\alpha$ in regulating BCR surface expression in mature B cells in vivo was confirmed by gene targeting (41). B cells with $Ig\alpha$ ITAM tyrosines mutated to phenylalanine displayed increased surface BCR expression in vivo (41). Although the mechanism for increased receptor expression in $Ig\alpha^{FF/FF}$ mice was not determined three alternatives were suggested. First, B cells with increased levels of surface BCR might be selected during development to compensate for the decreased signaling activity of the mutant BCR (41). Second, $Ig\alpha$ signaling might normally inhibit $Ig\beta$ function (36, 71). In this scheme uninhibited $Ig\beta$ signaling would account for increased BCR surface expression in $Ig\alpha^{FF/FF}$ B cells and increased BCR signaling in $Ig\alpha\Delta C$ B cells (36, 71). However, $Ig\beta\Delta C$ mutant B cells that lack the $Ig\beta$ cytoplasmic tail showed the same hyper-reactive phenotype as $Ig\alpha\Delta C$ B cells (35). Thus, the negative regulatory effect is not specific for either $Ig\alpha$ or $Ig\beta$ and is more likely due to inability of the single chain BCRs in $Ig\alpha\Delta C$ and $Ig\beta\Delta C$ B cells to recruit negative regulators of BCR function such as Src homology 2 domain-containing phosphatase 1 (35). Finally, $Ig\alpha$ signaling might directly control BCR surface expression or internalization (41). According to this hypothesis signals emanating from the $Ig\alpha$ ITAM would be required to set cell surface BCR levels and the absence of such signals in $Ig\alpha^{FF/FF}$ B cells would lead to increased expression and a proportional increase in BCR signaling. Our data is most consistent with the idea that $Ig\alpha$ functions as such a regulator as we find lower surface BCR levels when the BCR contains an additional $Ig\alpha$ cytoplasmic domain. This decrease in BCR expression is directly correlated with, and may be responsible for, the decreased signaling by BCRs with two $Ig\alpha$ ITAMs. Furthermore, phosphorylation of ITAM tyrosines is critical for regulating surface BCR levels because $Ig\beta_c \rightarrow \alpha_c/Ig\alpha^{FF/FF}$ BCRs with a lower total number of active $Ig\alpha$ ITAMs have increased BCR surface expression. Active feedback

regulation of cell surface receptor levels is a common mechanism for regulating cellular sensitivity to persistent stimuli. We speculate that Ig α performs this feedback function for regulating BCR surface expression in vivo.

Ig $\beta_c \rightarrow \alpha_c$ Mature B Cells Are Anergic. Subthreshold signaling leads to anergy in both T cells and B cells, possibly by partial stimulation of Ca²⁺-dependent pathways (72, 73). B cells stimulated with submitogenic doses of anti-BCR antibodies in vitro showed decreased surface BCR expression and became relatively anergic to further stimulation (74, 75). In the same way, anti-hen egg lysozyme-specific B cells became anergic when exposed to cognate antigen in vivo, and this was accompanied by decreased BCR surface expression (76). Finally, similar effects including decreased surface BCR expression and inability to secrete antibody were found in anti-double- (77) and anti-single-stranded DNA Ig transgenic mice (78). In Ig-transgenic models, subthreshold BCR signaling was associated with increased baseline Ca²⁺ levels (73,79), PKC δ -dependent changes in Ca²⁺ responses, and nuclear signaling pathways including NF κ B and JNK (79–82). In contrast, we found no differences in resting Ca²⁺ levels in Ig $\beta_c \rightarrow \alpha_c$ B cells. Anergy in this case may be due to an overall decrease in signaling as a result of low levels of cell surface receptor expression.

Ig $\beta_c \rightarrow \alpha_c$ B cells spontaneously display many of the features of anergic B cells, including decreased surface BCR expression and diminished responses to BCR cross-linking. In addition Ig $\beta_c \rightarrow \alpha_c$ B cells showed impaired polyclonal responses to T cell-independent and -dependent antigens in vivo. These changes in sensitivity to BCR cross-linking required signal transduction and Ig α tyrosine phosphorylation because they were partially reversed in Ig $\beta_c \rightarrow \alpha_c$ /Ig $\alpha^{FF/FF}$ B cells in which one of the two BCR ITAMs is inactivated. The major difference between Ig $\beta_c \rightarrow \alpha_c$ B cells and other anergic B cells is that they are long lived when compared with wild-type B cells (77, 83). Little is known about how B cell longevity is regulated but our experiments suggest that Ig α may play an important role in this process. We conclude that Ig α and Ig β signaling have distinct roles in regulating mature B cell physiology in vivo.

We thank Dr. Randolph Noelle for the CD8-gp39 fusion protein derived from insect cells. We thank the Nussenzweig lab for helpful discussions and Eva Besmar for her critical review of the manuscript. We appreciate the expertise of F. Isdell and M. Genova for flow cytometry.

A. Reichlin was supported by a National Institutes of Health grant RO3 and a March of Dimes Investigator award. M.C. Nussenzweig was supported by grants from the National Institutes of Health and is an Investigator at the Howard Hughes Medical Institute.

Submitted: 10 July 2003

Accepted: 2 February 2004

References

1. Rajewsky, K. 1996. Clonal selection and learning in the antibody system. *Nature*. 381:751–758.
2. Meffre, E., R. Casellas, and M.C. Nussenzweig. 2000. Antibody

- regulation in B cell development. *Nat. Immunol.* 1:379–385.
3. Hermanson, G.G., D. Eisenberg, P.W. Kincade, and R. Wall. 1988. B29: a member of the immunoglobulin gene superfamily exclusively expressed on B-lineage cells. *Proc. Natl. Acad. Sci. USA*. 85:6890–6894.
4. Sakaguchi, N., S. Kashiwamura, M. Kimoto, P. Thalmann, and F. Melchers. 1988. B lymphocyte lineage-restricted expression of mb-1, a gene with CD3-like structural properties. *EMBO J.* 7:3457–3464.
5. Shaw, A.C., R.N. Mitchell, Y.K. Weaver, T.J. Campos, A.K. Abbas, and P. Leder. 1990. Mutations of immunoglobulin transmembrane and cytoplasmic domains: effects on intracellular signaling and antigen presentation. *Cell*. 63:381–392.
6. Sanchez, M., Z. Misulovin, A.L. Burkhardt, S. Mahajan, T. Costa, R. Franke, J.B. Bolen, and M.C. Nussenzweig. 1993. Signal transduction by immunoglobulin is mediated through Ig α and Ig β . *J. Exp. Med.* 178:1049–1056.
7. Williams, G.T., C.J.G. Peaker, K.J. Patel, and M.S. Neuberger. 1994. The alpha/beta sheath and its cytoplasmic tyrosines are required for signaling by the B-cell antigen receptor but not for capping of for serine/threonine-kinase recruitment. *Proc. Natl. Acad. Sci. USA*. 91:474–478.
8. Reth, M. 1989. Antigen receptor tail clue. *Nature*. 338:383–384.
9. Kim, K.M., G. Alber, P. Weiser, and M. Reth. 1993. Differential signaling through the Ig-alpha and Ig-beta components of the B cell antigen receptor. *Eur. J. Immunol.* 23:911–916.
10. Flaswinkel, H., and M. Reth. 1994. Dual role of the tyrosine activation motif of the Ig-a protein during signal transduction via the B cell antigen receptor. *EMBO J.* 13:83–89.
11. Pao, L.I., S.J. Famiglietti, and J.C. Cambier. 1998. Asymmetrical phosphorylation and function of the immunoreceptor tyrosine-based activation motif tyrosines in B cell antigen receptor signal transduction. *J. Immunol.* 160:3305–3314.
12. Rowley, R., A.L. Burkhardt, H.G. Chao, G.R. Matsueda, and J.B. Bolen. 1995. Syk protein tyrosine kinase is regulated by tyrosine phosphorylated Ig alpha/Ig beta immunoreceptor tyrosine activation motif and autophosphorylation. *J. Biol. Chem.* 270:11590–11594.
13. Kurosaki, T., S.A. Johnson, L. Pai, K. Sada, H. Yamamura, and J.C. Cambier. 1995. Role of the Syk autophosphorylation site and SH2 domains in B cell antigen receptor signaling. *J. Exp. Med.* 182:1815–1823.
14. DeFranco, A. 1997. The complexity of signaling pathways activated by the BCR. *Curr. Opin. Immunol.* 9:296–308.
15. Cornall, R.J., A.M. Cheng, T. Pawson, and C.C. Goodnow. 2000. Role of Syk in B-cell development and antigen-receptor signaling. *Proc. Natl. Acad. Sci. USA*. 97:1713–1718.
16. Clark, M.R., K.S. Campbell, A. Kazlauskas, S.A. Johnson, M. Hertz, T.A. Potter, C. Pleiman, and J.C. Cambier. 1992. The B cell antigen receptor complex: association of Ig-alpha and Ig-beta with distinct cytoplasmic effectors. *Science*. 258: 123–126.
17. Clark, M., S.A. Johnson, and J.C. Cambier. 1994. Analysis of Ig α -tyrosine kinase interaction reveals two levels of binding specificity and tyrosine phosphorylated Ig-stimulation of Fyn activity. *EMBO J.* 13:1911–1919.
18. Pawson, T. and J.D. Scott. 1997. Signaling through scaffold, anchoring, and adaptor proteins. *Science* 278:2075–2080.
19. Hsueh, R., and R.H. Scheuermann. 2000. Tyrosine kinase activation in the decision between growth, differentiation, and death responses initiated from the B cell antigen receptor. *Adv. Immunol.* 75:283–316.
20. Fu, C., C.W. Turck, T. Kurosaki, and A.C. Chan. 1998.

- BLNK: a central linker protein in B cell activation. *Immunity*. 9:93–103.
21. Wienands, J., J. Schweikert, B. Wollscheid, H. Jumaa, P.J. Nielsen, and M. Reth. 1998. SLP-65 a new signaling component in B lymphocytes which requires expression of the antigen receptor for phosphorylation. *J. Exp. Med.* 188:791–795.
 22. Saijo, K., C. Schmedt, I.-H. Su, H. Karasuyama, C.A. Lowell, M. Reth, T. Adachi, A. Patke, A. Santana, and A. Tarakhovskiy. 2003. Essential role of Src-family protein tyrosine kinases in NF- κ B activation during B cell development. *Nat. Immunol.* 4:274–279.
 23. Turner, M., P.J. Mee, P.S. Costello, O. Williams, A.A. Price, L.P. Duddy, M.T. Furlong, R.L. Geahlen, and V.L. Tybulewicz. 1995. Perinatal lethality and blocked B-cell development in mice lacking the tyrosine kinase Syk. *Nature*. 378:298–302.
 24. Schweighoffer, E., L. Vanes, A. Mathiot, T. Nakamura, and V.L.J. Tybulewicz. 2003. Unexpected requirement for ZAP-70 in pre-B cell development and allelic exclusion. *Immunity*. 18:523–533.
 25. Hashimoto, S., A. Iwamatsu, M. Ishiai, T. Okawa, T. Yamadori, M. Matsushita, Y. Baba, T. Kishimoto, T. Kurosaki, and S. Tsukada. 1999. Identification of the SH2 binding protein of Bruton's tyrosine kinase as BLNK-functional significance of btk-SH2 domain in B-cell antigen receptor-coupled calcium signaling. *Blood*. 94:2357–2364.
 26. Engels, N., B. Wollscheid, and J. Wienands. 2001. Association of SLP-65/BLNK with the B cell antigen receptor through a non-ITAM tyrosine of Ig- α . *Eur. J. Immunol.* 31:2126–2134.
 27. Ishiai, M., M. Kurosaki, R. Pappu, K. Okawa, I. Ronko, C. Fu, M. Shibata, A. Iwamatsu, A.C. Chan, and T. Kurosaki. 1999. BLNK required for coupling Syk to PLC γ 2 and Rac1-JNK in B cells. *Immunity*. 10:117–125.
 28. Taddie, J., T.R. Hurley, B.S. Hardwick, and B.M. Sefton. 1994. Activation of B- and T-cells by the cytoplasmic domains of the B-cell antigen receptor proteins Ig- α and Ig β . *J. Biol. Chem.* 269:13529–13535.
 29. Luisiri, P., Y.J. Lee, B.J. Eisfelder, and M.R. Clark. 1996. Cooperativity and segregation of function within the Ig- α /beta heterodimer of the B cell antigen receptor complex. *J. Biol. Chem.* 271:5158–5163.
 30. Tseng, J., B.J. Eisfelder, and M.R. Clark. 1997. B-cell antigen receptor-induced apoptosis requires both Ig alpha and Ig beta. *Blood*. 89:1513–1520.
 31. Papavasiliou, F., Z. Misulovin, H. Suh, and M.C. Nussenzweig. 1995. The role of Ig β in precursor B cell transition and allelic exclusion. *Science*. 268:408–411.
 32. Papavasiliou, F., M. Jankovic, H. Suh, and M.C. Nussenzweig. 1995. The cytoplasmic domains of Ig α and Ig β can independently induce the pre-B cell transition and allelic exclusion. *J. Exp. Med.* 182:1389–1394.
 33. Teh, Y., and M.S. Neuberger. 1997. The immunoglobulin (Ig) α and Ig β cytoplasmic domains are independently sufficient to signal B cell maturation and activation in transgenic mice. *J. Exp. Med.* 185:1753–1768.
 34. Torres, R.M., H. Flaswinkel, M. Reth, and K. Rajewsky. 1996. Aberrant B cell development and immune response in mice with a compromised BCR complex. *Science*. 272:1804–1808.
 35. Reichlin, A., Y. Hu, E. Meffre, H. Nagaoka, S. Gong, M. Kraus, K. Rajewsky, and M.C. Nussenzweig. 2001. B cell development is arrested at the immature b cell stage in mice carrying a mutation in the cytoplasmic domain of immunoglobulin B. *J. Exp. Med.* 193:13–23.
 36. Torres, R.M., and K. Hafen. 1999. A negative regulatory role for Ig- α during B cell development. *Immunity*. 11: 527–536.
 37. Yagi, Y., Y. Ikawa, K. Yoshida, Y. Shigetani, N. Takeda, I. Mabuchi, T. Yamamoto, and S. Aizawa. 1990. Homologous recombination at c-fyn locus of mouse embryonic stem cells with use of diphtheria toxin A-fragment gene in negative selection. *Proc. Natl. Acad. Sci. USA*. 87:9918–9922.
 38. Costa, T., R.R. Franke, M. Sanchez, Z. Misulovin, and M.C. Nussenzweig. 1992. Functional reconstitution of an immunoglobulin antigen receptor in T cells. *J. Exp. Med.* 175:1669–1676.
 39. Foy, T.M., F.H. Durie, and R.J. Noelle. 1994. The expansive role of CD40 and its ligand, gp39, in immunity. *Semin. Immunol.* 6:259–266.
 40. Burkhardt, A., T. Costa, Z. Misulovin, B. Stealy, J.B. Bolen, and M.C. Nussenzweig. 1994. Ig alpha and Ig beta are functionally homologous to the signaling proteins of the T cell receptor. *Mol. Cell. Biol.* 14:1095–1103.
 41. Kraus, M., L. Pao, A. Reichlin, Y. Hu, B. Canono, J.C. Cambier, M.C. Nussenzweig, and K. Rajewsky. 2001. Interference with immunoglobulin (Ig) alpha immunoreceptor tyrosine activation motif (ITAM) phosphorylation modulates or blocks B cell development, depending on the availability of an Ig beta cytoplasmic tail. *J. Exp. Med.* 194:455–469.
 42. Hao, Z., and K. Rajewsky. 2001. Homeostasis of peripheral B cells in the absence of B cell influx from the bone marrow. *J. Exp. Med.* 194:1151–1164.
 43. Nagata, K., T. Nakamura, F. Kitamura, S. Kuramochi, S. Taki, K.S. Campbell, and H. Karasuyama. 1997. The Ig alpha/Ig beta heterodimer on mu-negative proB cells is competent for transducing signals to induce early B cell differentiation. *Immunity*. 7:559–570.
 44. Love, P., E.W. Shores, M.D. Johnson, M.E. Tremblay, J. Lee, A. Grinberg, S.P. Huang, A. Singer, and H. Westpahl. 1993. T cell development in mice that lack the z chain of the T cell receptor complex. *Science*. 261:918–921.
 45. Shores, E.W., M. Ono, T. Kawabe, C.L. Sommers, T. Tran, K. Lui, M.C. Udey, J. Ravetch, and P.E. Love. 1998. T cell development in mice lacking all t cell receptor z family members (z, n, and Fc ϵ R Ig). *J. Exp. Med.* 187:1093–1101.
 46. Shores, E.W., T. Tran, A. Grinberg, C.L. Sommers, H. Shen, and P.E. Love. 1997. Role of the multiple T cell receptor (TCR)-z chain signaling motifs in selection of the t cell repertoire. *J. Exp. Med.* 185:893–900.
 47. Siemasko, K., B.J. Eisfelder, C. Stebbins, S. Kabak, A.J. Sant, W. Song, and M.R. Clark. 1999. Ig alpha and Ig beta are required for efficient trafficking to late endosomes and to enhance antigen presentation. *J. Immunol.* 162:6518–6525.
 48. Siemasko, K., B.J. Skaggs, S. Kabak, E. Williamson, B.K. Brown, W. Song, and M.R. Clark. 2002. Receptor-facilitated antigen presentation requires the recruitment of B cell linker protein to Ig α . *J. Immunol.* 168:2127–2138.
 49. Goitsuka, R., Y. Fujimura, H. Mamada, A. Umeda, T. Morimura, K. Uetsuka, K. Doi, S. Tsuji, and D. Kitamura. 1998. BASH, a novel signaling molecule preferentially expressed in B cells of the bursa of Fabricius. *J. Immunol.* 161:5804–5808.
 50. Tan, J., S.C. Wong, S.K. Gan, S. Xu, and K.P. Lam. 2001. The adaptor protein BLNK is required for B cell antigen receptor-induced activation of nuclear factor- κ B and cell cycle entry and survival of B lymphocytes. *J. Biol. Chem.* 276: 20055–20063.

51. Tsuji, S., M. Okamoto, K. Yamada, N. Okamoto, R. Goitsuka, R. Arnold, F. Kefer, and D. Kitamura. 2001. B cell adaptor containing src homology 2 domain (BASH) links B cell receptor signaling to the activation of hematopoietic progenitor kinase 1. *J. Exp. Med.* 194:529–539.
52. Hayashi, K., R. Nittono, N. Okamoto, S. Tsuji, Y. Hara, R. Goitsuka, and D. Kitamura. 2000. The B cell-restricted adaptor BASH is required for normal development and antigen receptor-mediated activation of B cells. *Proc. Natl. Acad. Sci. USA.* 97:2755–2760.
53. Jumaa, H., B. Wollscheid, M. Mitterer, J. Wienands, M. Reth, and P.J. Nielsen. 1999. Abnormal development and function of B lymphocytes in mice deficient for the signaling adaptor protein SLP-65. *Immunity.* 11:547–554.
54. Flemming, A., T. Brummer, M. Reth, and H. Jumaa. 2003. The adaptor protein SLP-65 acts as a tumor suppressor that limits pre-B cell expansion. *Nat. Immunol.* 4:38–43.
55. Xu, S., J.E. Tan, E.P. Wong, A. Manickam, S. Ponniah, and K.P. Lam. 2000. B cell development and activation defects resulting in xid-like immunodeficiency in BLNK/SLP-65-deficient mice. *Int. Immunol.* 12:397–404.
56. Martin, F., and J.F. Kearney. 2000. Positive selection from newly formed to marginal zone B Cells depends on the rate of clonal production, CD19, and Btk. *Immunity.* 12:39–49.
57. Guinamard, R., M. Okigaki, J. Schlessinger, and J.V. Ravetch. 2000. Absence of marginal zone B cells in Pyk-2-deficient mice defines their role in the humoral response. *Nat. Immunol.* 1:31–36.
58. Cariappa, A., H.C. Liou, B.H. Horwitz, and S. Pillai. 2000. Nuclear factor kappa B is required for the development of marginal zone B lymphocytes. *J. Exp. Med.* 192:1175–1182.
59. Cariappa, A., M. Tang, C. Parnig, E. Nebelitskiy, M. Carroll, K. Georgopoulos, and S. Pillai. 2001. The follicular versus marginal zone B lymphocyte cell fate decision is regulated by Aiolos, Btk, and CD21. *Immunity.* 14:603–615.
60. Fukui, Y., O. Hashimoto, T. Sanui, T. Oono, H. Koga, M. Abe, A. Inayoshi, M. Noda, M. Oike, T. Shirai, and T. Sasazuki. 2001. Haematopoietic cell-specific CDM family protein DOCK2 is essential for lymphocyte migration. *Nature.* 412:826–831.
61. Girkontaite, I., K. Missy, V. Sakk, A. Harenberg, K. Tedford, T. Potzel, K. Pfeffer, and K.D. Fischer. 2001. Lsc is required for marginal zone B cells, regulation of lymphocyte motility and immune responses. *Nat. Immunol.* 2:855–862.
62. Fagarasan, S., N. Watanabe, and T. Honjo. 2000. Generation, expansion, migration and activation of mouse B1 cells. *Immunol. Rev.* 176:205–215.
63. Lam, K.P., and K. Rajewsky. 1999. B cell antigen receptor specificity and surface density together determine B-1 versus B-2 cell development. *J. Exp. Med.* 190:471–477.
64. Hayakawa, K., M. Asano, S.A. Shinton, M. Gui, D. Allman, C.L. Stewart, J. Silver, and R.R. Hardy. 1999. Positive selection of natural autoreactive B cells. *Science.* 285:113–116.
65. Hombach, J., T. Tsubata, L. Leclercq, H. Stappert, and M. Reth. 1990. Molecular components of the B-cell antigen receptor complex of the IgM class. *Nature.* 343:760–762.
66. Siemasko, K., B.J. Eisfelder, E. Williamson, S. Kabak, and M.R. Clark. 1998. Signals from the B lymphocyte antigen receptor regulate MHC class II containing late endosomes. *J. Immunol.* 160:5203–5208.
67. Jing, S., T. Spencer, K. Miller, C. Hopkins, and I.S. Trowbridge. 1990. Role of the human transferrin receptor cytoplasmic domain in endocytosis: localization of a specific signal sequence for internalization. *J. Cell Biol.* 110:283–294.
68. Chen, W., J.L. Goldstein, and M.S. Brown. 1990. NPXY a sequence often found in cytoplasmic tails is required for coated pit-mediated internalization of the low density lipoprotein receptor. *J. Biol. Chem.* 265:3116–3123.
69. Cassard, S., J. Salamero, D. Hanau, D. Spehner, J. Davoust, W.H. Fridman, and C. Bonnerot. 1998. A tyrosine-based signal present in Ig alpha mediates B cell receptor constitutive internalization. *J. Immunol.* 160:1767–1773.
70. Patel, K., and M.S. Neuberger. 1993. Antigen presentation by the B cell antigen receptor is driven by the a/b sheath and occurs independently of its cytoplasmic tyrosines. *Cell.* 74:939–946.
71. Kraus, M., K. Saijo, R.M. Torres, and K. Rajewsky. 1999. Ig-alpha cytoplasmic truncation renders immature B cells more sensitive to antigen contact. *Immunity.* 11:537–545.
72. Macián, F., F. Garcia-Cózar, S.H. Im, H.F. Horton, M.C. Byrne, and A. Rao. 2002. Transcriptional mechanisms underlying lymphocyte tolerance. *Cell.* 109:719–731.
73. Dolmetsch, R., R.S. Lewis, C.C. Goodnow, and J.I. Healy. 1997. Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration. *Nature.* 386:855–858.
74. Pike, B., A.W. Boyd, and G.J. Nossal. 1982. Clonal anergy: the universally anergic B lymphocyte. *Proc. Natl. Acad. Sci. USA.* 79:2013–2017.
75. Rudich, S., K.H. Roux, R.J. Winchester, and P.K. Mongini. 1988. Anti-IgM-mediated B cell signaling. Molecular analysis of ligand binding requisites for human B cell clonal expansion and tolerance. *J. Exp. Med.* 168:247–266.
76. Goodnow, C., J. Crosbie, H. Jorgensen, R.A. Brink, and A. Basten. 1989. Induction of self-tolerance in mature peripheral B lymphocytes. *Nature.* 342:340–341.
77. Mandik-Nayak, L., S.J. Seo, A. Eaton-Bassiri, D. Allman, R.R. Hardy, and J. Erikson. 2000. Functional consequences of the developmental arrest and follicular exclusion of anti-double-stranded DNA B cells. *J. Immunol.* 164:1161–1168.
78. Noorchashm, H., A. Bui, L. Hsiu-Ling, A. Eaton, L. Mandik-Nayak, C. Sokol, K.M. Potts, E. Pure, and J. Erikson. 1999. Characterization of anergic anti-DNA B cells: B cell anergy is a T cell-independent and potentially reversible process. *Int. Immunol.* 11:765–776.
79. Healy, J., R.E. Dolmetsch, L.A. Timmerman, J.G. Cyster, M.L. Thomas, G.R. Crabtree, R.S. Lewis, and C.C. Goodnow. 1997. Different nuclear signals are activated by the B cell receptor during positive versus negative signaling. *Immunity.* 6:419–428.
80. Weintraub, B., J.E. Jun, A.C. Bishop, K.M. Shokat, M.L. Thomas, and C.C. Goodnow. 2000. Entry of B cell receptor into signaling domains is inhibited in tolerant B cells. *J. Exp. Med.* 191:1443–1448.
81. Mecklenbrauker, I., K. Saijo, N.Y. Zheng, M. Leitges, and A. Tarakhovskiy. 2002. Protein kinase C controls self-antigen-induced B-cell tolerance. *Nature.* 416:860–864.
82. Miyamoto, A., K. Nakayama, I. Hiroyuki, S. Hirose, Y. Jiang, A. Masaaki, T. Tadasuke, H. Nagahama, S. Ohno, S. Hatakeyama, and K.I. Nakayama. 2002. Increased proliferation of B cells and auto-immunity in mice lacking protein Kinase C. *Nature.* 416:865–869.
83. Fulcher, D., and A. Basten. 1994. Reduced lifespan of anergic self reactive B cells in a double-transgenic model. *J. Exp. Med.* 179:125–134.