

Takashi Usui, Jan C. Preiss, Yuka Kanno, Zheng Ju Yao, Jay H. Bream, John J. O'Shea, and Warren Strober
 Vol. 203, No. 3, March 20, 2006. Pages 755–766.

The authors regret a labeling error in Fig. 1. In panel B, the first row, first column of the table should have read “–/–” and the second row, first column of the table should have read “+/+.” The corrected figure and its legend appear below.




Figure 1. Elevated IL-4 and GATA-3 and committed Th2 cells in T-bet^{-/-} mice. (A) Naive CD4⁺ T cells and whole CD4⁺ T cells from T-bet^{-/-}, T-bet^{+/+}, and wild-type mice were stimulated with anti-CD3/anti-CD28 and maintained under Th1 conditions (IL-12 only) and expanded with IL-2. On day 6, the cells were subjected to intracellular cytokine staining after restimulation with PMA/ionomycin. These data are representative of results obtained from three independent experiments. (B) Naive CD4⁺ T cells from T-bet^{-/-} and wild-type mice were stimulated with anti-CD3/anti-CD28 under Th1 conditions (IL-12 only) or neutral (no cytokines or antibodies added) conditions for 48 h, and then culture supernatants were harvested for IFN- γ

and IL-4 determination by ELISA. The data shown are averages \pm SE derived from three independent experiments. (C) Naive CD4⁺ T cells from T-bet^{-/-} and wild-type mice were stimulated with anti-CD3/anti-CD28 under Th1 conditions (IL-12 only) or neutral conditions and maintained for the time periods shown in the figure. At each time point, cells were harvested and total RNAs were isolated and subjected to real-time PCR analysis for GATA-3 mRNA quantification. Independently, total RNAs from freshly isolated naive and memory CD4⁺ T cells were prepared for day 9 samples. The data shown are averages \pm SE derived from three independent experiments.