

# Direct recognition of the mycobacterial glycolipid, trehalose dimycolate, by C-type lectin Mincle

Eri Ishikawa,<sup>1</sup> Tetsuaki Ishikawa,<sup>1,3</sup> Yasu S. Morita,<sup>4,6</sup> Kenji Toyonaga,<sup>1</sup> Hisakata Yamada,<sup>2</sup> Osamu Takeuchi,<sup>5,7</sup> Taroh Kinoshita,<sup>4,6</sup> Shizuo Akira,<sup>5,7</sup> Yasunobu Yoshikai,<sup>2</sup> and Sho Yamasaki<sup>1</sup>

<sup>1</sup>Division of Molecular Immunology, <sup>2</sup>Division of Host Defense, Medical Institute of Bioregulation, Kyushu University, Fukuoka 812-8582, Japan

<sup>3</sup>Cell Signaling, Chiba University School of Medicine, Chiba 260-8670, Japan

<sup>4</sup>Department of Immunoregulation, <sup>5</sup>Department of Host Defense, Research Institute for Microbial Diseases, <sup>6</sup>Laboratory of Immunoglycobiology, <sup>7</sup>Laboratory of Host Defense, WPI Immunology Frontier Research Center, Osaka University, Suita 565-0871, Japan

**Tuberculosis remains a fatal disease caused by *Mycobacterium tuberculosis*, which contains various unique components that affect the host immune system. Trehalose-6,6'-dimycolate (TDM; also called cord factor) is a mycobacterial cell wall glycolipid that is the most studied immunostimulatory component of *M. tuberculosis*. Despite five decades of research on TDM, its host receptor has not been clearly identified. Here, we demonstrate that macrophage inducible C-type lectin (Mincle) is an essential receptor for TDM. Heat-killed mycobacteria activated Mincle-expressing cells, but the activity was lost upon delipidation of the bacteria; analysis of the lipid extracts identified TDM as a Mincle ligand. TDM activated macrophages to produce inflammatory cytokines and nitric oxide, which are completely suppressed in Mincle-deficient macrophages. In vivo TDM administration induced a robust elevation of inflammatory cytokines in sera and characteristic lung inflammation, such as granuloma formation. However, no TDM-induced lung granuloma was formed in Mincle-deficient mice. Whole mycobacteria were able to activate macrophages even in MyD88-deficient background, but the activation was significantly diminished in Mincle/MyD88 double-deficient macrophages. These results demonstrate that Mincle is an essential receptor for the mycobacterial glycolipid, TDM.**

## CORRESPONDENCE

Sho Yamasaki:  
yamasaki@bioreg.kyushu-u.ac.jp

Abbreviations used: C:M, chloroform:methanol; CRD, carbohydrate recognition domain; HPTLC, high-performance thin-layer chromatography; iNOS, inducible NO synthase; LAM, lipoarabinomannan; LWI, lung weight index; Mincle, macrophage inducible C-type lectin; NO, nitric oxide; PIM, phosphatidylinositol mannoside; TDB, trehalose dibehenate; TDM, trehalose-6,6'-dimycolate; TLR, Toll-like receptor; TMM, trehalose monomycolate.

Many pathogens are directly recognized by pattern recognition receptors such as Toll-like receptors (TLRs), RIG-I-like helicases, or NOD-like receptors of the host cells (Akira et al., 2006), most of which sense the characteristic signatures of pathogens. Recently, some members of the C-type lectin family have also been identified as pattern recognition receptors for bacteria or fungi; however, the ligands of most C-type lectin receptors remain unidentified (Robinson et al., 2006).

Among these C-type lectin receptors is Mincle (macrophage inducible C-type lectin, also called Clec4e or Clec5f9), which is expressed in macrophages subjected to several types of stress (Matsumoto et al., 1999). Mincle possesses carbohydrate recognition domain

(CRD) within the extracellular region. We recently reported that Mincle is associated with an immunoreceptor tyrosine-based activation motif-containing Fc receptor  $\gamma$  chain (FcR $\gamma$ ) and functions as an activating receptor for damaged self- and non-self-pathogenic fungi (Yamasaki et al., 2008, 2009).

Tuberculosis is caused by *Mycobacterium tuberculosis* that infects one third of the world's population (Hunter et al., 2006). *M. tuberculosis* contains various unique components that affect the host immune system through both identified and unidentified receptors (Jo, 2008). Among these, the cord factor is the first immuno-

E. Ishikawa and T. Ishikawa contributed equally to this work.

© 2009 Ishikawa et al. This article is distributed under the terms of an Attribution-NonCommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.jem.org/misc/terms.shtml>). After six months it is available under a Creative Commons License (Attribution-NonCommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

stimulatory component to be identified that might elicit pulmonary inflammation, a characteristic feature of mycobacterial infection (Bloch, 1950; Yamaguchi et al., 1955). In 1956, the chemical structure of the cord factor was established as trehalose-6,6'-dimycolate (TDM; Noll et al., 1956), which is the most abundant glycolipid in the mycobacterial cell wall (Hunter et al., 2006). The long-chain lipids of TDM represent a structural component of the hydrophobic cell wall that is critical for the survival of mycobacteria within the phagosome of host cells (Indrigo et al., 2002). TDM has been shown to possess unique immunostimulatory activity, such as granulomagenesis, adjuvant activity for cell-mediated immune responses, humoral responses, and tumor regression (Hunter et al., 2006). Recently, Marco and TLR are proposed as TDM receptor (Bowdish et al., 2009), whereas other group suggested that FcR $\gamma$ -coupled receptor(s) are candidates (Werninghaus et al., 2009). Thus, the receptor for TDM is still controversial.

Some C-type lectins are involved in the recognition of *M. tuberculosis* (Jo, 2008). Mannose receptor expressed on macrophages is reported to mediate phagocytosis of mycobacteria (Kang et al., 2005). Another C-type lectin receptor, DC-SIGN, is known to recognize mycobacteria (Tailleux et al., 2003), but its binding results in the down-regulation of dendritic cell-mediated immune responses (Geijtenbeek et al., 2003). Dectin-1 is reported to recognize mycobacteria

through an unknown ligand (Yadav and Schorey, 2006; Rothfuchs et al., 2007). Other C-type lectins could potentially recognize mycobacteria, but the activating lectin receptors for mycobacteria in macrophages have not been clearly identified.


In this study, we show that Mincle is a direct receptor for mycobacterial TDM. We further demonstrate, through Mincle-deficient mice, that Mincle is an essential receptor for TDM-dependent inflammatory responses.

## RESULTS AND DISCUSSION

### Mincle can recognize mycobacteria

We first investigated whether Mincle recognizes mycobacteria through the use of NFAT-driven GFP reporter cells that express Mincle and its signaling subunit, the FcR $\gamma$  chain (Yamasaki et al., 2008). The heat-killed mycobacterial species *Mycobacterium smegmatis* and *Mycobacterium bovis* Bacille de Calmette et Guerin clearly activated NFAT-GFP in reporter cells expressing FcR $\gamma$  with Mincle (Fig. 1, A and B). Importantly, the virulent strain *M. tuberculosis* H37Rv also showed substantial ligand activity against Mincle (Fig. 1 C).

The recognition of mycobacteria by Mincle was shown to require the Glu-Pro-Asn (EPN) sequence, a putative mannose-binding motif within CRD, as the activity was eliminated by introducing a mutation of EPN into QPD



**Figure 1. Mincle recognizes mycobacterial species.** (A–C) NFAT-GFP reporter cells expressing FcR $\gamma$  only (FcR $\gamma$ ), Mincle + FcR $\gamma$ , and Mincle<sup>EPN→QPD</sup> + FcR $\gamma$  were co-cultured for 18 h with heat-killed *M. smegmatis* (A), *M. bovis* (B), or *M. tuberculosis* H37Rv (C). Induction of NFAT-GFP was analyzed by flow cytometry. (D) Reporter cells expressing Dectin-2 + FcR $\gamma$  were stimulated with *M. smegmatis*, *M. bovis*, and *Candida albicans* (*C. a.*). (E) Reporter cells expressing Mincle + FcR $\gamma$  were stimulated with 10<sup>7</sup> *M. smegmatis* together with anti-Mincle mAb and rat IgG1 as a control. (F) Cells were stimulated with wild-type *M. smegmatis* (WT), pimE-deficient *M. smegmatis* ( $\Delta$ pimE), and pimE-deficient *M. smegmatis* reconstituted with pimE ( $\Delta$ pimE + pimE). All data are means  $\pm$  SD for triplicate assays, and representative results from three independent experiments with similar results are shown.


(Mincle<sup>QPD</sup>; Fig. 1, B and C; Drickamer, 1992). However, the similar immunoreceptor tyrosine-based activation motif-coupled C-type lectin Dectin-2 (Sato et al., 2006) was not capable of recognizing mycobacteria (Fig. 1 D), suggesting selective recognition for Mincle. Indeed, activation of NFAT by mycobacteria was completely blocked in the presence of anti-Mincle mAb (Fig. 1 E).

As a putative mannose-binding motif within Mincle was essential for the recognition process (Fig. 1, B and C), we initially assumed that Mincle could recognize terminal  $\alpha$ 1,2 mannose residues of mycobacterial molecules, such as phosphatidylinositol mannoside (PIM), lipomannan, or lipoarabinomannan (LAM). To examine the contribution of PIM, we used mutant strains of mycobacteria lacking the terminal  $\alpha$ 1,2 mannose residues. *M. smegmatis*  $\Delta$ pimE, a deletion mutant of mannosyl-transferase PimE, lacks mature PIMs (Fig. S1; Morita et al., 2006). However,  $\Delta$ pimE still retained stimulatory activity at a comparable level to PimE-sufficient strains such as wild-type *M. smegmatis* and pimE-reconstituted  $\Delta$ pimE strain (Fig. 1 F). In addition, a mutant *M. smegmatis* strain deficient in  $\alpha$ 1,2 mannose of lipomannan and LAM could be recognized normally by Mincle, and purified LAM alone did not activate Mincle-expressing cells (unpublished data).

These results show that mycobacteria can be recognized by Mincle, but their  $\alpha$ 1,2 mannose-containing glycolipids appears not to be necessary for recognition to occur.

### Identification of mycobacterial glycolipids recognized by Mincle

Mycobacteria have a wealth of unique lipids on their cell walls, some of which presumably protect the bacteria from the host's defense system. To examine the contribution of other mycobacterial lipids as a potential candidate for a Mincle ligand, we extracted the lipids from *M. smegmatis* using various organic solvents (Fig. 2 A). *M. smegmatis* treated with chloroform:methanol (C:M) selectively lost their Mincle-stimulating activity (Fig. 2 B). Simultaneously, we analyzed the activity of extracted lipid fractions in plate-coated form and found that only the C:M phase after C:M extraction showed strong stimulatory activity (Fig. 2 C). We further analyzed this active fraction by means of high-performance thin-layer chromatography (HPTLC) and separated it into 22 subfractions to identify the active lipid components. Purified extracts from these subfractions showed strong ligand activity peaked at subfractions #7-9 (Fig. 2 D). Orcinol staining revealed a purple-red band at the position corresponding to subfractions #7-9 (Fig. 2 D, left lane), indicating that the ligand contains a sugar moiety. Furthermore, the purple-red band migrated to a position similar to that of purified TDM derived from *M. tuberculosis* (Fig. 2 D, right lane). From these characteristics, we hypothesized that TDM could be a candidate for the Mincle ligand. Subfraction #2 also showed weak activity, implying that trehalose monomycolate (TMM), a precursor of TDM biosynthesis, could also act as weak ligand, as previously suggested (Sueoka et al., 1995).



**Figure 2. Mincle recognizes mycobacterial trehalose-6,6'-dimycolate.** (A) Schematic diagram of delipidation of *M. smegmatis*. Delipidated bacteria (gray boxes) and lipid extracts (open boxes) were applied for ligand assays in B and C, respectively. (B and C) Heat-killed *M. smegmatis* treated with C:M, hexane, acetone, or 1-butanol (BuOH; B), and plate-coated lipid extract (C) were co-cultured with reporter cells expressing Mincle + Fc $\gamma$ R. (D) C:M phase of C:M extract was analyzed by HPTLC and divided into 22 subfractions. Each subfraction was coated onto a plate to stimulate reporter cells. Purified TDM was used as a reference (right lane). Arrowheads show the origin and solvent front. Data are means  $\pm$  SD for triplicate assays (B and C) or means for duplicate assays (D). Representative results from three independent experiments with similar results are shown.


**TDM as a Mincle ligand**

Indeed, we found that purified TDM, the structure of which is shown in Fig. 3 A, dramatically activated Mincle-expressing cells in plate-coated form (Fig. 3 B). The TDM analogue trehalose dibehenate (TDB), which is also used as a synthetic adjuvant, was a strong ligand for Mincle, as well (Fig. 3 C).

TDM consists of a trehalose moiety and two mycolate chains (Fig. 3 A), but purified mycolate did not itself activate Mincle-expressing cells (Fig. 3 D). To examine whether TMM also possesses ligand activity or not,

generated TMM by partial alkaline deacylation of TDM. As shown in Fig. 3 E, TMM could potentially activate Mincle-expressing cells, albeit less potent than TDM. Soluble trehalose had no stimulatory activity, and a large excess of trehalose did not block TDM-mediated NFAT activation (unpublished data). Thus, a combination of both the sugar and lipid moieties appears to be critical for the ligand activity of TDM.

Next, to verify the direct interaction between Mincle and TDM, we prepared soluble Mincle protein (Mincle-Ig). As



**Figure 3. Purified TDM is recognized by Mincle.** (A) Chemical structure of TDM. α-Mycolate (shown), methoxy-mycolate, and keto-mycolate are the major subclasses of mycolate found in *M. tuberculosis* TDM. (B and C) Reporter cells were stimulated with the indicated amount of plate-coated TDM (B) or TDB (C). (D) Reporter cells were stimulated with the indicated amount of TDM, methyl α-mycolate (α-mycolate), or methyl keto-mycolate (keto-mycolate). (E) Reporter cells were stimulated with the indicated amount of TDM and TMM. (F) ELISA-based detection of TDM by Mincle-Ig. hlgG1-Fc (Ig), Mincle-Ig, and Dectin2-Ig were incubated with 0.1 nmol/0.32 cm<sup>2</sup> of plate-coated TDM. Bound protein was detected with anti-hlgG-HRP followed by the addition of colorimetric substrate. (G) Effect of trehalose (100 μg/ml), EDTA (10 mM), rat IgG (10 μg/ml), and anti-Mincle mAb (10 μg/ml) on TDM recognition by Mincle-Ig. ELISA-based detection was performed as in E. (H) Reporter cells were stimulated with TDM, which was treated with trehalase as described in Materials and methods. Cells were also stimulated with plate-coated anti-Mincle mAb treated with trehalase as a negative control. All data are means ± SD for triplicate assays and representative results from three independent experiments with similar results are shown.

detected by anti-IgG, Mincle-Ig, but not other control proteins such as Ig or Dectin2-Ig, selectively binds to plate-coated TDM (Fig. 3 F). This biochemical data provides further proof that TDM is a direct Mincle ligand. This binding was blocked by EDTA and anti-Mincle mAb but not by excess trehalose, suggesting that Mincle recognizes specific glycolipid structures in a cation-dependent manner (Fig. 3 G).


It was recently proposed that TDM on mycobacterial cell wall could be converted into glucose monomycolate (GMM) in the host cell environment (Matsunaga et al., 2008). To test the hypothesis that this conversion may be of advantage to mycobacteria in escaping from Mincle-mediated recognition, we attempted to mimic the conversion in vitro by using trehalase, which hydrolyzed trehalose into two glucose units (Asano et al., 1996). Intriguingly, the trehalase treatment impaired the ligand activity of TDM (Fig. 3 H). Note that this enzymatic treatment did not grossly disrupt Mincle-mediated responses because stimulation by plate-coated anti-Mincle mAb was not affected by the trehalase treatment (Fig. 3 H). This finding is consistent with the idea that mycobacteria convert TDM into GMM upon infection into host, presum-

ably to escape from Mincle-mediated host immunity (Fig. S2; Matsunaga et al., 2008). Intriguingly, it has also been reported that as a possible counterdefense, host cells present this GMM on group1 CD1 molecules in human to provoke TCR-mediated acquired immune responses against mycobacteria (Moody et al., 1997; Matsunaga et al., 2008). Importantly, we found that TDM is recognized by human Mincle as well as murine Mincle (unpublished data).

Collectively, these results suggest that, among various mycobacterial components, Mincle is specific for the ester linkage of a fatty acid to trehalose.

### Mincle is essential for TDM-dependent macrophage activation

TDM also activated macrophages to produce nitric oxide (NO), which is critical for direct killing of mycobacteria. However, NO production was almost completely suppressed in Mincle<sup>-/-</sup> macrophages (Fig. 4 A). In contrast, LPS induced a similar response in Mincle<sup>-/-</sup> cells to that in WT cells. Inducible NO synthase (iNOS) is the enzyme responsible for production of NO. The transcriptional induction of




**Figure 4. Lack of TDM-mediated activation in Mincle-deficient macrophages.** (A) BMM $\phi$  from Mincle<sup>+/+</sup> and Mincle<sup>-/-</sup> were primed with IFN- $\gamma$  (10 ng/ml) and stimulated with plate-coated TDM or LPS (10 ng/ml) as a control. Culture supernatants were collected at 48 h and concentration of NO was measured. (B) IFN- $\gamma$ -primed BMM $\phi$  were stimulated with plate-coated TDM for 36 h as in A, and mRNA expression of iNOS was analyzed by real-time PCR. (C and D) IFN- $\gamma$ -primed BMM $\phi$  were stimulated with plate-coated TDM (C) or TDB (D) for 24 h. Culture supernatants were collected and concentrations of TNF and MIP-2 were determined by ELISA. All data are means  $\pm$  SD for triplicate assays and representative results from three independent experiments with similar results are shown.

iNOS by TDM stimulation was also completely suppressed in *Mincle*<sup>-/-</sup> mice (Fig. 4 B).

The crucial role of TNF has been reiterated by the recent observation of patients suffering from the reactivation of latent tuberculosis infection upon anti-TNF therapy for autoimmune diseases (Winthrop, 2006). TDM stimulation induced the production of TNF and MIP-2 (also called CXCL2), which are potent chemoattractant factors for in-

flammatory cells. However, these cytokines were not produced upon TDM stimulation in the absence of *Mincle* (Fig. 4 C). Similarly, TDB-induced cytokine production was also eliminated in *Mincle*<sup>-/-</sup> macrophages (Fig. 4 D).

We have previously reported that *Mincle* transduces signal through the FcR $\gamma$ -CARD9 signaling axis (Yamasaki et al., 2008). In line with these observations, it was recently reported that TDM and TDB activates macrophage in an



**Figure 5. *Mincle* is essential for TDM-induced inflammation in vivo.** (A) *Mincle*<sup>+/+</sup>, *Mincle*<sup>-/-</sup>, *FcR $\gamma$* <sup>-/-</sup> and *Myd88*<sup>-/-</sup> mice were injected intravenously with an oil-in-water emulsion containing TDM (150  $\mu$ g). Emulsion without TDM was injected as a vehicle control. At day 1 after injection, IL-6 and TNF concentrations in sera were determined by ELISA. Each symbol represents an individual mouse. (B) Lungs of TDM-injected mice were removed at day 7 and inflammatory intensity was evaluated by calculation of LWI. (C) Proinflammatory mediator mRNA levels in lungs at day 7 after TDM administration were evaluated by real-time PCR. Relative expression levels are shown as 2<sup>-Ct</sup>. Each symbol represents an individual mouse. Ct, cycle threshold. \*, P < 0.05; \*\*, P < 0.001. NS, not significant. (D) Histology of the lungs from untreated (control) and TDM-injected (TDM) mice was examined by hematoxylin-eosin staining at day 7. Bar, 0.1 mm. (E) Number of lung granulomas. Granuloma in lungs from mice injected with TDM was counted as described in Materials and methods. Data are means  $\pm$  SD for the mean number for at least three independent mice. Representative results from two independent experiments with similar results are shown.

FcR $\gamma$ - and CARD9-dependent manner (Werninghaus et al., 2009). Thus, Mincle is an essential receptor for macrophage activation elicited by TDM and TDB.

### TDM-induced lung granuloma formation requires Mincle

In vivo administration of TDM is capable of inducing inflammatory symptoms characteristic of tuberculosis (Yamaguchi et al., 1955; Hunter et al., 2006). Single injection of TDM into mice induced the robust production of inflammatory cytokines, such as IL-6 and TNF, in sera. However, these were completely eliminated in Mincle $^{-/-}$  mice (Fig. 5 A). FcR $\gamma$  is an essential signaling subunit for Mincle (Yamasaki et al., 2008). Indeed, FcR $\gamma$  $^{-/-}$  mice did not detectably respond to TDM in vivo (Fig. 5 A). In contrast, TDM was able to induce cytokine production in MyD88 $^{-/-}$  mice, suggesting that TLRs are not essential for TDM recognition (Fig. 5 A; Werninghaus et al., 2009), although potential roles of TLR2 and TLR4 have been proposed (Bowdish et al., 2009). TDM also induced inflammatory lung swelling as assessed by lung weight index (LWI), but this was totally dependent on the Mincle-FcR $\gamma$  axis (Fig. 5 B and Fig. S3). In concert with the acute lung inflammation, the up-regulation of mRNA for inflammatory cytokines/chemokines upon TDM treatment was severely impaired in Mincle $^{-/-}$  mice


(Fig. 5 C). In addition, TDM induced thymic atrophy in mice, and was also Mincle dependent (Fig. S4).

Granulomas are complex aggregates of immune cells, and are widely believed to constrain mycobacterial infection by physically surrounding the infecting bacteria (Adams, 1976). TDM alone dramatically induced granuloma formation, which is a characteristic of mycobacterial infection at day 7 after administration (Fig. 5 D). Strikingly, no granuloma formation was observed in the lungs of TDM-treated Mincle $^{-/-}$  mice (Fig. 5 D). Quantitative analysis on multiple sections revealed that granuloma was completely eliminated in Mincle $^{-/-}$  and FcR $\gamma$  $^{-/-}$  mice, but was induced normally in MyD88 $^{-/-}$  mice (Fig. 5 E). Thus, Mincle is a critical receptor for TDM-induced granuloma formation, most likely through the production of inflammatory cytokines/chemokines to recruit inflammatory cells (Welsh et al., 2008).

These results show that Mincle is an essential receptor for TDM-mediated inflammatory responses in vivo.

### Role of Mincle in response to whole mycobacteria

Finally we investigated the role of Mincle in response to whole mycobacteria using virulent strain, *M. tuberculosis* H37Rv. Heat-killed mycobacteria induced vigorous production of TNF and MIP-2 in macrophages, whereas Mincle



**Figure 6. Mincle is responsible for the TLR-independent pathway of anti-mycobacterium responses.** (A and B) BMM $\phi$  from WT, Mincle $^{-/-}$  (A), MyD88 $^{-/-}$ , and MyD88 $^{-/-}$ Mincle $^{-/-}$  (B) mice were stimulated with indicated number of heat-killed *M. tuberculosis* H37Rv. Cells were also stimulated with zymosan as a positive control. At day 2 after stimulation, production of TNF and MIP-2 in supernatants was determined. Data are means  $\pm$  SD for triplicate assays and representative results from three independent experiments with similar results are shown. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ .

deficiency did not detectably impair the production of TNF and only partially impaired MIP-2 production (Fig. 6 A). This is probably because whole mycobacteria possess many immunostimulatory components other than TDM, including TLR ligands (Jo, 2008). Indeed, cytokine production was markedly decreased, but was substantially induced in MyD88<sup>-/-</sup> macrophages (Fig. 6 B; Fremont et al., 2004). However, the level of these cytokines was significantly reduced in MyD88<sup>-/-</sup>Mincle<sup>-/-</sup> double-deficient macrophages when compared with MyD88<sup>-/-</sup> macrophages (Fig. 6 B). In contrast, the response to zymosan, which is known to be recognized by Dectin-1 and TLR2, in MyD88<sup>-/-</sup> macrophages was the same as that in MyD88<sup>-/-</sup>Mincle<sup>-/-</sup> cells (Fig. 6). These results suggest that Mincle plays a major role in TLR-independent recognition of mycobacteria. The remaining production of cytokines by mycobacteria observed in MyD88<sup>-/-</sup>Mincle<sup>-/-</sup> macrophages (Fig. 6 B) may reflect a possible contribution of the NOD-like receptor family NOD2, which was reported to recognize mycobacteria through muramyl dipeptide (Coulombe et al., 2009).

### Concluding remarks

TDM (also called cord factor) is an essential component that permits the survival of mycobacteria within the host cells (Indrigo et al., 2002), whereas it is barely present in vertebrates. It would therefore be a reasonable strategy for the host to recognize TDM as a signal of mycobacterial infection. In this study, we identified C-type lectin Mincle as an essential receptor for TDM.

It was demonstrated that cyclopropane within mycolate chain of TDM is critical for its immunostimulatory activity (Rao et al., 2005). However, TDB, which lacks the cyclopropane in the carbon chain, is still capable of activating macrophages and Mincle-expressing cells (Fig. 3 C and Fig. 4 D; Werninghaus et al., 2009). On the other hand, it has been suggested that specific configuration of TDM is necessary to exert its activity (Retzinger et al., 1981). Therefore the “kink” in the carbon chain of TDM may contribute to the optimal presentation of polar head to Mincle, rather than as a direct binding site for Mincle. It could be hypothesized that two Mincle receptors recognize one disaccharide head of TDM, presumably together with a proximal region of mycolate, although further structural analysis is needed to clarify this issue.

TDM has been known to induce lung granuloma in vivo, and we demonstrated that it is totally dependent on Mincle. We have recently found that Mincle also senses dead cells and recruits inflammatory cells (Yamasaki et al., 2008). Because the presence of necrotic cells in the center of a granuloma is a characteristic feature of tuberculosis (Adams, 1976), both TDM and dead cells may contribute cooperatively to the formation of granuloma through Mincle-mediated secretion of inflammatory cytokines/chemokines. The physiological role of Mincle during virulent infection is a critical issue that needs to be clarified and is now under investigation using Mincle<sup>-/-</sup> mice, as the contribution of TDM in the virulence of myco-

bacteria is still controversial (Hunter et al., 2006). Given its potent granuloma-inducing capacity, Mincle might also be involved in diseases characterized by granulomas.

Recent studies have suggested that T cells can respond to TDM, despite little evidence of CD1-mediated TDM presentation (Guidry et al., 2004; Otsuka et al., 2008). Mincle is also expressed in T cells upon activation (unpublished data), and some T cell populations use FcR $\gamma$  (Ohno et al., 1994). It is tempting to speculate that such T cells may directly recognize TDM to produce T cell-specific cytokines, such as IFN- $\gamma$  or IL-17, in a “TCR-independent” but “Mincle-dependent” manner.

We recently discovered that Mincle also recognizes the pathogenic fungus *Malassezia* (Yamasaki et al., 2009). Our current findings could suggest that Mincle uniquely recognizes glycolipid through the motifs that have generally been considered as mannose-binding motifs. We therefore speculate that the Mincle-*Malassezia* interaction might be mediated by unique fungal glycolipids similar to TDM. Interestingly, *Malassezia* species among other fungi uniquely require lipid for their growth (Schmidt, 1997).

TDM and its synthetic analogue, TDB, have been extensively studied because they are effective adjuvants (Azuma and Seya, 2001). It is believed that TDM accounts for part of the effect of CFA (Billiau and Matthys, 2001). Identification of the host receptor for TDM/TDB will provide valuable information related to the design of vaccine adjuvants, because rational screening of synthetic Mincle ligands is now feasible and could potentially lead to the development of an ideal synthetic adjuvant. Such a synthetic Mincle ligand could allow efficient vaccination against tuberculosis, other infectious diseases, and cancers.

### MATERIALS AND METHODS

**Mice.** Mincle-deficient mice were used as C57BL/6 and 129 mixed genetic background (Yamasaki et al., 2009). MyD88-deficient mice were purchased from Oriental Yeast. FcR $\gamma$ -deficient mice on a C57BL/6 background were provided by T. Saito (RIKEN, Yokohama, Japan; Park et al., 1998). C57BL/6 mice were obtained from Japan Clea or Kyudo. All mice were maintained in a filtered-air laminar-flow enclosure and given standard laboratory food and water ad libitum. Animal protocols were approved by the committee of Ethics on Animal Experiment, Faculty of Medical Sciences, Kyushu University.

**Bacteria.** *M. smegmatis* strain mc<sup>2</sup>155 were cultured in Middlebrook 7H9 broth as previously described (Morita et al., 2005). *M. bovis* Bacille de Calmette et Guerin were cultured in Middlebrook 7H9 supplemented with Middlebrook ADC enrichment and 0.05% Tween-80. *M. smegmatis*  $\Delta$ pimE, which is a mutant deficient in mannosyltransferase PimE, lacks  $\alpha$ 1,2-mannose moiety of PIM. *M. smegmatis* and *M. bovis* were heat-killed by pasteurization at 63°C for 40 min. Virulent strain *M. tuberculosis* H37Rv was autoclaved before use. *Candida albicans* (IFM No. 54349) was provided by T. Gono (Chiba University, Chiba, Japan).

**Delipidation.** *M. smegmatis* was delipidated with C:M (2:1), hexane, acetone, or 1-butanol (BuOH). Insoluble fractions were collected as delipidated bacteria. Soluble fractions were further partitioned by C:M:W (8:4:3; vol/vol) into lower organic phase (C:M) and upper aqueous phase (M:W). Upper aqueous phase (M:W) was further partitioned by 1-butanol:water (1:1; vol/vol) into upper butanol phase (BuOH) and lower aqueous phase (water).



Each fraction was dried, resuspended in DMSO relative to the original cell pellet weight, and tested as lipid extracts (Morita et al., 2005).

**Reagent.** TDM, Methyl  $\alpha$ -mycolate, Methyl keto-mycolate, and LAM were purchased from Nakalai tesque and TDB was obtained from Avanti Polar Lipids, Inc. D-Trehalose dihydrate (T3663), Trehalase (T8778), LPS (L4516), and zymosan (Z4250) were purchased from Sigma-Aldrich. For stimulation of reporter cells and bone marrow-derived macrophages (BMM $\phi$ ), TDM and TDB dissolved in chloroform at 1 mg/ml were diluted in isopropanol and added on 96-well plates at 20  $\mu$ l/well, followed by evaporation of the solvent as previously described (Ozeki et al., 2006). For generation of TMM, TDM was partially acylated with 0.4 M NaOH for 10 min, and the amount of TDM and TMM was determined by orcinol staining on HPTLC. Corresponding fraction to TDM and TMM was collected and used for assay.

**Cells.** 2B4-NFAT-GFP reporter cells expressing WT Mincle or Mincle<sup>QPD</sup> (E169Q/N171D) mutant and BMM $\phi$  were prepared as previously described (Yamasaki et al., 2008). BMM $\phi$  pretreated with 10 ng/ml IFN- $\gamma$  for 4 h and reporter cells were stimulated with various bacteria and bacterial cell wall components. Activation of NFAT-GFP was monitored by flow cytometry. The levels of cytokines were determined by ELISA. NO production was measured by Griess assay.

**Antibodies.** Anti-Mincle mAbs were established as described previously (Yamasaki et al., 2008) and clone 1B6 (IgG1,  $\kappa$ ) was used in this study. HRP-conjugated anti-human IgG (109-035-088) was from Jackson Immuno-Research Laboratories.

**Ig fusion protein.** The extracellular domain of Mincle (a.a. 46–214) and Dectin2 (a.a. 43–209) was fused to hIgG1 Fc region and prepared as described previously (Yamasaki et al., 2009).

**Trehalase treatment.** 0.8  $\mu$ g TDM was coated on a 96-well plate as described in the Reagents section, followed by incubation with 1 to 10 mU/ml of Trehalase in 10 mM Tris-HCl buffer (pH 5.9) at 37°C for 6 h.

**Quantitative RT-PCR.** Total RNA was isolated from stimulated BMM $\phi$  or lungs of TDM-administrated mice subjected to real-time PCR (Applied Biosystems). Sequence of gene-specific primers is available upon request.

**Administration of TDM.** TDM was prepared as oil-in-water emulsion consisting of mineral oil (9%), Tween-80 (1%), and saline (90%) as previously described (Numata et al., 1985). 100  $\mu$ l of emulsion containing 150  $\mu$ g of TDM was injected intravenously into 6–11-wk-old mice. Emulsion without TDM was injected as a vehicle control. At day 7, thymocyte number was calculated and lungs were weighed and fixed in 10% formaldehyde for hematoxylin-eosin staining. A part of lungs was frozen for quantitative RT-PCR. LWI was calculated as described (Guidry et al., 2004). The number of granulomas was determined by counting focal mononuclear cell infiltrations in randomized 10 microscopic fields (0.8 mm<sup>2</sup>) per mouse.

**Statistics.** An unpaired two-tailed Student's *t* test was used for all the statistical analyses.

**Online supplemental material.** Fig. S1 shows schematic structure of PIM in *M. smegmatis* mutant. Fig. S2 shows schematic representation of hypothetical evolutionary struggle between mycobacteria and host immunity. Fig. S3 shows lack of TDM-induced pulmonary inflammation in Mincle<sup>-/-</sup> mice. Fig. S4 shows impaired TDM-induced thymic atrophy in Mincle<sup>-/-</sup> mice. Online supplemental material is available at <http://www.jem.org/cgi/content/full/jem.20091750/DC1>.

We thank Y. Fukui, H. Hara, R. Kakutani, and Y. Miyake for discussion; A. Suzuki, and M. Tanaka, H. Koseki, and T. Hasegawa for embryonic engineering; T. Saito for providing Fc $\gamma$ -deficient mice; Y. Esaki, A. Oyamada and M. Sakuma for technical

assistance; and Y. Nishi and H. Yamaguchi for secretarial assistance. We also thank N. Kinoshita (Laboratory for Technical Support, Medical Institute of Bioregulation, Kyushu University) for histochemical staining.

This work was supported by Grant-in-Aid for Young Scientists (S) and Takeda Science Foundation.

The authors have no conflicting financial interests.

Submitted: 12 August 2009

Accepted: 3 November 2009

## REFERENCES

- Adams, D.O. 1976. The granulomatous inflammatory response. A review. *Am. J. Pathol.* 84:164–192.
- Akira, S., S. Uematsu, and O. Takeuchi. 2006. Pathogen recognition and innate immunity. *Cell.* 124:783–801. doi:10.1016/j.cell.2006.02.015
- Asano, N., A. Kato, and K. Matsui. 1996. Two subsites on the active center of pig kidney trehalase. *Eur. J. Biochem.* 240:692–698. doi:10.1111/j.1432-1033.1996.0692h.x
- Azuma, I., and T. Seya. 2001. Development of immunoadjuvants for immunotherapy of cancer. *Int. Immunopharmacol.* 1:1249–1259. doi:10.1016/S1567-5769(01)00055-8
- Billiau, A., and P. Matthys. 2001. Modes of action of Freund's adjuvants in experimental models of autoimmune diseases. *J. Leukoc. Biol.* 70:849–860.
- Bloch, H. 1950. Studies on the virulence of tubercle bacilli; isolation and biological properties of a constituent of virulent organisms. *J. Exp. Med.* 91:197–218. doi:10.1084/jem.91.2.197
- Bowdish, D.M., K. Sakamoto, M.J. Kim, M. Kroos, S. Mukhopadhyay, C.A. Leifer, K. Tryggvason, S. Gordon, and D.G. Russell. 2009. MARCO, TLR2, and CD14 are required for macrophage cytokine responses to mycobacterial trehalose dimycolate and *Mycobacterium tuberculosis*. *PLoS Pathog.* 5:e1000474. doi:10.1371/journal.ppat.1000474
- Coulombe, F., M. Divangahi, F. Veyrier, L. de Léséleuc, J.L. Gleason, Y. Yang, M.A. Kelliher, A.K. Pandey, C.M. Sasseti, M.B. Reed, and M.A. Behr. 2009. Increased NOD2-mediated recognition of N-glycolyl muramyl dipeptide. *J. Exp. Med.* 206:1709–1716. doi:10.1084/jem.20081779
- Drickamer, K. 1992. Engineering galactose-binding activity into a C-type mannose-binding protein. *Nature.* 360:183–186. doi:10.1038/360183a0
- Fremont, C.M., V. Yeremeev, D.M. Nicolle, M. Jacobs, V.F. Quesniaux, and B. Ryffel. 2004. Fatal *Mycobacterium tuberculosis* infection despite adaptive immune response in the absence of MyD88. *J. Clin. Invest.* 114:1790–1799.
- Geijtenbeek, T.B., S.J. Van Vliet, E.A. Koppel, M. Sanchez-Hernandez, C.M. Vandembroucke-Grauls, B. Appelmelk, and Y. Van Kooyk. 2003. Mycobacteria target DC-SIGN to suppress dendritic cell function. *J. Exp. Med.* 197:7–17. doi:10.1084/jem.20021229
- Guidry, T.V., M. Olsen, K.S. Kil, R.L. Hunter Jr., Y.J. Geng, and J.K. Actor. 2004. Failure of CD1D<sup>-/-</sup> mice to elicit hypersensitive granulomas to mycobacterial cord factor trehalose 6,6'-dimycolate. *J. Interferon Cytokine Res.* 24:362–371. doi:10.1089/107999004323142222
- Hunter, R.L., M.R. Olsen, C. Jagannath, and J.K. Actor. 2006. Multiple roles of cord factor in the pathogenesis of primary, secondary, and cavitary tuberculosis, including a revised description of the pathology of secondary disease. *Ann. Clin. Lab. Sci.* 36:371–386.
- Indrigo, J., R.L. Hunter Jr., and J.K. Actor. 2002. Influence of trehalose 6,6'-dimycolate (TDM) during mycobacterial infection of bone marrow macrophages. *Microbiology.* 148:1991–1998.
- Jo, E.K. 2008. Mycobacterial interaction with innate receptors: TLRs, C-type lectins, and NLRs. *Curr. Opin. Infect. Dis.* 21:279–286. doi:10.1097/QCO.0b013e3282f88b5d
- Kang, P.B., A.K. Azad, J.B. Torrelles, T.M. Kaufman, A. Beharka, E. Tibesar, L.E. Desjardin, and L.S. Schlesinger. 2005. The human macrophage mannose receptor directs *Mycobacterium tuberculosis* lipoarabinomannan-mediated phagosomal biogenesis. *J. Exp. Med.* 202:987–999. doi:10.1084/jem.20051239
- Matsumoto, M., T. Tanaka, T. Kaisho, H. Sanjo, N.G. Copeland, D.J. Gilbert, N.A. Jenkins, and S. Akira. 1999. A novel LPS-inducible C-type lectin is a transcriptional target of NF-IL6 in macrophages. *J. Immunol.* 163:5039–5048.

- Matsunaga, I., T. Naka, R.S. Talekar, M.J. McConnell, K. Katoh, H. Nakao, A. Otsuka, S.M. Behar, I. Yano, D.B. Moody, and M. Sugita. 2008. Mycolyltransferase-mediated glycolipid exchange in *Mycobacteria*. *J. Biol. Chem.* 283:28835–28841. doi:10.1074/jbc.M805776200
- Moody, D.B., B.B. Reinhold, M.R. Guy, E.M. Beckman, D.E. Frederique, S.T. Furlong, S. Ye, V.N. Reinhold, P.A. Sieling, R.L. Modlin, et al. 1997. Structural requirements for glycolipid antigen recognition by CD1b-restricted T cells. *Science*. 278:283–286. doi:10.1126/science.278.5336.283
- Morita, Y.S., R. Velasquez, E. Taig, R.F. Waller, J.H. Patterson, D. Tull, S.J. Williams, H. Billman-Jacobe, and M.J. McConville. 2005. Compartmentalization of lipid biosynthesis in mycobacteria. *J. Biol. Chem.* 280:21645–21652. doi:10.1074/jbc.M414181200
- Morita, Y.S., C.B. Sena, R.F. Waller, K. Kurokawa, M.F. Sernee, F. Nakatani, R.E. Haites, H. Billman-Jacobe, M.J. McConville, Y. Maeda, and T. Kinoshita. 2006. PimE is a polyprenol-phosphate-mannose-dependent mannosyltransferase that transfers the fifth mannose of phosphatidylinositol mannoside in mycobacteria. *J. Biol. Chem.* 281:25143–25155. doi:10.1074/jbc.M604214200
- Noll, H., H. Bloch, J. Asselineau, and E. Lederer. 1956. The chemical structure of the cord factor of *Mycobacterium tuberculosis*. *Biochim. Biophys. Acta.* 20:299–309. doi:10.1016/0006-3002(56)90289-X
- Numata, F., K. Nishimura, H. Ishida, S. Ukei, Y. Tone, C. Ishihara, I. Saiki, I. Sekikawa, and I. Azuma. 1985. Lethal and adjuvant activities of cord factor (trehalose-6,6'-dimycolate) and synthetic analogs in mice. *Chem. Pharm. Bull. (Tokyo)*. 33:4544–4555.
- Ohno, H., S. Ono, N. Hirayama, S. Shimada, and T. Saito. 1994. Preferential usage of the Fc receptor  $\gamma$  chain in the T cell antigen receptor complex by  $\gamma/\delta$  T cells localized in epithelia. *J. Exp. Med.* 179:365–369. doi:10.1084/jem.179.1.365
- Otsuka, A., I. Matsunaga, T. Komori, K. Tomita, Y. Toda, T. Manabe, Y. Miyachi, and M. Sugita. 2008. Trehalose dimycolate elicits eosinophilic skin hypersensitivity in mycobacteria-infected guinea pigs. *J. Immunol.* 181:8528–8533.
- Ozeki, Y., H. Tsutsui, N. Kawada, H. Suzuki, M. Kataoka, T. Kodama, I. Yano, K. Kaneda, and K. Kobayashi. 2006. Macrophage scavenger receptor down-regulates mycobacterial cord factor-induced proinflammatory cytokine production by alveolar and hepatic macrophages. *Microb. Pathog.* 40:171–176. doi:10.1016/j.micpath.2005.12.006
- Park, S.Y., S. Ueda, H. Ohno, Y. Hamano, M. Tanaka, T. Shiratori, T. Yamazaki, H. Arase, N. Arase, A. Karasawa, et al. 1998. Resistance of Fc receptor-deficient mice to fatal glomerulonephritis. *J. Clin. Invest.* 102:1229–1238. doi:10.1172/JCI3256
- Rao, V., N. Fujiwara, S.A. Porcelli, and M.S. Glickman. 2005. *Mycobacterium tuberculosis* controls host innate immune activation through cyclopropane modification of a glycolipid effector molecule. *J. Exp. Med.* 201:535–543. doi:10.1084/jem.20041668
- Retzinger, G.S., S.C. Meredith, K. Takayama, R.L. Hunter, and F.J. Kézdy. 1981. The role of surface in the biological activities of trehalose 6,6'-dimycolate. Surface properties and development of a model system. *J. Biol. Chem.* 256:8208–8216.
- Robinson, M.J., D. Sancho, E.C. Slack, S. LeibundGut-Landmann, and C. Reis e Sousa. 2006. Myeloid C-type lectins in innate immunity. *Nat. Immunol.* 7:1258–1265. doi:10.1038/ni1417
- Rothfuchs, A.G., A. Bafica, C.G. Feng, J.G. Egen, D.L. Williams, G.D. Brown, and A. Sher. 2007. Dectin-1 interaction with *Mycobacterium tuberculosis* leads to enhanced IL-12p40 production by splenic dendritic cells. *J. Immunol.* 179:3463–3471.
- Sato, K., X.L. Yang, T. Yudate, J.S. Chung, J. Wu, K. Luby-Phelps, R.P. Kimberly, D. Underhill, P.D. Cruz Jr., and K. Ariizumi. 2006. Dectin-2 is a pattern recognition receptor for fungi that couples with the Fc receptor gamma chain to induce innate immune responses. *J. Biol. Chem.* 281:38854–38866. doi:10.1074/jbc.M606542200
- Schmidt, A. 1997. *Malassezia furfur*: a fungus belonging to the physiological skin flora and its relevance in skin disorders. *Cutis*. 59:21–24.
- Sueoka, E., S. Nishiwaki, S. Okabe, N. Iida, M. Suganuma, I. Yano, K. Aoki, and H. Fujiki. 1995. Activation of protein kinase C by mycobacterial cord factor, trehalose 6-monomycolate, resulting in tumor necrosis factor-alpha release in mouse lung tissues. *Jpn. J. Cancer Res.* 86:749–755.
- Tailleux, L., O. Schwartz, J.L. Herrmann, E. Pivert, M. Jackson, A. Amara, L. Legres, D. Dreher, L.P. Nicod, J.C. Gluckman, et al. 2003. DC-SIGN is the major *Mycobacterium tuberculosis* receptor on human dendritic cells. *J. Exp. Med.* 197:121–127. doi:10.1084/jem.20021468
- Welsh, K.J., A.N. Abbott, S.A. Hwang, J. Indrigo, L.Y. Armitage, M.R. Blackburn, R.L. Hunter Jr., and J.K. Actor. 2008. A role for tumour necrosis factor-alpha, complement C5 and interleukin-6 in the initiation and development of the mycobacterial cord factor trehalose 6,6'-dimycolate induced granulomatous response. *Microbiology*. 154:1813–1824. doi:10.1099/mic.0.2008/016923-0
- Werninghaus, K., A. Babiak, O. Gross, C. Hölscher, H. Dietrich, E.M. Agger, J. Mages, A. Mocsai, H. Schoenen, K. Finger, et al. 2009. Adjuvanticity of a synthetic cord factor analogue for subunit *Mycobacterium tuberculosis* vaccination requires FcR $\gamma$ -Syk-Card9-dependent innate immune activation. *J. Exp. Med.* 206:89–97. doi:10.1084/jem.20081445
- Winthrop, K.L. 2006. Risk and prevention of tuberculosis and other serious opportunistic infections associated with the inhibition of tumor necrosis factor. *Nat. Clin. Pract. Rheumatol.* 2:602–610. doi:10.1038/ncprheum0336
- Yadav, M., and J.S. Schorey. 2006. The beta-glucan receptor dectin-1 functions together with TLR2 to mediate macrophage activation by mycobacteria. *Blood*. 108:3168–3175. doi:10.1182/blood-2006-05-024406
- Yamaguchi, M., Y. Ogawa, K. Endo, H. Takeuchi, S. Yasaka, S. Nakamura, and Y. Yamamura. 1955. Experimental formation of tuberculous cavity in rabbit lung. IV. Cavity formation by paraffine oil extract prepared from heat killed tubercle bacilli. *Kekkaku*. 30:521–524.
- Yamasaki, S., E. Ishikawa, M. Sakuma, H. Hara, K. Ogata, and T. Saito. 2008. Mincle is an ITAM-coupled activating receptor that senses damaged cells. *Nat. Immunol.* 9:1179–1188. doi:10.1038/ni.1651
- Yamasaki, S., M. Matsumoto, O. Takeuchi, T. Matsuzawa, E. Ishikawa, M. Sakuma, H. Tateno, J. Uno, J. Hirabayashi, Y. Mikami, et al. 2009. C-type lectin Mincle is an activating receptor for pathogenic fungus, *Malassezia*. *Proc. Natl. Acad. Sci. USA*. 106:1897–1902. doi:10.1073/pnas.0805177106