

Memory B cells, but not long-lived plasma cells, possess antigen specificities for viral escape mutants

Whitney E. Purtha,¹ Thomas F. Tedder,⁴ Syd Johnson,⁵
Deepta Bhattacharya,¹ and Michael S. Diamond^{1,2,3}

¹Department of Pathology and Immunology, ²Department of Medicine, and ³Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO 63110

⁴Department of Immunology, Duke University Medical Center, Durham, NC 27710

⁵MacroGenics, Inc., Rockville, MD 20850

Memory B cells (MBCs) and long-lived plasma cells (LLPCs) persist after clearance of infection, yet the specific and nonredundant role MBCs play in subsequent protection is unclear. After resolution of West Nile virus infection in mice, we demonstrate that LLPCs were specific for a single dominant neutralizing epitope, such that immune serum poorly inhibited a variant virus that encoded a mutation at this critical epitope. In contrast, a large fraction of MBC produced antibody that recognized both wild-type (WT) and mutant viral epitopes. Accordingly, antibody produced by the polyclonal pool of MBC neutralized WT and variant viruses equivalently. Remarkably, we also identified MBC clones that recognized the mutant epitope better than the WT protein, despite never having been exposed to the variant virus. The ability of MBCs to respond to variant viruses *in vivo* was confirmed by experiments in which MBCs were adoptively transferred or depleted before secondary challenge. Our data demonstrate that class-switched MBC can respond to variants of the original pathogen that escape neutralization of antibody produced by LLPC without a requirement for accumulating additional somatic mutations.

CORRESPONDENCE

Michael S. Diamond:
diamond@borcim.wustl.edu
OR
Deepta Bhattacharya:
deeptab@pathology.wustl.edu

Abbreviations used: BCR, B cell receptor; CDR, complementarity-determining region; CHIKV, Chikungunya virus; DIII, domain III; JEV, Japanese encephalitis virus; LDA, limiting dilution analysis; LLPC, long-lived plasma cell; LR, lateral ridge; MBC, memory B cell; WNV, West Nile virus.

Long-lived plasma cells (LLPCs) constitutively secrete antibody to neutralize antigen immediately upon reinfection, whereas memory B cells (MBCs) produce antibody only upon restimulation by specific antigen (Manz et al., 1997; Slifka et al., 1998). Although preexisting serum antibody titers correlate with vaccine efficacy, the importance of the MBC response in conferring protection to reinfection has remained controversial (Zinkernagel and Hengartner, 2006). As both LLPC and MBC have been retained through mammalian evolution, these cell types must have distinct functions that remain to be fully delineated.

Hapten-based studies have reported that the affinities of LLPC B cell receptors (BCRs) are higher than those of MBC (Smith et al., 1997). Studies using fixed BCR demonstrated that cells with high-affinity antigen receptors were recruited preferentially to the LLPC pool (Phan et al., 2006). In contrast, MBC can be formed even when BCR affinities are low (Dal Porto et al., 2002). Because of the reduced stringency

of selection, the repertoire of the MBC pool may be more diverse and reactive to pathogen variants than a more oligoclonal LLPC population. Experiments with influenza and HIV have supported these predictions, although several of these studies suggest that MBCs achieve broader reactivity primarily through somatic hypermutation and affinity maturation that occurs after secondary encounter with heterologous viruses (Wrammert et al., 2008, 2011; Galli et al., 2009; Scheid et al., 2009).

One caveat to these conclusions is the recent finding that isotype-switched MBCs, which dominate the initial antibody recall response, have a limited ability to form new germinal centers and thus may be incapable of undergoing further affinity maturation (Dogan et al., 2009; Pape et al., 2011). Based on these studies, we

© 2011 Purtha et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

hypothesized that MBC, but not LLPC, would recognize efficiently variant antigens before secondary encounter with heterologous viruses. To test this, we used a mouse model of West Nile virus (WNV) infection with a WT and variant virus that differ only by a single amino acid in a dominantly neutralizing epitope. Our results suggest that LLPCs function to prevent secondary infection with homologous viruses, whereas MBCs generated during the primary response can recognize pathogen variants that escape neutralization of preexisting serum antibody produced by LLPCs.

RESULTS AND DISCUSSION

In mice, both B cells and antibody are required for survival after WNV infection (Diamond et al., 2003). Strongly neutralizing mouse monoclonal and polyclonal antibodies bind to the lateral ridge (LR) epitope on domain III (DIII; DIII-LR, residues K307, T330, and T332) of the viral E glycoprotein (Beasley and Barrett, 2002; Oliphant et al., 2005, 2007). Virtually all antibodies specific for this epitope can be identified by comparing binding to WT and loss-of-function proteins with mutations at two of the critical contact residues (K307E and T330I). We measured a significant reduction (7.4-fold, $P < 0.01$) in serum antibody recognition of DIII-K307E/T330I in comparison with WT DIII at days 30–100 after infection in WT C57BL/6 mice (Fig. 1 A). Serum antibody titers against this epitope were sustained as late as 500 d after infection, a time at which short-lived plasma cell-derived antibody titers wane, suggesting that the LLPC population was skewed toward recognizing the DIII-LR epitope.

To measure the frequency of WNV-specific LLPCs, we interrogated bone marrow cells from infected mice for their ability to secrete IgG that recognized WT and variant DIII.

We detected DIII-specific LLPCs at 100 d after infection by ELISPOT (Fig. 1 B) and, similar to the ELISA data with serum, a substantial (9.3-fold, $P < 0.01$) reduction in binding to the DIII-LR epitope was observed. As an independent measure of LLPC specificity, we analyzed bone marrow cells for intracellular binding to a bivalent fusion protein composed of WNV DIII or DIII-K307E/T330I and human $\gamma 1$ Fc (DIII-Fc_{Hu}, DIII-K307E/T330I Fc_{Hu}). We identified a small population of plasma cells that recognized DIII that was not present in mice infected with an unrelated virus (Fig. 1, C and D). Importantly, there were notably fewer LLPCs that recognized DIII-K307E/T330I compared with DIII ($P < 0.005$). Thus, after WNV infection in C57BL/6 mice, a significant fraction of LLPC expressed BCRs that are restricted to a single immunodominant epitope.

Given the epitope specificity of the LLPC response, we hypothesized that the antibody derived from these cells would strongly neutralize homologous WNV but might be impaired at neutralizing a variant (WNV-K307E) with a single mutation introduced at the dominant epitope. Convalescent serum from either WNV-infected or vaccinated mice was tested in neutralization assays with the homologous WNV-WT or variant WNV-K307E. With serum collected 60 d after WNV-WT infection, we observed a reduction (8.5-fold, $P < 0.05$) in the EC₅₀ value for WNV-K307E compared with WNV-WT (Fig. 2, A and B). Similar results were observed with serum from mice 60 d after immunization with a formalin-inactivated vaccine, with less efficient neutralization of the variant WNV-K307E virus (Fig. 2, A and B).

To confirm these results, we performed protection experiments by transferring immune serum from vaccinated mice immediately before challenge with WNV-WT or WNV-K307E. Transfer of 1 μ l of immune serum, but not




Figure 1. LLPCs recognize the K307/T330, but not a mutant K307E/T330I epitope.

(A) Mice were infected with WNV, and serum was collected at the indicated time points after infection. Levels of DIII or DIII-K307E/T330I-specific IgG were measured by ELISA. Endpoint titers are expressed as the reciprocal serum dilution that was 3 SD above background. The data reflect 6–10 mice per time point from four experiments. (B) Bone marrow cells were collected at the indicated time points after WNV infection and DIII or DIII-K307E/T330I-specific plasma cells were enumerated by ELISPOT assay. The data reflect six to nine mice per time point from four experiments. (C) Bone marrow cells were collected from mice infected with WNV or Chikungunya virus (CHIKV) and antigen-specific plasma cells were enumerated by intracellular staining. Data are shown as percentage of total

B220⁺ CD138⁺ cells. The percentage of cells staining as DIII or DIII-K307E/T330I-specific LLPC in CHIKV-infected mice was considered background and subtracted (0.1%). The data reflect five mice from three experiments. Horizontal bars in A–C indicate the mean. (D) Representative flow cytometry plots showing DIII or DIII-K307E/T330I-specific plasma cells from mice at 60 d after infection with WNV or CHIKV. Statistical significance (***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$) was determined by an unpaired, two-tailed Student's *t* test.




Figure 2. LLPCs weakly neutralize a variant WNV.

(A) Mice were infected with WNV or WNV-K307E (left) or immunized with an inactivated WNV vaccine (right), and sera were collected at day 60. Neutralizing antibody titers were measured by plaque reduction assay. Means and SD are shown of six experiments. (B) EC₅₀ titers were calculated from neutralization curves of WNV-WT or WNV-K307E incubated with serum from WNV-infected or vaccinated mice. Means and SD are shown from three or six experiments. The dashed line represents the limit of detection of assay. (C) Mice were injected with 1 μ l WNV immune serum or PBS before infection with WNV or WNV-K307E or PBS. Data reflect 20–26 mice per group in three experiments. Survival differences were determined by the log-rank test. NS, not significant. (D) Mice were injected with 1 μ l of immune or PBS immediately before infection with WNV (left) or WNV-K307E (right) mice. Viral RNA in serum was measured by qRT-PCR. Means and SD are shown of two experiments with five mice per group. Statistical significance (**, $P < 0.01$; *, $P < 0.05$) was determined by an unpaired, two-tailed Student's *t* test.

a significant decrease in binding to the K307E/T330I DIII mutant relative to the WT DIII protein ($P > 0.1$).

As an independent measure of MBC frequency and specificity, we determined the relative number of virus-specific MBC by flow cytometry. Immune splenocytes were stained for MBC markers (CD19, CCR6, and CD80; Bhattacharya et al., 2007) and DIII-Fc_{Hu} or DIII-K307E/T330I-Fc_{Hu}. We defined

a population of isotype-switched CCR6⁺CD80⁺ MBCs specific to DIII in WNV immune mice, but not in mice infected with an irrelevant virus (Fig. 3, B and C; and Fig. S1 A). The antigen specificity was confirmed by sorting DIII⁺ MBCs, activating them in vitro, and testing the supernatant by ELISA (Fig. S1 B). DIII⁺-sorted MBCs recognized both DIII and DIII-K307E/T330I, whereas DIII⁻ MBC from WNV-immune mice failed to produce antibody that recognized DIII. All DIII-specific MBCs were within the CCR6⁺CD80⁺ gate, as adoptive transfer of isotype-switched CD80⁻CCR6⁻ cells from WNV-immune mice or CD80⁺CCR6⁺ cells from naive mice failed to produce anti-DIII antibody upon challenge with WNV-WT (Fig. S1 C). The kinetics of antigen-specific MBC accumulation and maintenance was consistent in both the LDA and flow cytometric experiments (Fig. 3, A and B). In contrast to the LDA, we observed a small decrease by flow cytometry in the frequency of MBCs that recognized DIII-K307E/T330I at days 30 and 60 after infection. Nonetheless, the LDA and flow cytometry experiments both suggest that compared with the LLPC pool, MBCs contain a much larger frequency of B cell clones that recognize the mutant DIII-K307E/T330I. We calculated the frequency of cells that recognize DIII-K307E/T330I relative to DIII between MBCs and LLPCs. At each time point, MBCs, whether enumerated by LDA or flow cytometry, were enriched in antibodies that recognized DIII-K307E/T330I compared with LLPCs (Fig. 3 D). Therefore, cells in the MBC compartment

saline, into naive 5-wk-old mice protected from WNV-WT challenge (100 vs. 5% survival, $P < 0.001$; Fig. 2 C). In comparison, the same immune serum did not improve survival significantly after challenge with the variant virus WNV-K307E (56 vs. 35% survival, $P > 0.05$). WNV-K307E was mildly attenuated in mice, as reported previously (Zhang et al., 2009). In mice receiving immune serum, viremia was reduced at 36 h (10-fold, $P < 0.001$) and 72 h (11-fold, $P < 0.05$) after infection with WNV-WT (Fig. 2 D). However, the same serum failed to reduce viremia after WNV-K307E infection ($P > 0.2$). Thus, LLPC-derived antibody was inefficient at controlling infection with a variant WNV that was altered at a single amino acid within the dominant neutralizing epitope.

To evaluate the frequency of MBCs specific to the neutralizing epitope, we used two approaches: (1) limiting dilution analysis (LDA) of supernatant from cultures of stimulated MBCs and (2) direct enumeration of antigen-specific MBCs by flow cytometry. At different time points after WNV-WT infection, CD19⁺ B cells from the spleen were diluted serially, antibody secretion was induced (Aman and Slifka, 2006), and supernatants were tested for reactivity to DIII or DIII-K307E/T330I. At day 30 after infection, there were, on average, 26 DIII-specific MBCs per 10⁶ B cells (Fig. 3 A). This frequency increased to 57 at day 60 but waned to 7 DIII-specific MBCs per 10⁶ B cells by day 300. Next, we compared the relative frequencies of DIII and DIII-K307E/T330I-specific MBCs. In contrast to antibody produced by LLPCs, we did not measure




Figure 3. MBCs recognize mutant DIII-K307E/T330I epitopes. (A) Mice were infected with WNV, splenocytes were collected at the indicated time points, and CD19⁺ B cells were isolated by positive selection. The frequency of DIII- or DIII-K307E/T330I-specific IgG⁺ MBC was assessed by LDA. Virus-specific IgG in supernatants from LDA cultures were measured by ELISA and frequencies of virus-specific MBC determined by linear regression analysis. The limit of detection was ~2 MBCs per 10⁶ B cells (dashed line), and data reflect 5–14 mice per group from four experiments. P-values were determined using an unpaired, two-tailed Student's *t* test. (B) The frequency of splenic DIII- or DIII-K307E/T330I-specific MBC was assessed by flow cytometry. The percentage of cells staining as DIII⁺ or DIII-K307E/T330I⁺ MBC in CHIKV-infected mice was considered background and subtracted (6.1 and 5.4 MBCs per 10⁶ CD19⁺ cells). Data reflect 7–11 mice per group from five experiments and are displayed as frequency of antigen-specific MBCs per 10⁶ CD19⁺ cells. P-values were determined using an unpaired, two-tailed Student's *t* test. (C) Flow cytometry plots were from B and show MBC recognizing DIII or DIII-K307E/T330I from WNV- or CHIKV-infected mice. (D) Ratio of frequency of DIII-K307E/T330I- to DIII-specific MBC or LLPC in individual mice as measured by flow cytometry, LDA, and ELISPOT. MBC frequencies were shown in A and B, and LLPC frequencies were from Fig 1 (B and C). P-values were determined by the Mann-Whitney test (***, *P* < 0.001; **, *P* < 0.01; *, *P* < 0.05). Horizontal bars indicate the mean.

were more poised, compared with LLPCs, to recognize variant DIII carrying mutations that cause functional escape from neutralization.

Given the broader antigen specificity of the MBC population, we asked whether a subset of MBCs encoded antibody that preferentially recognized the variant epitope that was not present during primary infection. Remarkably, wells that contained single B cell clones as isolated by LDA revealed MBC specificities that recognized the mutant DIII-K307E/T330I but bound WT DIII so poorly that it was below the limit of

assay detection (Fig. 4 A). To corroborate these findings, we isolated single DIII-specific or DIII-K307E/T330I-specific MBCs by cell sorting and expanded them in culture on feeder cells expressing BAFF and CD40L. Although DIII-K307E/T330I-specific MBC clones were isolated (Fig. 4 B), others produced antibody that recognized DIII and DIII-K307E/T330I equivalently (Fig. 4 C). Moreover, one clone (2F2) bound the variant DIII-K307E/T330I protein better than the WT protein. To confirm this observation, the heavy and light chain genes of 2F2 were cloned and expressed ectopically.




Figure 4. MBC clones can recognize variant epitope better than WT DIII. (A) Percentage of MBC clones that bind DIII, both DIII and DIII-K307E/T330I, or DIII-K307E/T330I by ELISA. Positive-scoring wells of the lowest dilution from LDA in A were assigned specificities based on binding to DIII and DIII-K307E/T330I. Data represent ~9.8 clones per mouse from 8–14 mice (low frequency of MBC at day 300 precluded analysis). (B and C) Individual WNV-specific MBCs were sorted and cultured. Supernatant was tested for binding to DIII or DIII-K307E/T330I by ELISA to identify DIII-LR-specific (B) or DIII-cross-reactive (C) MBC. Data represent four independent experiments. (D) Clone 2F2 was expressed ectopically in 293T cells and supernatant was tested for binding to DIII or DIII-K307E/T330I. Representative results are shown, and statistical significance was determined using a paired, two-tailed Student's *t* test using data from four independent experiments performed in duplicate (*, *P* < 0.01).

Notably, 293T cells transfected with 2F2 heavy and light chain genes produced antibody that recognized DIII-K307E/T330I better than WT DIII (Fig. 4 D). These results suggest that antigen specificities can be generated within the MBC compartment through mutation in the primary germinal center reaction, which preferentially recognizes variant epitopes.

To define the diversity within the MBC compartment, we analyzed the immunoglobulin gene sequences of individual WNV-specific MBCs. The analysis of nine cross-reactive clones revealed vast sequence variation within the complementarity-determining regions (CDRs), with no apparent consensus for gene family or amino acid usage (Tables S1 and S2). Among 11 DIII-LR-specific clones, we noted some conservation of the light chain V κ 19-25 gene segment. For comparison, sequencing data from the previously generated DIII-LR-specific monoclonal antibodies E16, E24, and E34 was included (Oliphant et al., 2005). Still, heavy chain usage was variable, even at contact residues between E16 and K307/T330 previously identified by crystallographic analysis (Nybakken et al., 2005). Although a similar analysis on LLPCs was attempted, the low frequency ($\sim 1\%$) of antigen-specific cells, low levels of cell surface BCR, and nonproliferative nature of these cells precluded isolation of individual WNV-DIII-specific LLPC clones. Still, the selection of MBCs with divergent immunoglobulin gene sequences is consistent with the broad reactivity of this population.

Because MBCs were better able to recognize DIII-K307E/T330I than LLPCs were, we hypothesized that antibody derived from these cells would more efficiently neutralize WNV-K307E. To assess this, we tested the ability of antibody-containing supernatant from polyclonally stimulated MBCs to neutralize WT or variant WNV. Although the total antibody concentration in MBC culture supernatant was lower than in immune serum (~ 500 -fold less, not depicted), resulting in less inhibition, antibody produced by MBC neutralized WNV and WNV-K307E equivalently ($P > 0.9$; Fig. 5 A). This contrasts with the relatively poor neutralizing activity of LLPC-derived antibody against WNV-K307E (Fig. 2 A).

We used two approaches to assess the ability of MBCs to respond to heterologous infection. We adoptively transferred CD19⁺CCR6⁺CD80⁺IgM⁺IgD⁺lin⁺ MBCs from vaccinated C57BL/6J-IgH^a mice into antibody allotype-mismatched IgH^b C57BL/6 mice and challenged recipients with either WNV-WT or WNV-K307E. Using IgG2a⁺-specific secondary antibodies, we detected MBC-derived antibody by ELISA. Transferred MBCs preferentially produced DIII-LR-specific antibodies when mice were challenged with WNV-WT (Fig. 5 B, left), suggesting that homotypic MBCs recognizing the DIII-LR epitope outcompete the cross-reactive MBCs in a secondary response, perhaps as a result of higher affinity for antigen. In contrast, MBCs produced antibodies that recognized both DIII and DIII-K307E/T330I equivalently when recipient mice were challenged with WNV-K307E (Fig. 5 B, right).

To evaluate the ability of MBCs to recognize naturally occurring viral variants without relying on adoptive transfer experiments, we used a CD20 antibody that depletes MBCs but




Figure 5. MBCs can respond to and neutralize variant virus.

(A) Neutralization curves of WNV-WT or WNV-K307E with supernatant from stimulated MBCs. Data reflects four independent experiments performed in duplicate. Error bars indicate SD. (B) IgH^a MBCs (CD19⁺IgM⁺IgD⁺lin⁺CCR6⁺CD80⁺) from WNV-vaccinated mice were sorted and transferred into allotypic IgH^b recipients and challenged with WNV-WT or WNV-K307E 1 d later. Specificity of the MBC-derived antibody at different time points was determined by ELISA using an IgG2a⁺-specific detection antibody. Data reflects eight mice per group and is the average of five experiments. (C) Mice were immunized with JEV-DIII and treated with B cell-depleting (CD20 mAb) or an isotype control antibody. Serum was tested before WNV infection (left) and at days 3 and 8 (right) after infection for IgG binding to WNV-DIII by ELISA. The data reflect a total of five to nine mice per group from three experiments. P-values were determined using an unpaired, two-tailed Student's *t* test (*, $P < 0.01$). NS, indicates not significant.

does not affect LLPCs (Uchida et al., 2004; Ahuja et al., 2008; DiLillo et al., 2008). In the absence of MBCs, mice would be predicted to respond more poorly to viral variants. To test this, mice were immunized with DIII of a related flavivirus, Japanese encephalitis virus (JEV). Although JEV and WNV share $\sim 75\%$ amino acid sequence identity (Sumiyoshi et al., 1987), the cross-reactive DIII-specific antibody generated after immunization with JEV DIII is not directed at the DIII-LR epitope and does not efficiently neutralize WNV (unpublished data). 60 d after immunization with JEV-DIII, mice were administered CD20 or isotype control mAb. After 2 wk, the depletion of CD19⁺ B cells was confirmed by analysis of peripheral blood.

Mice were rested for 13 wk to allow restoration of naive CD19⁺ B cells before challenge with WNV. Antibody response to an irrelevant hapten was normal in these mice, demonstrating that the naive B cell pool had fully recovered, whereas in vitro MBC cultures revealed the continued absence of JEV-DIII-specific MBC at this time point (unpublished data).

To evaluate the contribution of MBCs to heterologous challenge, JEV-DIII-immunized control or CD20 mAb-treated mice were infected with WNV. Serum was tested before WNV infection (Fig. 5 C, left), at day 3, and at day 8 (Fig. 5 C, right) after infection for binding to WNV-DIII. Importantly, before and at day 3 after infection, both control and CD20 mAb-treated mice had equivalent levels of anti-WNV-DIII IgG antibody that was derived exclusively from the initial JEV-DIII immunization (Fig. 5 C, $P > 0.6$). Thus, normal antibody titers derived from LLPCs were present in the CD20 mAb-treated mice. However, CD20 mAb-treated mice exhibited a reduced anti-WNV-DIII IgG response at day 8 in comparison with control mice ($P < 0.001$). Thus, MBC generated by JEV-DIII immunization produced anti-WNV DIII antibody after heterologous challenge despite the presence of nonneutralizing antibody derived from LLPCs.

Our data support a model in which broadly reactive MBCs are generated upon primary pathogen exposure, expanded without a requirement for further somatic hypermutation after secondary challenge, and preferentially neutralize variant viruses better than LLPC-derived serum antibody. It also remains possible that IgM⁺ memory cells undergo additional somatic hypermutation as they differentiate into IgG⁺ plasmablasts. Consistent with this, antibody-secreting cells were reported to have increased numbers of somatic mutations relative to resting IgG⁺ MBC (Wrammert et al., 2008). More comprehensive comparisons of these cell types from the same individual are required to fully address this issue.

Analysis of selected HIV-infected individuals with low levels of viremia revealed MBC specificities with a diverse repertoire (Scheid et al., 2009). Indeed, our immunoglobulin gene sequencing results of DIII-specific MBCs with WNV confirm this. Thus, although the frequency of any given individual MBC that produces highly potent neutralizing antibody may be low, the breadth of the repertoire allows for recognition of divergent epitopes and viruses, resulting in a capacity for broad neutralization. Our results support this concept, as the supernatant from stimulated MBCs neutralized both WT and variant WNV. Nonetheless, only a limited number of MBC clones that produce potentially neutralizing antibodies and recognize a diverse array of HIV strains have been characterized (Walker et al., 2009; Corti et al., 2010; Wu et al., 2010).

The ability of MBCs to target variant epitopes was revealed in our studies with MBC clones that preferentially recognized the mutant compared with the WT epitope in DIII. By LDA, we calculated that ~10% of our DIII-specific MBC response recognized variant protein significantly better than WT protein. Our observation of MBCs preferentially recognizing a protein variant is analogous to that observed with the anti-HIV gp120 antibody HJ16, which recognizes a mutant (D368R) gp120

more strongly than the WT protein (Corti et al., 2010). This particular antibody is potentially neutralizing across different clades, suggesting that antibodies with variant specificities may be functionally relevant in the context of the breadth of the response.

MATERIALS AND METHODS

Viruses and quantification of viral burden. The WNV strain (3000.0259) was isolated in New York in 2000 and passaged once in C6/36 *Aedes albopictus* cells. Mice were inoculated subcutaneously in the footpad with 10² PFU of WNV diluted in Hanks balanced salt solution and 1% heat inactivated fetal bovine serum. The WNV-K307E strain was isolated as an in vivo escape mutant of the neutralizing humanized mAb E16 from brains of *RAG1*^{-/-} mice (Zhang et al., 2009) and was passaged once in C6/36 cells. CHIKV (La Reunion 2006-OPY1) was generated from an infectious clone (gift of S. Higgs, University of Texas Medical Branch, Galveston, TX) and passaged once in C6/36 cells. For viral RNA quantification, RNA was purified from serum samples using a Qia-Amp RNA recovery kit (QIAGEN) and copies were determined by TaqMan real-time reverse transcription PCR as previously described (Diamond et al., 2003).

Immunizations and immunodepletion. The West Nile Innovator vaccine was obtained commercially (Valley Vet Supply). Mice were administered two doses (100 µl each) of vaccine by intraperitoneal injection at days 0 and 1. DIII (residues 301–406) of JEV was expressed in *E. coli* using a pET21 vector and purified by oxidative refolding as previously described (Oliphant et al., 2005). Mice were immunized with 25 µg JEV-DIII and 20 µg CpG (Integrated DNA Technologies) in alum subcutaneously. For depletion of B cells, mice were administered 50 µg of an antibody against CD20 (MB20-11; Uchida et al., 2004) by retroorbital injection. Depletion was confirmed by flow cytometric analysis of CD19⁺Ter119⁻ cells from peripheral blood 2 wk after antibody treatment, and B cell reconstitution was confirmed at least 13 wk later.

Mice. C57BL/6 (WT and *RAG1*^{-/-}) and C57BL/6J-Igh-1^a Thy1^a Gpi1^a mice were purchased from The Jackson Laboratory. All mice were bred in the animal facilities of the Washington University School of Medicine under pathogen-free conditions, and experiments were performed in compliance with Washington University Animal Studies guidelines. This study was performed in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The protocol was approved by the Institutional Animal Care and Use Committee at the Washington University School of Medicine (Assurance Number: A3381-01).

Flow cytometry and cell sorting. Single cell suspensions were generated from spleen or bone marrow. Lymphocytes were isolated by differential centrifugation (850 g, 10 min at room temperature) through a ficoll gradient composed of Histopaque-1119 and Histopaque-1077 (Sigma-Aldrich). Cells were resuspended in PBS containing 2% FBS (Hyclone) and incubated on ice with the directly conjugated antibodies. MAbs were purified from the following hybridomas (Bio-X-Cell): 2C11 (anti-CD3), GK1.5 (anti-CD4), 53-6.7 (anti-CD8), 8C5 (anti-Gr-1), TER119 (anti-Ter119), and 1D3 (anti-CD19). Antibodies were conjugated to Pacific Blue succinimidyl ester (Invitrogen) according to the manufacturer's instructions. The following antibodies were purchased from eBioscience: anti-B220 (6B2) conjugated to PE-Cy7; anti-CD19 (ID3) conjugated to Alexa Fluor 700; anti-CD80 (16-10A1) conjugated to PE; and anti-IgD (11-26) and anti-GL7 (GL-7) conjugated to biotin. The following antibodies were purchased from BD: anti-IgM (11/41) conjugated to PE-Cy5.5; anti-CCR6 (140706) conjugated to Alexa Fluor 647; anti-CD138 (281-2) conjugated to biotin; and anti-CD138 (281-2) conjugated to PE. Streptavidin conjugated to Qdot-605 was purchased from Invitrogen. Cells were stained at 4°C for 30 min.

For staining of antigen-specific MBCs, cells were incubated with 0.8 µg of a chimeric fusion protein (DIII-Fc_{Hu}) of WNV-DIII (or a mutant DIII-K307E/T330I) linked to a human Fc-γ1 heavy chain with a point mutation in the N-linked glycosylation site that abolishes Fc-γR and C1q binding (Tao and Morrison, 1989). This protein was generated as follows: WNV DIII-Fc

(DIII residues 586–705) proteins were expressed in CHO-S cells (Life Technologies). The respective N297Q and K307E/T330I mutations were introduced by site directed mutagenesis using the QuikChange method (Agilent Technologies) in the mammalian expression vector pCIneo (Promega) before transfection. The mutated DIII-Fc proteins were then purified from conditioned medium by sequential protein A affinity and size exclusion chromatography. After binding of DIII-Fc_{Hu}, cells were washed and incubated with an Alexa Fluor 488–conjugated goat anti-human IgG antibody (Invitrogen). For intracellular staining of antigen-specific LLPCs, bone marrow cells were depleted of CD4, CD3, CD19, GR-1, and Ter119 using anti-rat IgG magnetic beads (Dynabeads; Invitrogen). Remaining cells were stained for surface expression of CD8, CD19, and B220 and CD138. Cells were washed, fixed with 4% paraformaldehyde for 10 min, and permeabilized with saponin buffer (Hanks balanced salt solution containing 10 mM Hepes and 0.1% wt/vol saponin) for 10 min. Intracellular DIII-specific antibody was detected using the same method as described for staining of antigen-specific MBCs.

For adoptive transfer experiments, non-B cell lineage cells and IgM⁺, IgD⁺, CD138⁺, and GL-7⁺ cells were depleted with magnetic beads (Dynabeads). CD19⁺CCR6⁺CD80⁺ class-switched IgM⁺IgD⁺ cells were sorted into media (DME supplemented with 10% FBS, 10 mM Hepes, 10 mM nonessential amino acids, 10 mM sodium pyruvate, and penicillin and streptomycin) using a FACSAria II (BD). Cells were transferred adoptively by retroorbital injection at a fixed 3:1 donor to recipient ratio resulting in 30,000–140,000 total memory cells per recipient.

Serological analysis. WNV-specific antibody titers in serum were calculated as previously described (Oliphant et al., 2007). In brief, 96-well plates (Nalge Nunc) were coated with recombinant protein (5 µg/ml DIII or DIII-K307E/T330I) overnight at 4°C. After blocking for 1 h with PBS supplemented with 2% BSA and 0.05% Tween 20 at 37°C, diluted serum samples were added to wells and incubated for 1 h at room temperature. Plates were washed with PBS with 0.05% Tween 20 and incubated with 1 µg/ml biotinylated anti-IgG (Sigma-Aldrich) for 1 h. After washing, 2 µg/ml streptavidin-conjugated horseradish peroxidase (Invitrogen) was added to each well and plates were incubated for 1 h. Plates were developed using tetramethylbenzidine substrate (Dako) and quenched with H₂SO₄. The optical density was read at 450 nm and endpoint dilutions were calculated as 3 SD above the background (BSA-coated wells).

ELISPOT assay. Mixed cellulose filter plates (Millipore) were coated initially with a capture antibody (100 µl WNV E111 Fab fragment [Sukupolvi-Petty et al., 2007] at 40 µg/ml in PBS for 2 h at 37°C), washed twice with 200 µl PBS, and then 40 µg/ml of recombinant WNV DIII protein in PBS was added overnight at 4°C. Plates were washed twice with PBS with 0.1% Tween-20, twice with PBS, and blocked for 1 h at room temperature with DME (Sigma-Aldrich) supplemented with 10% FBS, penicillin and streptomycin, 10 mM Hepes, 50 µM β-mercaptoethanol, and 10 mM nonessential amino acids. Bone marrow cells were depleted of erythrocytes by differential centrifugation as described in Flow cytometry and cell sorting, resuspended at 2 × 10⁷ cells/ml, and serially diluted into wells. After 5 h at 37°C, plates were washed three times with PBS and three times with PBS + 0.1% Tween-20 (PBST). Wells were incubated with 1 µg/ml biotinylated anti-IgG (Sigma-Aldrich) diluted in PBST with 1% FBS and incubated overnight at 4°C. After washing with PBST, wells were incubated with streptavidin conjugated to 1.25 µg/ml HRP diluted in PBST with 1% FBS. Plates were washed a final time with PBST and PBS. Spots were developed using a chromogen substrate (AEC [3-amino-9 ethyl-carbazole]; Sigma-Aldrich), and the reaction was quenched by thoroughly rinsing wells with water. Spots were enumerated using a stereomicroscope.

Neutralization assay. Plaque reduction neutralization assays on BHK21-15 cells were performed after mixing serial dilutions of serum with a fixed amount (10² PFU) of WNV as previously described (Diamond et al., 2003). For MBC neutralization assays, MBCs were cultured as described in the next section, and pooled supernatant was concentrated using a centrifugal filter device (Millipore) before mixing with virus. The concentration of IgG was determined by

quantitative ELISA to normalize data between experiments. Neutralization data were fit to a variable slope to calculate the EC₅₀ values by linear regression.

MBC LDA cultures. WNV-specific MBC frequencies were determined by measuring the reactivity of supernatants by LDA to recombinant protein by ELISA and was adapted from published protocols (Amanna and Slifka, 2006). Single cell splenocyte suspensions were generated from WNV-infected mice. Erythrocytes were lysed (ACK lysing buffer; Invitrogen) and CD19⁺ B cells isolated by positive selection using magnetic beads (Miltenyi Biotec). B cells were resuspended in LDA medium (RPMI, 10% FBS, 1× penicillin-streptomycin, 1 mM sodium pyruvate, 0.1 mM nonessential amino acids, 10 mM Hepes, and 50 µM β-mercaptoethanol). B cells were cultured in twofold serial dilutions starting at 50,000–120,000 cells per well in 96-well flat-bottom plates in the presence of *Staphylococcus aureus* Cowan strain (SAC) lysate (diluted 1:10,000; Sigma-Aldrich), 1 µg/ml pokeweed mitogen lectin (Sigma-Aldrich), 10 µg/ml lipopolysaccharide (from *E. coli* 0.111:B4; Sigma-Aldrich), 1 µg/ml phosphorothioated murine CpG (5'-TCCAT-GACGTTCTCTGATGCT-3'; Integrated DNA Technologies), 16 ng/ml mL-2, 10 ng/ml mL-6, 17 ng/ml mL-10 (PeproTech), and 5,000 cells per well of irradiated syngeneic splenocytes. Plates were sealed with an adhesive lid to prevent evaporation and incubated at 37°C and 5% humidified CO₂ for 6 d. To calculate the frequency of WNV-specific MBC that produced IgG, half-area 96-well plates (Corning-Costar) were coated with 5 µg/ml WNV E protein overnight at 4°C. Plates were washed and blocked with PBS-T for 1 h at 37°C. Supernatants from LDA plates were added to the ELISA plates (15 µl per well), and detection of WNV-specific antibody was conducted as described in ELISPOT assay. Positive wells were defined as wells that scored twofold over the mean OD of negative control wells (wells containing irradiated splenocytes and supplements alone). Supernatant from cultured naive splenocytes did not score positive, and only LDA that contained at least two positive wells at the highest dilution were analyzed.

Single-cell MBC cultures and reexpression of immunoglobulin genes. NIH 3T3 cells (American Type Culture Collection) were retrovirally transduced with a murine stem cell retrovirus (Bhattacharya et al., 2002) expressing CD40L-IRES-GFP or BAFF-IRES-human CD4. Antigen-specific MBCs were sorted using a FACSAria II into individual wells of a flat-bottom 96-well plate containing LDA medium supplemented as described in the previous section. Each well was seeded with 5,000 mitomycin C-treated BAFF⁺CD40L⁺ NIH 3T3 cells. After 6 d, supernatant was harvested and tested for reactivity to DIII or DIII-K307E/T330I by ELISA and total RNA was extracted with Trizol (Invitrogen). cDNA was generated, and immunoglobulin heavy and light chain genes were amplified by PCR and sequenced as previously described (Tiller et al., 2009). Sequences were analyzed for immunoglobulin genes using IgBlast. Primers for specific Vh, Vκ, Jh, and Jκ segments were used to amplify variable regions for both heavy and light chain genes (Tiller et al., 2009), and these segments were cloned in-frame into pEF1α-myc-His expression vectors (Invitrogen) downstream of a consensus immunoglobulin leader sequence and upstream of either the mouse Cγ1 or Cκ constant region. Six individual clones of both heavy and light chain genes were resequenced, and individual heavy and light chain expression constructs containing the consensus sequence without point mutations were cotransfected into 293T cells by calcium phosphate transfection. Culture supernatant was harvested 3 d after transfection for ELISA assays.

Online supplemental material. Supplemental material shows the gating and sorting strategy of splenic MBC ex vivo (Fig. S1, A and B) and adoptive transfer studies in *RAG1*^{-/-} mice (Fig. S1C). Gene family usage (Table S1) and immunoglobulin CDR sequences (Table S2) from WNV-specific MBC clones also are described. Online supplemental material is available at <http://www.jem.org/cgi/content/full/jem.20110740/DC1>.

We thank M. Slifka and I. Amanna for technical advice, P. Pal for preparing the CHIKV stock, and M. Slifka critical comments on the manuscript.

This work was supported by grants from the National Institutes of Health (the Midwest and Southeastern Regional Centers of Excellence for Biodefense and Emerging Infectious Diseases Research; U54 AI057160, M.S. Diamond and D. Bhattacharya; U54 AI057157, T.F. Tedder; R01 AI56363, M.S. Diamond; and U01 AI082196, M.S. Diamond). This research was supported in part by Children's Discovery Institute grant PD-II-2011-116 (D. Bhattacharya).

S. Johnson is an employee of MacroGenics, which has licensed monoclonal antibodies against WNV from Washington University for possible commercial use. M. Diamond is a paid consultant for MacroGenics. The authors have no other competing financial interests.

Submitted: 13 April 2011

Accepted: 16 November 2011

REFERENCES

- Ahuja, A., S.M. Anderson, A. Khalil, and M.J. Shlomchik. 2008. Maintenance of the plasma cell pool is independent of memory B cells. *Proc. Natl. Acad. Sci. USA*. 105:4802–4807. <http://dx.doi.org/10.1073/pnas.0800555105>
- Amanna, I.J., and M.K. Slifka. 2006. Quantitation of rare memory B cell populations by two independent and complementary approaches. *J. Immunol. Methods*. 317:175–185. <http://dx.doi.org/10.1016/j.jim.2006.09.005>
- Beasley, D.W., and A.D. Barrett. 2002. Identification of neutralizing epitopes within structural domain III of the West Nile virus envelope protein. *J. Virol.* 76:13097–13100. <http://dx.doi.org/10.1128/JVI.76.24.13097-13100.2002>
- Bhattacharya, D., D.U. Lee, and W.C. Sha. 2002. Regulation of Ig class switch recombination by NF- κ B: retroviral expression of RelB in activated B cells inhibits switching to IgG1, but not to IgE. *Int. Immunol.* 14:983–991. <http://dx.doi.org/10.1093/intimm/14.6.983>
- Bhattacharya, D., M.T. Cheah, C.B. Franco, N. Hosen, C.L. Pin, W.C. Sha, and I.L. Weissman. 2007. Transcriptional profiling of antigen-dependent murine B cell differentiation and memory formation. *J. Immunol.* 179:6808–6819.
- Corti, D., J.P. Langedijk, A. Hinz, M.S. Seaman, F. Vanzetta, B.M. Fernandez-Rodriguez, C. Silacci, D. Pinna, D. Jarrossay, S. Balla-Jhaghoorsingh, et al. 2010. Analysis of memory B cell responses and isolation of novel monoclonal antibodies with neutralizing breadth from HIV-1-infected individuals. *PLoS ONE*. 5:e8805. <http://dx.doi.org/10.1371/journal.pone.0008805>
- Dal Porto, J.M., A.M. Haberman, G. Kelsoe, and M.J. Shlomchik. 2002. Very low affinity B cells form germinal centers, become memory B cells, and participate in secondary immune responses when higher affinity competition is reduced. *J. Exp. Med.* 195:1215–1221. <http://dx.doi.org/10.1084/jem.20011550>
- Diamond, M.S., B. Shrestha, A. Marri, D. Mahan, and M. Engle. 2003. B cells and antibody play critical roles in the immediate defense of disseminated infection by West Nile encephalitis virus. *J. Virol.* 77:2578–2586. <http://dx.doi.org/10.1128/JVI.77.4.2578-2586.2003>
- DiLillo, D.J., Y. Hamaguchi, Y. Ueda, K. Yang, J. Uchida, K.M. Haas, G. Kelsoe, and T.F. Tedder. 2008. Maintenance of long-lived plasma cells and serological memory despite mature and memory B cell depletion during CD20 immunotherapy in mice. *J. Immunol.* 180:361–371.
- Dogan, I., B. Bertocci, P. Vilmont, F. Delbos, J. Mègret, S. Storck, C.A. Reynaud, and J.C. Weill. 2009. Multiple layers of B cell memory with different effector functions. *Nat. Immunol.* 10:1292–1299. <http://dx.doi.org/10.1038/ni.1814>
- Galli, G., K. Hancock, K. Hoshler, J. DeVos, M. Praus, M. Bardelli, C. Malzone, F. Castellino, C. Gentile, T. McNally, et al. 2009. Fast rise of broadly cross-reactive antibodies after boosting long-lived human memory B cells primed by an MF59 adjuvanted prepandemic vaccine. *Proc. Natl. Acad. Sci. USA*. 106:7962–7967. <http://dx.doi.org/10.1073/pnas.0903181106>
- Manz, R.A., A. Thiel, and A. Radbruch. 1997. Lifetime of plasma cells in the bone marrow. *Nature*. 388:133–134. <http://dx.doi.org/10.1038/40540>
- Nybakken, G.E., T. Oliphant, S. Johnson, S. Burke, M.S. Diamond, and D.H. Fremont. 2005. Structural basis of West Nile virus neutralization by a therapeutic antibody. *Nature*. 437:764–769. <http://dx.doi.org/10.1038/nature03956>
- Oliphant, T., M. Engle, G.E. Nybakken, C. Doane, S. Johnson, L. Huang, S. Gorlatov, E. Mehlhop, A. Marri, K.M. Chung, et al. 2005. Development of a humanized monoclonal antibody with therapeutic potential against West Nile virus. *Nat. Med.* 11:522–530. <http://dx.doi.org/10.1038/nm1240>
- Oliphant, T., G.E. Nybakken, S.K. Austin, Q. Xu, J. Bramson, M. Loeb, M. Thorsby, D.H. Fremont, T.C. Pierson, and M.S. Diamond. 2007. Induction of epitope-specific neutralizing antibodies against West Nile virus. *J. Virol.* 81:11828–11839. <http://dx.doi.org/10.1128/JVI.00643-07>
- Pape, K.A., J.J. Taylor, R.W. Maul, P.J. Gearhart, and M.K. Jenkins. 2011. Different B cell populations mediate early and late memory during an endogenous immune response. *Science*. 331:1203–1207. <http://dx.doi.org/10.1126/science.1201730>
- Phan, T.G., D. Paus, T.D. Chan, M.L. Turner, S.L. Nutt, A. Basten, and R. Brink. 2006. High affinity germinal center B cells are actively selected into the plasma cell compartment. *J. Exp. Med.* 203:2419–2424. <http://dx.doi.org/10.1084/jem.20061254>
- Scheid, J.F., H. Mouquet, N. Feldhahn, M.S. Seaman, K. Velinzon, J. Pietzsch, R.G. Ott, R.M. Anthony, H. Zebroski, A. Hurley, et al. 2009. Broad diversity of neutralizing antibodies isolated from memory B cells in HIV-infected individuals. *Nature*. 458:636–640. <http://dx.doi.org/10.1038/nature07930>
- Slifka, M.K., R. Antia, J.K. Whitmire, and R. Ahmed. 1998. Humoral immunity due to long-lived plasma cells. *Immunity*. 8:363–372. [http://dx.doi.org/10.1016/S1074-7613\(00\)80541-5](http://dx.doi.org/10.1016/S1074-7613(00)80541-5)
- Smith, K.G., A. Light, G.J. Nossal, and D.M. Tarlinton. 1997. The extent of affinity maturation differs between the memory and antibody-forming cell compartments in the primary immune response. *EMBO J.* 16:2996–3006. <http://dx.doi.org/10.1093/emboj/16.11.2996>
- Sukupolvi-Petty, S., S.K. Austin, W.E. Purtha, T. Oliphant, G.E. Nybakken, J.J. Schlesinger, J.T. Roehrig, G.D. Gromowski, A.D. Barrett, D.H. Fremont, and M.S. Diamond. 2007. Type- and subcomplex-specific neutralizing antibodies against domain III of dengue virus type 2 envelope protein recognize adjacent epitopes. *J. Virol.* 81:12816–12826. <http://dx.doi.org/10.1128/JVI.00432-07>
- Sumiyoshi, H., C. Mori, I. Fuke, K. Morita, S. Kuhara, J. Kondou, Y. Kikuchi, H. Nagamatsu, and A. Igarashi. 1987. Complete nucleotide sequence of the Japanese encephalitis virus genome RNA. *Virology*. 161:497–510. [http://dx.doi.org/10.1016/0042-6822\(87\)90144-9](http://dx.doi.org/10.1016/0042-6822(87)90144-9)
- Tao, M.H., and S.L. Morrison. 1989. Studies of aglycosylated chimeric mouse-human IgG. Role of carbohydrate in the structure and effector functions mediated by the human IgG constant region. *J. Immunol.* 143:2595–2601.
- Tiller, T., C.E. Busse, and H. Wardemann. 2009. Cloning and expression of murine Ig genes from single B cells. *J. Immunol. Methods*. 350:183–193. <http://dx.doi.org/10.1016/j.jim.2009.08.009>
- Uchida, J., Y. Lee, M. Hasegawa, Y. Liang, A. Bradney, J.A. Oliver, K. Bowen, D.A. Steeber, K.M. Haas, J.C. Poe, and T.F. Tedder. 2004. Mouse CD20 expression and function. *Int. Immunol.* 16:119–129. <http://dx.doi.org/10.1093/intimm/16.2.dxh009>
- Walker, L.M., S.K. Phogat, P.Y. Chan-Hui, D. Wagner, P. Phung, J.L. Goss, T. Wrinn, M.D. Simek, S. Fling, J.L. Mitcham, et al. 2009. Broad and potent neutralizing antibodies from an African donor reveal a new HIV-1 vaccine target. *Science*. 326:285–289. <http://dx.doi.org/10.1126/science.1178746>
- Wrammert, J., K. Smith, J. Miller, W.A. Langley, K. Kokko, C. Larsen, N.Y. Zheng, I. Mays, L. Garman, C. Helms, et al. 2008. Rapid cloning of high-affinity human monoclonal antibodies against influenza virus. *Nature*. 453:667–671. <http://dx.doi.org/10.1038/nature06890>
- Wrammert, J., D. Koutsouanos, G.M. Li, S. Edupuganti, J. Sui, M. Morrissey, M. McCausland, I. Skountzou, M. Hornig, W.I. Lipkin, et al. 2011. Broadly cross-reactive antibodies dominate the human B cell response against 2009 pandemic H1N1 influenza virus infection. *J. Exp. Med.* 208:181–193. <http://dx.doi.org/10.1084/jem.20101352>
- Wu, X., Z.Y. Yang, Y. Li, C.M. Hogerkorp, W.R. Schief, M.S. Seaman, T. Zhou, S.D. Schmidt, L. Wu, L. Xu, et al. 2010. Rational design of envelope identifies broadly neutralizing human monoclonal antibodies to HIV-1. *Science*. 329:856–861. <http://dx.doi.org/10.1126/science.1187659>
- Zhang, S., M.R. Vogt, T. Oliphant, M. Engle, E.I. Bovshik, M.S. Diamond, and D.W. Beasley. 2009. Development of resistance to passive therapy with a potentially neutralizing humanized monoclonal antibody against West Nile virus. *J. Infect. Dis.* 200:202–205. <http://dx.doi.org/10.1086/599794>
- Zinkernagel, R.M., and H. Hengartner. 2006. Protective ‘immunity’ by pre-existent neutralizing antibody titers and preactivated T cells but not by so-called ‘immunological memory’. *Immunol. Rev.* 211:310–319. <http://dx.doi.org/10.1111/j.0105-2896.2006.00402.x>