

S1P and the birth of platelets

Timothy Hla, Sylvain Galvani, Shahin Rafii, and Ralph Nachman

Recent work has highlighted the multitude of biological functions of sphingosine 1-phosphate (S1P), which include roles in hematopoietic cell trafficking, organization of immune organs, vascular development, and neuroinflammation. Indeed, a functional antagonist of S1P₁ receptor, FTY720/Gilenya, has entered the clinic as a novel therapeutic for multiple sclerosis. In this issue of the *JEM*, Zhang et al. highlight yet another function of this lipid mediator: thrombopoiesis. The S1P₁ receptor is required for the growth of proplatelet strings in the bloodstream and the shedding of platelets into the circulation. Notably, the sharp gradient of S1P between blood and the interstitial fluids seems to be essential to ensure the production of platelets, and S1P appears to cooperate with the CXCL12–CXCR4 axis. Pharmacologic modulation of the S1P₁ receptor altered circulating platelet numbers acutely, suggesting a potential therapeutic strategy for controlling thrombocytopenic states. However, the S1P₄ receptor may also regulate thrombopoiesis during stress-induced accelerated platelet production. This work reveals a novel physiological action of the S1P/S1P₁ duet that could potentially be harnessed for clinical translation.

S1P and immune/hematopoietic cell trafficking

The lipid mediator S1P has received significant attention as an extracellular factor that signals via G protein–coupled receptors to regulate lymphocyte trafficking and vascular function (Blaho and Hla, 2011; Cyster and Schwab, 2012; Obinata and Hla, 2012). It is now established that an S1P gradient exists between vascular and nonvascular compartments (Schwab et al., 2005; Hla et al., 2008). High levels of S1P are found in blood and lymph, whereas the actions of degradative enzymes in tissue compartments such as secondary lymphoid organs keep its concentration low. This S1P gradient is necessary for the directional egress of lymphocytes from secondary lymphoid organs and the thymus into the circulatory system. Interference with the gradient by pharmacologic or genetic manipulation of

S1P lyase (Schwab et al., 2005), genetic knockout of sphingosine kinases (Pham et al., 2010), S1P transporter (Spns2; Fukuhara et al., 2012), or the lipid phosphate phosphatase-3 (LPP3; Bréart et al., 2011) results in the attenuation of lymphocyte egress, and lymphopenia. Similar mechanisms may also be operative in the trafficking of dendritic cells, NKT cells, and hematopoietic progenitor cells (Massberg et al., 2007; Cyster and Schwab, 2012). These findings attest to the generality of the S1P gradient-dependent trafficking paradigm for lymphocytes and other hematopoietic cells.

Detailed investigations of S1P receptors have also shown that S1P₁ receptor on immune cells is required for ligand-dependent egress (Grigorova et al., 2009; Allende et al., 2010). The rate of egress is determined by the net effect between retention signals and egress signals. In the case of lymph node–resident T cells, a key retention signal is determined by the chemokine CCL21 signaling via CCR7 (Pham et al., 2008). Activation of S1P₁ by S1P appears to be the only egress signal identified to date. Intravital two-photon fluorescence microscopy studies have shown that S1P₁ signaling on immune

cells allows probing of the endothelial lining of cortical sinuses with cellular processes that ultimately allows productive egress. The endothelial S1P₁ receptor appears to be dispensable for egress, whereas plasma membrane residence of S1P₁ on lymphocytes is one of the key factors that determine egress rates (Thangada et al., 2010). It is likely that such mechanisms are applicable to many situations in which hematopoietic cells traffic into the S1P-rich environments via trans-endothelial egress.

S1P and platelets

Early work identified that S1P is released by activated platelets stimulated with thrombin or ADP (Yatomi et al., 1995, 1997). Platelets carry endothelial cell-protective cargo (trophogens) such as platelet-derived growth factor (PDGF), vascular endothelial growth factor, CXCL12, fibroblast growth factor (FGF), and stem cell factor, among others, and thrombocytopenic states lead to vascular endothelial dysfunction and breach of vascular barrier (Nachman and Rafii, 2008). S1P seems to “nourish” the endothelium, supporting the integrity of the vascular bed by activating endothelial S1P₁ receptors. In addition, platelets also express S1P receptors; however, their role in platelet biology has remained elusive.

Zhang et al. (2012) report that S1P signaling via its multifunctional receptor S1P₁ is important in platelet production from megakaryocytes. Even though multiple S1P receptors are expressed in megakaryocytes (i.e., S1P_{1,2,4}), S1P₁ is unique in that it is required for two specific events important in platelet formation and release.

T. Hla and S. Galvani are at the Center for Vascular Biology, Department of Pathology and Laboratory Medicine; S. Rafii is at the Ansary Stem Cell Center, Department of Genetic Medicine; and R. Nachman is at the Department of Medicine Weill Cornell Medical College, Cornell University, New York, NY 10065.

CORRESPONDENCE
T. Hla: th2002@med.cornell.edu

The positioning of megakaryocytes to the endothelial lining of the bone marrow sinusoids via the VCAM1/VLA4 adhesion pair is known to be critical for thrombopoiesis (Hamada et al., 1998; Majka et al., 2000; Avecilla et al., 2004; Schulze et al., 2006). In S1P₁ knockout megakaryocytes, the positioning itself was not altered; however, the directional migration of proplatelet-containing cytoplasmic extensions into the circulatory compartment was inhibited. These data, coupled with in vitro studies using S1P gradients, suggest that compartmentalized S1P₁ signaling is important for directional growth of proplatelet-containing megakaryocyte processes. S1P₁ signaling is also required for the shedding of proplatelets in a Rac-dependent manner. Because it is well established that S1P₁ couples to the G_i-dependent Rac activation (Lee et al., 2001), the findings suggest that active signaling by this S1P receptor is required to complete the final stages of thrombopoiesis (Fig. 1). The importance of this pathway was demonstrated in the hematopoietic-specific *S1pr1* knockout mice, which showed severe thrombocytopenia.

These findings also highlight the cooperative action of different GPCRs in megakaryocytes in ensuring optimal thrombopoiesis. Previous studies have determined that endothelial cell expression of CXCL12 and its action on megakaryocytes via CXCR4 GPCR is important for the interaction and positioning of the mature megakaryocytes in their proper vascular niche (Avecilla et al., 2004). Indeed, provision of CXCL12 and FGF-4 (another endothelial-active cytokine; Konishi et al., 1996) was able to support platelet formation even in thrombopoietin knockout mice. Thus, CXCR4 supports megakaryocyte interaction and positioning at the vascular niche, whereas S1P₁ supports polarized proplatelet process formation and release into the circulation.

The intracellular signaling mechanisms used by CXCR4 in megakaryocytes to allow interaction with endothelial cells are not well understood. The CXCR4 receptor can activate multiple G proteins, such as G_i, G_q, and G_{12/13} (Alkhatib, 2009). How such pathways lead to endothelial–

megakaryocyte interactions is not well understood. However, S1P₁ is known to activate the G_i pathway exclusively (Windh et al., 1999). This results in activation of Rac-dependent cortical actin assembly (Lee et al., 2001). In addition to inducing actin cytoskeleton rearrangement, the S1P₁–Rac pathway also potently induces microtubule dynamics (Paik et al., 2004; Obinata and Hla, 2012). Therefore, S1P₁-dependent Rac activation is critical for process extension and the release of proplatelets. Indeed, a small molecule inhibitor of Rac potently blocked platelet release.

A previous study examined S1P₄, another megakaryocyte-expressed S1P receptor that possesses different signaling properties (Golffier et al., 2010). This receptor is strongly induced in megakaryocyte differentiation, but upon gene deletion, platelet numbers were not altered. However, stress-induced thrombo-

poiesis was slightly delayed in *S1pr4* KO mice in this study, suggesting a possible function under accelerated platelet generation. Further, a significant number of *S1pr4* KO megakaryocytes exhibited abnormal cellular morphology characterized by cytoplasmic vacuolation and nuclear ploidy changes. In contrast, *S1pr4* KO megakaryocytes did not exhibit alterations in proplatelet generation in vitro. Thus, S1P₄ may also have a role in thrombopoiesis, even though its exact significance in physiological and stress-induced thrombopoiesis needs further elucidation.

Recent studies also show that S1P₁ is intimately involved in flow-dependent signal transduction in the endothelium (Jung et al., 2012). In vascular endothelial cells, S1P₁ is necessary for shear stress-induced signaling events, which culminate in the stabilization of newly formed vascular networks (Gaengel et al., 2012; Jung et al., 2012). Notably, S1P₁ GPCR




Figure 1. S1P₁ receptor on megakaryocytes is required for thrombopoiesis. The S1P₁ receptor, which activates the G_i protein, Ras GTPase, PI-3-kinase (PI3K), and phospholipase C (PLC) pathways, regulates the formation of proplatelet-containing cytoplasmic protrusions and release of platelet fragments. CXCR4 expression is required to position mature megakaryocytes in the appropriate vascular niche for platelet formation, and S1P₁ receptor is essential for process formation and proplatelet release. Both actin-based and microtubule cytoskeleton changes may be required for such events, which are likely to require both plasma-derived S1P and shear forces exerted by blood flow. The S1P₄ receptor may also regulate thrombopoiesis because it is also highly expressed in megakaryocytes. However, endothelial S1P₁ is essential for vascular stability and homeostasis.

can signal in response to laminar shear stress in a ligand-independent manner. Either the S1P₁ GPCR itself contains a mechanosensitive domain or is capable of associating with a mechanosensor, thus promoting signal transduction in a ligand-independent manner (Jung et al., 2012). Therefore, it is possible that proplatelet release from the transendothelial processes into the circulation requires S1P-dependent and flow-dependent mechanisms.

S1P therapeutics and potential modulation of platelet biology

The ability of the S1P–S1P₁ axis to regulate immune cell trafficking was harnessed in the novel treatment paradigm for multiple sclerosis (MS), in which autoreactive immune cells migrate into the central nervous system (CNS) and destroy myelin-containing axons, leading to astrogliosis and neuronal deficit. Treatment of MS patients with the S1P₁ receptor antagonist Fingolimod (a.k.a., FTY720/Gilenya) resulted in the disruption of normal trafficking patterns as indicated by the reduced numbers of circulating central memory type T cells that are IL-17⁺ (LaMontagne et al., 2006; Brinkmann et al., 2010; Cohen et al., 2010). In mouse models of experimental autoimmune encephalomyelitis, S1P₁ receptor inhibitors induced profound lymphopenia and reduced penetration of inflammatory cells into the CNS (Chun and Hartung, 2010). Fingolimod and related S1P₁ receptor-targeting drugs are functional antagonists; even though they act as agonists upon initial binding to S1P₁, they induce irreversible receptor internalization, resulting in a reduced plasma membrane residence of S1P₁ (Oo et al., 2011). Zhang et al. (2012) show that FTY720 administration causes a rapid increase in platelet numbers in mice, suggesting that acute agonistic action of FTY720 on the megakaryocyte S1P₁ receptors induced platelet release. Thus, it may be possible to therapeutically regulate platelet deficiencies by targeting S1P₁. However, because this pathway influences immune cell trafficking and vascular endothelial cell function, mechanism-based potential adverse events should be considered for translational approaches.

Several clinical conditions are associated with thrombocytopenia. In various infectious conditions, such as sepsis and Dengue hemorrhagic fever, platelet counts are markedly reduced and pose a significant risk for hemorrhage. Thus, in many hematological malignancies, as well as after administration of bone marrow cytotoxic therapies, the ability to increase platelet counts acutely may be useful. Thus, activation of this pathway with a long-lasting agonist of S1P₁ may be beneficial not only in increasing platelet counts but also in preserving endothelial function. It is important to note that current S1P₁ receptor modulators act as functional antagonists because of their ability to induce irreversible receptor internalization (Oo et al., 2011). In this scenario, such compounds are unlikely to be effective inducers of platelet formation. Thus, a new generation of S1P₁ agonists will need to be developed to therapeutically harness this system. Because the S1P pathway is involved in the terminal steps of thrombopoiesis, it is likely that S1P₁ agonists will have to be used in conjunction with other inducers of thrombopoiesis, for example, thrombopoietin, a key megakaryocyte differentiation factor.

These recent findings of Zhang et al. (2012) have highlighted a novel function of the lipid mediator S1P that signals through its multifunctional receptor S1P₁. This pathway may be potentially useful in therapeutic modulation of thrombocytopenia. However, because of the multitude of biological systems that S1P₁ regulates, any therapeutic strategy will need to consider possible adverse events. Novel agents that selectively target S1P₁ receptors in a cell- or tissue-specific manner will likely be needed to fully harness this potential translational opportunity.

This work is supported by National Institutes of Health grants HL67330, HL89934 and HL70694 (TH).

REFERENCES

Alkhatib, G. 2009. The biology of CCR5 and CXCR4. *Curr. Opin. HIV AIDS.* 4:96–103. <http://dx.doi.org/10.1097/COH.0b013e328324bbec>

Allende, M.L., G. Tuymetova, B.G. Lee, E. Bonifacino, Y.P. Wu, and R.L. Proia. 2010. S1P₁ receptor directs the release of immature B cells from bone marrow into blood. *J. Exp. Med.* 207:1113–1124. <http://dx.doi.org/10.1084/jem.20092210>

Avecilla, S.T., K. Hattori, B. Heissig, R. Tejada, F. Liao, K. Shido, D.K. Jin, S. Dias, F. Zhang, T.E. Hartman, et al. 2004. Chemokine-mediated interaction of hematopoietic progenitors with the bone marrow vascular niche is required for thrombopoiesis. *Nat. Med.* 10:64–71. <http://dx.doi.org/10.1038/nm973>

Blaho, V.A., and T. Hla. 2011. Regulation of mammalian physiology, development, and disease by the sphingosine 1-phosphate and lysophosphatidic acid receptors. *Chem. Rev.* 111:6299–6320. <http://dx.doi.org/10.1021/cr200273u>

Bréart, B., W.D. Ramos-Perez, A. Mendoza, A.K. Salous, M. Gobert, Y. Huang, R.H. Adams, J.J. Lafaille, D. Escalante-Alcalde, A.J. Morris, and S.R. Schwab. 2011. Lipid phosphate phosphatase 3 enables efficient thymic egress. *J. Exp. Med.* 208:1267–1278. <http://dx.doi.org/10.1084/jem.20102551>

Brinkmann, V., A. Billich, T. Baumruker, P. Heinig, R. Schmouder, G. Francis, S. Aradhya, and P. Burstin. 2010. Fingolimod (FTY720): discovery and development of an oral drug to treat multiple sclerosis. *Nat. Rev. Drug Discov.* 9:883–897. <http://dx.doi.org/10.1038/nrd3248>

Chun, J., and H.P. Hartung. 2010. Mechanism of action of oral fingolimod (FTY720) in multiple sclerosis. *Clin. Neuropharmacol.* 33:91–101. <http://dx.doi.org/10.1097/WNF.0b013e3181cbf825>

Cohen, J.A., F. Barkhof, G. Comi, H.P. Hartung, B.O. Khatri, X. Montalban, J. Pelletier, R. Capra, P. Gallo, G. Izquierdo, et al; TRANSFORMS Study Group. 2010. Oral fingolimod or intramuscular interferon for relapsing multiple sclerosis. *N. Engl. J. Med.* 362:402–415. <http://dx.doi.org/10.1056/NEJMoa0907839>

Cyster, J.G., and S.R. Schwab. 2012. Sphingosine-1-phosphate and lymphocyte egress from lymphoid organs. *Annu. Rev. Immunol.* 30:69–94. <http://dx.doi.org/10.1146/annurev-immunol-020711-075011>

Fukuhara, S., S. Simmons, S. Kawamura, A. Inoue, Y. Orba, T. Tokudome, Y. Sunden, Y. Arai, K. Moriwaki, J. Ishida, et al. 2012. The sphingosine-1-phosphate transporter Spns2 expressed on endothelial cells regulates lymphocyte trafficking in mice. *J. Clin. Invest.* 122:1416–1426. <http://dx.doi.org/10.1172/JCI60746>

Gaengel, K., C. Niaudet, K. Hagikura, B.L. Siemsen, L. Muhl, J.J. Hoffmann, L. Ebarasi, S. Nyström, S. Ryomo, L.L. Chen, et al. 2012. The Sphingosine-1-Phosphate Receptor S1PR1 Restricts Sprouting Angiogenesis by Regulating the Interplay between VE-Cadherin and VEGFR2. *Dev. Cell.* 23:587–599. <http://dx.doi.org/10.1016/j.devcel.2012.08.005>

Golfiger, S., S. Kondo, T. Schulze, T. Takeuchi, G. Vassileva, A.H. Achtmann, M.H. Gräler, S.J. Abbondanzo, M. Wiekowski, E. Kremmer, et al. 2010. Shaping of terminal megakaryocyte differentiation and proplatelet development by sphingosine-1-phosphate receptor S1P4. *FASEB J.* 24:4701–4710. <http://dx.doi.org/10.1096/fj.09-141473>

Grigorova, I.L., S.R. Schwab, T.G. Phan, T.H. Pham, T. Okada, and J.G. Cyster. 2009.

Cortical sinus probing, S1P1-dependent entry and flow-based capture of egressing T cells. *Nat. Immunol.* 10:58–65. <http://dx.doi.org/10.1038/ni.1682>

Hamada, T., R. Möhle, J. Hesselgesser, J. Hoxie, R.L. Nachman, M.A. Moore, and S. Rafii. 1998. Transendothelial migration of megakaryocytes in response to stromal cell-derived factor 1 (SDF-1) enhances platelet formation. *J. Exp. Med.* 188:539–548. <http://dx.doi.org/10.1084/jem.188.3.539>

Hla, T., K. Venkataraman, and J. Michaud. 2008. The vascular S1P gradient-cellular sources and biological significance. *Biochim. Biophys. Acta.* 1781:477–482. <http://dx.doi.org/10.1016/j.bbapap.2008.07.003>

Jung, B., H. Obinata, S. Galvani, K. Mendelson, B.S. Ding, A. Skoura, B. Kinzel, V. Brinkmann, S. Rafii, T. Evans, and T. Hla. 2012. Flow-Regulated Endothelial S1P Receptor-1 Signaling Sustains Vascular Development. *Dev. Cell.* 23:600–610. <http://dx.doi.org/10.1016/j.devcel.2012.07.015>

Konishi, H., T. Ochiya, Y. Yasuda, H. Sakamoto, T. Muto, T. Sugimura, and M. Terada. 1996. HST-1/FGF-4 stimulates proliferation of megakaryocyte progenitors synergistically and promotes megakaryocyte maturation. *Oncogene.* 13:9–19.

LaMontagne, K., A. Littlewood-Evans, C. Schnell, T. O'Reilly, L. Wyder, T. Sanchez, B. Probst, J. Butler, A. Wood, G. Liau, et al. 2006. Antagonism of sphingosine-1-phosphate receptors by FTY720 inhibits angiogenesis and tumor vascularization. *Cancer Res.* 66:221–231. <http://dx.doi.org/10.1158/0008-5472.CAN-05-2001>

Lee, M.J., S. Thangada, J.H. Paik, G.P. Sapkota, N. Ancellin, S.S. Chae, M. Wu, M. Morales-Ruiz, W.C. Sessa, D.R. Alessi, and T. Hla. 2001. Akt-mediated phosphorylation of the G protein-coupled receptor EDG-1 is required for endothelial cell chemotaxis. *Mol. Cell.* 8:693–704. [http://dx.doi.org/10.1016/S1097-2765\(01\)00324-0](http://dx.doi.org/10.1016/S1097-2765(01)00324-0)

Majka, M., A. Janowska-Wieczorek, J. Ratajczak, M.A. Kowalska, G. Vilaine, Z.K. Pan, M. Honczarenko, L.A. Marquez, M. Poncz, and M.Z. Ratajczak. 2000. Stromal-derived factor 1 and thrombopoietin regulate distinct aspects of human megakaryopoiesis. *Blood.* 96:4142–4151.

Massberg, S., P. Schaerli, I. Knezevic-Maramica, M. Köllnberger, N. Tubo, E.A. Moseman, I.V. Huff, T. Junt, A.J. Wagers, I.B. Mazo, and U.H. von Andrian. 2007. Immunosurveillance by hematopoietic progenitor cells trafficking through blood, lymph, and peripheral tissues. *Cell.* 131:994–1008. <http://dx.doi.org/10.1016/j.cell.2007.09.047>

Nachman, R.L., and S. Rafii. 2008. Platelets, petechiae, and preservation of the vascular wall. *N. Engl. J. Med.* 359:1261–1270. <http://dx.doi.org/10.1056/NEJMra0800887>

Obinata, H., and T. Hla. 2012. Sphingosine 1-phosphate in coagulation and inflammation. *Semin. Immunopathol.* 34:73–91. <http://dx.doi.org/10.1007/s00281-011-0287-3>

Oo, M.L., S.H. Chang, S. Thangada, M.T. Wu, K. Rezaul, V. Blaho, S.I. Hwang, D.K. Han, and T. Hla. 2011. Engagement of S1P₁-degradative mechanisms leads to vascular leak in mice. *J. Clin. Invest.* 121:2290–2300. <http://dx.doi.org/10.1172/JCI45403>

Paik, J.H., A. Skoura, S.S. Chae, A.E. Cowan, D.K. Han, R.L. Proia, and T. Hla. 2004. Sphingosine 1-phosphate receptor regulation of N-cadherin mediates vascular stabilization. *Genes Dev.* 18:2392–2403. <http://dx.doi.org/10.1101/gad.1227804>

Pham, T.H., T. Okada, M. Matloubian, C.G. Lo, and J.G. Cyster. 2008. S1P1 receptor signaling overrides retention mediated by G alpha i-coupled receptors to promote T cell egress. *Immunity.* 28:122–133. <http://dx.doi.org/10.1016/j.immuni.2007.11.017>

Pham, T.H., P. Baluk, Y. Xu, I. Grigorova, A.J. Bankovich, R. Pappu, S.R. Coughlin, D.M. McDonald, S.R. Schwab, and J.G. Cyster. 2010. Lymphatic endothelial cell sphingosine kinase activity is required for lymphocyte egress and lymphatic patterning. *J. Exp. Med.* 209:2165–2181.

207:17–27. <http://dx.doi.org/10.1084/jem.20091619>

Schulze, H., M. Korpal, J. Hurov, S.W. Kim, J. Zhang, L.C. Cantley, T. Graf, and R.A. Shvidasani. 2006. Characterization of the megakaryocyte demarcation membrane system and its role in thrombopoiesis. *Blood.* 107:3868–3875. <http://dx.doi.org/10.1182/blood-2005-07-2755>

Schwab, S.R., J.P. Pereira, M. Matloubian, Y. Xu, Y. Huang, and J.G. Cyster. 2005. Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P gradients. *Science.* 309:1735–1739. <http://dx.doi.org/10.1126/science.1113640>

Thangada, S., K.M. Khanna, V.A. Blaho, M.L. Oo, D.S. Im, C. Guo, L. Lefrancois, and T. Hla. 2010. Cell-surface residence of sphingosine 1-phosphate receptor 1 on lymphocytes determines lymphocyte egress kinetics. *J. Exp. Med.* 207:1475–1483. <http://dx.doi.org/10.1084/jem.20091343>

Windh, R.T., M.J. Lee, T. Hla, S. An, A.J. Barr, and D.R. Manning. 1999. Differential coupling of the sphingosine-1-phosphate receptors Edg-1, Edg-3, and H218/Edg-5 to the G(i), G(q), and G(12) families of heterotrimeric G proteins. *J. Biol. Chem.* 274:27351–27358. <http://dx.doi.org/10.1074/jbc.274.39.27351>

Yatomi, Y., F. Ruan, S. Hakomori, and Y. Igarashi. 1995. Sphingosine-1-phosphate: a platelet-activating sphingolipid released from agonist-stimulated human platelets. *Blood.* 86:193–202.

Yatomi, Y., Y. Igarashi, L. Yang, N. Hisano, R. Qi, N. Asazuma, K. Satoh, Y. Ozaki, and S. Kume. 1997. Sphingosine 1-phosphate, a bioactive sphingolipid abundantly stored in platelets, is a normal constituent of human plasma and serum. *J. Biochem.* 121:969–973. <http://dx.doi.org/10.1093/oxfordjournals.jbchem.a021681>

Zhang, L., M. Orban, M. Lorenz, V. Barocke, D. Braun, N. Urtz, C. Schulz, M. von Bruhl, A. Tirniceriu, F. Gaertner, et al. 2012. A novel role for sphingosine 1-phosphate receptor S1pr1 in mouse thrombopoiesis. *J. Exp. Med.* 209:2165–2181.