

The common mouse protozoa *Tritrichomonas muris* alters mucosal T cell homeostasis and colitis susceptibility

Nichole K. Escalante,¹ Paul Lemire,² Mayra Cruz Tleugabulova,¹ David Prescott,^{1,2} Arthur Mortha,³ Catherine J. Streutker,² Stephen E. Girardin,² Dana J. Philpott,^{1*} and Thierry Mallevaey^{1*}

¹Department of Immunology and ²Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario M5S 1A8, Canada
³Department of Oncological Science, Icahn School of Medicine at Mount Sinai, New York, NY 10029

The mammalian gastrointestinal tract hosts a diverse community of microbes including bacteria, fungi, protozoa, helminths, and viruses. Through coevolution, mammals and these microbes have developed a symbiosis that is sustained through the host's continuous sensing of microbial factors and the generation of a tolerant or pro-inflammatory response. While analyzing T cell–driven colitis in nonlittermate mouse strains, we serendipitously identified that a nongenetic transmissible factor dramatically increased disease susceptibility. We identified the protozoan *Tritrichomonas muris* as the disease-exacerbating element. Furthermore, experimental colonization with *T. muris* induced an elevated Th1 response in the cecum of naive wild-type mice and accelerated colitis in *Rag1*^{−/−} mice after T cell transfer. Overall, we describe a novel cross-kingdom interaction within the murine gut that alters immune cell homeostasis and disease susceptibility. This example of unpredicted microbial priming of the immune response highlights the importance of studying trans-kingdom interactions and serves as a stark reminder of the importance of using littermate controls in all mouse research.

INTRODUCTION

Microorganisms within the mammalian gastrointestinal tract have developed complex relationships with the host, such that the host provides a nutrient-rich microbial niche while the microbes provide vitamins, metabolites, and other nutrients otherwise inaccessible to the host (Hooper et al., 2012). To maintain this symbiosis, the host has evolved many microbial sensors, such as Toll-like receptors and Nod-like receptors, that stimulate the immune system to secrete factors such as mucous, antimicrobial peptides, and IgA, to keep microbes in check (Philpott et al., 2014; Caballero and Pamer, 2015). In turn, these microbial signals are necessary for the proper development of the mucosal immune system (Honda and Littman, 2012; Hooper et al., 2012).

Although it is clear that the presence of certain bacteria (Ivanov et al., 2009; Atarashi et al., 2013), viruses (Pfeiffer and Virgin, 2016), and helminths (Elliott and Weinstock, 2012) within the microbiota can impact intestinal homeostasis, the potential immune-shaping role of other kingdoms, such as Protista, has been less well studied. Protozoa are unicellular eukaryotes known to cause human diseases such as malaria, giardiasis, and trichomoniasis (Lindsay et al., 2008). In wild and laboratory mice, several protozoa have been documented to be disease causing, whereas others, such as *Entamoeba muris* and *Tritrichomonas muris*, are considered nonpatho-

genic members of the murine microbiome (Baker, 2008). However, recent studies suggest that *T. muris* and a related organism, *Tritrichomonas musculis*, have the potential to be pathogenic as the presence of these protists in the gut microbiota promote type 2 immune responses (Howitt et al., 2016) and Th1 inflammation (Chudnovskiy et al., 2016), respectively. These observations suggest that even though murine health is not overtly altered, protozoa colonization has the potential to modify the immune response in disease models.

While analyzing colitis induced using a T cell transfer model in two nonlittermate mouse lines, we unexpectedly observed that the exacerbated disease susceptibility of one of the lines was transmissible by cohousing and was dominant in littermates. We further identified the protozoan parasite *T. muris* to be the pro-colitogenic confounding factor in this T cell transfer model of colitis. Indeed, mice infected with *T. muris* developed an increased IFN- γ ⁺ CD4 T cell response and accelerated epithelial damage within the colon. We have also found that colonizing mice chronically with isolated *T. muris* altered the baseline number of Th1 T cells in the mouse intestine. This work reveals an unexpected critical role of a murine intestinal protozoan parasite in exacerbating colitis. This also highlights the need for monitoring parasite infections in specific pathogen–free animal facilities and argues for the standard use of littermate controls in all murine research.

*D.J. Philpott and T. Mallevaey contributed equally to this paper as co-senior authors.

Correspondence to Dana J. Philpott: dana.philpott@utoronto.ca

A. Mortha's present address is Dept. of Immunology, University of Toronto, Toronto, Ontario M5S 1A8, Canada.

Abbreviations used: LP, lamina propria; MLN, mesenteric lymph node; MNV, murine norovirus; qPCR, quantitative real-time PCR.

© 2016 Escalante et al. This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

RESULTS AND DISCUSSION

***Rip2*^{-/-}*Rag1*^{-/-} mice have accelerated T cell transfer colitis pathology**

With the aim of studying the role of NOD1 and NOD2 signaling in regulating non-T/B cells during colitis, we generated a *Rip2*^{-/-}*Rag1*^{-/-} mouse line for use in the T cell transfer model of colitis. Indeed, RIP2 is an essential signaling adapter molecule downstream of both NOD1 and NOD2 (Magalhaes et al., 2011). After transfer of wild-type CD4⁺CD45RB^{High} naive T cells, *Rip2*^{-/-}*Rag1*^{-/-} mice rapidly lost weight compared with nonlittermate *Rag1*^{-/-} mice and were euthanized at week 4 because of their poor health (Fig. 1 A). Upon examination, *Rip2*^{-/-}*Rag1*^{-/-} mice had larger spleens and short, thick colons, consistent with an increase in cellularity and significant colon pathology (Fig. 1, B–E). Preparation of colon lamina propria (LP) cells revealed a decreased frequency but a significant absolute increase in the number of IFN- γ ⁺CD4⁺ T cells (Fig. 1, F and G). There was also an increased number of TNF α ⁺CD4⁺ T cells but no increase in IL-17A⁺CD4⁺ T cells (Fig. 1, F and G). Analysis of colon explant cultures revealed an elevated protein level of IFN- γ and IL-12/IL-23p40 (Fig. 1 H). These results demonstrated that, in comparison with nonlittermate *Rag1*^{-/-} mice, *Rip2*^{-/-}*Rag1*^{-/-} mice had an increased pro-inflammatory response and exacerbated colon pathology after naive T cell transfer.

***Rip2*^{-/-}*Rag1*^{-/-} mice are not protected by regulatory cells**

Co-transfer of regulatory CD4⁺CD45RB^{Low} T cells in *Rag1*^{-/-} mice has been shown to protect mice from CD4⁺CD45RB^{High}-mediated colon pathology (Ostanin et al., 2009). Co-transfer of CD4⁺CD45RB^{Low} T cells into *Rip2*^{-/-}*Rag1*^{-/-} mice, however, did not protect the mice from colitis (Fig. 2 A). These mice still developed increased cellularity in the spleen and increased CD4⁺ T cell infiltration of the colon LP and mesenteric lymph node (MLN; Fig. S1, A and B). Increased numbers of IFN- γ ⁺CD4⁺ T cells and TNF α ⁺CD4⁺ T cells were also still observed in the colon, along with an increase in colon IFN- γ and IL-12/IL-23p40 protein (Fig. 2, B–D; and Fig. S1, C and D). Various other cytokines, such as IL-1 β , IL-6, IL-10, IL-12p70, IL-13, KC, MIP-1 α , and MIP-2 were also increased in *Rip2*^{-/-}*Rag1*^{-/-} colons, regardless of the T cells transferred (Fig. 2 E). This lack of protection was not caused by a lack of CD4⁺Foxp3⁺ regulatory T cells (T reg cells) as there was a similar ratio of CD4⁺Foxp3⁺ to CD4⁺Foxp3⁻ T cells in both *Rag1*^{-/-} and *Rip2*^{-/-}*Rag1*^{-/-} mice when CD4⁺CD45RB^{Low} T cells were co-transferred (Fig. 2 C). These results suggest that the early increase in pro-inflammatory mediators in the *Rip2*^{-/-}*Rag1*^{-/-} mice may be blocking the ability of the CD4⁺CD45RB^{Low} T cells to suppress inflammation.

Exacerbated *Rip2*^{-/-}*Rag1*^{-/-} pathology is transmissible

To determine whether the accelerated colitis seen in *Rip2*^{-/-}*Rag1*^{-/-} mice was driven by *Rip2* deficiency or was caused by other factors such as a dysbiotic microbiota,

Rip2^{-/-}*Rag1*^{-/-} and *Rag1*^{-/-} mice were co-caged at weaning and 5 wk later injected with CD4⁺CD45RB^{High} T cells. Strikingly, 4 wk after transfer, co-caged *Rag1*^{-/-} mice developed equally accelerated pathology as the *Rip2*^{-/-}*Rag1*^{-/-} mice (Fig. 3 A), suggesting that the disease-inducing factor could be horizontally transferred. Littermate mice were then generated, and the resulting *Rip2*^{+/+}*Rag1*^{-/-} mice also developed accelerated colitis and had similar pro-inflammatory cytokine levels as their *Rip2*^{-/-}*Rag1*^{-/-} littermates (Fig. 3, B and C). These results indicate that microbial factors, rather than *Rip2* genotype, were likely driving the exacerbated colitis in *Rip2*^{-/-}*Rag1*^{-/-} mice.

To determine the potential pro-colitogenic microbial factors, *Rip2*^{-/-}*Rag1*^{-/-} and *Rag1*^{-/-} nonlittermate and littermate mice were screened for viral infections. Interestingly, *Rip2*^{-/-}*Rag1*^{-/-} nonlittermate and all littermates, regardless of genotype, were positive for murine norovirus (MNV) infection (IDEXX testing by quantitative real-time PCR [qPCR]), whereas nonlittermate *Rag1*^{-/-} mice were negative. MNV is commonly found in the gastrointestinal tract of laboratory mice, and *Rag1*^{-/-} mice have been reported to be persistently infected after exposure (Karst et al., 2003). Because previous studies had shown that MNV infection may exacerbate both chemical- and bacterial-induced colitis in mice (Lencioni et al., 2008; Cadwell et al., 2010), we investigated the hypothesis that MNV may exacerbate colitis in *Rag1*^{-/-} mice during T cell transfer colitis. To test this, we infected *Rag1*^{-/-} mice with MNV CW3 or MNV CR6 and 2 wk later compared them with their noninfected littermates during T cell transfer colitis. Although MNV levels were still detectable by qPCR 4 wk after CD4⁺CD45RB^{High} T cell transfer, no differences were observed in pathology or weight loss, suggesting that MNV infection is not a contributing factor to T cell transfer colitis (Fig. 3, D–F).

***T. muris* infection exacerbates T cell transfer colitis**

Rip2^{-/-}*Rag1*^{-/-} and *Rag1*^{-/-} nonlittermate mice were also screened for parasitic infection. All mice were found to be negative for *E. muris*, *Giardia muris*, and *Spironucleus muris*. Notably, however, the original *Rip2*^{-/-}*Rag1*^{-/-} mouse line and littermate mice were all found to be positive for the protozoa *T. muris*, whereas *Rag1*^{-/-} nonlittermates were not. Indeed, large numbers of these protozoa could be observed by microscope in the cecal contents of *Rip2*^{-/-}*Rag1*^{-/-} but not *Rag1*^{-/-} nonlittermate mice (Fig. 4 A). Giemsa staining revealed the characteristic three anterior flagella of *T. muris* (Fig. 4 B). qPCR screening of our colony revealed that 18 out of 24 separate mouse strains were *T. muris* colonized. *T. muris* is transferred between mice by ingestion of infected fecal pellets containing pseudocysts, but transmission between cages is limited when sterile technique is strictly followed (Baker, 2008). Co-caging of mice led to transmission of the parasitic infection by 2 wk (not depicted). We next tested the hypothesis that *T. muris* infection could be the confounding factor that accelerated colitis in our T cell transfer experiments.




Figure 1. Nonlittermate *Rip2*^{−/−}*Rag1*^{−/−} mice develop accelerated colitis. Nonlittermate *Rip2*^{−/−}*Rag1*^{−/−} and *Rag1*^{−/−} mice were injected i.p. with 0.5×10^6 $CD45RB^{high}$ $CD4$ T cells. (A) Mice were weighed weekly, and the percentage of initial body weight was calculated. Data from four experiments were pooled, $n = 15$ mice per group. (B) Mice were sacrificed at week 4. Colons were fixed, H&E stained, and scored for colitis severity. Data from three experiments were pooled. (C) Representative spleens and colons were imaged. (D and E) Spleen, colon LP, and MLNs were harvested at week 4, and isolated cells were quantified by flow cytometry. (F and G) Colon cells were restimulated with plate-bound anti-CD3/anti-CD28 antibody and analyzed by flow cytometry to determine the total (F) and frequency (G) of cytokine-positive cells. Data from three experiments were pooled with 5–15 mice per group. (H) Proximal colon explants were cultured for 18 h, and supernatants were analyzed by ELISA for cytokine production. Data from two experiments were pooled with four to seven samples per group. (A, B, and D–H) Mean \pm SEM is shown with * $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$; **** $P \leq 0.0001$ using an unpaired Student's *t* test.

T. muris was isolated from the cecal contents of *Rip2*^{−/−}*Rag1*^{−/−} mice (Fig. S2 A) and orally gavaged into *Rag1*^{−/−} mice. 2 wk later, $CD4^+CD45RB^{high}$ T cells were

transferred into the *T. muris*–infected *Rag1*^{−/−} mice and their noninfected littermate controls. By 4 wk after T cell transfer, the *T. muris*–infected mice lost significantly more





Figure 2. Nonlittermate Rip2^{-/-}Rag1^{-/-} mice are not protected from pathology by regulatory CD45RB^{Low} T cells. Nonlittermate Rip2^{-/-}Rag1^{-/-} and Rag1^{-/-} mice were injected i.p. with 0.5×10^6 CD45RB^{High} with or without 0.25×10^6 CD45RB^{Low} CD4 T cells and sacrificed at week 4. (A) Colons were fixed, H&E stained, and scored for colitis severity. Data from three experiments were pooled. (B and C) Spleen, colon LP, and MLNs were harvested, and isolated cells were counted, restimulated with plate bound anti-CD3/anti-CD28 antibody, and analyzed by flow cytometry to determine tissue Foxp3⁺ (C) and colon IFN γ ⁺ T cell numbers (B). Data from two experiments were pooled with 4–12 mice per group. (D) Proximal colon explants were cultured for 18 h, and supernatants were analyzed by ELISA for cytokine production. Data from two experiments were pooled with three to seven samples per group. (A–D) Mean \pm SEM is shown with *, P \leq 0.05; **, P \leq 0.01; ***, P \leq 0.001 using a one-way ANOVA and Tukey's post-hoc analysis. (E) Proximal colon explants were cultured for 18 h, and supernatants were analyzed by Bio-Plex assay for cytokine production. Data are one representative of three experiments with two samples per group. *, samples below minimum level of detection; #, samples above maximal level of detection.



weight than their mock-challenged siblings (Fig. 4 C). The *T. muris*-infected mice had significantly increased loss of crypt structure, epithelial damage ($P = 0.0545$), and lymphocyte infiltration (chronic inflammation score; Fig. 4, D–F). IFN- γ and IL-12/IL-23p40 protein levels were also elevated in the colons of the infected mice, whereas IL-18 was not significantly increased (Fig. 4 G). Together, these data show that *T. muris* infection contributed to an elevated Th1 pro-inflammatory environment in the colon that resulted in increased colon pathology during T cell transfer colitis.

Chronic *T. muris* infection alters baseline T cell homeostasis

To determine whether *T. muris* could be altering the steady-state level of immune activation in the gut of immunocompetent mice during natural infection, we created a line of chronically infected C57BL/6 mice by infecting a pair of C57BL/6 breeders. Baseline T cell numbers and cytokine production in the offspring of these infected mice were compared with pups from noninfected breeders (littermates to the infected breeders). Interestingly, infected mice had increased frequencies and a tendency for numbers ($P = 0.0617$) of IFN- γ^+ CD4 T cells in their cecal LP (Fig. 5 A). In contrast, frequencies of IL-17A $^+$ CD4 T cells were decreased in the colon LP (Fig. 5 B). Ror γ t $^+$ Foxp3 $^+$ T reg cell

Figure 3. Co-caged and littermate mice develop equally exacerbated T cell–induced colitis. (A and B) Co-caged (A) and littermate (B) *Rip2^{-/- Rag1^{-/-}}* and *Rip2^{+/+ Rag1^{-/-}}* mice were injected i.p. with 0.5×10^6 CD45RB $^{\text{High}}$ CD4 T cells and sacrificed at week 4. Colons were fixed, H&E stained, and scored for colitis severity. Data from three experiments were pooled. Mean \pm SEM is shown, and an unpaired Student's *t* test was performed. (C) Proximal colon explants from week 4 after transfer littermate mice were cultured for 18 h, and supernatants were analyzed by ELISA for cytokine production. Data from two experiments were pooled with five to six samples per group. Mean \pm SEM is shown, and a Student's *t* test was performed. (D–F) *Rag1^{-/-}* mice were orally gavaged with 10^6 PFU MNV CR6 or MNV CW3 and 2 wk later injected i.p. with 0.5×10^6 CD45RB $^{\text{High}}$ CD4 T cells. (D) Body weight was monitored, and mice were sacrificed at week 4. (E) Colons were fixed, H&E stained, and scored for colitis severity. (F) Fecal pellets were confirmed positive by qPCR. Data are from one experiment with three to four mice per group. Mean \pm SEM is shown, and a one-way ANOVA was performed.

frequencies and numbers were similar between infected and noninfected mice, but Ror γ t $^+$ Foxp3 $^+$ T reg cell frequencies were decreased (Fig. 5, A and B). IFN- γ protein levels were below the assay level of detection in cultured colon explant supernatants, and IL-18 was not significantly different, but interestingly, IL-12/IL-23p40 protein was elevated in *T. muris* colonized mice (Fig. 5 C). This baseline increase in IL-12/IL-23p40 protein and IFN- γ -producing CD4 T cells, accompanied by the decrease in T reg cells, indicated a shift to a more pro-inflammatory rather than tolerant mucosal environment in the presence of *T. muris*.

Although we focused on a Th1 immune response in the large intestine, the wide array of cytokines found to be increased in the *Rip2^{-/- Rag1^{-/-}}* colons during T cell transfer colitis suggested that *T. muris* may be acting as a general stimulator of the mucosal immune system and not specifically favoring a Th1 response. This would agree with the study from Howitt et al. (2016) that outlined the induction of a type 2 response in the small intestine upon *T. muris* colonization. Interestingly, we also observed a significant increase in type 2–inducing tuft cells in the small intestines of our chronically infected wild-type mice, suggesting *T. muris* responses may differ in distinct gut regions (Fig. 5 D). Recently, Chudnovskiy et al. (2016) reported that acute infection of mice




Figure 4. *T. muris* accelerates T cell–induced colitis. (A) Fresh cecal contents were diluted in phosphate-buffered saline and visualized under a microscope. (B) *T. muris* from the cecal contents of *Rip2*^{−/−}*Rag1*^{−/−} mice were stained with Giemsa and visualized. (C–G) *T. muris* was isolated from the cecum of *Rip2*^{−/−}*Rag1*^{−/−} mice, and 10⁶ protozoans were orally gavaged into *Rag1*^{−/−} mice. 2 wk later, mice were injected i.p. with 0.5 × 10⁶ CD45RB^{High} CD4 T cells. (C) Body weight was measured, and mice were sacrificed at week 4. (D–F) Colons were fixed, H&E stained, and scored for colitis severity. Data from three experiments were pooled with 12 mice per group. Bars: (A) 100 μ m; (B) 10 μ m; (F) 200 μ m. (G) Proximal colon explants from week 4 after T cell transfer *T. muris*-infected and control mice were cultured for 18 h, and supernatants were analyzed by ELISA for cytokine production. Data from two experiments were pooled with 9–10 samples per group. (C–E and G) Mean \pm SEM is shown with *, P ≤ 0.05; **, P ≤ 0.01 using an unpaired Student's *t* test.

with a related protozoan *T. musculis* induced IL-18–driven Th1 and Th17 responses that altered the outcome of bacterial-induced colitis, T cell–driven colitis, and colorectal tumor formation. Although our study also demonstrates increased Th1 responses after protozoa infection, IL-18 and IL-17 levels were not significantly increased. This difference may be the result of chronic versus acute infection or the result of dif-

ferent protozoa. Indeed, the 28s rRNA primers used in our study cannot differentiate between *T. muris* and *T. musculis*. Collectively, these studies are further evidence that *T. muris* and related intestinal parasites could alter disease outcomes in many other mouse models.

Because of the presence of *T. muris* in our *Rip2*^{−/−}*Rag1*^{−/−} mice, we were unable to determine whether




Figure 5. *T. muris* infection chronically increases cecal IFN- γ ⁺ T cells. (A–D) C57BL/6 breeders were orally gavaged with 10^6 isolated protozoans. (A and B) Ceca (A) and colons (B) of infected pups and control pups from noninfected breeders were harvested. Cells were isolated and restimulated for 4 h with PMA and ionomycin, in the presence of protein transport inhibitor cocktail, before flow cytometry analysis. Data from two experiments were pooled with three to six mice per group. (C) Proximal colon explants were cultured for 24 h, and supernatants were analyzed by ELISA for IL-12/IL-23p40 and IL-18 protein. Data from two experiments were pooled with six to seven samples per group. (D) Small intestines were fixed, stained for tufts cells (DCLK1⁺DAPI⁺), and quantified. Data from two experiments were pooled with six mice per group. (A–D) Mean \pm SEM is shown with *, P \leq 0.05 using an unpaired Student's *t* test.

NOD1 and 2 signaling could alter colitis onset in the T cell transfer model of colitis. A previous study has described a pro-colitogenic bacterial composition in *Nod2*^{−/−} and

Rip2^{−/−} mice; however, no details about the colonization of these mice with protozoa were given (Couturier-Maillard et al., 2013). Although the relationship between *T. muris*

and bacteria within the gut is unknown, the large number of protozoa present in the cecal contents of *T. muris* colonized mice suggests a likely impact on bacterial composition. Thus, along with searching for potential *T. muris*–specific antigens, the potential of a *T. muris*–induced bacterial dysbiosis could also be explored.

T. muris screening is not standard in animal facilities; therefore, the current prevalence of *T. muris* in laboratory mice is unknown. Treatment with metronidazole has been reported to rid mice of *T. muris* (Roach et al., 1988; Howitt et al., 2016); however, antibiotic treatment also alters bacterial composition, which could have other confounding effects on a given phenotype. Our experience suggests that cross-fostering newborn pups with *T. muris*–free dams is a potential option. Although *T. muris* may now be flagged as a potential confounding factor in intestinal disease models, whether it impacts systemic disease remains to be determined. Moreover, there still remains many other unidentified protozoans or microbes that may also play immunomodulatory roles in mouse models, underscoring the importance of proper littermate controlled experiments.

Pentatrichomonas hominis and *Dientamoeba fragilis* are two species of trichomonads known to sporadically colonize the human gastrointestinal tract (Maritz et al., 2014). Although these trichomonads were originally described to be commensal protists, the correlation between the presence of these organisms in patients presenting gastrointestinal symptoms and the work presented here calls for a reevaluation of their pathogenicity (Meloni et al., 2011; Stark et al., 2016). Overall, our findings highlight the need for a better understanding of cross-kingdom interactions between host and protozoa within the gastrointestinal tract and emphasize the importance of controlling for microbial factors in mouse models of disease using littermate controls.

MATERIALS AND METHODS

Animals

All animals were housed in specific pathogen–free conditions at the University of Toronto, Division of Comparative Medicine. *Rag1*^{−/−} and C57BL/6 mice were purchased from The Jackson Laboratory and subsequently bred in house. *Rip2*^{−/−} mice were obtained from R.A. Flavell (Yale University School of Medicine, New Haven, CT; Kobayashi et al., 2002). *Rip2*^{−/−}*Rag1*^{−/−} mice were generated by crossing *Rag1*^{−/−} mice with in-house *Rip2*^{−/−} mice. All animal experiments were approved by the animal care committee, University of Toronto.

T cell transfer colitis

$CD4^+CD45RB^{High}$ and $CD4^+CD45RB^{Low}$ T cells were sorted from *T. muris*–negative C57BL/6 spleens. 0.5×10^6 $CD4^+CD45RB^{High}$ with or without 0.25×10^6 $CD4^+CD45RB^{Low}$ T cells were injected i.p. into 8-wk-old *Rag1*^{−/−} or *Rip2*^{−/−}*Rag1*^{−/−} mice. Mice were weighed twice a week to obtain a mean weekly weight. Mice were sacrificed at 4 wk

after T cell transfer, and colons were fixed with 10% formalin and hematoxylin and eosin (H&E) stained for pathological scoring or used for LP preps and flow cytometry. Colon pathology was scored by a blinded pathologist as previously described (Ostanin et al., 2009).

Tissue preparations

Colon LP cells were isolated as previously described (Good-year et al., 2014). In brief, colons were washed in pre-stripping buffer (HBSS with 5 mM DTT, penicillin, streptomycin, and FBS) to remove mucous and fecal contents before incubation in epithelial stripping buffer (HBSS with 5 mM EDTA, penicillin, streptomycin, and FBS). Colons were then washed in HBSS + 10 mM HBSS, minced, and digested in digestion buffer (HBSS with 10 mM Hepes, penicillin, streptomycin, 20 µg/ml DNase, and 0.2U/ml Liberase TM). Colon pieces were titrated with an 18GA needle, filtered, and washed. Spleen and MLNs were mashed through a filter, and cells were isolated by centrifugation. Red blood cells were lysed using ammonium chloride.

Flow cytometry

For intracellular cytokine staining, cells were restimulated for 18 h on plates coated with anti-CD3 and anti-CD28 antibody or for 4 h with PMA and ionomycin. Brefeldin A and Monensin (Protein transport inhibitor cocktail; eBioscience) were added during the last 4 h of restimulation. Dead cells were marked using the LIVE/DEAD Fixable Aqua Dead Cell Stain kit (Molecular Probes), and Fc receptors were blocked using mouse CD16/CD32 antibody (eBioscience). Surface staining was performed with anti-mouse CD4 (GK1.5), CD45RB (C363-16A), and TCR β (H57-597) from eBioscience. Cells were fixed and permeabilized using Foxp3/Transcription Factor staining buffer set (eBioscience) before staining with anti-mouse IFN- γ (XMG1.2), TNF α (MP6-XT22), IL-17A (eBio17B7), and Foxp3 (FJK-16s) from eBioscience. Samples were analyzed on a FACSCanto II or LSR Fortessa X20 (BD).

Colon explants, ELISAs, and Bio-Plex assay

Proximal colon explants were weighed, rinsed in RPMI, and cultured in 500 µl complete RPMI for 18–24 h as previously described (Hue et al., 2006). Supernatants were centrifuged and used for cytokine measurement by IL-12/IL-23p40 (eBioscience; sensitivity 2 pg/ml), IFN- γ (eBioscience; sensitivity 15 pg/ml), and IL-18 (MBL; sensitivity 12.5 pg/ml) ELISA or Bio-Plex Mouse Cytokine 23-plex Assay (Bio-Rad Laboratories) according to the manufacturer’s instructions. Results were normalized to tissue weight.

Immunofluorescence staining

Small intestines were collected, formalin fixed, and paraffin embedded. Paraffin sections (5 µm) were deparaffinized in xylene and rehydrated in an ethanol gradient. Antigen retrieval was performed in 10 mM sodium citrate buffer, pH

6.0, at 95°C for 20 min. Sections were stained with rabbit polyclonal anti-DCLK1 (1:1,000; Abcam) overnight at 4°C, followed by staining with Alexa Fluor 647-conjugated anti-rabbit secondary antibody (1:4,000; Molecular Probes), and counterstaining with DAPI (1:10,000; Sigma-Aldrich). Images were acquired with the Axio Scan Z.1 Slide Scanner (ZEISS). Quantification was performed blind by counting the total number of tuft cells per crypt-villus with the Zen Blue 2 software (ZEISS).

MNV infection

MNV CR6 and CW3 stocks and plasmids were provided by H.W.Virgin (Washington University School of Medicine, St. Louis, MO). MNV was propagated, quantified, and detected by qPCR as previously described (Hwang et al., 2014). Mice were mock infected or infected with 10⁶ PFU MNV and confirmed to be positive by qPCR (IDEXX Bioresearch or in house as previously described).

T. muris isolation and infection

Cecal contents of *Rip2^{-/-}Rag1^{-/-}* mice were resuspended in PBS and filtered twice through a 70-μm cell strainer. Flow through was centrifuged, resuspended, and refiltered twice. Protozoa were double sorted based on size, granularity, and violet autofluorescence on an Influx (BD; Fig. S2 A) and visualized under a microscope (BX43; Olympus) to confirm protozoa isolation. Eubacteria 16s rDNA levels in sort samples were determined by qPCR as previously described (Robertson et al., 2013) to measure the reduction of bacterial contamination (Fig. S2 B). 10⁶ protozoa in PBS or protozoa-free PBS was orally gavaged into *Rag1^{-/-}* or C57BL/6 mice. 2 wk after infection, mice were confirmed to be *T. muris* positive and used for T cell transfer colitis or as breeders. *T. muris* testing was performed by IDEXX Bioresearch, cecal content wet smear, and/or by qPCR for the 28s rRNA as previously described (Howitt et al., 2016). For qPCR, fecal DNA was isolated using a NucleoSpin Soil kit (MACHER EY-NAGEL) and amplified using Power SYBR green master mix (Invitrogen). For microscopic visualization, *T. muris* were fixed with paraformaldehyde, adhered to slides by cytospin, and stained with Giemsa.

Statistical analysis

Unpaired Student's *t* tests were performed when two groups were compared. A one-way ANOVA was used for comparison of more than two groups, followed by a post-hoc Tukey's multiple comparisons test if needed.

Online supplemental material

Supporting data for Fig. 2 (CD4⁺CD45RB^{High+Low} T cells into *Rip2^{-/-}Rag1^{-/-}* vs. *Rag1^{-/-}* mice), *T. muris* sorting strategy, and measurement of bacterial contamination are provided. Fig. S1 shows *Rip2^{-/-}Rag1^{-/-}* are not protected from pathology by regulatory CD45RB^{Low} T cells. Fig. S2 shows the sorting strategy for *T. muris*.

ACKNOWLEDGMENTS

The authors of this paper would like to thank Dionne White and Joanna Warzyszynska for their expert FACS assistance, Elisabeth Foerster for her microscopy expertise, and the University of Toronto Department of Comparative Medicine animal facility for their assistance.

This work was supported by grants from the Canadian Institutes of Health Research (to D.J. Philpott) and the Crohn's and Colitis Foundation of Canada (to T. Mallevaey). N.K. Escalante was supported by a Vanier Canada Graduate Scholarship. The authors declare no competing financial interests.

Author contributions: Conceptualization, N.K. Escalante, D.J. Philpott, T. Mallevaey, and S.E. Girardin; methodology, N.K. Escalante, A. Mortha, D.J. Philpott, T. Mallevaey, and S.E. Girardin; formal analysis, N.K. Escalante; investigation, N.K. Escalante, P. Lemire, M. Cruz Tleugabulova, D. Prescott, and C.J. Streutker; writing—original draft, N.K. Escalante; writing—review and editing, D.J. Philpott and S.E. Girardin; visualization, N.K. Escalante; supervision, D.J. Philpott, T. Mallevaey, and S.E. Girardin; project administration, N.K. Escalante, D.J. Philpott, and T. Mallevaey; funding acquisition, D.J. Philpott and T. Mallevaey.

Submitted: 21 October 2016

Revised: 4 November 2016

Accepted: 7 November 2016

REFERENCES

- Atarashi, K., T. Tanoue, K. Oshima, W. Suda, Y. Nagano, H. Nishikawa, S. Fukuda, T. Saito, S. Narushima, K. Hase, et al. 2013. Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature*. 500:232–236. <http://dx.doi.org/10.1038/nature12331>
- Baker, D.G. 2008. Parasites of rats and mice. In *Flynn's Parasites of Laboratory Animals*. Second edition. D.G. Baker, editor. Blackwell Publishing Ltd., Ames, IA. 303–397.
- Caballero, S., and E.G. Pamer. 2015. Microbiota-mediated inflammation and antimicrobial defense in the intestine. *Annu. Rev. Immunol.* 33:227–256. <http://dx.doi.org/10.1146/annurev-immunol-032713-120238>
- Cadwell, K., K.K. Patel, N.S. Maloney, T.C. Liu, A.C.Y. Ng, C.E. Storer, R.D. Head, R. Xavier, T.S. Stappenbeck, and H.W. Virgin. 2010. Virus-plus-susceptibility gene interaction determines Crohn's disease gene Atg16L1 phenotypes in intestine. *Cell*. 141:1135–1145. <http://dx.doi.org/10.1016/j.cell.2010.05.009>
- Chudnovskiy, A., A. Mortha, V. Kana, A. Kennard, J.D. Ramirez, A. Rahman, R. Remark, I. Mogno, R. Ng, S. Gnajtic, et al. 2016. Host-protozoan interactions protect from mucosal infections through activation of the inflammasome. *Cell*. 167:444–456.e14. <http://dx.doi.org/10.1016/j.cell.2016.08.076>
- Couturier-Maillard, A., T. Secher, A. Rehman, S. Normand, A. De Arcangelis, R. Haesler, L. Huot, T. Grandjean, A. Bressenot, A. Delanoye-Crespin, et al. 2013. NOD2-mediated dysbiosis predisposes mice to transmissible colitis and colorectal cancer. *J. Clin. Invest.* 123:700–711. <http://dx.doi.org/10.1172/JCI62236>
- Elliott, D.E., and J.V. Weinstock. 2012. Helminth-host immunological interactions: prevention and control of immune-mediated diseases. *Ann. N. Y. Acad. Sci.* 1247:83–96. <http://dx.doi.org/10.1111/j.1749-6632.2011.06292.x>
- Goodyear, A.W., A. Kumar, S. Dow, and E.P. Ryan. 2014. Optimization of murine small intestine leukocyte isolation for global immune phenotype analysis. *J. Immunol. Methods*. 405:97–108. <http://dx.doi.org/10.1016/j.jim.2014.01.014>
- Honda, K., and D.R. Littman. 2012. The microbiome in infectious disease and inflammation. *Annu. Rev. Immunol.* 30:759–795. <http://dx.doi.org/10.1146/annurev-immunol-020711-074937>
- Hooper, L.V., D.R. Littman, and A.J. Macpherson. 2012. Interactions between the microbiota and the immune system. *Science*. 336:1268–1273. <http://dx.doi.org/10.1126/science.1223490>

Howitt, M.R., S. Lavoie, M. Michaud, A.M. Blum, S.V. Tran, J.V. Weinstock, C.A. Gallini, K. Redding, R.F. Margolskee, L.C. Osborne, et al. 2016. Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in the gut. *Science*. 351:1329–1333. <http://dx.doi.org/10.1126/science.aaf1648>

Hue, S., P. Ahern, S. Buonocore, M.C. Kullberg, D.J. Cua, B.S. McKenzie, F. Powrie, and K.J. Maloy. 2006. Interleukin-23 drives innate and T cell-mediated intestinal inflammation. *J. Exp. Med.* 203:2473–2483. <http://dx.doi.org/10.1084/jem.20061099>

Hwang, S., B. Alhatlani, A. Arias, S.L. Caddy, C. Christodoulou, J.B. Cunha, E. Emmott, M. Gonzalez-Hernandez, A. Kolawole, J. Lu, et al. 2014. Murine norovirus: propagation, quantification, and genetic manipulation. *Curr. Protoc. Microbiol.* 33:1–61: 61.

Ivanov, I.I., K. Atarashi, N. Manel, E.L. Brodie, T. Shima, U. Karaoz, D. Wei, K.C. Goldfarb, C.A. Santee, S.V. Lynch, et al. 2009. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell*. 139:485–498. <http://dx.doi.org/10.1016/j.cell.2009.09.033>

Karst, S.M., C.E. Wobus, M. Lay, J. Davidson, and H.W. Virgin IV. 2003. STAT1-dependent innate immunity to a Norwalk-like virus. *Science*. 299:1575–1578. <http://dx.doi.org/10.1126/science.1077905>

Kobayashi, K., N. Inohara, L.D. Hernandez, J.E. Galán, G. Núñez, C.A. Janeway, R. Medzhitov, and R.A. Flavell. 2002. RICK/Rip2/CARDIAK mediates signalling for receptors of the innate and adaptive immune systems. *Nature*. 416:194–199. <http://dx.doi.org/10.1038/416194a>

Lencioni, K.C., A. Seamons, P.M. Treuting, L. Maggio-Price, and T. Brabb. 2008. Murine norovirus: an intercurrent variable in a mouse model of bacteria-induced inflammatory bowel disease. *Comp. Med.* 58:522–533.

Lindsay, D.S., C.A. Sundermann, and D.G. Baker. 2008. Biology of the protozoa. In *Flynn's Parasites of Laboratory Animals*. Second edition. D.G. Baker, editor. Blackwell Publishing Ltd., Ames, IA. 15–25.

Magalhaes, J.G., J. Lee, K. Geddes, S. Rubino, D.J. Philpott, and S.E. Girardin. 2011. Essential role of Rip2 in the modulation of innate and adaptive immunity triggered by Nod1 and Nod2 ligands. *Eur. J. Immunol.* 41:1445–1455. <http://dx.doi.org/10.1002/eji.201040827>

Maritz, J.M., K.M. Land, J.M. Carlton, and R.P. Hirt. 2014. What is the importance of zoonotic trichomonads for human health? *Trends Parasitol.* 30:333–341. <http://dx.doi.org/10.1016/j.pt.2014.05.005>

Meloni, D., C. Mantini, J. Goustille, G. Desoubeaux, Z. Maakaroun-Vermesse, J. Chandenier, N. Gantois, C. Duboucher, P.L. Fiori, E. Dei-Cas, et al. 2011. Molecular identification of *Pentatrichomonas hominis* in two patients with gastrointestinal symptoms. *J. Clin. Pathol.* 64:933–935. <http://dx.doi.org/10.1136/jcp.2011.089326>

Ostanin, D.V., J. Bao, I. Koboziev, L. Gray, S.A. Robinson-Jackson, M. Kosloski-Davidson, V.H. Price, and M.B. Grisham. 2009. T cell transfer model of chronic colitis: concepts, considerations, and tricks of the trade. *Am. J. Physiol. Gastrointest. Liver Physiol.* 296:G135–G146. <http://dx.doi.org/10.1152/ajpgi.90462.2008>

Pfeiffer, J.K., and H.W. Virgin. 2016. Transkingdom control of viral infection and immunity in the mammalian intestine. *Science*. 351:aad5872. <http://dx.doi.org/10.1126/science.aad5872>

Philpott, D.J., M.T. Sorbara, S.J. Robertson, K. Croitoru, and S.E. Girardin. 2014. NOD proteins: regulators of inflammation in health and disease. *Nat. Rev. Immunol.* 14:9–23. <http://dx.doi.org/10.1038/nri3565>

Roach, P.D., P.M. Wallis, and M.E. Olson. 1988. The use of metronidazole, tinidazole and dimetridazole in eliminating trichomonads from laboratory mice. *Lab. Anim.* 22:361–364. <http://dx.doi.org/10.1258/002367788780746287>

Robertson, S.J., J.Y. Zhou, K. Geddes, S.J. Rubino, J.H. Cho, S.E. Girardin, and D.J. Philpott. 2013. Nod1 and Nod2 signaling does not alter the composition of intestinal bacterial communities at homeostasis. *Gut Microbes*. 4:222–231. <http://dx.doi.org/10.4161/gmic.24373>

Stark, D., J. Barratt, D. Chan, and J.T. Ellis. 2016. *Dientamoeba fragilis*, the neglected trichomonad of the human bowel. *Clin. Microbiol. Rev.* 29:553–580. <http://dx.doi.org/10.1128/CMR.00076-15>