

ARTICLE

Regulatory roles of IL-10-producing human follicular T cells

Pablo F. Cañete¹, Rebecca A. Sweet¹, Paula Gonzalez-Figueroa¹, Ilenia Papa¹, Naganari Ohkura³, Holly Bolton⁴, Jonathan A. Roco¹, Marta Cuenca¹, Katharine J. Bassett¹, Ismail Sayin⁵, Emma Barry⁶, Angel Lopez⁶, David H. Canaday⁵, Michael Meyer-Hermann⁷, Claudio Doglioni⁸, Barbara Fazekas de St Groth⁴, Shimon Sakaguchi³, Matthew C. Cook^{1,2}, and Carola G. Vinuesa¹

Mucosal lymphoid tissues such as human tonsil are colonized by bacteria and exposed to ingested and inhaled antigens, requiring tight regulation of immune responses. Antibody responses are regulated by follicular helper T (T_{FH}) cells and FOXP3⁺ follicular regulatory T (T_{FR}) cells. Here we describe a subset of human tonsillar follicular T cells identified by expression of T_{FH} markers and CD25 that are the main source of follicular T (T_F) cell-derived IL-10. Despite lack of FOXP3 expression, CD25⁺ T_F cells resemble T reg cells in high CTLA4 expression, low IL-2 production, and their ability to repress T cell proliferation. CD25⁺ T_F cell-derived IL-10 dampens induction of B cell class-switching to IgE. In children, circulating total IgE titers were inversely correlated with the frequencies of tonsil CD25⁺ T_F cells and IL-10-producing T_F cells but not with total T reg cells, T_{FR} , or IL-10-producing T cells. Thus, CD25⁺ T_F cells emerge as a subset with unique T and B cell regulatory activities that may help prevent atopy.

Downloaded from http://upress.org/jem/article-pdf/216/8/1843/1766359/jem_20190493.pdf by guest on 09 February 2026

Introduction

High-affinity antibodies are critical for long-lived host defense after infection or vaccination. Conversely, dysregulation of antibody responses is the basis of both serious autoimmune diseases and allergy (Vinuesa et al., 2016). It is clear that specialized B cell lymphoma 6 protein (BCL6)-driven B helper follicular T (T_{FH}) cells are essential in supporting and regulating the quality and longevity of antibody responses (Crotty, 2011; Vinuesa et al., 2016). T_{FH} cells first interact with antigen-specific B cells at the borders between T cell zones and B cell follicles, driving B cells to differentiate in extrafollicular foci as short-lived plasmablasts (Lee et al., 2011), and then after repeated cycles of division and mutation within germinal centers (GCs). T_{FH} cells also drive GC B cell differentiation into long-lived plasma cells and memory B cells. Limiting T_{FH} cell help appears to be crucial for long-lived high-affinity antibody responses (Victora et al., 2010), and aberrant accumulation of T_{FH} cells has been shown to promote selection of GC B cells and lead to autoantibodies (Vinuesa et al., 2005; Linterman et al., 2009; Simpson et al., 2010) and IgE⁺ B cells (Yang et al., 2016). The BCL6⁺ follicular T (T_F) cell population also contains regulatory cells made up of a thymic-derived and peripherally induced forkhead box P3 (FOXP3)-expressing population known as follicular regulatory T cells

(T_{FR} ; Chung et al., 2011; Linterman et al., 2011; Wollenberg et al., 2011). In mice, T_{FR} cells have been shown to repress GC B cells and antibody responses (Sage et al., 2016). Regulatory CD25⁺ T cells and follicular FOXP3⁺ T cells have also been reported in humans (Lim et al., 2004; Carreras et al., 2006; Chung et al., 2011) and circulating follicular FOXP3⁺ regulatory populations have been described (Fonseca et al., 2017; Wing et al., 2017). Nevertheless, the nature of T_{FR} cells in human tonsil, the most accessible human secondary lymphoid tissue, remains uncharacterized. CD25⁺ T_F cells have been reported in human tonsils but are not considered to carry out regulatory roles based on their lack of FOXP3 expression (Li and Pauza, 2015), even though functional studies are lacking.

Anaphylaxis and other acute allergic reactions are growing in incidence and are poorly understood problems causing increasing morbidity and mortality (Yue et al., 2018). These reactions are driven by cross-linking of IgE bound to the high-affinity IgE receptor (Fc ϵ RI) on mast cells and basophils, which leads to the release of inflammatory and vasoactive mediators (Galli et al., 2008). Allergic pathology is often located at epithelial and mucosal sites and consists of type 2 immune responses, in which signature cytokines IL-4 and IL-13 are derived

¹Department of Immunology and Infectious Disease and Centre for Personalised Immunology, The John Curtin School of Medical Research, The Australian National University, Canberra, Australia; ²Department of Immunology, Canberra Hospital, Canberra, Australia; ³Laboratory of Experimental Immunology, World Premier International Research Center Immunology Frontier Research Center, Osaka University, Osaka, Japan; ⁴Discipline of Pathology, School of Medical Sciences, Charles Perkins Centre, University of Sydney, New South Wales, Australia; ⁵Division of Infectious Diseases and HIV Medicine, Case Western Reserve University, Cleveland, OH; ⁶Cytokine Receptor Laboratory, Centre for Cancer Biology, Adelaide, Australia; ⁷Department of Systems Immunology and Braunschweig Integrated Centre of Systems Biology, Helmholtz Centre for Infection Research, Braunschweig, Germany; ⁸Department Pathology, San Raffaele Scientific Institute, Università Vita-Salute, Milan, Italy.

Correspondence to Carola G. Vinuesa: carola.vinuesa@anu.edu.au.

© 2019 Cañete et al. This article is available under a Creative Commons License (Attribution 4.0 International, as described at <https://creativecommons.org/licenses/by/4.0/>).

from type 2 innate lymphoid (ILC2) cells, basophils, or CD4⁺ helper T cells (Voehringer et al., 2006; Licona-Limón et al., 2013; Hammad and Lambrecht, 2015). These signature cytokines drive B cells to undergo class switch recombination (CSR) to IgE. There is evidence that IgE-producing plasma cells can arise both upon T cell priming of B cell differentiation along the extrafollicular route and upon interaction with T cells within epithelial lesions as a result of sequential CSR in IgG memory B cells that arose in GCs (Erazo et al., 2007; Xiong et al., 2012; He et al., 2013). IgE⁺ B cells can also be found in GCs (He et al., 2013), and several lines of evidence have suggested that T_{FH} cells contribute to IgE production (Glatman Zaretsky et al., 2009; King and Mohrs, 2009; Reinhardt et al., 2009; Coquet et al., 2015; Ballesteros-Tato et al., 2016). Recently, a dependence of T_{FH} cells has been confirmed for mouse IgE responses induced by airborne antigens (Kobayashi et al., 2017). IgE responses appear to be tightly regulated, particularly in the context of GC responses, with several B cell-intrinsic mechanisms shown to limit IgE production (Yang et al., 2012; He et al., 2013; Butt et al., 2015). To date, despite an established role of regulatory T (T reg) cells in IgE repression (Josefowicz et al., 2012a), whether T_{FR} cells contribute to IgE regulation remains unclear.

Here, we describe unique regulatory functions of human tonsillar follicular CD25⁺ T cells, which are the main producers of IL-10 among human T_F cells. Despite exhibiting similarities to mouse T_{FR} cells both phenotypically and functionally, CD25⁺ T_F cells in tonsil are FOXP3 negative. Strikingly, these cells exert a strong suppression on human T_{FH} cells and expression of key molecules BCL6, CD40L, and IL-21, while not directly suppressing plasma cell differentiation of B cells. CD25⁺ T_F cell-derived IL-10 dampened induction of Ig switching to IgE. In children, serum total IgE titers were inversely correlated with the frequencies of tonsil CD25⁺ T_F cells and IL-10-producing T_F cells, but not with total T reg cells, T_{FR} cells, or IL-10-producing T cells. Thus, CD25⁺ T_F cells emerge as a subset with unique T and B cell regulatory activities that may help prevent atopy.

Results

Identification of tonsillar CD25⁺ FOXP3⁻ T_F cells that express abundant IL-10

In an effort to identify the human equivalent of mouse T_{FR} cells, we stained cells from the most accessible human secondary lymphoid tissue, tonsil, for markers of T_F cells and T reg cells. Total T reg cells were identified by expression of CD25 in the absence of CD127 (Seddiki et al., 2006; Fig. 1 a). The majority of CD25⁺ CD127⁻ T reg cells found within the nonfollicular T effector gate (C-X-C motif chemokine receptor 5 [CXCR5]^{int} and programmed cell death protein 1 [PD-1]^{int}) expressed FOXP3, and as such constitute the conventional T reg cell population (Fig. 1, a and b). A CD25⁺ CD127⁻ subset was also identified within the follicular CXCR5^{hi} PD-1^{hi} population, which we refer to as “CD25⁺ T_F” cells (Fig. 1 a). We speculated that these might correspond to the T_{FR} cell population identified in mice. To our surprise, the majority (~95%) of these tonsillar CD25⁺ CD127⁻ T_F cells lacked FOXP3 expression (P < 0.0001; Fig. 1 b). We also identified follicular CD25⁺ FOXP3⁺ and CD25⁻ FOXP3⁺ T cells

(Fig. S3 a). CD25⁺ FOXP3⁻ T_F cells were also present in human mesenteric lymph nodes, although these were less frequent than those seen in the tonsil and found at proportions comparable to CD25⁺ FOXP3⁺ T cells (P = 0.2236; Fig. S1).

We were intrigued by the lack of FOXP3 expression among the majority of CD25⁺ T_F cells and considered that CD25 expression may be either identifying activated T_{FH} cells or a T reg cell subset independent of FOXP3 expression. FOXP3-negative T reg cells have been described in humans and mice, and among these are type 1 regulatory (Tr1) cells that produce IL-10 and are identified by expression of the lymphocyte activating gene 3 (LAG3; Gagliani et al., 2013). We therefore investigated the relative ability of the different tonsil T cell subsets to produce IL-10. Strikingly, staining for IL-10 revealed that the FOXP3⁻ CD25⁺ T_F cell population was the subset containing the largest fraction of IL-10-producing T cells in human tonsil (Fig. 1 c): 20–30% expressed IL-10 compared with 3–12% of conventional T reg cells (P < 0.0001) and barely any T_{FH} cells (P < 0.0001; Fig. 1 c). Unlike IL-10-producing Tr1 cells, IL-10-producing T_F cells did not express LAG3 (Fig. 1 d), suggesting that CD25⁺ T_F cells may not be Tr1 cells with a follicular phenotype.

Having demonstrated that only a fraction of CD25⁺ T_F cells expressed IL-10, we asked whether IL-10⁺ and IL-10⁻ T_F cells were fundamentally different subsets. Transcriptional profiling using Affymetrix RNA microarrays of IL-10-producing versus nonproducing T_F cells revealed highly comparable transcriptomes (Fig. 2 a), with only a few differentially expressed transcripts including IL-10, C-C motif chemokine 4-like (CCL4L2), T-lymphocyte maturation-associated protein (MAL), microRNA424 (MIR424; reported to be an activator of TGF- β signaling; Li et al., 2014), lymphoid enhancer binding factor 1 (LEF1; a transcription factor important for T_{FH} differentiation; Choi et al., 2015), and the surface receptor cell adhesion molecule 1 (CADM1), which has been previously reported to mark a population of adult T cell leukemia/lymphoma (Komohara et al., 2017). To evaluate whether CADM1 expression could identify the IL-10-producing CD25⁺ T_F cells, we stained human tonsils with an anti-CADM1 antibody together with T_F- and T reg cell-specific markers. While CADM1 expression failed to detect all IL-10-producing cells within the CD25⁺ T_F population (Fig. 2 b, upper panels), it did exclude most FOXP3-expressing T_F cells (Fig. 2 b, lower panels). Altogether, these results suggest that IL-10-producing and nonproducing CD25⁺ T_F cells are closely related, with IL-10 probably being produced upon CD25⁺ T_F cell activation.

CD25⁺ T_F cells suppress T cell proliferation

To investigate whether CD25⁺ T_F cells were simply activated T_{FH} cells that had up-regulated CD25, as previously proposed (Li and Pauza, 2015) and as opposed to T reg cells, we tested their ability to suppress T cell proliferation in vitro. Unlike typical T_{FH} cells (CD25⁻ CD127⁻), human CD25⁺ T_F cells and T reg cells were equally effective in suppressing conventional (CD25⁻ CD4⁺) T cell proliferation (Fig. 2 c), as previously described for mouse T_{FR} cells (Chung et al., 2011; Linterman et al., 2011). While it is unlikely that the small contaminating number of FOXP3⁺ cells among tonsillar CD25⁺ T_F cells (i.e., the equivalent to mouse T_{FR}




Figure 1. Identification of IL-10-producing CD25⁺ FOXP3⁻ human T_F cells in human tonsils. **(a)** Flow cytometric plots showing gating strategy to identify the indicated populations. **(b)** Flow cytometric plots and quantification ($n = 8$) showing percentage of FOXP3⁺ cells and FOXP3 mean fluorescence intensity (MFI) within the indicated subset according to panel a. Data are representative of 10 independent experiments. **(c)** Flow cytometric plots and quantification of PMA/ionomycin-stimulated tonsillar cell suspensions showing IL-10 expression in the indicated subset ($n = 9$). Data are representative of 10 experiments. **(d)** Flow cytometric plots and quantification showing LAG3 and CD25 expression in total T_F cells ($n = 14$; left), and IL-10 and LAG3 in CD25⁺ T_F cells ($n = 5$; right). Data are representative of five independent experiments. In all graphs, bars represent medians, and each dot represents a single tonsil donor. ****, $P \leq 0.0001$, nonparametric Mann-Whitney U test.

cells) is responsible for the overall suppressive effect of this population in the cell titration response, we sought to exclude the contribution from such T_{FR} cells. We used CADM1 to sort FOXP3⁻ CD25⁺ T_F cells. CADM1⁺ and CADM1⁻ T_F cells suppressed T cell proliferation as effectively as T_{reg} cells (Fig. 2 d), demonstrating that CD25⁺ T_F cells exert T_{reg} cell functions even in the absence of FOXP3 expression. Given that CADM1 does not distinguish IL-10⁺ versus IL-10⁻ CD25⁺ T_F cells, it is unlikely that IL-10 mediates suppression of T cell proliferation. Indeed, IL-10 blockade in these cultures did not affect this regulatory property of CD25⁺ T_F cells (data not shown). Altogether, these results suggest that human CD25⁺ CD127⁻ T_F cells are not simply activated T_{FR} cells, but can behave like T_{reg} cells.

To gain insights into the ontogeny of CD25⁺ T_F cells, and specifically ask whether these cells may have expressed FOXP3

as thymic T_{reg} cells (tT_{reg} cells) and later down-regulated it, we investigated the methylation of the conserved noncoding sequence 2 (CNS2) locus of the FOXP3 promoter. Demethylation of this locus is critical for the stable FOXP3 expression that characterizes tT_{reg} cells; Zheng et al., 2010). The CNS2 locus was >90% methylated in human CD25⁺ T_F cells (Fig. 3 a), suggesting a peripheral rather than thymic origin. Similarly, compared with conventional nonfollicular T_{reg} cells, CD25⁺ T_F cells expressed 40% less HELIOS (Fig. 3 b, $P = 0.0002$), a known target of FOXP3 (Fu et al., 2012) that is abundant in, although not exclusive of, tT_{reg} cells (Thornton et al., 2010; Szurek et al., 2015). Together, these data demonstrate the existence of human CD25⁺ FOXP3⁻ T_{FR} cells that are unlikely to originate from tT_{reg} cells and appear to be enriched in tonsillar tissue.




Figure 2. Human CD25⁺ T_F cells repress T cell proliferation. **(a)** Volcano plot showing Affymetrix RNA microarrays comparing gene expression between IL-10-producing and nonproducing T_F cells ($n = 3$). **(b)** Flow cytometric plots showing coexpression of CADM1 with IL-10 or FOXP3 in PMA/ionomycin-stimulated tonsillar cell suspensions in the indicated subset ($n = 5$). Data are representative of two independent experiments. **(c and d)** Flow cytometric plots and quantification of proliferating CD4⁺CD25⁻ responder T cells according to dilution of the cytoplasmic fluorescent dye CTV after 3 d of α -CD3 and α -CD28 stimulation in the presence or absence of T_{reg} cells ($n = 10$), CD25⁺ T_F cells ($n = 8$), and CD25⁻ T_{FH} cells ($n = 5$; c) or T_{reg} cells ($n = 3$), CADM1⁺CD25⁺ T_F cells ($n = 3$), and CADM1⁻CD25⁺ T_F cells ($n = 3$; d). Bars represent medians, and each dot represents the mean value of cultures set up in triplicate from a single donor. Data are representative of five (c) and two (b and d) independent experiments. ns, not significant; **, $P \leq 0.01$; ***, $P \leq 0.001$, nonparametric Mann-Whitney U test (c, left panel) and two-way ANOVA (c, right panel, and d).




Figure 3. Human CD25⁺ T_F cells may be peripherally induced and express high CTLA4 but low IL-2 production. **(a)** Bisulfite sequencing of the 11 CpG islands of the FOXP3 CNS2 locus with 12 representative clones per population and per donor (filled circle, methylated; open circle, demethylated). **(b)** Flow cytometric plots and quantification showing HELIOS expression in the indicated cell subsets ($n = 8$). Data are representative of three independent experiments. **(c)** Flow cytometric plots and quantification ($n = 8$) showing CTLA4 expression in the indicated cell subset. Data are representative of five independent experiments. **(d and e)** Flow cytometric plots and quantification showing gating strategy (d) and IL-2 expression of PMA/ionomycin-stimulated tonsillar cell suspensions in the indicated subset ($n = 8$; e). Data are representative of two independent experiments. In all graphs, bars represent medians, and each dot represents a single donor. **, $P \leq 0.01$; ***, $P \leq 0.001$, nonparametric Mann-Whitney *U* test. FSCA, forward scatter.

We next investigated whether, in spite of absent FOXP3 expression, CD25⁺ T_F cells display the two key characteristics of conventional T_{reg} cells known to be crucial for their suppressor function: high cytotoxic T lymphocyte-associated protein 4 (CTLA4) expression and low IL-2 production. CTLA4 is one of

the most important targets of FOXP3 in T_{reg} cells, and high amounts of CTLA4 expression plus absence of IL-2 production is a good marker of T_{reg} cell function even in the absence of FOXP3 expression (Yamaguchi et al., 2013). We found that CD25⁺ T_F cells had the highest amount of CTLA4 among tonsil

T cell subsets (Fig. 3 c), and IL-2 production was detected in only ~5% of human tonsil CD25⁺ T_F cells (Fig. 3, d and e). These results further suggest that CD25⁺ T_F cells are indeed a T reg cell subset.

Human CD25⁺ T_F cells resemble mouse T_{FR} cells

To investigate the extent to which human CD25⁺ T_F cells resemble mouse T_{FR} cells, we obtained the transcriptional signature of CD25⁺ T_F cells. As shown before, we noted that among T_F cells, CD25⁺ T_F cells did not overlap with LAG3⁺ T_F cells. Thus, we FACS-purified naive T cells and the three major T_F cell subsets according to CD25 and LAG3 expression, CD25⁺ LAG3⁻ T_F cells, CD25⁻ LAG3⁺ T_F cells, and CD25⁻ LAG3⁻ T_{FH} cells (Fig. 4 a), and performed RNA sequencing (RNA-seq). Paired analyses from three different tonsil donors revealed that CD25⁺ T_F cells were fundamentally different from naive, T_{FH}, and LAG3⁺ T_{FH} cells (Fig. 4 b) and were remarkably similar to the phenotype described for mouse T_{FR} cells (FOXP3⁺ T_F cells). Indeed, human CD25⁺ T_F cells expressed key molecules required for T_{FH} development, including BCL6, and showed the highest expression of transcripts associated with effector T reg cells including CTLA4, glucocorticoid-induced TNFR-related protein (GITR), PR domain zinc finger protein 1 (PRDM1), Runt-related transcription factor 2 (RUNX2), C-C chemokine receptor type 5 (CCR5), and IL-10 (Fig. 4 c; Fu et al., 2012). Similar to mouse T_{FR} cells, human CD25⁺ T_F cells expressed the lowest amount of the key B cell helper molecule CD40L but abundant IL-21 transcript and protein (Fig. S2, a and b), which is low in mouse T_{FR} cells but has also been shown to be expressed in T_{FR} cells from macaques (Chowdhury et al., 2015).

Flow cytometric analysis of protein expression validated our transcriptomic analyses and confirmed the similarities between CD25⁺ T_F cells and T reg cells (Fig. 4 d). Also, minimal differences were observed between the small fraction of tonsil follicular FOXP3⁺ CD25⁺ and the more abundant FOXP3⁻ CD25⁺ T_F cells. The latter expressed more IL-10 and BCL6, whereas FOXP3⁺ cells expressed more HELIOS and GITR (both direct targets of FOXP3; Fig. S3). Together, these results suggest that human CD25⁺ T_F cells have a gene expression profile that allows migration to facilitate T:B cell encounters and T reg cell function as well as abundant expression of IL-10.

Human CD25⁺ T_F cells regulate T_{FH} cells and B cell IgE production

In mice, both B and T_{FH} cells have been suggested to be targets of T_{FR} cell suppression (Sage and Sharpe, 2016). To test whether CD25⁺ T_F cells could regulate T_{FH} cell function, we analyzed T_{FH} cells after 3 d in culture with autologous memory B cells and CD25⁺ T_F cells in the presence of *Staphylococcus* enterotoxin B (SEB). Addition of CD25⁺ T_F cells suppressed T_{FH} cell proliferation (Fig. 5 a). In addition, we observed that CD25⁺ T_F:T_{FH} co-cultures reduced the percentage of CD40L-expressing T_{FH} cells ($P < 0.0001$) and also reduced CD40L expression per T_{FH} cell by 50% ($P < 0.0001$; Fig. 5 b), as well as IL-21 production ($P < 0.0001$) and BCL6 expression ($P = 0.0010$; Fig. 5 b). Reduction of each of these T_{FH} cell molecules is known to limit T_{FH} cell help for B cells (Vinuesa et al., 2016). Similar results were obtained in

cultures in which B cells were not included and T_{FH} cells were activated with αCD3/αCD28 antibodies (Fig. 5 c), suggesting that CD25⁺ T_F cells act directly on T_{FH} cells to suppress their function. These regulatory effects were not IL-10 mediated, as addition of an IL-10 blocking antibody did not rescue CD25⁺ T_F cell-mediated repression of T_{FH} cell proliferation or expression of CD40L, IL-21, or BCL6 (Fig. S4 a).

Next, we compared the ability of human T_{FH} and CD25⁺ T_F cells to induce B cell responses. Human memory B cells were cocultured with autologous T_{FH} or CD25⁺ T_F cells for 7 d in the presence of SEB. Coculture of B cells with T_{FH} cells successfully induced cell division, differentiation of plasma cells (Fig. 5 d), and secretion of IgG and IgE (Fig. 5 e). Coculture with CD25⁺ T_F cells induced plasma cell differentiation, although to a lesser extent (Fig. 5 d, left), and modest B cell proliferation (Fig. 5 d, right) and resulted in a 5.2-fold decrease in IgE secretion and a 2.7-fold decrease in IgG compared with T_{FH} cocultures (Fig. 5 e). Addition of T_{FH} cells to the CD25⁺ T_F:B cell cocultures at equal ratios rescued plasma cell production, B cell proliferation, and IgG production, but not IgE production (Fig. 5 d and e). Together, these data suggest that human CD25⁺ T_F cells are able to regulate T_{FH} cell function and B cell IgE production.

Human CD25⁺ T_F cell-derived IL-10 represses epsilon germline transcription

Early reports from human in vitro studies showed that IL-10 could suppress class-switching to IgE but not IgG (Jeannin et al., 1998; Akdis and Blaser, 2001). Since IL-10 can have additional effects on B cells such as promoting plasma cell differentiation (Arpin et al., 1995), we sought to separate the observed action of CD25⁺ T_F cells on plasma cell induction (Fig. 5 d) from possible repressive effects on Ig switching to IgE by looking at the earliest event that occurs in cells undergoing CSR: production of germline transcripts (GLTs). To investigate direct CSR events to IgE from IgM⁺ B cells as opposed to sequential CSR, in which an initial IgM-to-IgG1 CSR event is followed by switching to IgE, we limited our culture to 25 h. This is sufficient to induce the production of εGLTs, but is not sufficient for naive B cells to undergo two cell divisions, thus preventing sequential CSR from occurring (Erazo et al., 2007). Naive CD19⁺ IgD⁺ human B cells were purified and cocultured with T_{FH} cells and stimulated with IL-4, IL-13, and SEB in the presence or absence of CD25⁺ T_F cells. Although addition of CD25⁺ T_F cells did not have a statistically significant effect on γGLT induction, it consistently reduced εGLTs in T_{FH}:B cell cocultures (Fig. 5 f). While there was high variability across individuals, paired statistics revealed a reduction in εGLT across 18 human donors ($P = 0.0006$). CD25⁺ T_F cells repressed εGLTs in IgM⁺ but not in IgG⁺ memory B cells (Fig. S4 b), suggesting that CD25⁺ T_F cells preferentially repress direct as opposed to sequential induction of CSR to IgE, or that CD25⁺ T_F cells preferentially interact with IgM⁺ B cells (naive or memory).

IL-10 has been shown to suppress εGLTs in human B cell cultures (Jeannin et al., 1998). We therefore evaluated the contribution of IL-10 to CD25⁺ T_F cell-mediated suppression of εGLTs. Blocking the IL-10 receptor (Fig. S4 c) did not have a profound effect on γGLT but rescued εGLTs in CD25⁺ T_F:B cell




Figure 4. Human CD25⁺ T_F cells' transcriptome resembles that of murine T_{FR} cells. **(a)** Flow cytometric plots showing the gating strategy used to FACS-purify each of the indicated cell subsets. **(b)** Heatmap analysis of RNA-seq data showing differentially expressed genes in CD25⁺ T_F cells compared with the indicated T cell populations (\log_2 value of counts per million) extracted from the tonsils of three individuals. **(c)** Selected transcripts from panel b in the indicated subsets (RNA counts per million [CPM]). **(d)** Flow cytometric plots and quantification ($n = 8$) of the indicated proteins. Data are representative of at least three independent experiments. In all graphs, bars represent medians; each dot represents a single tonsil donor. ***, $P \leq 0.001$, nonparametric Mann-Whitney U test.

cocultures (Fig. 5 g) in 8 out of 10 donors. Taken together, these data suggest that IL-10 contributes to suppression of IgE production by human CD25⁺ T_F cells.

IL-10-producing CD25⁺ T_F cells inversely correlate with circulating IgE in the serum

To gain some insights into the possible clinical relevance of CD25⁺ T_F cells, we collected blood from pediatric tonsil donors at the time of tonsillectomy and produced a cohort of matched serum and tonsil samples from 50 individuals. We then investigated associations between total IgE in serum and CD25⁺ T_F cells in tonsils. Strikingly, we observed an inverse correlation

between the frequency of CD25⁺ T_F cells, measured as either a percentage of T_{reg} cells or of CD4⁺ T cells, and the amount of IgE in serum (Fig. 6 a). The more frequent CD25⁺ T_F cells were, the less abundant total IgE in serum, suggesting that CD25⁺ T_F cells may indeed regulate IgE production. Although the goodness of fit in this model was somewhat weak ($r = 0.5107$), the deviation from zero in the correlation was highly significant, demonstrating a negative correlation between the two variables. Similar results were obtained when we correlated the amount of total IgE and the frequency of IL-10-producing T_F cells (Fig. 6 a). Compared with CD25⁺ T_F cells that lack FOXP3 expression, no negative correlation was observed between serum IgE and the




Figure 5. Human CD25⁺ T_F cells repress T_{FH} cells and induction of IgE switching. (a and b) Flow cytometric plots (a) and quantification ($n = 8$; b) of CTV-labeled- T_{FH} cells, cocultured with memory B cells with or without CD25⁺ T_F cells, showing expression of the indicated proteins after 3 d. Upper panels are representative data for a single donor, and lower panels are pooled values from different donors and experiments. Data in each panel is representative of at least five independent experiments. **(c)** Flow cytometric plots and quantification of CTV-labeled T_{FH} cells, cocultured with or without CD25⁺ T_F cells, showing expression of the indicated proteins after 3 d of α -CD3 and α -CD28 stimulation. **(d)** Quantification of plasma cell (CD27⁺ CD38⁺) differentiation ($n = 25$) or proliferation ($n = 12$) from memory B cells cocultured with T_{FH} cells, CD25⁺ T_F cells, or both in the presence of SEB (500 ng/ml), IL-4, and IL-13 (40 ng/ml) for 7 d. **(e)** IgG ($n = 10$) or IgE ($n = 14$) in coculture supernatants of cocultures as in panel d. Data were normalized to values from cultures without T cells. Data in d and e were pooled from eight independent experiments. **(f and g)** Quantification of γ GLTs (top) or ϵ GLTs via qPCR in naive B cells incubated with T_{FH} cells with or without CD25⁺ T_F cells ($n = 19$; f), and naive B cells cocultured with CD25⁺ T_F cells in the presence or absence of an IL-10 blocking antibody ($n = 10$; g). GLT expression values were calculated using the $\Delta\Delta CT$ method and were normalized to RPL13A levels, then normalized to the untreated control. Data were pooled from five independent experiments. Bars represent means of three technical replicates (b and c) or medians (d–g), each dot represents a single tonsil donor, and error bars represent standard deviations. ns, not significant; *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$; ****, $P \leq 0.0001$, two-tailed Student's t test (b, top panels, and c–g) and nonparametric paired Wilcoxon test (b, bottom panels). All data are representative of at least three independent experiments. MFI, mean fluorescence intensity; RU, relative units.




Figure 6. Tonsillar CD25⁺ T_F cells inversely correlate to total IgE in the serum. (a and b)

Pearson correlation analyses between serum total IgE and the frequency of the indicated cell subset in tonsil ($n = 49$). Data are representative of two independent experiments that were pooled together. **(c)** Histogram showing the frequency distribution of serum total IgE (log₁₀ ng/ml) from 49 children (mean = 2.07). **(d)** Quantification of the frequency of the indicated cell subset from tonsil donors with high (>100 ng/ml) or low (<100 ng/ml) total serum IgE titers. Bars represent medians, and dots represent individual tonsil donors ($n = 49$). Data are representative of two independent experiments that were pooled together. ***, $P \leq 0.001$; ****, $P \leq 0.0001$, nonparametric Mann-Whitney test U test.

frequency of FOXP3⁺ T_{FR} cells; in fact, a positive correlation was seen (Fig. 6 b, left). No correlations were detected between serum IgE and total T reg cells or even total IL-10-producing CD4⁺ T cells (Fig. 6 b, right), suggesting that CD25⁺ T_F cells are the cells specialized in preventing unwanted IgE production.

Given that IL-10 has been reported to cooperate with IL-4 in inducing B cell production of IgG4, and that the latter correlates with allergy desensitization and tolerance, we asked whether CD25⁺ T_F cells could also regulate IgG4 responses (Akdis and Akdis, 2014). Pearson correlation analysis revealed no relationship between circulating IgG4 and the frequency of CD25⁺ T_F cells or IL-10-producing T_F cells (Fig. S5 a). A positive correlation was observed between the frequency of T reg cells and IgG4 (Fig. S5 b), possibly suggesting a division of labor between T reg cells and CD25⁺ T_F cells in promoting tolerogenic IgG4 versus repressing IgE responses.

Finally, we asked whether the frequency of CD25⁺ T_F cells differed between those donors who exhibited high amounts of IgE versus those with low titers. The clinical value for normal total IgE in children ranges from 2 to 200 IU, or 2.4 to 480 ng/ml (Martins et al., 2014). An average of total serum IgE in healthy

individuals of 100 ng/ml has been reported (Gould, 1998) and was the average value found in our tonsil/serum cohort (Fig. 6 c). We selected 100 ng/ml as a cutoff to separate tonsil donors with high versus low IgE titers and thus interrogated whether the frequency of CD25⁺ T_F cells differed in both groups. CD25⁺ T_F cells, as a percentage of T reg cells or CD4 T cells, were significantly less frequent in donors with high IgE titers (62% and 42% median decrease, respectively; Fig. 6 d). Similarly, the frequency of IL-10-producing T_F cells was more abundant in donors with low IgE titers. Altogether, these results show an inverse correlation between CD25⁺ T_F cells and serum IgE, suggesting that CD25⁺ T_F cells are physiological regulators of IgE production.

Discussion

Here we have described a novel regulatory function of a unique human T_F cell subset that is abundant in tonsil, CD25⁺ T_F cells, and is the major IL-10-producing subset among T_F cells. CD25⁺ T_F cells are identified by high expression of the GC T_{FH} cell markers PD-1 and CXCR5 together with T reg cell markers

(CD25^{hi} CD127^{lo} CTLA4^{hi} IL-2^{lo}) and high IL-10 production in the absence of FOXP3 expression. Several lines of evidence indicate that this subset is functionally related to FOXP3-expressing T reg cells. First, our transcriptomic and proteomic characterization of CD25⁺ T_F cells has revealed that they closely resemble mouse T_{FR} cells, except for the absence of FOXP3 expression and IL-21 production. A CD25⁺ nonfollicular T reg cell population that also lacks FOXP3 expression and produces IL-10 has been previously described (Facciotti et al., 2016). These cells regulate B cell responses and, when dysregulated, are associated with autoimmune diseases. Second, besides high CTLA4 expression, CD25⁺ T_F cells express low levels of IL-2. These observations are consistent with previous evidence that T cells engineered to express CTLA4 and not IL-2 adopt full T reg cell-like activity even in the absence of FOXP3 (Yamaguchi et al., 2013). Absence of demethylation in the FOXP3 CNS2 promoter, together with low HELIOS expression, suggests that this subset may be induced in the periphery, arising from conventional naive CD4⁺ T cells or from T_{FH} cells themselves.

Functionally, we show that human CD25⁺ T_F cells exert suppressive effects on both T and B cells. With respect to T cells, CD25⁺ T_F cells repress total T cell and T_{FH} cell proliferation and function. CD25⁺ T_F cells dampen T_{FH} cell help through direct down-regulation of CD40L, IL-21, and BCL6 on T_{FH} cells. CD25⁺ T_F cells also exert regulatory effects on B cells. First, CD25⁺ T_F cells appear to dampen B cell CSR to IgE. The latter occurs in an IL-10-dependent manner. Indeed, CD25⁺ T_F cells reduce ϵ GLT production in T_{FH}:B cell cocultures, and this effect is rescued by IL-10 blockade. This is interesting, because IL-10 acts very early, within 25 h of naive (unswitched) B cell activation, to suppress CSR by inhibiting ϵ GLT production. Our experimental system shows this effect is a direct consequence of limiting direct CSR from μ to ϵ , rather than inhibiting γ GLT production and subsequent sequential class-switching. It is still unclear whether CD25⁺ T_F cells predominantly target naive, GC B, or memory B cells in vivo. T_{FH} cells influence B cell activation at multiple stages, from the earliest B cell priming stages at the T:B border where Ig CSR is initially triggered (Jacob et al., 1991; Toellner et al., 1996), to selection in GCs, and reactivation as memory B cells (Vinuesa et al., 2016). Thus, CD25⁺ T_F cells are also likely to interact with and influence naive and memory B cell isotype switching.

Interestingly, and as shown for mouse IL-10-producing T_F cells, CD25⁺ T_F cells induced some plasma cell differentiation, although to a lesser extent than T_{FH} cells. This contrasts with mouse T_{FR} cells that effectively repress B cell differentiation (Sage et al., 2016; Botta et al., 2017; Maceiras et al., 2017). One obvious difference between mouse (FOXP3⁺) T_{FR} cells and the human tonsillar FOXP3⁻ CD25⁺ T_F cells described here is the production of IL-21 by the latter but not the former. IL-21 is known to promote plasma cell differentiation in mice (Ozaki et al., 2002) and humans (Ettinger et al., 2005), so it is possible that this cytokine contributes to plasma cell differentiation, potentially in conjunction with CD25⁺ T_F cell-derived IL-10. Although we could not see effects of IL-10R blockade on plasma cell differentiation in 7-d cultures (unpublished data), it is possible that the conditions were not optimal for durable IL-10

blockade in these assays. Interestingly, unlike in mice, in which IL-21 inhibits IgE formation (Ozaki et al., 2002), IL-21 is a potent inducer of IgE in CD40L-stimulated human B cells (Berglund et al., 2013). Thus, there appears to be dissociation between the effects of CD25⁺ T_F cells on plasma cell formation, possibly conferred by IL-21 with or without contribution of IL-10, and CSR conferred by IL-10. CD25⁺ T_F cells emerge as a unique regulatory cell that controls switching independently of B cell differentiation.

Thymus-derived T reg cells that stably express FOXP3 are selected on the basis of their ability to recognize self-antigen in the thymus (Hsieh et al., 2012). T reg cells give rise to T_{FR} cells in immunized mice (Chung et al., 2011; Linterman et al., 2011; Wollenberg et al., 2011; Sage et al., 2014; Wing et al., 2014; Aloulou et al., 2016), making them effective suppressors of responses against self. Recent reports have also suggested mouse T_{FR} cells can be induced in the periphery in responses to both self and foreign antigens (Aloulou et al., 2016). The human CD25⁺ T_F cells described in this study lack FOXP3 expression and appear not to be thymus derived. The fact that CD25⁺ T_F cells are so abundant in tonsils, which are exposed to oral and inhaled antigens, but are scarce in other lymphoid tissues that are not exposed to such antigens, makes them particularly good candidates to suppress responses to harmless foreign antigens. Furthermore, our observation that CD25⁺ T_F cells repress CSR to IgE and are associated with lower circulating IgE titers in children undergoing tonsillectomy suggests that this subset may be important to prevent IgE-mediated allergic reactions to inhaled or ingested antigens.

Besides a role in preventing immune responses and proallergenic IgE to innocuous antigens, the chronic and ongoing nature of GC reactions in human tonsils makes this subset highly suitable to control excessive and harmful inflammation to foreign antigens, thus serving as a natural homeostatic mechanism to curtail T_{FH} cell responses. Equivalent T reg cell subsets have been described that are specialized in repressing T_H2 and T_H1 lineages (Jankovic et al., 2007; Altin et al., 2012; Josefowicz et al., 2012b). There is evidence that T_{FR} cells can be induced against both self and foreign antigens (Aloulou et al., 2016), but it remains unclear whether self- or innocuous environmental antigens such as allergens are more conducive to T_{FR} or CD25⁺ T_F cell development.

In summary, our work unveils a novel subset of T_{FR} cells in humans, characterized by abundant production of IL-10, which regulates several aspects of T and B cell responses and prevents induction of switching to IgE. Importantly, low numbers of CD25⁺ T_F cells are associated with elevated circulating IgE levels. We therefore predict that deficiencies in homeostasis or function of this cell subset could underpin susceptibility to allergic and anaphylactic reactions induced by inhaled and ingested antigens. Should this be the case, these findings may open up new avenues for boosting CD25⁺ T_F cells to reduce the risk of allergy.

Materials and methods

Human tonsil and lymph node cells

Human tonsils were obtained from children undergoing routine tonsillectomy. Tonsillar lymphocyte single-cell suspensions were prepared by mechanical disruption of the tissue followed by cell separation using Ficoll-Hypaque (GE Healthcare Life Sciences) gradient and frozen until further use, except for RNA-seq, for which fresh samples were used. Human mesenteric lymph nodes were obtained as discarded tissue from nonmalignancy gastrointestinal surgery. Informed consent was obtained from all subjects. All experiments with human samples were approved by the Australian National University's Human Experimentation Ethics Committee and the Australian Capital Territory Health Human Research Ethics Committee.

Flow cytometry

Tonsillar lymphocytes were stained with the following anti-human antibodies: anti-CADM1 biotin (3E1; MBL), anti-CD4 APC Cy7 (RPA-T4; BD Biosciences), anti-CD8 FITC (RPA-T8; BD Biosciences) or PE (SK1; BD Biosciences), anti-CD19 FITC (SJ25C1; BioLegend) or PE Cy7 (SJ25C1; BD Biosciences), anti-CXCR5 Alexa Fluor 488 or 647 or PerCP Cy5.5 (RF8B2; BD Biosciences) or PerCP/Cy5.5 (J252D4; BioLegend), anti-CD45 RA (HI100; BioLegend), anti-CD45RO (UCHL1; BioLegend), anti-CTLA-4 PE (BNI3; BD Biosciences) or PE Cy7 (L3D10; BioLegend), anti-PD-1 PE (MIH4; eBioscience) or BV605 or BV421 (EH12.2H7; BioLegend) or unconjugated (J105; eBioscience), anti-CD127 FITC (11-1278; eBioscience) or BV421 (A019D5; BioLegend), anti-CD25 biotin (BC96; eBioscience or BioLegend) or PE-Cy7 (BC96; BD Biosciences or BioLegend) or APC (2A3; BD Biosciences), anti-CD127 FITC (A019D5; eBioscience) or BV421 (A109D5; BioLegend) or BV510 (HIL-7R-M21; BD Biosciences), anti-FOXP3 Alexa Fluor 647 (259D; eBioscience), anti-GITR PE (110416; R&D Systems), anti-CTLA-4 PE-CF-594 (BNI3; BD Biosciences) or PE Cy7 (L3D10; BioLegend), anti-CD40L Pacific Blue or FITC (24-31; BioLegend), anti-BCL6 Alexa Fluor 647 or PE-Cy7 (K112-91; BD Biosciences), anti-BLIMP-1 Alexa Fluor 647 (646702; R&D Systems), anti-CXCL13 APC (53610; R&D Systems), anti-Helios PE (22F6; BioLegend), anti-IL-10 PE (JES3-19F1; BioLegend) or APC (JES3-19F1; BD Biosciences), anti-IL-10R PE (3F9; iCyt), anti-IL-21 PE (3A3-N21; BD Biosciences), anti-IL2 Alexa Fluor 488 (MQ1-17H12; BioLegend), anti-CD3 APC (HIT3a; BD Biosciences) or Alexa Fluor 700 (UCHT1; BD Biosciences) or Pacific Blue (OKT3; BioLegend) or BV510 (OKT3; BioLegend), anti-CD27 FITC or APC (M-T271; BD Biosciences), anti-CD38 FITC (HIT2; BD Biosciences) or PE (HB7; BD Biosciences), anti-LAG3 PE or FITC (2319-L3-050; R&D Systems), and anti-pSTAT3 Alexa Fluor 647 (4/P-STAT3; BD Biosciences). Intracellular staining was performed using the FOXP3/Transcription Factor Staining Buffer Set (eBioscience) or Cytofix/Cytoperm (BD Biosciences) according to the manufacturer's instructions. LAG3 was stained at 37°C for 15 min in the dark.

Mesenteric lymph nodes were stained with the following anti-human antibodies: anti-CD3 BUV395 (UCHT1; BD

Biosciences), anti-CD4 PerCP (RPA-T4; BioLegend), anti-CD8 APC-C7 (RPA-T8; BioLegend), anti-CD19 APC-Cy7 (HIB19; BioLegend), anti-CD45RA PE-TR (MEM-56; Invitrogen), anti-CXCR5 BV510 (RF8B2; BD Biosciences), anti-PD-1 BV421 (EH12.2H7; BioLegend), anti-CD127 BV650 (HIL-7R-M21; BD Biosciences), anti-CD25 PE-C7 (BC96; BioLegend), anti-FOXP3 PE (259D/C7; BD Biosciences), and anti-CD27 BV711 (L128; BD Biosciences). Intracellular staining was performed using the FOXP3/Transcription Factor Staining Buffer Set according to the manufacturer's instructions.

CpG methylation analysis by bisulfite sequencing

Genomic DNA was prepared using the NucleoSpin Tissue XS kit (Macherey-Nagel). After sodium bisulfite treatment (Methyl-Easy Xceed, Human Genetic Signatures), modified DNA was amplified by PCR and subcloned into PCR2.1-TOPO Vector (Invitrogen). PCR primers used were 5'-TTGGGTTAACGTTGTTGTTGTTGAGGATAG-3' and 5'-ATCTAAACCCATTATCACAAACCC-3'. The colonies (16–48 colonies/region) were directly amplified using the Illustra TempliPhi Amplification Kit (GE Healthcare) and sequenced.

In vitro stimulation

Intracellular cytokine staining was performed following 4–6 h of PMA (50 ng/ml) and ionomycin (500 ng/ml) stimulation with GolgiStop (BD Biosciences) or Brefeldin A (BioLegend) in RPMI 1640 medium supplemented with 10% FCS, 2 mM L-glutamine, 100 U penicillin-streptomycin, 0.1 mM nonessential amino acids, 100 mM Hepes, and 44 μM 2-mercaptoethanol. 200,000 naïve B cells were FACS-purified and stimulated with 50 ng/ml recombinant human IL-10 (Peprotech) for 30 min with or without 5 μg/ml of anti-IL-10 blocking antibody (3F9; BioLegend).

Secreted cytokine surface capture

Live CD4⁺ human tonsillar lymphocytes were stained with the anti-human IL-10 catch reagent (Miltenyi) at 9 million cells per 100 μl. After 10 min on ice, the cells were diluted 1:20 in RPMI supplemented as above. Cells were stimulated with PMA (100 ng/ml) and ionomycin (500 ng/ml) and incubated for 2 h in a 37°C incubator with 5% CO₂, rotating slowly using a MacsMix (Miltenyi). Cells were then stained as above using the anti-human IL-10 detection antibody (Miltenyi).

Microarray RNA analysis

IL-10-positive and -negative CD25⁺ T_F cells were FACS-purified from three human subjects. mRNA was extracted, and samples were sent to the Ramaciotti Centre for Genomics (Sydney, Australia) for analysis with Affymetrix GeneChip Human Gene 2.0 ST microarrays. The resulting nine CEL files were imported into Partek Genomics Suite (version 6.6) using the RMA algorithm with RMA background correction, quantile normalization, and median polish probe set summarization. Differentially expressed probe sets were identified in Partek GS using a two-factor ANOVA model with the factor SubjectID. The data are deposited under the National Center for Biotechnology Information Gene Expression Omnibus accession no. GSE79887.

Autologous human tonsillar T_{FH} :B cell cocultures and IgE detection

20,000 human FACS-purified T_{FH} and/or CD25 $^{+}$ T_F cells were cocultured with 100,000 GC B (CD4 $^{-}$ CD19 $^{+}$ CD38 $^{+}$ CD27 int) cells or memory B (CD4 $^{-}$ CD19 $^{+}$ CD38 $^{-}$ CD27 $^{+}$) cells in the presence of SEB (500 ng/ml; Sigma-Aldrich) for 3, 5, or 7 d. For IgE-inducing conditions, recombinant IL-4 and IL-13 (Peprotech) were used at 40 ng/ml. Culture supernatants, or human serum, were used to measure IgE, IgG, and IgG4 using the Cytometric Bead Array flex sets (558682, 558679, and 558678; BD Bioscience) according to the manufacturer's instructions. For proliferation assays, cells were stained with CellTrace Violet (CTV; Thermo Fisher Scientific) according to manufacturer's instructions. IL-10R blockade was achieved using anti-IL-10R blocking antibody (3F9; BioLegend) or isotype control (RTK2758; BioLegend) at 5 μ g/ml. For ϵ GLT detection, 100,000 naive B cells (IgD $^{+}$ CD19 $^{+}$ cells) were incubated with recombinant IL-4 and IL-13 together with SEB with or without T_{FH} or CD25 $^{+}$ T_F cells for 24–25 h. RNA was extracted using phenol/chloroform extraction, and cDNA was synthesized using SuperScript III (Invitrogen), according to the manufacturer's instructions. The following transcripts were measured via qPCR and analyzed using the $\Delta\Delta CT$ method. ϵ GLTs were detected using the following primers (forward: 5'-TGCATC CACAGGCACCAAAT-3', reverse: 5'-ATCACCGGCTCCGGGAAG TA-3) and normalized to *RPL13A* expression levels (forward: 5'-CTCAAGGTGTTGACGGCATCC-3', reverse: 5'-TACTTCCAGCCA ACCTCGTGAG-3'). γ GLTs were measured by using the following primers (forward: 5'-TCCAAGCCAACAGGGCAGGACACACCC AGAG-3', reverse: 5'-AAGTAGTCCTTGACCAGGCAG-3').

Human T reg cell suppression assays

10,000 CTV-labeled FACS-purified responder T cells (CD4 $^{+}$ CD3 $^{-}$ CD25 $^{-}$) were cocultured in the presence or absence of serially diluted follicular or nonfollicular T reg cells starting with 20,000 cells. Cells were stimulated with α CD3/ α CD28 microbeads (Miltenyi) at a 1:1 ratio of beads to T responder cells. After 3 d, cells were stained with 7-aminoactinomycin D and then analyzed by flow cytometry.

RNA-seq and analysis

The different subsets of human T_F cells were FACS-purified from three fresh tonsils. mRNA was extracted and sent to the Australian Cancer Research Foundation Biomolecular Resource Facility, The John Curtin School of Medical Research, The Australian National University (ANU; Canberra, Australia), for library construction using the TruSeq Stranded mRNA LT Sample Prep Kit (Illumina). Library samples were sequenced on a Hi-Seq2000 machine with a coverage of 25 million reads. The data were then sent to the Genome Discovery Unit (ANU Bioinformatics Consulting Unit) for analysis. Initial quality control checks with FastQC revealed that none of the 12 samples were problematic. All reads were aligned to the *Homo sapiens* genome reference sequence using TopHat version 2.0.13 with default parameters. Read counts were then generated for each gene in each sample using featureCounts version 1.4.6-p1 by using annotated gene locations. Differential expression analysis was performed using the edgeR package, version 3.10. Read counts

per gene were normalized by trimmed mean of M-values. Because edgeR uses the negative binomial distribution as its basic model for differential expression data, dispersion estimates were obtained using the quantile-adjusted conditional maximum likelihood method for single-factor experiments. Then, the quantile-adjusted conditional maximum likelihood-based exact test for the negative binomial distribution was performed to test for differentially expressed genes in our groups of samples. We used a Benjamini-Hochberg adjusted P value threshold of 0.05 to identify significantly differentially regulated genes. Data are deposited under National Center for Biotechnology Information Sequence Read Archive accession no. SRP072739.

Statistical analyses

All data were analyzed with nonparametric Mann-Whitney *U* test, except for some human cell culture experiments, in which paired Student's *t* tests or two-way ANOVAs were used. Paired analyses were performed using the Wilcoxon test. All statistical analyses were performed with Prism software (version 6, GraphPad). Statistically significant differences are indicated as *, $P \leq 0.05$; **, $P \leq 0.01$; ***, $P \leq 0.001$; ****, $P \leq 0.0001$; and ns, not significant.

Online supplemental material

Fig. S1 shows the frequency of CD25 $^{+}$ cells in human lymph node tissue. Fig. S2 shows IL-21 production by the different T_F cell subsets. Fig. S3 compares T reg cell- and T_{FH} cell-associated protein expression among CD25 $^{+}$ T_F cells and the more classical FOXP3 $^{+}$ T_{FR} cells. Fig. S4 shows that CD25 $^{+}$ T_F -mediated repression of T_{FH} cells is IL-10 independent, as well as repression of ϵ GLTs in IgG $^{+}$ and IgM $^{+}$ memory B cells. Fig. S5 shows Pearson correlation analyses between the frequency of CD25 $^{+}$ T_F cells in tonsil versus IgG4 serum titers.

Acknowledgments

The authors thank Anastasia Wilson and Ann-Maree Hatch for assistance with obtaining tonsil samples and obtaining clinical data; the Australian Cancer Research Foundation Biomolecular Resource Facility and the Imaging and Cytometry Facility (John Curtin School of Medical Research), especially H. Vohra and M. Devoy for FACS sorting and S. Ohms for help with microarray analysis; and B. Kaehler and A. Chua for RNA-seq analysis.

C.G. Vinuesa was supported by the Australian National Health and Medical Research Council fellowship (APP1117812), program and project grants (APP1113577), and a Human Frontier Science Program program grant (RGPO033/2015).

The authors declare no competing financial interests.

Author contributions: P.F. Cañete and R.A. Sweet designed, planned, and conducted experiments; P. Gonzalez-Figueroa and I. Papa provided substantial intellectual input and performed additional experiments; N. Ohkura and S. Sakaguchi performed bisulfate sequencing; K.J. Bassett performed additional experiments; I. Sayin and D.H. Canaday analyzed human mesenteric lymph nodes; J.A. Roco and M. Cuenca provided substantial intellectual input; E. Barry, A. Lopez, H. Bolton, and B. Fazekas de St Groth provided additional reagents and samples; C. Doglioni

and M. Meyer-Hermann conducted additional experiments and analyses; M.C. Cook provided substantial intellectual input and helped write the manuscript; C.G. Vinuesa conceived, designed, and supervised the study; P.F. Cañete and C.G. Vinuesa wrote the manuscript.

Submitted: 19 March 2019

Revised: 2 May 2019

Accepted: 20 May 2019

References

Akdis, C.A., and M. Akdis. 2014. Mechanisms of immune tolerance to allergens: role of IL-10 and Tregs. *J. Clin. Invest.* 124:4678–4680. <https://doi.org/10.1172/JCI78891>

Akdis, C.A., and K. Blaser. 2001. Mechanisms of interleukin-10-mediated immune suppression. *Immunology*. 103:131–136. <https://doi.org/10.1046/j.1365-2567.2001.01235.x>

Aloulou, M., E.J. Carr, M. Gador, A. Bignon, R.S. Liblau, N. Fazilleau, and M.A. Linterman. 2016. Follicular regulatory T cells can be specific for the immunizing antigen and derive from naïve T cells. *Nat. Commun.* 7: 10579. <https://doi.org/10.1038/ncomms10579>

Altin, J.A., C.C. Goodnow, and M.C. Cook. 2012. IL-10+ CTLA-4+ Th2 inhibitory cells form in a Foxp3-independent, IL-2-dependent manner from Th2 effectors during chronic inflammation. *J. Immunol.* 188:5478–5488. <https://doi.org/10.4049/jimmunol.1102994>

Arpin, C., J. Déchanet, C. Van Kooten, P. Merville, G. Grouard, F. Brière, J. Banchereau, and Y.J. Liu. 1995. Generation of memory B cells and plasma cells in vitro. *Science*. 268:720–722. <https://doi.org/10.1126/science.7537388>

Ballesteros-Tato, A., T.D. Randall, F.E. Lund, R. Spolski, W.J. Leonard, and B. León. 2016. T Follicular Helper Cell Plasticity Shapes Pathogenic T Helper 2 Cell-Mediated Immunity to Inhaled House Dust Mite. *Immunity*. 44:259–273. <https://doi.org/10.1016/j.jimmuni.2015.11.017>

Berglund, L.J., D.T. Avery, C.S. Ma, L. Moens, E.K. Deenick, J. Bustamante, S. Boisson-Dupuis, M. Wong, S. Adelstein, P.D. Arkwright, et al. 2013. IL-21 signalling via STAT3 primes human naïve B cells to respond to IL-2 to enhance their differentiation into plasmablasts. *Blood*. 122:3940–3950. <https://doi.org/10.1182/blood-2013-06-506865>

Botta, D., M.J. Fuller, T.T. Marquez-Lago, H. Bachus, J.E. Bradley, A.S. Weinmann, A.J. Zajac, T.D. Randall, F.E. Lund, B. León, and A. Ballesteros-Tato. 2017. Dynamic regulation of T follicular regulatory cell responses by interleukin 2 during influenza infection. *Nat. Immunol.* 18: 1249–1260. <https://doi.org/10.1038/ni.3837>

Butt, D., T.D. Chan, K. Bourne, J.R. Hermes, A. Nguyen, A. Statham, L.A. O'Reilly, A. Strasser, S. Price, P. Schofield, et al. 2015. FAS Inactivation Releases Unconventional Germinal Center B Cells that Escape Antigen Control and Drive IgE and Autoantibody Production. *Immunity*. 42: 890–902. <https://doi.org/10.1016/j.jimmuni.2015.04.010>

Carreras, J., A. Lopez-Guillermo, B.C. Fox, L. Colomo, A. Martinez, G. Roncador, E. Montserrat, E. Campo, and A.H. Banham. 2006. High numbers of tumor-infiltrating FOXP3-positive regulatory T cells are associated with improved overall survival in follicular lymphoma. *Blood*. 108: 2957–2964. <https://doi.org/10.1182/blood-2006-04-018218>

Choi, Y.S., J.A. Gullicksrud, S. Xing, Z. Zeng, Q. Shan, F. Li, P.E. Love, W. Peng, H.H. Xue, and S. Crotty. 2015. LEP-1 and TCF-1 orchestrate T(FH) differentiation by regulating differentiation circuits upstream of the transcriptional repressor Bcl6. *Nat. Immunol.* 16:980–990. <https://doi.org/10.1038/ni.3226>

Chowdhury, A., P.M. Del Rio Estrada, G.K. Tharp, R.P. Trible, R.R. Amara, A. Chahroudi, G. Reyes-Teran, S.E. Bosingher, and G. Silvestri. 2015. Decreased T Follicular Regulatory Cell/T Follicular Helper Cell (TFH) in Simian Immunodeficiency Virus-Infected Rhesus Macaques May Contribute to Accumulation of TFH in Chronic Infection. *J. Immunol.* 195: 3237–3247. <https://doi.org/10.4049/jimmunol.1402701>

Chung, Y., S. Tanaka, F. Chu, R.I. Nurieva, G.J. Martinez, S. Rawal, Y.H. Wang, H. Lim, J.M. Reynolds, X.H. Zhou, et al. 2011. Follicular regulatory T cells expressing Foxp3 and Bcl-6 suppress germinal center reactions. *Nat. Med.* 17:983–988. <https://doi.org/10.1038/nm.2426>

Coquet, J.M., M.J. Schuijs, M.J. Smyth, K. Deswarte, R. Beyaert, H. Braun, L. Boon, G.B. Karlsson Hedestam, S.L. Nutt, H. Hammad, and B.N. Lambrecht. 2015. Interleukin-21-Producing CD4(+) T Cells Promote Type 2 Immunity to House Dust Mites. *Immunity*. 43:318–330. <https://doi.org/10.1016/j.jimmuni.2015.07.015>

Crotty, S. 2011. Follicular helper CD4 T cells (TFH). *Annu. Rev. Immunol.* 29: 621–663. <https://doi.org/10.1146/annurev-immunol-031210-101400>

Erazo, A., N. Kutchukhidze, M. Leung, A.P. Christ, J.F. Urban Jr., M.A. Curotto de Lafaille, and J.J. Lafaille. 2007. Unique maturation program of the IgE response in vivo. *Immunity*. 26:191–203. <https://doi.org/10.1016/j.jimmuni.2006.12.006>

Ettinger, R., G.P. Sims, A.-M. Fairhurst, R. Robbins, Y.S. da Silva, R. Spolski, W.J. Leonard, and P.E. Lipsky. 2005. IL-21 induces differentiation of human naïve and memory B cells into antibody-secreting plasma cells. *J. Immunol.* 175:7867–7879. <https://doi.org/10.4049/jimmunol.175.12.7867>

Facciotti, F., N. Gagliani, B. Häringer, J.S. Alfen, A. Penatti, S. Maglie, M. Paroni, A. Iseppon, M. Moro, M.C. Crosti, et al. 2016. IL-10-producing forkhead box protein 3-negative regulatory T cells inhibit B-cell responses and are involved in systemic lupus erythematosus. *J. Allergy Clin. Immunol.* 137:318–321.e5. <https://doi.org/10.1016/j.jaci.2015.06.044>

Fonseca, V.R., A. Agua-Doce, A.R. Maceiras, W. Pierson, F. Ribeiro, V.C. Romão, A.R. Pires, S.L. da Silva, J.E. Fonseca, A.E. Sousa, et al. 2017. Human blood T_{fr} cells are indicators of ongoing humoral activity not fully licensed with suppressive function. *Sci. Immunol.* 2:eaan1487. <https://doi.org/10.1126/sciimmunol.aan1487>

Fu, W., A. Ergun, T. Lu, J.A. Hill, S. Haxhinasto, M.S. Fassett, R. Gazit, S. Adoro, L. Glimcher, S. Chan, et al. 2012. A multiply redundant genetic switch 'locks in' the transcriptional signature of regulatory T cells. *Nat. Immunol.* 13:972–980. <https://doi.org/10.1038/ni.2420>

Gagliani, N., C.F. Magnani, S. Huber, M.E. Gianolini, M. Pala, P. Licona-Limon, B. Guo, D.R. Herbert, A. Bulfone, F. Trentini, et al. 2013. Coexpression of CD49b and LAG-3 identifies human and mouse T regulatory type 1 cells. *Nat. Med.* 19:739–746. <https://doi.org/10.1038/nm.3179>

Galli, S.J., M. Tsai, and A.M. Piliponsky. 2008. The development of allergic inflammation. *Nature*. 454:445–454. <https://doi.org/10.1038/nature07204>

Glatman Zaretsky, A., J.J. Taylor, I.L. King, F.A. Marshall, M. Mohrs, and E.J. Pearce. 2009. T follicular helper cells differentiate from Th2 cells in response to helminth antigens. *J. Exp. Med.* 206:991–999. <https://doi.org/10.1084/jem.20090303>

Gould, H.J. B.R. 1998. Immunoglobulin E—an overview. In *Encyclopedia of Immunology*, 2nd Ed. Academic Press, Cambridge, MA. 1202–1208. <https://doi.org/10.1006/rwei.1999.0312>

Hammad, H., and B.N. Lambrecht. 2015. Barrier Epithelial Cells and the Control of Type 2 Immunity. *Immunity*. 43:29–40. <https://doi.org/10.1016/j.jimmuni.2015.07.007>

He, J.-S., M. Meyer-Hermann, D. Xiangying, L.Y. Zuan, L.A. Jones, L. Ramakrishna, V.C. de Vries, J. Dolpady, H. Aina, S. Joseph, et al. 2013. The distinctive germinal center phase of IgE+ B lymphocytes limits their contribution to the classical memory response. *J. Exp. Med.* 210: 2755–2771. <https://doi.org/10.1084/jem.20131539>

Hsieh, C.S., H.M. Lee, and C.W. Lio. 2012. Selection of regulatory T cells in the thymus. *Nat. Rev. Immunol.* 12:157–167. <https://doi.org/10.1038/nri3155>

Jacob, J., R. Kassir, and G. Kelsoe. 1991. In situ studies of the primary immune response to (4-hydroxy-3-nitrophenyl)acetyl. I. The architecture and dynamics of responding cell populations. *J. Exp. Med.* 173:1165–1175. <https://doi.org/10.1084/jem.173.5.1165>

Jankovic, D., M.C. Kullberg, C.G. Feng, R.S. Goldszmid, C.M. Collazo, M. Wilson, T.A. Wynn, M. Kamanaka, R.A. Flavell, and A. Sher. 2007. Conventional T-bet(+)/Foxp3(-) Th1 cells are the major source of host-protective regulatory IL-10 during intracellular protozoan infection. *J. Exp. Med.* 204:273–283. <https://doi.org/10.1084/jem.20062175>

Jeannin, P., S. Lecoanet, Y. Delneste, J.-F. Gauchat, and J.-Y. Bonnefoy. 1998. IgE versus IgG4 production can be differentially regulated by IL-10. *J. Immunol.* 160:3555–3561.

Josefowicz, S.Z., L.F. Lu, and A.Y. Rudensky. 2012a. Regulatory T cells: mechanisms of differentiation and function. *Annu. Rev. Immunol.* 30: 531–564. <https://doi.org/10.1146/annurev.immunol.25.022106.141623>

Josefowicz, S.Z., R.E. Niec, H.Y. Kim, P. Treuting, T. Chinen, Y. Zheng, D.T. Umetsu, and A.Y. Rudensky. 2012b. Extrathymically generated regulatory T cells control mucosal TH2 inflammation. *Nature*. 482:395–399. <https://doi.org/10.1038/nature10772>

King, I.L., and M. Mohrs. 2009. IL-4-producing CD4+ T cells in reactive lymph nodes during helminth infection are T follicular helper cells. *J. Exp. Med.* 206:1001–1007. <https://doi.org/10.1084/jem.20090313>

Kobayashi, T., K. Iijima, A.L. Dent, and H. Kita. 2017. Follicular helper T cells mediate IgE antibody response to airborne allergens. *J. Allergy Clin. Immunol.* 139:300–313.e7. <https://doi.org/10.1016/j.jaci.2016.04.021>

Komohara, Y., C. Ma, H. Yano, C. Pan, H. Horlad, Y. Saito, K. Ohnishi, Y. Fujiwara, Y. Okuno, K. Nosaka, et al. 2017. Cell adhesion molecule-1 (CADM1) expressed on adult T-cell leukemia/lymphoma cells is not involved in the interaction with macrophages. *J. Clin. Exp. Hematop.* 57: 15–20. <https://doi.org/10.3960/jsh.17003>

Lee, S.K., R.J. Rigby, D. Zotos, L.M. Tsai, S. Kawamoto, J.L. Marshall, R.R. Ramiscal, T.D. Chan, D. Gatto, R. Brink, et al. 2011. B cell priming for extrafollicular antibody responses requires Bcl-6 expression by T cells. *J. Exp. Med.* 208:1377–1388. <https://doi.org/10.1084/jem.20102065>

Li, H., and C.D. Pauza. 2015. CD25(+) Bcl6(low) T follicular helper cells provide help to maturing B cells in germinal centers of human tonsil. *Eur. J. Immunol.* 45:298–308. <https://doi.org/10.1002/eji.201444911>

Li, Y., W. Li, Z. Ying, H. Tian, X. Zhu, J. Li, and M. Li. 2014. Metastatic heterogeneity of breast cancer cells is associated with expression of a heterogeneous TGF β -activating miR424-503 gene cluster. *Cancer Res.* 74:6107–6118. <https://doi.org/10.1158/0008-5472.CAN-14-0389>

Licona-Limón, P., L.K. Kim, N.W. Palm, and R.A. Flavell. 2013. TH2, allergy and group 2 innate lymphoid cells. *Nat. Immunol.* 14:536–542. <https://doi.org/10.1038/ni.2617>

Lim, H.W., P. Hillsamer, and C.H. Kim. 2004. Regulatory T cells can migrate to follicles upon T cell activation and suppress GC-Th cells and GC-Th cell-driven B cell responses. *J. Clin. Invest.* 114:1640–1649. <https://doi.org/10.1172/JCI200422325>

Linterman, M.A., R.J. Rigby, R.K. Wong, D. Yu, R. Brink, J.L. Cannons, P.L. Schwartzberg, M.C. Cook, G.D. Walters, and C.G. Vinuesa. 2009. Follicular helper T cells are required for systemic autoimmunity. *J. Exp. Med.* 206:561–576. <https://doi.org/10.1084/jem.20081886>

Linterman, M.A., W. Pierson, S.K. Lee, A. Kallies, S. Kawamoto, T.F. Rayner, M. Srivastava, D.P. Divekar, L. Beaton, J.J. Hogan, et al. 2011. Foxp3⁺ follicular regulatory T cells control the germinal center response. *Nat. Med.* 17:975–982. <https://doi.org/10.1038/nm.2425>

Maceiras, A.R., S.C.P. Almeida, E. Mariotti-Ferrandiz, W. Chaara, F. Jebbawi, A. Six, S. Hori, D. Klatzmann, J. Faro, and L. Graca. 2017. T follicular helper and T follicular regulatory cells have different TCR specificity. *Nat. Commun.* 8:15067. <https://doi.org/10.1038/ncomms15067>

Martins, T.B., M.E. Bandhauer, A.M. Bunker, W.L. Roberts, and H.R. Hill. 2014. New childhood and adult reference intervals for total IgE. *J. Allergy Clin. Immunol.* 133:589–591. <https://doi.org/10.1016/j.jaci.2013.08.037>

Ozaki, K., R. Spolski, C.G. Feng, C.F. Qi, J. Cheng, A. Sher, H.C. Morse III, C. Liu, P.L. Schwartzberg, and W.J. Leonard. 2002. A critical role for IL-21 in regulating immunoglobulin production. *Science.* 298:1630–1634. <https://doi.org/10.1126/science.1077002>

Reinhardt, R.L., H.E. Liang, and R.M. Locksley. 2009. Cytokine-secreting follicular T cells shape the antibody repertoire. *Nat. Immunol.* 10: 385–393. <https://doi.org/10.1038/ni.1715>

Sage, P.T., and A.H. Sharpe. 2016. T follicular regulatory cells. *Immunol. Rev.* 271:246–259. <https://doi.org/10.1111/imr.12411>

Sage, P.T., A.M. Paterson, S.B. Lovitch, and A.H. Sharpe. 2014. The co-inhibitory receptor CTLA-4 controls B cell responses by modulating T follicular helper, T follicular regulatory, and T regulatory cells. *Immunity.* 41:1026–1039. <https://doi.org/10.1016/j.immuni.2014.12.005>

Sage, P.T., N. Ron-Harel, V.R. Juneja, D.R. Sen, S. Maleri, W. Sungnak, V.K. Kuchroo, W.N. Haining, N. Chevrier, M. Haigis, and A.H. Sharpe. 2016. Suppression by T_F cells leads to durable and selective inhibition of B cell effector function. *Nat. Immunol.* 17:1436–1446. <https://doi.org/10.1038/ni.3578>

Seddiki, N., B. Santner-Nanan, J. Martinson, J. Zaunders, S. Sasson, A. Landay, M. Solomon, W. Selby, S.I. Alexander, R. Nanan, et al. 2006. Expression of interleukin (IL)-2 and IL-7 receptors discriminates between human regulatory and activated T cells. *J. Exp. Med.* 203: 1693–1700. <https://doi.org/10.1084/jem.20060468>

Simpson, N., P.A. Gatenby, A. Wilson, S. Malik, D.A. Fulcher, S.G. Tangye, H. Manku, T.J. Vyse, G. Roncador, G.A. Huttley, et al. 2010. Expansion of circulating T cells resembling follicular helper T cells is a fixed phenotype that identifies a subset of severe systemic lupus erythematosus. *Arthritis Rheum.* 62:234–244. <https://doi.org/10.1002/art.25032>

Szurek, E., A. Cebula, L. Wojciech, M. Pietrzak, G. Rempala, P. Kisielow, and L. Ignatowicz. 2015. Differences in Expression Level of Helios and Neuropilin-1 Do Not Distinguish Thymus-Derived from Extrathymically-Induced CD4⁺Foxp3⁺ Regulatory T Cells. *PLoS One.* 10:e0141161. <https://doi.org/10.1371/journal.pone.0141161>

Thornton, A.M., P.E. Korty, D.Q. Tran, E.A. Wohlfert, P.E. Murray, Y. Belkaid, and E.M. Shevach. 2010. Expression of Helios, an Ikarus transcription factor family member, differentiates thymic-derived from peripherally induced Foxp3⁺ T regulatory cells. *J. Immunol.* 184:3433–3441. <https://doi.org/10.4049/jimmunol.0904028>

Toellner, K.M., A. Gulbranson-Judge, D.R. Taylor, D.M. Sze, and I.C. MacLennan. 1996. Immunoglobulin switch transcript production in vivo related to the site and time of antigen-specific B cell activation. *J. Exp. Med.* 183:2303–2312. <https://doi.org/10.1084/jem.183.5.2303>

Victoria, G.D., T.A. Schwickert, D.R. Fooksman, A.O. Kamphorst, M. Meyer-Hermann, M.L. Dustin, and M.C. Nussenzweig. 2010. Germinal center dynamics revealed by multiphoton microscopy with a photoactivatable fluorescent reporter. *Cell.* 143:592–605. <https://doi.org/10.1016/j.cell.2010.10.032>

Vinuesa, C.G., S.G. Tangye, B. Moser, and C.R. Mackay. 2005. Follicular B helper T cells in antibody responses and autoimmunity. *Nat. Rev. Immunol.* 5:853–865. <https://doi.org/10.1038/nri1714>

Vinuesa, C.G., M.A. Linterman, D. Yu, and I.C. MacLennan. 2016. Follicular Helper T Cells. *Annu. Rev. Immunol.* 34:335–368. <https://doi.org/10.1146/annurev-immunol-041015-055605>

Voehringer, D., T.A. Reese, X. Huang, K. Shinkai, and R.M. Locksley. 2006. Type 2 immunity is controlled by IL-4/IL-13 expression in hematopoietic non-eosinophil cells of the innate immune system. *J. Exp. Med.* 203: 1435–1446. <https://doi.org/10.1084/jem.20052448>

Wing, J.B., W. Ise, T. Kurosaki, and S. Sakaguchi. 2014. Regulatory T cells control antigen-specific expansion of Tfh cell number and humoral immune responses via the coreceptor CTLA-4. *Immunity.* 41:1013–1025. <https://doi.org/10.1016/j.immuni.2014.12.006>

Wing, J.B., Y. Kitagawa, M. Locci, H. Hume, C. Tay, T. Morita, Y. Kidani, K. Matsuda, T. Inoue, T. Kurosaki, et al. 2017. A distinct subpopulation of CD25⁺ T-follicular regulatory cells localizes in the germinal centers. *Proc. Natl. Acad. Sci. USA.* 114:E6400–E6409. <https://doi.org/10.1073/pnas.1705551114>

Wollenberg, I., A. Agua-Doce, A. Hernández, C. Almeida, V.G. Oliveira, J. Faro, and L. Graca. 2011. Regulation of the germinal center reaction by Foxp3⁺ follicular regulatory T cells. *J. Immunol.* 187:4553–4560. <https://doi.org/10.4049/jimmunol.1101328>

Xiong, H., J. Dolpady, M. Wabl, M.A. Curotto de Lafaille, and J.J. Lafaille. 2012. Sequential class switching is required for the generation of high affinity IgE antibodies. *J. Exp. Med.* 209:353–364. <https://doi.org/10.1084/jem.20111941>

Yamaguchi, T., A. Kishi, M. Osaki, H. Morikawa, P. Prieto-Martin, K. Wing, T. Saito, and S. Sakaguchi. 2013. Construction of self-recognizing regulatory T cells from conventional T cells by controlling CTLA-4 and IL-2 expression. *Proc. Natl. Acad. Sci. USA.* 110:E2116–E2125. <https://doi.org/10.1073/pnas.1307185110>

Yang, Z., B.M. Sullivan, and C.D. Allen. 2012. Fluorescent in vivo detection reveals that IgE(+) B cells are restrained by an intrinsic cell fate predisposition. *Immunity.* 36:857–872. <https://doi.org/10.1016/j.immuni.2012.02.009>

Yang, Z., M.J. Robinson, X. Chen, G.A. Smith, J. Taunton, W. Liu, and C.D.C. Allen. 2016. Regulation of B cell fate by chronic activity of the IgE B cell receptor. *eLife.* 5:e21238. <https://doi.org/10.7554/eLife.21238>

Yue, D., A. Ciccolini, E. Avilla, and S. Waserman. 2018. Food allergy and anaphylaxis. *J. Asthma Allergy.* 11:111–120. <https://doi.org/10.2147/JAA.S162456>

Zheng, Y., S. Josefowicz, A. Chaudhry, X.P. Peng, K. Forbush, and A.Y. Rudensky. 2010. Role of conserved non-coding DNA elements in the Foxp3 gene in regulatory T-cell fate. *Nature.* 463:808–812. <https://doi.org/10.1038/nature08750>