

REVIEW

Antibody–drug conjugates in solid tumors: Efficacy, challenges, and opportunities

Nakul M. Shah¹ and Funda Meric-Bernstam²

Antibody–drug conjugates (ADCs) have become an increasingly important component of the therapeutic landscape of many solid tumors. Currently, there are eight ADCs approved for solid tumors, with hundreds being developed and in clinical trials. Initially being approved in the advanced or metastatic settings, ADCs are also being incorporated as neoadjuvant or adjuvant therapies. In this review, we discuss the important components of ADC design in the context of clinical successes and failures. We further evaluate mechanisms of intrinsic and acquired resistance and strategies to overcome these barriers. Finally, we discuss the landscape of potential combination partners to increase efficacy of ADC therapies.

Introduction

Antibody–drug conjugates (ADCs) have revolutionized the treatment landscape for patients with solid tumors, precipitating a surge in research, development, and innovation. They have been coined “biological missiles,” combining the precise targeting of a monoclonal antibody with the potent cytotoxicity of the attached payload (Yao et al., 2026; Fu et al., 2022; Veneziani et al., 2024). There are currently eight Food and Drug Administration (FDA)-approved ADCs in solid tumors (Wang et al., 2025b; Camidge et al., 2024), and numerous ADCs are currently in development and in clinical trials (Zhou et al., 2024a). ADCs were initially developed more than four decades ago, and they suffered many setbacks until the first ADC for solid tumors was approved in 2013; hundreds of ADCs have been discontinued (Colombo et al., 2024). Important innovations in target discovery, antibody engineering, linker design, and payloads led to the development of currently successful ADC programs. In this review, we highlight characteristics of approved ADCs that have led to their success, describe pitfalls and resistance mechanisms that limit their efficacy, and explore strategies to overcome these barriers. We first discuss important design considerations of ADCs and their impact on the success and failure of clinical programs. We further explore mechanisms of intrinsic and acquired resistance. Finally, in the context of these limitations, we discuss how combination strategies with ADCs and future innovations may address current limitations and continue to expand the efficacy of these promising agents.

Current ADC targets

Broadly, ADCs consist of an antibody to a tumor-enriched target that is linked to a therapeutic payload. However, each aspect of

the ADC design—the antibody (and target), conjugation strategy, linker type, and payload—must be carefully considered in developing a successful agent. Trastuzumab emtansine was the first ADC to receive FDA approval for a solid tumor in February 2013 (Verma et al., 2012), and there have been seven additional ADCs that have FDA approval for solid tumors targeting six cell surface proteins. In this section, we will briefly discuss each target and each approved ADC.

Human EGF receptor 2 (HER2)

HER2 is a member of the EGF receptor (EGFR) family of receptor tyrosine kinases. Heterodimerization with other members of the EGFR family leads to autophosphorylation of tyrosine residues and activates various pathways that lead to cell proliferation, survival, and tumorigenesis. When overexpressed in preclinical models, it leads to increases in mitogenic and cell cycle signaling (Zhou and Agazie, 2012). HER2 is expressed across a diverse range of cancer types (Uzunpirmak et al., 2023) and is the only target that has a therapy with tumor-agnostic approval (Shah and Meric-Bernstam, 2025).

Trastuzumab emtansine (T-DM1)

T-DM1 is a HER2-targeting IgG1 monoclonal antibody with a noncleavable linker that is attached to mertansine (DM1), a maytansine derivative that is a potent microtubule-targeting compound and membrane impermeable (Lambert and Chari, 2014; Erickson et al., 2006). Lysine conjugation is used in production, and the drug-to-antibody ratio (DAR) is ~3.5 (Poon et al., 2013). T-DM1 is approved for HER2-positive (HER2 3+ or HER2 2+/ISH positive) metastatic breast cancer for patients that have previously received trastuzumab and a taxane

¹Division of Cancer Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX, USA; ²Department of Investigational Cancer Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, TX, USA.

Correspondence to Funda Meric-Bernstam: fmeric@mdanderson.org.

© 2026 Shah and Meric-Bernstam. This article is distributed under the terms as described at <https://rupress.org/pages/terms102024/>.

(Verma et al., 2012). Trastuzumab deruxtecan (T-DXd) showed superiority in this setting in a subsequent trial and is now the preferred agent (Cortés et al., 2022). T-DM1 is also approved in the adjuvant setting for patients with HER2-positive breast cancer with residual disease after receiving neoadjuvant trastuzumab and taxane-based regimens (von Minckwitz et al., 2019). Recently, T-DXd has also shown superiority to T-DM1 for high-risk residual disease (Loibl et al., 2026). Notable high-grade toxicities of T-DM1 seen in trials include thrombocytopenia, liver toxicity, and cardiac events (von Minckwitz et al., 2019).

Trastuzumab deruxtecan (T-DXd)

T-DXd is a HER2-targeting IgG1 monoclonal antibody with an enzyme-cleavable linker that is attached to DXd, a potent camptothecin derivative that is a topoisomerase 1 inhibitor that is membrane permeable (Ogitani et al., 2016b). Interchain cysteine conjugation is used in production, and the DAR is ~7.7 (Ogitani et al., 2016a). T-DXd has the most approved indications of any ADC that spans across histologies and levels of HER2 expression. T-DXd first achieved accelerated approval for HER2-positive (HER2 3+ or HER2 2+/ISH positive) unresectable or metastatic breast cancer for patients previously treated with T-DM1. Subsequently, in the DESTINY-Breast03 phase 3 trial, T-DXd showed improved clinical outcomes compared with T-DM1 and achieved FDA approval in patients with unresectable/metastatic breast cancer with previous treatment of trastuzumab and taxane or those that had recurred within 6 mo of neoadjuvant treatment (Cortés et al., 2022). Recently, T-DXd plus pertuzumab have been FDA approved as first-line therapy for patients with advanced or metastatic HER2+ breast cancer (Tolaney et al., 2026). T-DXd has also had recent approvals in the perioperative setting. For patients with HER2-positive breast cancer with high-risk residual disease, defined as nonoperable disease at presentation or operable disease that is axillary-node positive, T-DXd has been approved showing superiority to T-DM1 after neoadjuvant therapy with anti-HER2 therapy and taxane-based chemotherapy (Geyer et al., 2025). Furthermore, T-DXd is also approved in the neoadjuvant setting for patients with stage II or III HER2-positive breast cancer, followed by the combination of a taxane, trastuzumab, and pertuzumab (Harbeck et al., 2026).

In hormone-positive breast cancer, T-DXd has further been approved in HER2-low (HER2 2+/ISH⁻ or HER2 1+) breast cancer that has received prior chemotherapy in metastatic setting or had recurrence within 6 mo of adjuvant chemotherapy (Modi et al., 2022). In addition, most recently, it has also been approved in the HER2 low and ultralow setting (HER2 0 with >0% and ≤10% invasive cancer cells showing incomplete membrane staining) after endocrine therapy (Bardia et al., 2024a).

Outside of breast cancer, T-DXd also received accelerated approval in HER2-positive gastric cancer (Shitara et al., 2020) and non-small cell lung cancer (NSCLC) with HER2 mutations (Goto et al., 2023). T-DXd is also the only ADC with tumor agnostic approval for patients with HER2-positive (IHC 3+) solid tumors based on data from three trials, including the DESTINY-PanTumor-02 basket trial that included multiple histologies (Raghav et al., 2024; Smit et al., 2024; Meric-Bernstam et al.,

2024). Important high-grade toxicities recognized across clinical trials include cardiotoxicity, interstitial lung disease, cytopenias, and gastrointestinal (GI) toxicities (Pathak et al., 2025).

Trophoblast cell surface antigen-2 (TROP2)

TROP2 is a cell surface glycoprotein that activates numerous pathways associated with cell proliferation, invasion, and metastasis. Consistent with its involvement in a variety of cancer-associated signaling pathways, overexpression of TROP2 has been reported and linked to poor prognosis in many tumors of epithelial origin, including pancreatic, gastric, and cervical cancers (Zhao et al., 2015; Zeng et al., 2016; Ning et al., 2013; Liu et al., 2013; Fong et al., 2008). Gene expression analyses have shown upregulation of TROP2 in cancerous tissues compared with their matching normal tissues across a broad spectrum of solid tumors (Trerotola et al., 2013). Through these findings, TROP2 has become a highly attractive therapeutic target on the cell surface.

Sacituzumab govitecan (SG)

SG is a TROP2-targeting IgG1 monoclonal antibody with a pH-sensitive linker that is attached to SN-38, the active metabolite of irinotecan that is a potent topoisomerase 1 inhibitor that is membrane permeable (Cardillo et al., 2015). Interchain cysteine conjugation is used in production, and the DAR is ~7.6 (Cardillo et al., 2015). SG first received FDA approval in patients with relapsed or refractory metastatic triple-negative breast cancer (TNBC) (Bardia et al., 2021a). Subsequently, SG was approved in patients with hormone receptor (HR)-positive and HER2-negative breast cancer that have received at least one previous line of endocrine therapy, a taxane, and a CDK4/6 inhibitor and at least two previous lines of chemotherapy (Rugo et al., 2023). Of note, SG initially received accelerated approval in metastatic urothelial cancer based on promising phase II activity (Loriot et al., 2024); however, lack of survival benefit and increased rates of treatment-related adverse events, mostly from neutropenia, resulting in death in the confirmatory trial led to withdrawal (Powles et al., 2025). ADCs are not necessarily always less toxic than systemic chemotherapy, and prophylaxis of significant toxicities needs to be carefully considered. Major adverse events with SG include neutropenia (high-risk patients should have G-CSF prophylaxis), nausea, and diarrhea (Schlam et al., 2023).

Datopotamab DXd (Dato-DXd)

Dato-DXd is a TROP2-targeting IgG1 monoclonal antibody with an enzyme-sensitive linker that is attached to the topoisomerase I inhibitor payload DXd (Okajima et al., 2021). Interchain cysteine conjugation is used in production, and the DAR is ~4 (Okajima et al., 2021). For patients with HR+/HER2⁻ inoperable/metastatic breast cancer that had progressed on endocrine therapy and one to two lines of chemotherapy, Dato-DXd received accelerated approval based on superiority compared with chemotherapy in this setting (Bardia et al., 2025). For patients with metastatic TNBC ineligible for immunotherapy, first-line Dato-DXd has also recently been approved (Dent et al., 2026). Currently, Dato-DXd is also in neoadjuvant and adjuvant therapy

trials for breast cancer (Bardia et al., 2024b; Shatsky et al., 2024). Dato-DXd also received accelerated approval for patients with EGFR-mutated NSCLC that had received previous EGFR-directed therapy and platinum-based chemotherapy (Ahn et al., 2025; Sands et al., 2025). Important toxicities that require monitoring and prophylaxis include stomatitis and ocular surface events (Heist et al., 2024).

Nectin cell adhesion molecule 4 (NECTIN4)

NECTIN4 is canonically involved in cell adhesion. In normal conditions, expression of NECTIN4 is restricted to the embryo and placenta with low-level expression in most adult tissues (Li et al., 2024; Challita-Eid et al., 2016). NECTIN4 IHC staining showed positivity in numerous tumor types that include bladder, breast, pancreatic, lung, ovarian, head and neck, and esophageal cancer (Challita-Eid et al., 2016). In transformation of epithelial cells, NECTIN4 plays an important role in allowing for anchorage-independent growth seen in early stages of progression, and its knockdown can prevent tumor growth (Pavlova et al., 2013).

Enfortumab vedotin (EV)

EV is a NECTIN4-targeting IgG1 monoclonal antibody with an enzyme-sensitive linker that is attached to vedotin, a monomethyl auristatin E (MMAE) microtubule-disrupting payload that is membrane permeable. Interchain cysteine conjugation is used in production, and the DAR is ~4 (Challita-Eid et al., 2016). EV is approved for locally advanced or metastatic urothelial cancer in combination with pembrolizumab as first-line therapy (Thomas et al., 2024). Recently, EV plus pembrolizumab also received approval as perioperative treatment for patients with muscle invasive bladder cancer that are not eligible for cisplatin (Center for Drug Evaluation and Research, 2025). Important toxicities to monitor for include neuropathy, skin toxicities, and GI toxicities (Lacouture et al., 2022).

Folate receptor 1 (FOLR1)

FOLR1 is a glycosylphosphatidylinositol-anchored protein that is involved in folate transportation and is capable of reducing folates. In normal tissues, FOLR1 is expressed on the apical surfaces of the lung, kidney, choroid plexus, and placenta (Varaganti et al., 2023). It is overexpressed in multiple cancer types that include ovarian cancer, uterine cancer, pancreatic cancer, biliary tract cancer, testicular germ cell tumors, and glioblastoma (Nawaz and Kipreos, 2022; Liu et al., 2025b). Though some studies have shown association with poor prognosis (Liu et al., 2020), these results are not consistent (Köbel et al., 2014). The exact role of FOLR1 in tumorigenesis has not been established (Nawaz and Kipreos, 2022).

Mirvetuximab soravtansine

Mirvetuximab soravtansine is a FOLR1-targeting IgG1 monoclonal antibody with an enzyme-sensitive linker that is attached to DM4, a maytansine derivative that is a potent microtubule targeting compound that is membrane permeable (Ab et al., 2015; Ponte et al., 2016). Lysine conjugation is used in production, and the DAR is ~3.5 (Tu et al., 2024). In the MIRASOL phase

3 trial, patients with high-grade serous ovarian cancer and high FOLR1 expression ($\geq 75\%$ tumor cells with $\geq 2+$ staining) were randomized to receive mirvetuximab soravtansine or chemotherapy, and mirvetuximab soravtansine showed improved progression-free survival (PFS) and overall survival (OS) and received FDA approval for this indication (Moore et al., 2023). Major toxicities include ocular surface events, GI toxicities, and neuropathy (Lang et al., 2025).

Tissue factor (TF)

TF is an important part of the coagulation cascade and is the receptor and cofactor for FVIIa. After tissue injury or trauma, The TF-FVIIa complex can trigger the coagulation cascade (Grover and Mackman, 2018). In cancer, it is associated with angiogenesis and invasion and can lead to activation of multiple proliferative pathways through its binding and cleavage activities (Ahmadi et al., 2023; Hassan et al., 2023). It is upregulated in many cancer types (de Bono et al., 2023).

Tisotumab vedotin

Tisotumab vedotin is a TF-targeting IgG1 monoclonal antibody with an enzyme-sensitive linker that is attached to vedotin, a MMAE microtubule-disrupting payload that is membrane permeable. Interchain cysteine conjugation is used in production, and the DAR is ~4 (Breij et al., 2014). In the phase 3 innovaTV-301 trial, tisotumab vedotin displayed improved PFS compared with the investigator's choice of chemotherapy for patients with recurrent or metastatic cervical cancer as second- or third-line therapy and received FDA approval. Major toxicities of tisotumab vedotin include ocular surface events, anemia, and GI toxicities (Vergote et al., 2024).

MET

The hepatocyte growth factor receptor (MET) is a tyrosine kinase that is physiologically involved in wound healing and embryonic development (Chmielowiec et al., 2007; Zhao et al., 2022; Latimer and Jessen, 2008). It activates many proliferative pathways through activation from binding its ligand or ligand-independent activation of MET through interaction with partners such as integrins and fibronectin (Huang et al., 2025; Mitra et al., 2011). MET is expressed in various normal adult tissues and is overexpressed across various solid tumor (Li et al., 2020b). MET amplification can also derive resistance to therapies targeting other receptor tyrosine kinases such as EGFR inhibitors (Chmielecki et al., 2023).

Telisotuzumab vedotin

Telisotuzumab vedotin is a MET-targeting IgG1 monoclonal antibody with an enzyme-sensitive linker that is attached to vedotin, a MMAE microtubule-disrupting payload that is membrane permeable. Interchain cysteine conjugation is used in production, and the DAR is ~3 (Wang et al., 2017b). In the phase 2 LUMINOSITY trial, patients with non-squamous, EGFR-wild type NSCLC with previous treatment and cMET overexpression ($\geq 50\%$ tumor cells with 3+ expressions) were treated with telisotuzumab vedotin and showed an objective response rate (ORR) of 35% and duration of response (DOR) of 9.0 mo. Based on these

“Build your own” Antibody-drug conjugate




Figure 1. **ADC design menu.** Illustration highlighting the major components of ADC design used in currently approved and investigational ADCs: (1) target, (2) IgG subtype, (3) conjugation method, (4) linker, and (5) payload. Copyright used with the permission of The Board of Regents of the University of Texas System through The University of Texas MD Anderson Cancer Center.

results, telisotuzumab vedotin received accelerated FDA approval for this setting. The major adverse events observed were neuropathy and pneumonitis (Camidge et al., 2024).

Design of ADCs

Each component of the ADC is a crucial decision that determines important characteristics, including its target specificity, off-target toxicity, stability, therapeutic window, potency, bystander effect, and immune system activation (Fig. 1). Furthermore, even with the same target antibody, ADCs can have large differences in efficacy and indication as seen with T-DM1 and T-DXd. There are many lessons that can be learned from both the aforementioned clinical successes but also the many setbacks. Here, we describe each major component of an ADC, the general characteristics that have defined successful agents, competing interests that need to be balanced, and upcoming innovations. It is important to emphasize, however, that ADCs are complex

therapeutic agents, and key determinants of pharmacokinetics, efficacy, and toxicity, are not fully understood.

Antibody

The first decision is to choose an ideal target to then generate an antibody against. Though most current ADC targets are found on the surface of cancer cells, therapies are emerging that target secreted proteins (Cela et al., 2025) or proteins found in the tumor-microenvironment (Storey et al., 2025; Gallant et al., 2024; Hooper et al., 2022). Ideal surface proteins are believed to have high expression, surface abundance, tumor-specificity, and turnover to promote internalization once the ADC is bound to its target. Though high cell surface density is desirable (Maecker et al., 2023), increased payload potency and bystander effect can allow for efficacy at lower thresholds (Akay et al., 2025). There is not a strict cutoff for normal tissue expression; however, on-target/off-tumor toxicity can often correlate with high expression found in alternative tissues such as HER2

expression in the heart (Pondé et al., 2016) and lungs (Liu et al., 2022) and TROP2 expression in the oral mucosa (Stepan et al., 2011). Multiple ADCs had to halt clinical development due to unexpectedly high rates of on-target/off-tumor effects such as severe skin toxicities with bivatuzumab mertansine, targeting CD44v6 (Riechelmann et al., 2008) that is expressed in the skin, and exudative gastritis with BR96-doxorubicin, targeting the Lewis Y antigen that is expressed in the GI tract (Gottlich et al., 2023). However, expression in a normal tissue does not always predict toxicity (Bosi et al., 2023), and many of the common toxicities seen reflect payload therapeutic class toxicities rather than ones specific for ADCs (i.e., neuropathy with microtubule inhibitors) (Nguyen et al., 2023). This is most apparent in the difference of toxicities between Dato-DXd and SG, two TROP2-targeting ADCs with TOPli payloads. TROP2 is expressed in the salivary glands and mucosal tissues (Stepan et al., 2011), yet Dato-DXd causes much higher rates of stomatitis than SG. This may be due to the higher potency of DXd; other investigational TROP2 ADCs with higher potency payloads than SG also have higher rates of stomatitis (Meric-Bernstam et al., 2025). Furthermore, TROP2 is also not expressed in blood cells (Wang et al., 2025a), yet SG has very high rates of cytopenias. This is likely due to the relatively more unstable linker for SG leading to release of payload in circulation (Nguyen et al., 2023; Colombo et al., 2026). Expression of ADC targets in normal tissues clearly does not completely predict toxicities and is influenced by linker, payload, and tissue-intrinsic susceptibility to the payload. Nonetheless, most targets of successful ADCs have high expression in tumors and relatively low expression in majority of normal tissues.

Targets with important roles in tumorigenesis or survival may have advantages. The most clinically successful ADC, T-DXd, targets a well-known cancer driver in HER2 that was associated with a more aggressive disease with poor prognosis until the advent of HER2-targeting antibodies (Slamon et al., 1987). Many ADC targets have been associated with tumorigenesis and promoting cancer cell survival. There can also be therapeutic effect from disrupting function and downstream signaling of the target; there has been experimental evidence with HER2 (Nagata et al., 2004), TF (Breij et al., 2014), and MET (Wang et al., 2016). Selection of cancer drivers can also mitigate against clonal selection, which leads to antigen downregulation, loss, or mutation in resistance (Li et al., 2025; Chang et al., 2023).

Next is to decide on antibody design and binding affinity. Major characteristics to account for include target specificity, immunogenicity, and serum stability. Though, logically, it seems that higher affinity would be better, too high of an affinity can lead to unintended off-target effects. In preclinical models, MET-targeting ADCs with too high an affinity led to increased toxicity in the liver and other normal tissues, but decreased affinity had less off-target toxicity while maintaining therapeutic effect on cancer cells (Datta-Mannan et al., 2024). High affinity can also lead to concentration of the antibody at the vascular-tumor interface and reduced tumor penetration (Adams et al., 2001). This is known as the binding site barrier that can limit spatial penetration of the ADC (Wei et al., 2024). Initial ADCs used murine antibodies that led to immune reaction and limited efficacy before humanized constructs were made (Chari et al., 2014). There

are also four human IgG subclasses (IgG1, IgG2, IgG3, and IgG4) that can be chosen to construct ADCs (Vidarsson et al., 2014; Yu et al., 2020). A potential determinant of immune effects of antibodies that include antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) is the binding affinity of the crystallizable fragment (Fc) of the antibody to the Fcγ receptors (FcγR) on immune effector cells. IgG3 and IgG1 have the highest complement activation and Fc-binding affinity (Yu et al., 2020; Chu et al., 2021). The majority of ADCs are constructed with IgG1 subclass antibodies. Though Fc interaction can promote ADCC, it can also lead to off-target toxicity through internalization through the Fc receptor in immune and other normal cells and also promote antibody sequestration. Antibody engineering and Fc modification can modulate the immune activation and off-target toxicities of these antibodies (Hoffmann et al., 2017; Moquist et al., 2024). Successful antibody development requires finding optimal binding affinity balancing target selectivity and tumor penetration and engineering to modulate immune effects.

Conjugation

An important parameter of ADCs is the DAR, which is the amount of payload that is attached to each antibody. Effective ADC design balances low potency of a low DAR with increased toxicity of a high DAR (Pettinato, 2021). The conjugation method used to attach the linker to the antibody is an important determination of the DAR and the distribution of the payload on the antibody molecule. The antibody-linker stability is also an important determinant of the pharmacokinetics of ADCs that is often overlooked; many currently approved ADCs will release the payload-linker complex from the antibody with a half-life of ~7 days (Colombo et al., 2026). Interestingly, the majority of these payload-linker complexes end up attached to circulating albumin; the consequences of which have not been well studied (Colombo and Rich, 2022). There are two major categories of conjugation methods: nonspecific (or stochastic) and site specific. In nonspecific conjugation, the linker attachment can be to multiple sites on the antibody, and thus the exact site used and DAR can be variable. One of the most common methods that has been used on commercial antibodies is random lysine conjugation, where the linkers are attached to solvent-accessible lysine groups on the antibody. Early ADCs, such as gemtuzumab ozogamicin, using this conjugation technique had highly variable DARs that limited its therapeutic window (Joubert et al., 2020; Fan et al., 2025). However, with more modern manufacturing conditions, it has been shown that DAR can remain stable (Wu et al., 2019). Another example of this is using the four disulfide bonds found in IgG1 antibodies, one between each light and heavy chain pair and two between the heavy chains. The eight cysteines at these locations can be targeted for conjugation, but most current methods will lead to variability in which ones are conjugated between ADC molecules (Fu et al., 2022). However, newer ADCs such as T-DXd can have DARs that are almost 8, and thus, nearly every interchain cysteine is conjugated (Ogitani et al., 2016a). In site-specific conjugation, alternative methods introduce specific modifications to the antibody to then be able to control which site is being conjugated and have more precise control over the DAR.

Potential benefits include increasing ADC stability and homogeneity. This includes introducing specific cysteines, noncanonical amino acids, and modified glycans to then react with the linker (Walsh et al., 2021). The full spectrum of conjugation technologies is reviewed elsewhere (Matsuda et al., 2025; Fan et al., 2025). Novel methodologies also allow development of ADCs with dual payloads (Zhou et al., 2024b; Wen et al., 2025). Though site-specific ADCs have shown better preclinical parameters and offer more precise tuning of DAR, clinical evidence of their advantage is yet to be fully established (Fan et al., 2025).

Linker

The linker that connects the payload to the conjugation site also has important properties that determine stability, cleavability, therapeutic index, and bystander effects from the ADC. It is important to separate linker stability from linker cleavability. Linker stability refers to premature release of payload regardless of environment. Cleavability, on the other hand, is a feature; the linker being designed to release the payload based on presence of a trigger in the tumor microenvironment or target cancer cell. Initial ADCs were limited by unstable linkers that led to premature release in the plasma and systemic toxicity (Chari et al., 2014; Ducry and Stump, 2010). However, increasing stability does not necessarily mean universal less toxicity or more efficacy. A recently published insightful review discusses the balance of linker stability; unstable linkers lead to premature payload release, but high stability can also increase certain toxicities; with MMAE ADCs, higher stability of the linker is associated with increased corneal toxicity (Colombo et al., 2026). Cantuzumab mertansine (Helft et al., 2004) and DCDS0780A (Herrera et al., 2022) are two examples of unsuccessful ADCs designed with more stable linkers that ultimately displayed high rates of corneal events. Preclinical testing of increasingly stable linkers has shown that intermediate stability having the optimal efficacy; not the most stable construct (Kellogg et al., 2011; Su and Zhang, 2021). This is confounded by the fact that when linkers are more stable, they also may be more difficult to cleave in the TME, lowering the concentration of payload in the tumor. All but one FDA-approved ADC for solid tumors have cleavable linkers. One example is linkers susceptible to cathepsins, a family of proteases that can play a role in tumor invasion through breakdown of the extracellular matrix (Yadati et al., 2020). Cathepsin B and cathepsin L are highly expressed in multiple tumor types and are believed to contribute to ADC linker cleavage (Liu et al., 2024; Sudhan and Siemann, 2015; Tsao et al., 2025). Another example is acid-labile linkers sensitive to the low pH found in the tumor microenvironment (Hosonuma and Yoshimura, 2023) and in lysosomes (Yim and Mizushima, 2020). Linkers can also be susceptible to redox, oftentimes having a disulfide bond that is susceptible to high glutathione concentrations intracellularly (Franco and Cidrowski, 2009). Non-cleavable linkers are made to be stable and can prevent premature release of the payload before the ADC is broken down internally within the lysosome of the cell. T-DM1 is an example of an ADC with a non-cleavable linker (Peddi and Hurvitz, 2013). Advantages include greater plasma stability and lower off-target activity. However, this also comes at a tradeoff with efficacy

(Ogitani et al., 2016b). When T-DM1 is broken down, the payload and linker remain attached, and this complex is not membrane permeable. Thus, there is no “bystander effect,” where the payload can permeate to nearby cells that may not have high expression of the target (Ogitani et al., 2016b). There are many other linker design considerations, including the addition of spacers to help optimize flexibility, length, and hydrophobicity (Matsuda et al., 2025; Lei et al., 2025; Su et al., 2021). Most successful ADCs have intermediately stable and cleavable linkers, and additional study of tumor-specific TME characteristics may help expand ADC efficacy to tumors with low target expression.

Payload

The final component of the ADC is the payload, and this is classically a cytotoxic agent. There are two main classes of drugs in currently approved ADCs for solid tumors: topoisomerase I inhibitors and anti-microtubule agents. The majority of topoisomerase I inhibitor systemic therapies and payloads are derivatives of camptothecin, an alkaloid found in plants. Examples include SN-38, exatecan, and DXd. Anti-microtubule payloads include DM1, DM4, MMAE, and monomethyl auristatin F. There are many important considerations when choosing and deciding on a payload. First is balancing on-target killing and off-target toxicity. A key advantage of precision delivery of ADCs is the ability to conjugate agents with high potency. Early studies with systemic exatecan showed promising activity but a limited therapeutic window due to bone marrow toxicity that prevented its development into systemic therapy (Ajani et al., 2005). However, DXd, a payload derived from exatecan, has been successfully incorporated into multiple ADCs with an improved therapeutic window (Modi et al., 2020, 2022; Bardia et al., 2024a; Ahn et al., 2025). Potency also has been balanced; pyrrolobenzodiazepine dimers are highly cytotoxic, but many ADCs with pyrrolobenzodiazepine (PBD) payloads have been discontinued due to high toxicities (Hartley, 2021). Membrane permeability of the payload can also determine the bystander effect. When DM1 is released from T-DM1, it is polar and thus does not diffuse into neighboring cells that limits this effect (Ogitani et al., 2016b). Previous treatment lines and development of cross-resistance is also an important consideration. Multiple studies in breast cancer have shown that PFS is generally lower with a second ADC with the same payload class (Huppert et al., 2025; Nezirevic et al., 2025). Given the predominance of topoisomerase I inhibitor and anti-microtubule agents in current ADC development (Colombo et al., 2024), diversification of payloads will likely be important for ADC efficacy in pretreated patients (Su et al., 2021). Many novel payloads that include DNA-damage toxins (Calabretta et al., 2022), RNA toxins (Orlik et al., 2022; Pahl et al., 2018), radioisotopes (Lawal et al., 2025), protein degraders (Nakazawa et al., 2024), and immune modulators (Sega et al., 2025; Li et al., 2023) have been developed and are being tested in early-phase trials.

Determinants of ADC efficacy

ADCs have received FDA approval across various indications in solid tumors and have led to impressive improvements in

outcomes in various cancer types. However, the efficacy is not universal, and patient tumors can have inherent resistance or acquired resistance. Here, we discuss the current understanding of ADC efficacy and resistance.

Expression and histology

Given that ADC binding and internalization is thought to be essential for its therapeutic mechanism, expression of the target would presumably be an important determinant of efficacy. However, there is not always a clear relationship between expression levels and clinical outcome. For example, in breast cancer, T-DXd has shown efficacy and is approved across a wide range of expression, from ultralow staining to IHC 3⁺-positive staining. In addition, in the phase 2 DAISY trial, responses were seen in patients with no detectable HER2 expression (29.7%) (Mosele et al., 2023). Intriguingly, recent studies indicate that extracellular proteases in the tumor microenvironment can lead to release of the DXd payload (Tsao et al., 2025). The tumor microenvironment is also acidic, and ADCs with acid-labile linkers, such as SG, could also have payload release without internalization (Tolaney et al., 2025b). On the other hand, many approvals of ADCs do not require checking for expression that includes SG, Dato-DXd, EV, and tisotumab vedotin. Studies evaluating relationship between expression of TROP2 determined by both RNA levels and IHC staining have not shown conclusive relationship to clinical outcomes with SG in TNBC (Bardia et al., 2021b) or HR⁺/HER2⁻ breast cancer (Bardia et al., 2023), though higher expression does have a positive trend with improved outcomes (Bardia et al., 2021b). The phase 2 ADC-MATCH trial is assessing if patients with solid tumor that have high expression of HER2, NECTIN4, or TROP2 will benefit from matched ADC therapies regardless of histology (NCI10397). The discrepancy between expression and efficacy suggests our current tools, relying mostly on IHC of a single sample, may not be accurately quantifying heterogeneous expression (Dernbach et al., 2025; Wang et al., 2024) across all tumor sites or that target-independent mechanisms of efficacy, such as payload sensitivity and extracellular payload release, may be playing an unappreciated role in clinical outcomes that need further study.

Mechanisms of inherent and acquired resistance

Patient tumors can harbor inherent resistance to ADCs, leading to limited upfront efficacy, or they can develop resistance that eventually can lead to progression. There are five major steps to the therapeutic activity of ADCs, and at each one there can be factors that limit efficacy: (1) antigen binding, (2) internalization, (3) lysosomal degradation, (4) payload release and transport, and (5) payload cytotoxic activity.

The first step in ADC activity is binding to the target on the surface of cancer cells. As discussed previously, expression and clinical activity do not always correlate, but in general, a trend toward improved activity with higher expression is observed. Lower expressing tumors or those with heterogeneity in expression may harbor inherent resistance to ADCs. For example, in a study of neoadjuvant therapy with T-DM1 with pertuzumab, expression heterogeneity in tumor samples was associated with lower rates of response (Filho et al., 2021). During treatment,

expression of the target can decrease or be lost. For example, a study of 34 patients treated with T-DXd with pre-posttreatment biopsies showed that 34% had HER2 loss, and 29.4% of patients had HER2 score decrease. An additional study showed HER2 loss in 52% (26/50) patients treated with T-DXd (Chen et al., 2025). Mutations on the binding surface of the target can also occur that lead to decreased binding as has been seen with HER2 (Chen et al., 2025). Antigen-specific mechanisms of resistance may be mitigated with newer ADCs that have bystander effect as well as bispecific ADCs that bind to multiple targets on the cell surface (Gu et al., 2024).

The next essential step is internalization of the ADC-antigen complex. This can be done through various endocytic pathways that include clathrin-mediated and clathrin-independent; the latter includes many alternative pathways, including caveolin-mediated endocytosis (Rennick et al., 2021). Antigens differ in the pathways used for internalization; for example, it is believed that HER2 is internalized with both clathrin-mediated and caveolin-mediated endocytosis (Hammood et al., 2021). Alterations in these pathways could lead to resistance. For example, endophilin A2 (also named SH3GL1), a protein that is important for clathrin-independent internalization (Renard et al., 2015), promotes internalization of HER2 (Baldassarre et al., 2017). Silencing its expression in cell lines leads to impaired internalization and resistance to HER2 antibody-based therapies (Baldassarre et al., 2017). Preclinical studies have also suggested that increased expression of caveolin-1 is associated with decreased sensitivity to HER2-directed therapies in gastric cancer (Pereira et al., 2022); however, in breast cancer cell lines, increased caveolin-1 expression was associated with increased sensitivity (Chung et al., 2015). In addition, clear evidence of changes in the endocytosis machinery happening in ADC-resistant patient tumors has yet to be reported. Overall, additional research is needed to establish resistance mechanisms affecting ADC-target internalization and develop strategies for enhanced internalization.

Once internalized, ADCs are typically trafficked to the acidic environment of the lysosome, where they are degraded, and the linker is cleaved to release the payload. The acidic environment of the lysosome is essential for the proper function of its protease enzymes used to cleave enzyme-cleavable linker. The low pH is also important for acid-labile linkers. In breast cancer and gastric cancer cell lines with induced T-DM1 resistance, lysosomal alkalization and impaired enzyme activity has been observed (Ríos-Luci et al., 2017; Wang et al., 2017a). In addition, inhibiting acidification of lysosomes in gastric cell lines has been shown to reduce T-DM1 activity in gastric cancer (Wang et al., 2017a). Thus, lysosomal dysfunction could be a mechanism to develop ADC resistance, though a therapeutic target to reverse this dysfunction has not been established.

A key aspect of the function of both topoisomerase-1 inhibitors and microtubule disruptors is transport from the lysosome to their site of activity in the nucleus and cytoplasm, respectively. For ADCs with payloads that are not cleavable, transport of the payload out of the lysosome is an essential step. A study found that SLC46A3 was essential for transport of the catabolites of T-DM1 into the cytoplasm, and its silencing

increased the concentration of catabolites in the lysosome (Hamblett et al., 2015). Further study in in vitro and in vivo models with acquired T-DM1 resistance showed that diminished SLC46A3 expression could also be a mechanism of acquired resistance (Kinneer et al., 2018). In addition, expression of multiple efflux transporters of the ATP-binding cassette class has been shown to correlate with outcomes and be a mechanisms of resistance to ADCs. Examples include MDR1 or P-gp (encoded by *ABCB1*), MRP1 (encoded by *ABCC1*), MRP2 (encoded by *ABCC2*), and BCRP (encoded by *ABCG2*). Studies of cell lines with acquired T-DM1 resistance revealed increased *ABCC1* expression as a common mechanism (Loganzo et al., 2015; Li et al., 2018). Recent large-scale transcriptomic efforts have also revealed trends toward higher expression of MDR1, MRP1, and BCRP in patients with acquired resistance to ADCs (Alkassis et al., 2025; Sledge et al., 2025). In addition, high expression of MDR1 was associated with shorter duration of treatment and OS with T-DXd treatment in breast cancer patients (Alkassis et al., 2025). Choosing payloads with membrane permeability, bystander effect, and poor substrates for efflux transporters can help mitigate this effect (Roth et al., 2026).

The final step is the effect of the payload, and currently, this consists of either a topoisomerase 1 inhibitor or an anti-microtubule agent, and there can be development of resistance to the cytotoxic effect of the payload. In multiple studies of ADCs, samples at progression have shown development of alterations in the *TOP1* gene that may confer resistance to the *TOP1i* payload (Shitara et al., 2024; Abelman et al., 2025; Coates et al., 2021). In a study of breast cancer patients treated with *TOP1i* ADCs and with pre- and post-progression ctDNA analysis, 12.9% of patients were found to have *TOP1* mutations and had poor outcomes with subsequent ADCs with *TOP1i* payloads (Abelman et al., 2025). Though alterations in expression of tubulin proteins and mutations have been explored as mechanisms of resistance to systemic anti-microtubule agents such as taxanes (Mozzetti et al., 2005; Uba et al., 2023; Maloney et al., 2020), their exact role in conferring resistance to ADC payloads has not been established. However, resistant cells have been shown to avoid cell death in preclinical breast cancer models treated with T-DM1 by having defective cyclin B1 that plays an important role in mitotic catastrophe (Sabbaghi et al., 2017). Multiple resistance mechanisms can also develop in tumors from the same patient (Coates et al., 2021). New strategies conjugating multiple payloads on the same ADC are being developed to help overcome resistance (Khosravifarsani et al., 2025).

Combination therapies with ADCs

Though ADCs have shown potent activity as single agents, virtually all patients with widely metastatic solid tumors will eventually progress. ADCs often have a favorable therapeutic window and toxicity profile that could make them favorable for combination therapies. Rational combinations that synergize with ADCs could improve clinical outcomes. Currently, EV plus pembrolizumab is the only approved ADC combination for solid tumors; however, there are many promising ADC combination therapies with strong preclinical and clinical data that are being

developed. We categorize them based on their role in the pathway of ADC efficacy (Fig. 2).

Regulation of surface target

Many therapeutic agents modulate the expression of ADC targets. In breast cancer, our group has shown that treatment with decitabine, a DNA methyltransferase inhibitor, can lead to up-regulation and increased expression of TROP2 (Zhao et al., 2023). SG and decitabine show in vitro synergy (Zhao et al., 2023). In addition, we also observed that decitabine increases the expression of SLFN11. SLFN11 is a DNA/RNA helicase, and its expression is often correlated with sensitivity to therapies such as *TOP1i* (Murai et al., 2019; Metzner et al., 2022). Valemetostat, an EZH1/EZH2 inhibitor, has also been shown to increase the expression of HER2 and SLFN11 and enhance the antitumor activities of T-DXd and Dato-DXd (Honma et al., 2025). There is an ongoing trial assessing these combinations (NCT06244485). There has also been evidence that statin therapy can increase HER2 density on the cancer cell surface (Rao et al., 2023).

Target internalization

Therapies can also enhance the internalization of the ADC target. This has been most consistently seen with the irreversible pan-HER2 inhibitor neratinib. Preclinical studies showed that neratinib treatment led to increased ubiquitination and internalization of HER2, and this also led to increased T-DM1 internalization and efficacy in lung cancer cell lines and PDX models (Li et al., 2020a). This was also seen in gastric cancer models (Zidel et al., 2025). The phase 2 Translational Breast Cancer Research Consortium trial tested the combination of neratinib and T-DM1 in HER2⁺ breast cancer patients with brain metastasis and showed promising intracranial activity of the combination even in setting of previous T-DM1 exposure (Freedman et al., 2024). A phase I trial testing the combination of T-DXd and neratinib in solid tumors with HER2 IHC 3⁺. HER2 amplification or HER2-activating mutations showed responses in pancreatic, gastroesophageal, and ovarian cancer during dose escalation (Davis et al., 2025).

DNA damage response (DDR) pathway inhibitor

For ADCs with *TOP1i* payloads, combination with inhibitors of the DDR pathway inhibitors can be synergistic and lead to synthetic lethality. The *TOP1* enzyme recognizes supercoiled DNA and creates a single-strand break, unwinds the DNA, and then repairs the single-strand break. *TOP1* inhibitors trap the *TOP1* after it creates the single-strand break, and this is known as a *TOP1* cleavage complex (*TOP1cc*). This leads to accumulation of *TOP1cc*'s and single-strand breaks (Pommier, 2006). Poly (ADP-ribose) polymerases (PARPs) are critical components to multiple DNA repair processes. PARPs play an important role in recognition of early DNA lesions, including the *TOP1cc* to regulate repair (Chowdhuri and Das, 2021). PARP inhibitors (PARPi) have shown synergy with *TOP1* inhibitors, including topotecan in preclinical models (Smith et al., 2005; Znojek et al., 2014). Furthermore, PARPi enhance the cytotoxicity of SN-38, the payload of SG (Tahara et al., 2014). Multiple trials have tested the combination of ADC and PARPi, which was initially limited by




Figure 2. **ADC combination therapies.** Illustration highlighting seven categories of ADC combination therapies that are currently being developed. Copyright used with the permission of The Board of Regents of the University of Texas System through The University of Texas MD Anderson Cancer Center.

toxicity. To overcome toxicity, Bardia et al. tested sequential dosing of SG and PARPi in TNBC preclinical models, and the combination showed synergy in enhancing the efficacy of SG. These data were used to rationalize a phase I trial with SG and talazoparib, where sequential dosing had better PFS and safety than concurrent administration (Bardia et al., 2024c). A subsequent phase 2 trial of SG and talazoparib in patients with TNBC showed an ORR of 30.1% (Abelman et al., 2024). Furthermore, accumulation of TOP1cc also increases replication stress. An important mediator of response to replication stress is the ataxia telangiectasia and Rad3-related protein (ATR). ATR helps to coordinate repair while also leading to cell cycle arrest. Inhibition of ATR in addition to TOP1i can lead to progression of the cell cycle in the presence of DNA damage that can lead to mitotic catastrophe and cell death (Cliby et al., 2002). A siRNA screen of nearly 7,000 genes in a breast cancer cell line for synthetic lethality with the TOP1i camptothecin identified ATR as a top candidate (Jossé et al., 2014). The combination of T-DXd and the ATR inhibitor, ceralasertib, showed promising efficacy in the phase I DASH trial with ORR of 57% in HER2 3⁺ metastatic solid tumors and ORR of 27% in HER2 1⁺/2⁺ metastatic solid tumors (Raghav et al., 2025). In a phase I trial of SG with the ATR inhibitor, berzosertib, in solid tumors, there were no dose-limiting

toxicities or clinically relevant adverse events and responses were seen in 2/12 patients (Abel et al., 2023). Combination of ATRi and Dato-DXd has also been shown to resensitize gastric cell lines with acquired resistance to Dato-DXd (Przybyla et al., 2023).

Cell cycle control

A key step in effective DNA-damage repair, especially after treatment with DNA-damaging agents, is cell cycle arrest. Wee1 is an important cell cycle checkpoint that regulates the transition from G₂ to mitosis. When there is DNA damage, Wee1 leads to cell cycle arrest that allows time for DNA repair (Thangaretnam et al., 2025). Inhibition of Wee1 has been shown to prematurely force cells into mitosis leading to abnormal mitosis and ultimately cell death (Aarts et al., 2012). Preclinical testing has shown single-agent efficacy of Wee1 inhibitors such as azenosertib (Ma et al., 2025; Huang et al., 2021) and adavosertib (Vuaroqueaux et al., 2022) across a wide range of tumor types. A key mediator of sensitivity to Wee1 inhibition has been shown to be amplification of cyclin E1 (CCNE1). CCNE1 regulates G₁ to S transition, and amplification is associated with worse prognosis and increased replication stress. Adavosertib has shown promising single agent activity in early-phase trials in uterine serous

carcinoma (Liu et al., 2025a) and *CCNE1*-amplified solid tumors (Fu et al., 2023). *CCNE1* amplification can also mediate resistance to HER2-directed therapies, and *CCNE1* and *HER2* co-amplification is common (DiPeri et al., 2023). DiPeri et al. showed that *CCNE1* amplification can also decrease sensitivity to T-DXd. Combination treatment of T-DXd plus adavosertib had synergy in both HER2 low (IHC 1⁺/2⁺) and HER2 high (IHC 3⁺) tumors in PDX model experiments (DiPeri et al., 2023). Cell line experiments have also shown synergy between both TOP2 α and ADCs with TOP2 α payloads and Wee1 inhibitors (Ma et al., 2024). There are ongoing trials testing ADC and Wee1 inhibitor combinations (Doerfler et al., 2025).

Cell death control

Inhibitors of apoptosis proteins (IAPs) play an important role in regulating apoptosis. They can prevent programmed cell death and are overexpressed in many cancer types (Ye et al., 2025). IAPs have been shown to confer drug resistance to various chemotherapeutic agents (Ye et al., 2025). FL118 is a camptothecin analog that has also been shown to inhibit expression of IAPs (Ling et al., 2015). It has shown efficacy in topotecan- and irinotecan-resistant cell lines (Ling et al., 2015). Given these findings, further research is warranted to assess the combination of IAP inhibitors and TOP2 α ADCs.

Immune cell activation

The interplay between ADCs and the immune system plays an important role in their efficacy, and many effective combinations of ADCs and checkpoint inhibitor therapies have been developed. Potent cytotoxic payloads of ADCs can lead to immunogenic cell death (Schulte et al., 2025; Iwata et al., 2018; Müller et al., 2014). Upon ADC-mediated cytotoxicity, cancer cells release damage-associated molecular patterns into the TME, which are recognized by immature dendritic cells (DCs) via toll-like receptors. This interaction, along with direct stimulation exerted by the payload, boosts the maturation of DCs that migrate in lymph nodes to activate naïve T cells (Fuentes-Antrás et al., 2023). Activated CD8 T cells can subsequently infiltrate the tumor and lead to increased cytotoxicity (Iwata et al., 2018). In addition, ADCs can activate the immune system through ADCC, ADCP, and complement-dependent cellular cytotoxicity (Chang et al., 2023). As described above, the combination of EV and pembrolizumab has been approved in metastatic urothelial cancer, and many additional ADC plus immunotherapy combinations have shown promising results. As first-line therapy for patients with metastatic TNBC, SG plus pembrolizumab displayed superior PFS (HR-0.65), DOR (16.5 vs. 9.2 mo), and a trend toward improved OS (immature) when compared with chemotherapy plus pembrolizumab in the phase 3 ASCENT-04 trial (Tolaney et al., 2025a). In the cohort of metastatic TNBC patients treated with first-line Dato-DXd in combination with durvalumab in the BEGONIA trial, the confirmed ORR was 79% and the response was irrespective of PD-L1 expression; 87% of patients had tumors with low PD-L1 expression. Median PFS was 13.8 mo (Schmid et al., 2023). There are multiple ongoing trials assessing ADC and immunotherapy combinations.

Antibody therapy combinations

Cancer cells can display multiple cell surface targets, and this can allow for ADC combination therapies with other antibody-based therapies and ADCs. For example, there is significant co-expression of TROP2 and NECTIN4 in metastatic urothelial cancer (Fan et al., 2022). The phase I double-antibody-drug trial of SG plus EV showed promising results, indicating that ADCs can be used in combination. For patients with metastatic urothelial cancer, the response rate was 70% across the dose ranges in the trial, but there was significant toxicity with 48% of patients requiring dose reductions and occurrence of one grade 5 toxicity (McGregor et al., 2024). The recently approved combination of T-DXd and pertuzumab shows combining ADC with antibody targeting a different target domain can be successful (Tolaney et al., 2026). In addition, ADCs with unique payloads may have favorable toxicity profiles to allow for more successful ADC combination therapies. Finally, bispecific ADCs that target multiple cell surface receptors are also in clinical development. They can be engineered to only have potent activity when expression of both antigens is high, and this significantly limit off-target effects (Luo et al., 2025). Overall, there are many innovations in ADC design that may lead to effective ADC combinations in the future.

Conclusions

With the pace of current research and innovation, ADCs are poised to enter the therapeutic toolkit for a wide range of tumor types. There are over 400 ADCs currently in development globally (Wang et al., 2025b), and in 2024 alone, there were over 100 first-in-human trials of ADCs (Crescioli and Reichert, 2025). Though they hold great promise, there are many limitations that need to be considered. There are many resistance mechanisms, both inherent and acquired, that can limit their efficacy. Furthermore, lack of payload diversity and cross-resistance may greatly dampen the possibility of sequential ADC therapy. ADCs also have significant toxicities that need to be considered (Cheng et al., 2025). As use of ADCs spreads, it will be important to educate clinicians on recognizing and managing these toxicities. Even with these limitations, the numerous innovations in ADC design and potential combination partners offer hope for greater efficacy and improved patient outcomes in the future.

Acknowledgments

This work was supported in part by a Yosemite-American Cancer Society Award (YACS-25-1519593-01-YACS), METAvivor Award, Nellie B. Connally Research Funds, National Center for Advancing Translational Sciences grant UM1 TR0045906, and the National Institutes of Health/National Cancer Institute award number P30 CA016672 (University of Texas MD Anderson Cancer Center).

Nakul M. Shah: conceptualization, investigation, and writing—original draft, review, and editing. Funda Meric-Bernstam: conceptualization, funding acquisition, investigation, project administration, resources, supervision, validation, and writing—review and editing.

Disclosures: N.M. Shah reported equity in Aarvik Therapeutics. F. Meric-Bernstam reported personal fees and grants from AstraZeneca Pharmaceuticals; personal fees from Becton Dickinson, Biocartis NV, Boehringer Ingelheim International GmbH, Calibr (a division of Scripps Research), CrossBridge Bio, and Cybrexa Therapeutics; nonfinancial support from the Cholangiocarcinoma Foundation; personal fees and grants from Daiichi Sankyo; personal fees, "other," and nonfinancial support from Dava Oncology; grants and personal fees from Debiopharm International; personal fees from DEM BioPharma Inc. and EcoRI Capital; personal fees and grants from EFFECTOR Therapeutics; personal fees from Elevation Oncology, Exelixis, go Therapeutics, and GT Aperion; personal fees and grants from Guardant Health Inc; nonfinancial support from European Society for Medical Oncology (ESMO); nonfinancial support and "other" from Physician Education Resource (PER); personal fees from Harbinger Health, Illumen Therapeutics, Kivu Biosciences, LOXO-Oncology, Incyte Corporation, Jazz Pharmaceuticals, LigaChem Biosciences (formally LegoChem), Menarini Group, ModeX Therapeutics, Inc., Molecular Templates, OnCusp Therapeutics, Precede Biosciences Inc, Protai Bio, Ribometrix, and SystImmune; grants from Jazz Pharmaceuticals; grants and personal fees from Zymeworks; and personal fees from Tacalyx GmbH, TEMPUS, TORL BioTherapeutics, Vir Biotechnology, Seagen (formerly Seattle Genetics), Sutro Biopharma Inc, Theratechnologies Inc, and Zentalis Pharmaceuticals outside the submitted work.

Submitted: 15 December 2025

Revised: 15 April 2026

Accepted: 25 June 2026

References

- Aarts, M., R. Sharpe, I. Garcia-Murillas, H. Gevensleben, M.S. Hurd, S.D. Shumway, C. Toniatti, A. Ashworth, and N.C. Turner. 2012. Forced mitotic entry of S-phase cells as a therapeutic strategy induced by inhibition of WEE1. *Cancer Discov.* 2:524–539. <https://doi.org/10.1158/2159-8290.CD-11-0320>
- Ab, O., K.R. Whiteman, L.M. Bartle, X. Sun, R. Singh, D. Tavares, A. LaBelle, G. Payne, R.J. Lutz, J. Pinkas, et al. 2015. IMGN853, a folate receptor- α (FR α)-Targeting antibody-drug conjugate, exhibits potent targeted antitumor activity against FR α -expressing tumors. *Mol. Cancer Ther.* 14: 1605–1613. <https://doi.org/10.1158/1535-7163.MCT-14-1095>
- Abel, M.L., N. Takahashi, C. Peer, C.E. Redon, S. Nichols, R. Vilimas, M.-J. Lee, S. Lee, M. Shelat, R. Kattappuram, et al. 2023. Targeting replication stress and chemotherapy resistance with a combination of sacituzumab govitecan and berzosertib: A phase I clinical trial. *Clin. Cancer Res.* 29: 3603–3611. <https://doi.org/10.1158/1078-0432.CCR-23-0536>
- Abelman, R.O., L. Spring, A. Niemierko, E. Abraham, M. McNeice, J. Goff, A. Valenti, S.A. Wander, S.J. Isakoff, B. Moy, et al. 2024. Sequential combination of sacituzumab govitecan and talazoparib in metastatic triple negative breast cancer (mTNBC): Results from a phase II study. *J. Clin. Oncol.* 42:1102. https://doi.org/10.1200/JCO.2024.42.16_suppl.1102
- Abelman, R.O., B. Wu, H. Barnes, A. Medford, B. Norden, A. Putur, E. Bitman, W. Thant, T. Liu, C. Weipert, et al. 2025. TOP1 mutations and cross-resistance to antibody-drug conjugates in patients with metastatic breast cancer. *Clin. Cancer Res.* 31:1966–1974. <https://doi.org/10.1158/1078-0432.CCR-24-2771>
- Adams, G.P., R. Schier, A.M. McCall, H.H. Simmons, E.M. Horak, R.K. Alpaugh, J.D. Marks, and L.M. Weiner. 2001. High affinity restricts the localization and tumor penetration of single-chain Fv antibody Molecules. *Cancer Res.* 61:4750–4755.
- Ahmadi, S.E., A. Shabannezhad, A. Kahrizi, A. Akbar, S.M. Safdari, T. Hoseinnehzad, M. Zahedi, S. Sadeghi, M.G. Mojarad, and M. Safa. 2023. Tissue factor (coagulation factor III): A potential double-edge molecule to be targeted and re-targeted toward cancer. *Biomark. Res.* 11:60. <https://doi.org/10.1186/s40364-023-00504-6>
- Ahn, M.-J., K. Tanaka, L. Paz-Ares, R. Cornelissen, N. Girard, E. Pons-Tostivint, D. Vicente Baz, S. Sugawara, M. Cobo, M. Pérol, et al. 2025. Datopotamab deruxtecan versus docetaxel for previously treated advanced or metastatic non-small cell lung cancer: The randomized, open-label phase III TROPION-Lung01 study. *J. Clin. Oncol.* 43:260–272. <https://doi.org/10.1200/JCO.24-01544>
- Ajani, J.A., C. Takimoto, C.R. Becerra, A. Silva, L. Baez, A. Cohn, P. Major, M. Kamida, K. Feit, and R. De Jager. 2005. A phase II clinical and pharmacokinetic study of intravenous exatecan mesylate (DX-8951f) in patients with untreated metastatic gastric cancer. *Invest. New Drugs* 23: 479–484. <https://doi.org/10.1007/s10637-005-2907-z>
- Akay, S., M. Emiroglu, C. Kelten Talu, and O.U. Unal. 2025. Comparison of clinicopathological characteristics for HER2-Null, HER2-Ultralow and HER2-low breast cancer: A single-center study. *Medicina (Kaunas)*. 61: 719. <https://doi.org/10.3390/medicina61040719>
- Alkassiss, S., M. Lipsyc-Sharf, C. Traverso, A.J. Dugan, B. Yilma, Y. Yuan, F. Meric-Bernstam, H.S. Rugo, J. O'Shaughnessy, M.F. Rimawi, et al. 2025. Transcriptomic biomarkers of therapeutic response to antibody-drug conjugates in metastatic breast cancer: A comprehensive multi-center study. *J. Clin. Oncol.* 43:1041. https://doi.org/10.1200/JCO.2025.43.16_suppl.1041
- Baldassarre, T., P. Truesdell, and A.W. Craig. 2017. Endophilin A2 promotes HER2 internalization and sensitivity to trastuzumab-based therapy in HER2-positive breast cancers. *Breast Cancer Res.* 19:110. <https://doi.org/10.1186/s13058-017-0900-z>
- Bardia, A., S.A. Hurvitz, S.M. Tolaney, D. Loirat, K. Punie, M. Oliveira, A. Brufsky, S.D. Sardesai, Kevin Kalinsky, A.B. Zelnak, et al. 2021a. Sacituzumab govitecan in metastatic triple-negative breast cancer. *N. Engl. J. Med.* 384:1529–1541. <https://doi.org/10.1056/NEJMoa2028485>
- Bardia, A., S.M. Tolaney, K. Punie, D. Loirat, M. Oliveira, K. Kalinsky, A. Zelnak, P. Aftimos, F. Dalenc, S. Sardesai, et al. 2021b. Biomarker analyses in the phase III ASCENT study of sacituzumab govitecan versus chemotherapy in patients with metastatic triple-negative breast cancer. *Ann. Oncol.* 32:1148–1156. <https://doi.org/10.1016/j.annonc.2021.06.002>
- Bardia, A., H.S. Rugo, J. Cortés, S.M. Tolaney, P. Schmid, M. Motwani, O.K. Yoon, J. Boice, L. Zhuo, and F. Marmé. 2023. Trop-2 mRNA expression and association with clinical outcomes with sacituzumab govitecan (SG) in patients with HR+/HER2- metastatic breast cancer (mBC): Biomarker results from the phase 3 TROPiCS-02 study. *J. Clin. Oncol.* 41: 1082. https://doi.org/10.1200/JCO.2023.41.16_suppl.1082
- Bardia, A., X. Hu, R. Dent, K. Yonemori, C.H. Barrios, J.A. O'Shaughnessy, H. Wildiers, J.-Y. Pierga, Q. Zhang, C. Saura, et al. 2024a. Trastuzumab deruxtecan after endocrine therapy in metastatic breast cancer. *N. Engl. J. Med.* 391:2110–2122. <https://doi.org/10.1056/NEJMoa2407086>
- Bardia, A., L. Pusztai, K. Albain, E.M. Ciruelos, S.-A. Im, D. Hershman, K. Kalinsky, C. Isaacs, D. Loirat, L. Testa, et al. 2024b. TROPION-Breast03: A randomized phase III global trial of datopotamab deruxtecan ± durvalumab in patients with triple-negative breast cancer and residual invasive disease at surgical resection after neoadjuvant therapy. *Ther. Adv. Med. Oncol.* 16:17588359241248336. <https://doi.org/10.1177/17588359241248336>
- Bardia, A., S. Sun, N. Thimmiah, J.T. Coates, B. Wu, R.O. Abelman, L. Spring, B. Moy, P. Ryan, M.N. Melkonyan, et al. 2024c. Antibody-drug conjugate sacituzumab govitecan enables a sequential TOP1/PARP inhibitor therapy strategy in patients with breast cancer. *Clin. Cancer Res.* 30: 2917–2924. <https://doi.org/10.1158/1078-0432.CCR-24-0428>
- Bardia, A., K. Jhaveri, S.-A. Im, S. Pernas, M. De Laurentiis, S. Wang, N. Martínez Jañez, G. Borges, D.W. Cescon, M. Hattori, et al. 2025. Datopotamab deruxtecan versus chemotherapy in previously treated inoperable/metastatic hormone receptor-positive human epidermal growth factor receptor 2-negative breast cancer: Primary results from TROPION-Breast01. *J. Clin. Oncol.* 43:285–296. <https://doi.org/10.1200/JCO.24.00920>
- Bosi, C., Á. Bartha, B. Galbardi, G. Notini, M.M. Naldini, L. Licata, G. Viale, M. Mariani, B. Pistilli, H.R. Ali, et al. 2023. Pan-cancer analysis of antibody-drug conjugate targets and putative predictors of treatment response. *Eur. J. Cancer* 195:113379. <https://doi.org/10.1016/j.ejca.2023.113379>
- Breij, E.C.W., B.E.C.G. de Goeij, S. Verploegen, D.H. Schuurhuis, A. Amir-khosravi, J. Francis, V.B. Miller, M. Houtkamp, W.K. Bleeker, D. Satijn,

- and P.W.H.I. Parren. 2014. An antibody–drug conjugate that targets tissue factor exhibits potent therapeutic activity against a broad range of solid tumors. *Cancer Res.* 74:1214–1226. <https://doi.org/10.1158/0008-5472.CAN-13-2440>
- Calabretta, E., M. Hamadani, P.L. Zinzani, P. Caimi, and C. Carlo-Stella. 2022. The antibody–drug conjugate loncastuximab tesirine for the treatment of diffuse large B-cell lymphoma. *Blood.* 140:303–308. <https://doi.org/10.1182/blood.2021014663>
- Camidge, D.R., J. Bar, H. Horinouchi, J. Goldman, F. Moiseenko, E. Filippova, I. Cicin, T. Ciuleanu, N. Daaboul, C. Liu, et al. 2024. Telisotuzumab vedotin monotherapy in patients with previously treated c-Met protein-overexpressing advanced nonsquamous EGFR-wildtype non-small cell lung cancer in the phase II LUMINOSITY trial. *J. Clin. Oncol.* 42: 3000–3011. <https://doi.org/10.1200/JCO.24.00720>
- Cardillo, T.M., S.V. Govindan, R.M. Sharkey, P. Trisal, R. Arrojo, D. Liu, E.A. Rossi, C.-H. Chang, and D.M. Goldenberg. 2015. Sacituzumab govitecan (IMMU-132), an anti-Trop-2/SN-38 antibody–drug conjugate: Characterization and efficacy in pancreatic, gastric, and other cancers. *Bioconj. Chem.* 26:919–931. <https://doi.org/10.1021/acs.bioconjchem.5b00223>
- Cela, I., E. Capone, A. Pece, G. Lovato, P. Simeone, M. Colasante, A. Lamolinara, A. Piro, M. Iezzi, P. Lanuti, et al. 2025. LGALS3BP antibody–drug conjugate enhances tumor-infiltrating lymphocytes and synergizes with immunotherapy to restrain neuroblastoma growth. *J. Transl. Med.* 23:431. <https://doi.org/10.1186/s12967-025-06434-1>
- Center for Drug Evaluation and Research. 2025. FDA approves pembrolizumab with enfortumab vedotin-efv for muscle invasive bladder cancer. *Food and Drug Administration.* <https://www.fda.gov/drugs/resources-information-approved-drugs/fda-approves-pembrolizumab-enfortumab-vedotin-efv-muscle-invasive-bladder-cancer> (accessed April 4, 2026)
- Challita-Eid, P.M., D. Satpayev, P. Yang, Z. An, K. Morrison, Y. Shostak, A. Raitano, R. Nadell, W. Liu, D.R. Lortie, et al. 2016. Enfortumab vedotin antibody–drug conjugate targeting Nectin-4 Is a highly potent therapeutic agent in multiple preclinical cancer models. *Cancer Res.* 76: 3003–3013. <https://doi.org/10.1158/0008-5472.CAN-15-1313>
- Chang, H.L., B. Schwettmann, H.L. McArthur, and I.S. Chan. 2023. Antibody–drug conjugates in breast cancer: Overcoming resistance and boosting immune response. *J. Clin. Invest.* 133:e127156. <https://doi.org/10.1172/JCI172156>
- Chari, R.V.J., M.L. Miller, and W.C. Widdison. 2014. Antibody–drug conjugates: An emerging concept in cancer therapy. *Angew. Chem. Int. Ed. Engl.* 53:3796–3827. <https://doi.org/10.1002/anie.201307628>
- Chen, W., A. Gupta, N. Mai, S. Nag, J.S. Lau, S. Singh, J.D. Chodera, B. Liu, E. de Stanchina, F. Pareja, et al. 2025. Trastuzumab Deruxtecan (T-DXd) resistance via loss of HER2 expression and binding. *Cancer Discov.* 16: 235–249. <https://doi.org/10.1158/2159-8290.CD-25-0647>
- Cheng, Y., J. Lu, C. Zhang, W. Yan, P. Zhu, Q. Qin, and L. Gong. 2025. Overview of antibody–drug conjugates nonclinical and clinical toxicities and related contributing factors. *Antib. Ther.* 8:124–144. <https://doi.org/10.1093/abt/tbaf004>
- Chmielecki, J., J.E. Gray, Y. Cheng, Y. Ohe, F. Imamura, B.C. Cho, M.-C. Lin, M. Majem, R. Shah, Y. Rukazenzov, et al. 2023. Candidate mechanisms of acquired resistance to first-line osimertinib in EGFR-mutated advanced non-small cell lung cancer. *Nat. Commun.* 14:1070. <https://doi.org/10.1038/s41467-023-35961-y>
- Chmielewicz, J., M. Borowiak, M. Morkel, T. Stradal, B. Munz, S. Werner, J. Wehland, C. Birchmeier, and W. Birchmeier. 2007. c-Met is essential for wound healing in the skin. *J. Cell Biol.* 177:151–162. <https://doi.org/10.1083/jcb.200701086>
- Chowdhuri, S.P., and B.B. Das. 2021. Top1-PARP1 association and beyond: From DNA topology to break repair. *NAR Cancer.* 3:zcab003. <https://doi.org/10.1093/narcan/zcab003>
- Chu, T.H., E.F. Patz, and M.E. Ackerman. 2021. Coming together at the hinges: Therapeutic prospects of IgG3. *mAbs.* 13:1882028. <https://doi.org/10.1080/19420862.2021.1882028>
- Chung, Y.-C., J.-F. Kuo, W.-C. Wei, K.-J. Chang, and W.-T. Chao. 2015. Caveolin-1 Dependent endocytosis enhances the chemosensitivity of HER-2 positive breast cancer cells to trastuzumab Emtrastine (T-DM1). *PLoS One.* 10:e0133072. <https://doi.org/10.1371/journal.pone.0133072>
- Cliby, W.A., K.A. Lewis, K.K. Lilly, and S.H. Kaufmann. 2002. S phase and G2 arrests induced by topoisomerase I poisons are dependent on ATR kinase function. *J. Biol. Chem.* 277:1599–1606. <https://doi.org/10.1074/jbc.M106287200>
- Coates, J.T., S. Sun, I. Leshchiner, N. Thimmiah, E.E. Martin, D. McLoughlin, B.P. Danysh, K. Slowik, R.A. Jacobs, K. Rhrissorrakrai, et al. 2021. Parallel genomic alterations of antigen and payload targets mediate polyclonal acquired clinical resistance to sacituzumab govitecan in triple-negative breast cancer. *Cancer Discov.* 11:2436–2445. <https://doi.org/10.1158/2159-8290.CD-21-0702>
- Colombo, R., and J.R. Rich. 2022. The therapeutic window of antibody drug conjugates: A dogma in need of revision. *Cancer Cell.* 40:1255–1263. <https://doi.org/10.1016/j.ccell.2022.09.016>
- Colombo, R., P. Tarantino, J.R. Rich, P.M. LoRusso, and E.G.E. de Vries. 2024. The Journey of antibody–drug conjugates: Lessons learned from 40 years of development. *Cancer Discov.* 14:2089–2108. <https://doi.org/10.1158/2159-8290.CD-24-0708>
- Colombo, R., S.D. Seredick, S.D. Barnscher, and J.R. Rich. 2026. Re-evaluating antibody–drug conjugate linker stability: Assessment, interpretation, and clinical translation. *Ann. Oncol.* 0:902–919. <https://doi.org/10.1016/jannonc.2026.04.008>
- Cortés, J., S.-B. Kim, W.-P. Chung, S.-A. Im, Y.H. Park, R. Hegg, M.H. Kim, L.-M. Tseng, V. Petry, C.-F. Chung, et al. 2022. Trastuzumab deruxtecan versus trastuzumab emtansine for breast cancer. *N. Engl. J. Med.* 386: 1143–1154. <https://doi.org/10.1056/NEJMoa2115022>
- Crescioli, S., and J.M. Reichert. 2025. Innovative antibody therapeutic development in China compared with the USA and Europe. *Nat. Rev. Drug Discov.* 25:169–170. <https://doi.org/10.1038/d41573-025-00185-w>
- Datta-Mannan, A., H. Choi, Z. Jin, L. Liu, J. Lu, D.J. Stokell, A.T. Murphy, K.W. Dunn, M.M. Martinez, and Y. Feng. 2024. Reducing target binding affinity improves the therapeutic index of anti-MET antibody–drug conjugate in tumor bearing animals. *PLoS One.* 19:e0293703. <https://doi.org/10.1371/journal.pone.0293703>
- Davis, A.A., H. Park, E.E. Dumbrava, N.V. Uboha, A. Kalyan, F. Dayyani, M. Arora, M.A. Villalona-Calero, H.S. Mahdi, R.A. Patel, et al. 2025. Abstract CT071: Phase I trial of trastuzumab deruxtecan in combination with neratinib in solid tumors with HER2 alterations (NCI 10495). *Cancer Res.* 85:CT071. <https://doi.org/10.1158/1538-7445.AM2025-CT071>
- de Bono, J.S., J.R. Harris, S.M. Burm, A. Vanderstichele, M.A. Houtkamp, S. Aarass, R. Riisnaes, I. Figueiredo, D. Nava Rodrigues, R. Christova, et al. 2023. Systematic study of tissue factor expression in solid tumors. *Cancer Rep.* 6:e1699. <https://doi.org/10.1002/cnr2.1699>
- Dent, R., Z. Shao, P. Schmid, J. Cortes, D.W. Cescon, S. Saji, K.H. Jung, T. Bachelot, S. Wang, E.M. Ramirez, et al. 2026. Datopotamab deruxtecan in patients with untreated, advanced triple-negative breast cancer (TROPION-Breast02): A randomised, open-label, international, phase III trial. *Ann. Oncol.* <https://doi.org/10.1016/j.annonc.2026.03.008>
- Dernbach, G., M.-L. Eich, M.P. Dragomir, P. Anders, N. Jurczok, C. Stief, P. Jurmeister, T. Schlomm, F. Klauschen, D. Horst, et al. 2025. Spatial expression of HER2, NECTIN4, and TROP-2 in muscle-invasive bladder cancer and metastases: Implications for pathological and clinical management. *Mod. Pathol.* 38:100753. <https://doi.org/10.1016/j.modpat.2025.100753>
- DiPeri, T.P., K.W. Evans, M.G. Raso, M. Zhao, Y.Q. Rizvi, X. Zheng, B. Wang, B.P. Kirby, K. Kong, M. Kahle, et al. 2023. Adavosertib enhances anti-tumor activity of trastuzumab deruxtecan in HER2-expressing cancers. *Clin. Cancer Res.* 29:4385–4398. <https://doi.org/10.1158/1078-0432.CCR-23-0103>
- Doerfler, A.M., E.E. Dumbrava, Y. Yuan, C. Pickett-Gies, S. Gore, S.N. Westin, J.A. Ajani, A.-M. Cimo, S. Fu, C. Lee, et al. 2025. Abstract B009: Trial in progress: Phase 1 study of trastuzumab deruxtecan (DS-8201a) in combination with azenosertib (ZN-c3) in HER2-expressing/amplified gastric/gastroesophageal junction cancer and other solid tumors with HER2 expression. *Mol. Cancer Ther.* 24:B009. <https://doi.org/10.1158/1535-7163.TARG-25-B009>
- Ducry, L., and B. Stump. 2010. Antibody–drug conjugates: Linking cytotoxic payloads to monoclonal antibodies. *Bioconj. Chem.* 21:5–13. <https://doi.org/10.1021/bc9002019>
- Erickson, H.K., P.U. Park, W.C. Widdison, Y.V. Kovtun, L.M. Garrett, K. Hoffman, R.J. Lutz, V.S. Goldmacher, and W.A. Blättler. 2006. Antibody–maytansinoid conjugates are activated in targeted cancer cells by lysosomal degradation and linker-dependent intracellular processing. *Cancer Res.* 66:4426–4433. <https://doi.org/10.1158/0008-5472.CAN-05-4489>
- Fan, Y., Q. Li, Q. Shen, Z. Liu, Z. Zhang, S. Hu, W. Yu, Z. He, Q. He, and Q. Zhang. 2022. Head-to-head comparison of the expression differences of NECTIN-4, TROP-2, and HER2 in urothelial carcinoma and its histologic variants. *Front. Oncol.* 12:858865. <https://doi.org/10.3389/fonc.2022.858865>
- Fan, Q., H. Chen, G. Wei, D. Wei, Z. Wang, L. Zhang, J. Wang, and M. Zhu. 2025. A review of conjugation technologies for antibody drug conjugates. *Antib. Ther.* 8:157–170. <https://doi.org/10.1093/abt/tbaf010>

- Filho, O.M., G. Viale, S. Stein, L. Trippa, D.A. Yardley, I.A. Mayer, V.G. Abramson, C.L. Arteaga, L.M. Spring, A.G. Waks, et al. 2021. Impact of HER2 heterogeneity on treatment response of early-stage HER2-positive breast cancer: Phase II neoadjuvant clinical trial of T-DM1 combined with pertuzumab. *Cancer Discov.* 11:2474–2487. <https://doi.org/10.1158/2159-8290.CD-20-1557>
- Fong, D., G. Spizzo, J.M. Gostner, G. Gastl, P. Moser, C. Krammel, S. Gerhard, M. Rasse, and K. Laimer. 2008. TROP2: A novel prognostic marker in squamous cell carcinoma of the oral cavity. *Mod. Pathol.* 21:186–191. <https://doi.org/10.1038/modpathol.3801001>
- Franco, R., and J.A. Cidlowski. 2009. Apoptosis and glutathione: Beyond an antioxidant. *Cell Death Differ.* 16:1303–1314. <https://doi.org/10.1038/cdd.2009.107>
- Freedman, R.A., H.M. Heiling, T. Li, D. Trapani, N. Tayob, K.L. Smith, R. Davis, A.M. Peresleite, M.K. DeMeo, C. Cotter, et al. 2024. Neratinib and ado-trastuzumab emtansine for pretreated and untreated human epidermal growth factor receptor 2 (HER2)-positive breast cancer brain metastases: Translational breast cancer research consortium trial 022. *Ann. Oncol.* 35:993–1002. <https://doi.org/10.1016/j.annonc.2024.07.245>
- Fu, Z., S. Li, S. Han, C. Shi, and Y. Zhang. 2022. Antibody drug conjugate: The “biological missile” for targeted cancer therapy. *Signal Transduct. Target. Ther.* 7:93. <https://doi.org/10.1038/s41392-022-00947-7>
- Fu, S., S. Yao, Y. Yuan, R.A. Previs, A.D. Elias, R.D. Carvajal, T.J. George, Y. Yuan, L. Yu, S.N. Westin, et al. 2023. Multicenter Phase II trial of the WEE1 inhibitor adavosertib in refractory solid tumors harboring CCNE1 amplification. *J. Clin. Oncol.* 41:1725–1734. <https://doi.org/10.1200/JCO.22.00830>
- Fuentes-Antrás, J., S. Genta, A. Vijenthira, and L.L. Siu. 2023. Antibody-drug conjugates: In search of partners of choice. *Trends Cancer.* 9:339–354. <https://doi.org/10.1016/j.trecan.2023.01.003>
- Gallant, J.P., H.M. Hintz, G.S. Gunaratne, M.T. Breneman, E.E. Recchia, J.L. West, K.L. Ott, E. Heninger, A.E. Jackson, N.Y. Luo, et al. 2024. Mechanistic characterization of cancer-associated fibroblast depletion via an antibody-drug conjugate targeting fibroblast activation protein. *Cancer Res. Commun.* 4:1481–1494. <https://doi.org/10.1158/2767-9764.CRC-24-0248>
- Geyer, C.E., Y.H. Park, Z.-M. Shao, C.-S. Huang, C.H.E. Barrios, J. Abraham, A. Prat, N. Ninkura, M. Untch, S.-A. Im, et al. 2025. LBA1 Trastuzumab Deruxtecan (T-DXd) vs Trastuzumab Emtansine (T-DM1) in Patients (Pts) with high-risk human epidermal growth factor receptor 2-positive (HER2+) primary Breast Cancer (BC) with residual invasive disease after Neoadjuvant Therapy (Tx): interim analysis of DESTINY-Breast05. *Ann. Oncol.* 36:S1556–S1557. <https://doi.org/10.1016/j.annonc.2025.09.021>
- Goto, K., Y. Goto, T. Kubo, K. Ninomiya, S.-W. Kim, D. Planchard, M.-J. Ahn, E.F. Smit, A.J. de Langen, M. Pérol, et al. 2023. Trastuzumab deruxtecan in patients with HER2-mutant metastatic non-small-cell lung cancer: Primary results from the randomized, Phase II DESTINY-Lung02 trial. *J. Clin. Oncol.* 41:4852–4863. <https://doi.org/10.1200/JCO.23.01361>
- Gottlich, H.C., R. Nabavizadeh, M. Dumbra, R.R. Pessoa, A.M. Mahmoud, I. Garg, J. Orme, B.A. Costello, J. Cheville, and F. Lucien. 2023. Emerging antibody-drug conjugate therapies and targets for metastatic renal cell carcinoma. *Kidney Cancer.* 7:161–172. <https://doi.org/10.3233/KCA-230012>
- Grover, S.P., and N. Mackman. 2018. Tissue factor. *Arterioscler. Thromb. Vasc. Biol.* 38:709–725. <https://doi.org/10.1161/ATVBAHA.117.309846>
- Gu, Y., Z. Wang, and Y. Wang. 2024. Bispecific antibody drug conjugates: Making 1+1>2. *Acta Pharm. Sin. B.* 14:1965–1986. <https://doi.org/10.1016/j.apsb.2024.01.009>
- Hamblett, K.J., A.P. Jacob, J.L. Gurgel, M.E. Tometsko, B.M. Rock, S.K. Patel, R.R. Milburn, S. Siu, S.P. Ragan, D.A. Rock, et al. 2015. SLC46A3 is required to transport catabolites of noncleavable antibody maytansine conjugates from the lysosome to the cytoplasm. *Cancer Res.* 75:5329–5340. <https://doi.org/10.1158/0008-5472.CAN-15-1610>
- Hammood, M., A.W. Craig, and J.V. Leyton. 2021. Impact of endocytosis mechanisms for the receptors targeted by the currently approved Antibody-Drug Conjugates (ADCs)—A necessity for future adc research and development. *Pharmaceuticals.* 14:674. <https://doi.org/10.3390/ph14070674>
- Harbeck, N., S. Modi, L. Pusztai, S. Ohno, J. Wu, S.-B. Kim, A. Yoshida, A. Fabi, X. Cao, R. Joseph, et al. 2026. Neoadjuvant trastuzumab deruxtecan alone or followed by paclitaxel, trastuzumab, and pertuzumab for high-risk HER2-positive early breast cancer (DESTINY-Breast11): a randomised, open-label, multicentre, phase III trial. *Ann. Oncol.* 37:166–179. <https://doi.org/10.1016/j.annonc.2025.10.019>
- Hartley, J.A. 2021. Antibody-drug conjugates (ADCs) delivering pyrrolone-benzodiazepine (PBD) dimers for cancer therapy. *Expert Opin. Biol. Ther.* 21:931–943. <https://doi.org/10.1080/14712598.2020.1776255>
- Hassan, N., J. Efling, L. Kiesel, G. Bendas, and M. Götte. 2023. The tissue factor pathway in cancer: Overview and role of heparan sulfate proteoglycans. *Cancers.* 15:1524. <https://doi.org/10.3390/cancers15051524>
- Heist, R.S., J. Sands, A. Bardia, T. Shimizu, A. Lisberg, I. Krop, N. Yamamoto, T. Kogawa, S. Al-Hashimi, S.S.M. Fung, et al. 2024. Clinical management, monitoring, and prophylaxis of adverse events of special interest associated with datopotamab deruxtecan. *Cancer Treat. Rev.* 125:102720. <https://doi.org/10.1016/j.ctrv.2024.102720>
- Helft, P.R., R.L. Schilsky, F.J. Hoke, D. Williams, H.L. Kindler, E. Sprague, M. DeWitte, H.K. Martino, J. Erickson, L. Pandite, et al. 2004. A phase I study of cantuzumab mertansine administered as a single intravenous infusion once weekly in patients with advanced solid tumors. *Clin. Cancer Res.* 10:4363–4368. <https://doi.org/10.1158/1078-0432.CCR-04-0088>
- Herrera, A.F., M.R. Patel, J.M. Burke, R. Advani, B.D. Cheson, J.P. Sharman, E. Penuel, A.G. Polson, C.D. Liao, C. Li, et al. 2022. Anti-CD79B Antibody-drug conjugate DCDS0780A in patients with B-Cell non-hodgkin lymphoma: Phase 1 dose-escalation study. *Clin. Cancer Res.* 28:1294–1301. <https://doi.org/10.1158/1078-0432.CCR-21-3261>
- Hoffmann, R.M., B.G.T. Coumbe, D.H. Josephs, S. Mele, K.M. Ilieva, A. Cheung, A.N. Tutt, J.F. Spicer, D.E. Thurston, S. Crescioli, and S.N. Karagiannis. 2017. Antibody structure and engineering considerations for the design and function of Antibody Drug Conjugates (ADCs). *Oncoimmunology.* 7:e1395127. <https://doi.org/10.1080/2162402X.2017.1395127>
- Honma, D., A. Toyota, Y. Hama, Y. Kamai, D. Okajima, Y. Hizukuri, M. Narahara, Y. Yamamoto, E. Nosaka, K. Usami, et al. 2025. Abstract 3790: Preclinical assessment of valemetostat, a dual inhibitor of EZH2 and EZH1, combined with trastuzumab deruxtecan and datopotamab deruxtecan for multiple solid tumors. *Cancer Res.* 85:3790. <https://doi.org/10.1158/1538-7445.AM2025-3790>
- Hooper, A.T., K. Marquette, C.-P.B. Chang, J. Golas, S. Jain, M.-H. Lam, M. Guffroy, M. Leal, H. Falahatpisheh, D. Mathur, et al. 2022. Anti-extra domain B splice variant of fibronectin antibody-drug conjugate eliminates tumors with enhanced efficacy when combined with checkpoint blockade. *Mol. Cancer Ther.* 21:1462–1472. <https://doi.org/10.1158/1535-7163.MCT-22-0099>
- Hosonuma, M., and K. Yoshimura. 2023. Association between pH regulation of the tumor microenvironment and immunological state. *Front. Oncol.* 13:1175563. <https://doi.org/10.3389/fonc.2023.1175563>
- Huang, P.Q., B.C. Boren, S.G. Hegde, H. Liu, A.K. Unni, S. Abraham, C.D. Hopkins, S. Paliwal, A.A. Samatar, J. Li, and K.D. Bunker. 2021. Discovery of ZN-c3, a highly potent and selective Wee1 inhibitor undergoing evaluation in clinical trials for the treatment of cancer. *J. Med. Chem.* 64:13004–13024. <https://doi.org/10.1021/acs.jmedchem.1c01121>
- Huang, X., R. Fang, Y. Pang, Z. Zhang, J. Huang, Y. Li, T. Yuan, Y. Zeng, Z. Yao, S. Vega-Rubin-de-Celis, et al. 2025. HHLA2 activates c-Met and identifies patients for targeted therapy in hepatocellular carcinoma. *J. Exp. Clin. Cancer Res.* 44:153. <https://doi.org/10.1186/s13046-025-03407-6>
- Huppert, L.A., R. Mahtani, S. Fisch, N. Dempsey, S. Premji, A. Raimonde, S. Jacob, L. Quintal, M. Melisko, J. Chien, et al. 2025. Multicenter retrospective cohort study of the sequential use of the antibody-drug conjugates (ADCs) trastuzumab deruxtecan (T-DXd) and sacituzumab govitecan (SG) in patients with HER2-low metastatic breast cancer (MBC). *Npj Breast Cancer.* 11:34. <https://doi.org/10.1038/s41523-025-00748-5>
- Iwata, T.N., C. Ishii, S. Ishida, Y. Ogitani, T. Wada, and T. Agatsuma. 2018. A HER2-targeting antibody-Drug Conjugate, Trastuzumab Deruxtecan (DS-8201a), enhances antitumor immunity in a mouse model. *Mol. Cancer Ther.* 17:1494–1503. <https://doi.org/10.1158/1535-7163.MCT-17-0749>
- Jossé, R., S.E. Martin, R. Guha, P. Ormanoglu, T.D. Pfister, P.M. Reaper, C.S. Barnes, J. Jones, P. Charlton, J.R. Pollard, et al. 2014. ATR inhibitors VE-821 and VX-970 sensitize cancer cells to topoisomerase I inhibitors by disabling DNA replication initiation and fork elongation responses. *Cancer Res.* 74:6968–6979. <https://doi.org/10.1158/0008-5472.CAN-13-3369>
- Joubert, N., A. Beck, C. Dumontet, and C. Denevault-Sabourin. 2020. Antibody-drug conjugates: The last decade. *Pharmaceuticals.* 13:245. <https://doi.org/10.3390/ph13090245>
- Kellogg, B.A., L. Garrett, Y. Kovtun, K.C. Lai, B. Leece, M. Miller, G. Payne, R. Steeves, K.R. Whiteman, W. Widdison, et al. 2011. Disulfide-linked

- antibody-maytansinoid conjugates: Optimization of in vivo activity by varying the steric hindrance at carbon atoms adjacent to the disulfide linkage. *Bioconjug. Chem.* 22:717–727. <https://doi.org/10.1021/bc100480a>
- Khosravifarsani, M., F.N. Njotu, D.A. Fon, and H. Fonge. 2025. Maximizing therapeutic potential and safety: Exploring multi/dual-payload antibody conjugates as cancer theranostics. *Adv. Drug Deliv. Rev.* 222:115608. <https://doi.org/10.1016/j.addr.2025.115608>
- Kinneer, K., J. Meekin, A.C. Tiberghien, Y.-T. Tai, S. Phipps, C.M. Kiefer, M.C. Rebelatto, N. Dimasi, A. Moriarty, K.P. Papadopoulos, et al. 2018. SLC46A3 as a potential predictive biomarker for antibody–drug conjugates bearing noncleavable linked maytansinoid and pyrrolobenzodiazepine warheads. *Clin. Cancer Res.* 24:6570–6582. <https://doi.org/10.1158/1078-0432.CCR-18-1300>
- Köbel, M., J. Madore, S.J. Ramus, B.A. Clarke, P.D.P. Pharoah, S. Deen, D.D. Bowtell, K. Odunsi, U. Menon, C. Morrison, et al. 2014. Evidence for a time-dependent association between FOLR1 expression and survival from ovarian carcinoma: Implications for clinical testing. An ovarian tumour tissue analysis consortium study. *Br. J. Cancer.* 111:2297–2307. <https://doi.org/10.1038/bjc.2014.567>
- Lacouture, M.E., A.B. Patel, J.E. Rosenberg, and P.H. O'Donnell. 2022. Management of dermatologic events associated with the Nectin-4-directed antibody–drug conjugate enfortumab vedotin. *The Oncologist.* 27:e223–e232. <https://doi.org/10.1093/oncolo/oyac001>
- Lambert, J.M., and R.V.J. Chari. 2014. Ado-trastuzumab Emtrastine (T-DM1): An antibody–drug conjugate (ADC) for HER2-positive breast cancer. *J. Med. Chem.* 57:6949–6964. <https://doi.org/10.1021/jm500766w>
- Lang, S.M., F. Vingopoulos, I. Beshar, S. Somasegar, E.V. Adams, S.A. Sasse, T. Ghezelayagh, E.C. McClung, A. Karam, O. Dorigo, et al. 2025. Early experience and assessment of real-world toxicities with mirvetuximab soravtansine in a heavily pretreated patient cohort with ovarian cancer. *Gynecol. Oncol. Rep.* 59:101738. <https://doi.org/10.1016/j.gore.2025.101738>
- Latimer, A.J., and J.R. Jessen. 2008. hgf/c-met expression and functional analysis during zebrafish embryogenesis. *Dev. Dyn.* 237:3904–3915. <https://doi.org/10.1002/dvdy.21794>
- Lawal, I.O., S. Abubakar, H. Ndlovu, A. Ismaila, and M.M. Sathekge. 2025. Radiolabeled Antibody–drug conjugates in the treatment of solid tumors. *Semin. Nucl. Med.* <https://doi.org/10.1053/j.semnuclmed.2025.08.002>
- Lei, Y., M. Zheng, P. Chen, C. Seng Ng, T. Peng Loh, and H. Liu. 2025. Linker design for the antibody drug conjugates: A comprehensive review. *ChemMedChem.* 20:e202500262. <https://doi.org/10.1002/cmdc.202500262>
- Li, G., J. Guo, B.-Q. Shen, D.B. Yadav, M.X. Sliwowski, L.M. Crocker, J.A. Lacap, and G.D.L. Phillips. 2018. Mechanisms of acquired resistance to trastuzumab emtansine in breast cancer cells. *Mol. Cancer Ther.* 17:1441–1453. <https://doi.org/10.1158/1535-7163.MCT-17-0296>
- Li, B.T., F. Michelini, S. Misale, E. Cocco, L. Baldino, Y. Cai, S. Shifman, H.-Y. Tu, M.L. Myers, C. Xu, et al. 2020a. HER2-mediated internalization of cytotoxic agents in ERBB2 amplified or mutant lung cancers. *Cancer Discov.* 10:674–687. <https://doi.org/10.1158/2159-8290.CD-20-0215>
- Li, J., K. Hu, L. Zhou, J. Huang, S. Zeng, Z. Xu, and Y. Yan. 2020b. Spectrum of Mesenchymal-epithelial transition aberrations and potential clinical implications: Insights from integrative pancancer analysis. *Front. Oncol.* 10:560615. <https://doi.org/10.3389/fonc.2020.560615>
- Li, B.T., M.D. Pegram, K.-W. Lee, M. Sharma, J. Lee, A.I. Spira, G.J. Hanna, Y.-K. Kang, D.W. Rasco, K.N. Moore, et al. 2023. A phase 1/2 study of a first-in-human immune-stimulating antibody conjugate (ISAC) BDC-1001 in patients with advanced HER2-expressing solid tumors. *J. Clin. Oncol.* 41:2538. https://doi.org/10.1200/JCO.2023.41.16_suppl.2538
- Li, K., Y. Zhou, M. Zang, X. Jin, and X. Li. 2024. Therapeutic prospects of nectin-4 in cancer: Applications and value. *Front. Oncol.* 14:1354543. <https://doi.org/10.3389/fonc.2024.1354543>
- Li, S., X. Zhao, K. Fu, S. Zhu, C. Pan, C. Yang, F. Wang, K.K.W. To, and L. Fu. 2025. Resistance to antibody–drug conjugates: A review. *Acta Pharm. Sin.* 15:737–756. <https://doi.org/10.1016/j.apsb.2024.12.036>
- Ling, X., X. Liu, K. Zhong, N. Smith, J. Prey, and F. Li. 2015. FL118, a novel camptothecin analogue, overcomes irinotecan and topotecan resistance in human tumor xenograft models. *Am. J. Transl. Res.* 7:1765–1781.
- Liu, T., Y. Liu, X. Bao, J. Tian, Y. Liu, and X. Yang. 2013. Overexpression of TROP2 predicts poor prognosis of patients with cervical cancer and promotes the proliferation and invasion of cervical cancer cells by regulating ERK signaling pathway. *PLoS One.* 8:e75864. <https://doi.org/10.1371/journal.pone.0075864>
- Liu, Y., T. Lian, and Y. Yao. 2020. A systematic review and meta-analysis of higher expression of folate receptor alpha (FOLR1) predicts poor cancer prognosis. *Biomarkers.* 25:367–374. <https://doi.org/10.1080/1354750X.2020.1771420>
- Liu, X., Y. Geng, J. Liang, A.L. Coelho, C. Yao, N. Deng, Y. Wang, K. Dai, G. Huang, T. Xie, et al. 2022. HER2 drives lung fibrosis by activating a metastatic cancer signature in invasive lung fibroblasts. *J. Exp. Med.* 219:e20220126. <https://doi.org/10.1084/jem.20220126>
- Liu, F., T. Zhou, S. Zhang, Y. Li, Y. Chen, Z. Miao, X. Wang, G. Yang, Q. Li, L. Zhang, and Y. Liu. 2024. Cathepsin B: The dawn of tumor therapy. *Eur. J. Med. Chem.* 269:116329. <https://doi.org/10.1016/j.ejmech.2024.116329>
- Liu, J.F., N. Colombo, A.M. Oza, J.-S. Frenel, B.R. Corr, M.M. Rubinstein, N.S. Nevadunsky, S. Lheureux, L. Gaba, L. González Cortijo, et al. 2025a. ADAGIO: A phase IIb, open-label, single-arm, multicenter study assessing the efficacy and safety of Adavosertib (AZD1775) as treatment for recurrent or persistent uterine serous carcinoma. *J. Clin. Oncol.* 43:2897–2907. <https://doi.org/10.1200/JCO-24-01606>
- Liu, Y., X. Chen, T. Evan, B. Esapa, A. Chenoweth, A. Cheung, and S.N. Karagiannis. 2025b. Folate receptor alpha for cancer therapy: An antibody and antibody–drug conjugate target coming of age. *MAbs.* 17:2470309. <https://doi.org/10.1080/19420862.2025.2470309>
- Loganzo, F., X. Tan, M. Sung, G. Jin, J.S. Myers, E. Melamud, F. Wang, V. Diesl, M.T. Follettie, S. Musto, et al. 2015. Tumor cells chronically treated with a trastuzumab–maytansinoid antibody–drug conjugate develop varied resistance mechanisms but respond to alternate treatments. *Mol. Cancer Ther.* 14:952–963. <https://doi.org/10.1158/1535-7163.MCT-14-0862>
- Loibl, S., Y.H. Park, Z. Shao, C.-S. Huang, C. Barrios, J. Abraham, A. Prat, N. Niikura, S.-A. Im, W. Li, et al. 2026. Trastuzumab Deruxtecan in residual HER2-positive early breast cancer. *N. Engl. J. Med.* 394:845–857. <https://doi.org/10.1056/NEJMoa2514661>
- Loriot, Y., D.P. Petrylak, A. Rezazadeh Kalebasty, A. Fléchon, R.K. Jain, S. Gupta, M. Bupathi, P. Beuzebec, P. Palmbo, A.V. Balar, et al. 2024. TROPY-U-01, a phase II open-label study of sacituzumab govitecan in patients with metastatic urothelial carcinoma progressing after platinum-based chemotherapy and checkpoint inhibitors: Updated safety and efficacy outcomes. *Ann. Oncol.* 35:392–401. <https://doi.org/10.1016/j.annonc.2024.01.002>
- Luo, M., X. Wang, G. Yu, J. Ji, L. Li, and F. Song. 2025. Development of a bispecific antibody–drug conjugate targeting EpCAM and CLDN3 for the treatment of multiple solid tumors. *Exp. Hematol. Oncol.* 14:33. <https://doi.org/10.1186/s40164-025-00624-9>
- Ma, J., X. Guo, C. Lee, F. Meric-Bernstam, O. Harismendy, A. Jubb, D. Kim, and M.R. Lackner. 2024. 35P the selective WEE1 inhibitor azenosertib shows synergistic anti-tumor effects in combination with topoisomerase I inhibitor-based antibody drug conjugates. *Ann. Oncol.* 35:S225–S226. <https://doi.org/10.1016/j.annonc.2024.08.044>
- Ma, J., W. Liu, J. Li, D. Kim, S. Kim, A. Levy, Z. Cai, K.D. Bunker, A. Recio-Boiles, J.M. Segar, et al. 2025. Azenosertib is a potent and selective WEE1 kinase inhibitor with broad antitumor activity across a range of solid tumors. *Mol. Cancer Ther.* 24:1171–1185. <https://doi.org/10.1158/1535-7163.MCT-24-1194>
- Maecker, H., V. Jonnalagadda, S. Bhakta, V. Jammalamadaka, and J.R. Junutula. 2023. Exploration of the antibody–drug conjugate clinical landscape. *MAbs.* 15:2229101. <https://doi.org/10.1080/19420862.2023.2229101>
- Maloney, S.M., S.M. Spring, L.V. Morejon-Lasso, and J.R. Prospero. 2020. Mechanisms of taxane resistance. *Cancers (Basel).* 12:3323. <https://doi.org/10.3390/cancers12113323>
- Matsuda, Y., J.R. Chang, and B.A. Mendelsohn. 2025. Advanced antibody–drug conjugates design: Innovation in linker chemistry and site-specific conjugation technologies. *ChemBiochem.* 26:2500305. <https://doi.org/10.1002/cbic.202500305>
- McGregor, B.A., G.P. Sonpavde, L. Kwak, M.M. Regan, X. Gao, H. Hvidsten, C.M. Mantia, X.X. Wei, J.E. Berchuck, S.A. Berg, et al. 2024. The double antibody drug conjugate (DAD) phase I trial: Sacituzumab govitecan plus enfortumab vedotin for metastatic urothelial carcinoma*. *Ann. Oncol.* 35:91–97. <https://doi.org/10.1016/j.annonc.2023.09.3114>
- Meric-Bernstam, F., V. Makker, A. Oaknin, D.-Y. Oh, S. Banerjee, A. González-Martín, K.H. Jung, I. Ługowska, L. Manso, A. Manzano, et al. 2024. Efficacy and safety of trastuzumab deruxtecan in patients with HER2-expressing solid tumors: Primary results from the DESTINY-PanTumor02 Phase II trial. *J. Clin. Oncol.* 42:47–58. <https://doi.org/10.1200/JCO.23.02005>
- Meric-Bernstam, F., A. Bardia, P. Bossi, G. Bianchini, F. Gatlin, R.V. Lalla, B. Melosky, N. Niikura, T.A. Yap, S.S. Kim, et al. 2025. Prophylaxis, clinical management, and monitoring of datopotamab deruxtecan-associated oral mucositis/stomatitis. *Oncol.* 30:oyaf031. <https://doi.org/10.1093/oncolo/oyaf031>

- Metzner, F.J., S.J. Wenzl, M. Kugler, S. Krebs, K.-P. Hopfner, and K. Lammens. 2022. Mechanistic understanding of human SLFN11. *Nat. Commun.* 13: 5464. <https://doi.org/10.1038/s41467-022-33123-0>
- Mitra, A.K., K. Sawada, P. Tiwari, K. Mui, K. Gwin, and E. Lengyel. 2011. Ligand independent activation of c-Met by fibronectin and $\alpha 5 \beta 1$ -integrin regulates ovarian cancer invasion and metastasis. *Oncogene*. 30:1566–1576. <https://doi.org/10.1038/onc.2010.532>
- Modi, S., C. Saura, T. Yamashita, Y.H. Park, S.-B. Kim, K. Tamura, F. Andre, H. Iwata, Y. Ito, J. Tsurutani, et al. 2020. Trastuzumab deruxtecan in previously treated HER2-positive breast cancer. *N. Engl. J. Med.* 382: 610–621. <https://doi.org/10.1056/NEJMoa1914510>
- Modi, S., W. Jacot, T. Yamashita, J. Sohn, M. Vidal, E. Tokunaga, J. Tsurutani, N.T. Ueno, A. Prat, Y.S. Chae, et al. 2022. Trastuzumab deruxtecan in previously treated HER2-low advanced breast cancer. *N. Engl. J. Med.* 387:9–20. <https://doi.org/10.1056/NEJMoa2203690>
- Moore, K.N., A. Angelergues, G.E. Konecny, Y. García, S. Banerjee, D. Lorusso, J.-Y. Lee, J.W. Moroney, N. Colombo, A. Roszak, et al. 2023. Mirvetuximab soravtansine in FR α -positive, platinum-resistant ovarian cancer. *N. Engl. J. Med.* 389:2162–2174. <https://doi.org/10.1056/NEJMoa2309169>
- Moquist, P.N., X. Zhang, C.I. Leiske, N.M.-L. Eng-Duncan, W. Zeng, N.A. Bindman, S.W. Wo, A. Wong, C.M. Henderson, K. Crowder, et al. 2024. Reversible chemical modification of antibody effector function mitigates unwanted systemic immune activation. *Bioconjug. Chem.* 35: 855–866. <https://doi.org/10.1021/acs.bioconjchem.4c00212>
- Mosele, F., E. Deluche, A. Lusque, L. Le Bescond, T. Filleron, Y. Pradat, A. Ducoulombier, B. Pistilli, T. Bachelot, F. Viret, et al. 2023. Trastuzumab deruxtecan in metastatic breast cancer with variable HER2 expression: The phase 2 DAISY trial. *Nat. Med.* 29:2110–2120. <https://doi.org/10.1038/s41591-023-02478-2>
- Mozzetti, S., C. Ferlini, P. Concolino, F. Filippetti, G. Raspaglio, S. Prislei, D. Gallo, E. Martinelli, F.O. Ranelletti, G. Ferrandina, and G. Scambia. 2005. Class III β -tubulin overexpression is a prominent mechanism of paclitaxel resistance in ovarian cancer patients. *Clin. Cancer Res.* 11: 298–305. <https://doi.org/10.1158/1078-0432.CCR11.1>
- Müller, P., K. Martin, S. Theurich, J. Schreiner, S. Savic, G. Terszowski, D. Lardinois, V.A. Heinzelmann-Schwarz, M. Schlaak, H.-M. Kvasnicka, et al. 2014. Microtubule-depolymerizing agents used in antibody–drug conjugates induce antitumor immunity by stimulation of dendritic cells. *Cancer Immunol. Res.* 2:741–755. <https://doi.org/10.1158/2326-6066.CIR-13-0198>
- Murai, J., A. Thomas, M. Miettinen, and Y. Pommier. 2019. Schlafen 11 (SLFN11), a restriction factor for replicative stress induced by DNA-targeting anti-cancer therapies. *Pharmacol. Ther.* 201:94–102. <https://doi.org/10.1016/j.pharmthera.2019.05.009>
- Nagata, Y., K.-H. Lan, X. Zhou, M. Tan, F.J. Esteva, A.A. Sahin, K.S. Klos, P. Li, B.P. Monia, N.T. Nguyen, et al. 2004. PTEN activation contributes to tumor inhibition by trastuzumab, and loss of PTEN predicts trastuzumab resistance in patients. *Cancer Cell.* 6:117–127. <https://doi.org/10.1016/j.ccr.2004.06.022>
- Nakazawa, Y., M. Miyano, S. Tsukamoto, H. Kogai, A. Yamamoto, K. Iso, S. Inoue, Y. Yamane, Y. Yabe, H. Umihara, et al. 2024. Delivery of a BET protein degrader via a CEACAM6-targeted antibody–drug conjugate inhibits tumour growth in pancreatic cancer models. *Nat. Commun.* 15: 2192. <https://doi.org/10.1038/s41467-024-46167-1>
- Nawaz, F.Z., and E.T. Kipreos. 2022. Emerging roles for folate receptor FOLR1 in signaling and cancer. *Trends Endocrinol. Metab. TEM.* 33:159–174. <https://doi.org/10.1016/j.tem.2021.12.003>
- Nezirevic, S., C. Anders, S. Dent, R. Bansal, L.Z. Yang, A. Erkanli, and H. Moore. 2025. Evaluation of efficacy and safety of sequential antibody drug conjugates (ADCs) in human epidermal growth factor 2 (HER2)-negative metastatic breast cancer. *Breast Cancer Res. Treat.* 214:329–337. <https://doi.org/10.1007/s10549-025-07818-z>
- Nguyen, T.D., B.M. Bordeau, and J.P. Balthasar. 2023. Mechanisms of ADC toxicity and strategies to increase ADC tolerability. *Cancers.* 15:713. <https://doi.org/10.3390/cancers15030713>
- Ning, S., S. Guo, J. Xie, Y. Xu, X. Lu, and Y. Chen. 2013. TROP2 Correlates with microvessel density and poor prognosis in hilar cholangiocarcinoma. *J. Gastrointest. Surg.* 17:360–368. <https://doi.org/10.1007/s11605-012-2105-1>
- Ogitani, Y., T. Aida, K. Hagihara, J. Yamaguchi, C. Ishii, N. Harada, M. Soma, H. Okamoto, M. Oitate, S. Arakawa, et al. 2016a. DS-8201a, a novel HER2-targeting ADC with a Novel DNA topoisomerase I inhibitor, demonstrates a promising antitumor efficacy with differentiation from T-DMI. *Clin. Cancer Res.* 22:5097–5108. <https://doi.org/10.1158/1078-0432.CCR-15-2822>
- Ogitani, Y., K. Hagihara, M. Oitate, H. Naito, and T. Agatsuma. 2016b. By-stander killing effect of DS-8201a, a novel anti-human epidermal growth factor receptor 2 antibody–drug conjugate, in tumors with human epidermal growth factor receptor 2 heterogeneity. *Cancer Sci.* 107: 1039–1046. <https://doi.org/10.1111/cas.12966>
- Okajima, D., S. Yasuda, T. Maejima, T. Karibe, K. Sakurai, T. Aida, T. Toki, J. Yamaguchi, M. Kitamura, R. Kamei, et al. 2021. Datopotamab deruxtecan, a Novel TROP2-directed antibody–drug conjugate, demonstrates potent antitumor activity by efficient drug delivery to tumor cells. *Mol. Cancer Ther.* 20:2329–2340. <https://doi.org/10.1158/1535-7163.MCT-21-0206>
- Orlik, C., F. Ebeling, K. Decker, I. Dranova, A. Pálfi, C. Mueller, T. Hechler, A. Pahl, and M. Kulke. 2022. Abstract 1754: Amatoxin-based antibody–drug conjugates induce immunogenic cell death and improve the anti-tumor efficacy of immune checkpoint inhibitors in humanized mouse models. *Cancer Res.* 82:1754. <https://doi.org/10.1158/1538-7445.AM2022-1754>
- Pahl, A., C. Lutz, and T. Hechler. 2018. Amanitins and their development as a payload for antibody–drug conjugates. *Drug Discov. Today Technol.* 30: 85–89. <https://doi.org/10.1016/j.ddtec.2018.08.005>
- Pathak, N., M. Di Iorio, D.M. Gimenez, Y. Berner-Wygoda, J. Savill, A. Aljuhani, A. Mittal, V. Kumar, and E. Amir. 2025. Adverse effect of trastuzumab deruxtecan in solid tumours: A systematic review and meta-analysis. *Crit. Rev. Oncol. Hematol.* 213:104787. <https://doi.org/10.1016/j.critrevonc.2025.104787>
- Pavlova, N.N., C. Pallasch, A.E.H. Elia, C.J. Braun, T.F. Westbrook, M. Hermann, and S.J. Elledge. 2013. A role for PVRL4-driven cell–cell interactions in tumorigenesis. *Elife.* 2:e00358. <https://doi.org/10.7554/eLife.00358>
- Peddi, P.F., and S.A. Hurvitz. 2013. Trastuzumab emtansine: The first targeted chemotherapy for treatment of breast cancer. *Future Oncol. Lond. Engl.* 9:319–326. <https://doi.org/10.2217/fon.13.710.2217/fon.13.7>
- Pereira, P.M.R., K. Mandleywala, S. Monette, M. Lumish, K.M. Tully, S.S. Panikar, M. Cornejo, A. Mauguen, A. Ragupathi, N.C. Keltee, et al. 2022. Caveolin-1 temporal modulation enhances antibody drug efficacy in heterogeneous gastric cancer. *Nat. Commun.* 13:2526. <https://doi.org/10.1038/s41467-022-30142-9>
- Pettinato, M.C. 2021. Introduction to antibody–drug conjugates. *Antibodies.* 10: 42. <https://doi.org/10.3390/antib10040042>
- Pommier, Y. 2006. Topoisomerase I inhibitors: Camptothecins and beyond. *Nat. Rev. Cancer* 6:789–802. <https://doi.org/10.1038/nrc1977>
- Pondé, N.F., M. Lambertini, and E. de Azambuja. 2016. Twenty years of anti-HER2 therapy-associated cardiotoxicity. *ESMO Open.* 1:e000073. <https://doi.org/10.1136/esmoopen-2016-000073>
- Ponte, J.F., O. Ab, L. Lanieri, J. Lee, J. Coccia, L.M. Bartle, M. Themeles, Y. Zhou, J. Pinkas, and R. Ruiz-Soto. 2016. Mirvetuximab Soravtansine (IMGN853), a folate receptor alpha-targeting antibody–drug conjugate, potentiates the activity of standard of care therapeutics in ovarian cancer models. *Neoplasia.* 18:775–784. <https://doi.org/10.1016/j.neo.2016.11.002>
- Poon, K.A., K. Flagella, J. Beyer, J. Tibbitts, S. Kaur, O. Saad, J.-H. Yi, S. Girish, N. Dybdal, and T. Reynolds. 2013. Preclinical safety profile of trastuzumab emtansine (T-DM1): Mechanism of action of its cytotoxic component retained with improved tolerability. *Toxicol. Appl. Pharmacol.* 273:298–313. <https://doi.org/10.1016/j.taap.2013.09.003>
- Powles, T., S. Tagawa, C. Vulsteke, M. Gross-Goupil, S.H. Park, A. Necchi, M. De Santis, I. Duran, R. Morales-Barrera, J. Guo, et al. 2025. Sacituzumab govitecan in advanced urothelial carcinoma: TROPICS-04, a phase III randomized trial. *Ann. Oncol.* 36:561–571. <https://doi.org/10.1016/j.annonc.2025.01.011>
- Przybyla, A., C. Rane, G. Patel, D. Bhavsar, S. Wen, D. Clark, Y.J. Kim, A. Lau, Y. Wallez, J. Mettetal, et al. 2023. Abstract LB227: Leveraging novel Dato-DXd resistance models to inform biomarker discovery and rational combinations to combat drug resistance. *Cancer Res.* 83:LB227. <https://doi.org/10.1158/1538-7445.AM2023-LB227>
- Raghav, K., S. Siena, A. Takashima, T. Kato, M. Van den Eynde, F. Pietrantonio, Y. Komatsu, H. Kawakami, M. Peeters, T. Andre, et al. 2024. Trastuzumab deruxtecan in patients with HER2-positive advanced colorectal cancer (DESTINY-CRC02): Primary results from a multicentre, randomised, phase 2 trial. *Lancet Oncol.* 25:1147–1162. [https://doi.org/10.1016/S1470-2045\(24\)00380-2](https://doi.org/10.1016/S1470-2045(24)00380-2)
- Raghav, K., Y. Yuan, V. Florou, M.A. Shah, B. Bashir, S. Kazmi, S. Ulahannan, F. Dayyani, H.J. Lenz, J.W. Riess, et al. 2025. 939P Phase I, dose-escalation study evaluating safety & efficacy of Trastuzumab Deruxtecan (T-DXd) in combination with ATR inhibitor (AZD6738,

- Ceralasertib) in advanced solid tumors with HER2 expression (DASH). *Ann. Oncol.* 36:S615–S615. <https://doi.org/10.1016/j.annonc.2025.08.1508>
- Rao, Y., Z. Samuels, L.M. Carter, S. Monette, S.S. Panikar, P.M.R. Pereira, and J.S. Lewis. 2023. Statins enhance the efficacy of HER2-targeting radioligand therapy in drug-resistant gastric cancers. *Proc. Natl. Acad. Sci. USA*. 120:e2220413120. <https://doi.org/10.1073/pnas.2220413120>
- Renard, H.-F., M. Simunovic, J. Lemièrre, E. Boucrot, M.D. Garcia-Castillo, S. Arumugam, V. Chambon, C. Lamaze, C. Wunder, A.K. Kenworthy, et al. 2015. Endophilin-A2 functions in membrane scission in clathrin-independent endocytosis. *Nature*. 517:493–496. <https://doi.org/10.1038/nature14064>
- Rennick, J.J., A.P.R. Johnston, and R.G. Parton. 2021. Key principles and methods for studying the endocytosis of biological and nanoparticle therapeutics. *Nat. Nanotechnol.* 16:266–276. <https://doi.org/10.1038/s41565-021-00858-8>
- Riechelmann, H., A. Sauter, W. Golze, G. Hanft, C. Schroen, K. Hoermann, T. Erhardt, and S. Gronau. 2008. Phase I trial with the CD44v6-targeting immunoconjugate bivatuzumab mertansine in head and neck squamous cell carcinoma. *Oral Oncol.* 44:823–829. <https://doi.org/10.1016/j.oraloncology.2007.10.009>
- Ríos-Luci, C., S. García-Alonso, E. Díaz-Rodríguez, M. Nadal-Serrano, J. Arribas, A. Ocaña, and A. Pandiella. 2017. Resistance to the antibody–drug conjugate T-DM1 is based in a reduction in lysosomal proteolytic activity. *Cancer Res.* 77:4639–4651. <https://doi.org/10.1158/0008-5472.CAN-16-3127>
- Roth, J.S., H. Guo, L. Chen, M. Shen, O. Gbadegesin, R.W. Robey, M.M. Gottesman, and M.D. Hall. 2026. Identification of antibody–drug conjugate payloads that are substrates of ATP-binding cassette drug efflux transporters. *Cancer Drug Resist.* 9:2. <https://doi.org/10.20517/cdr.2025.151>
- Rugo, H.S., A. Bardia, F. Marmé, J. Cortés, P. Schmid, D. Loirat, O. Trédan, E. Ciruelos, F. Dalenc, P.G. Pardo, et al. 2023. Overall survival with sacituzumab govitecan in hormone receptor-positive and human epidermal growth factor receptor 2-negative metastatic breast cancer (TROPICS-02): A randomised, open-label, multicentre, phase 3 trial. *The Lancet*. 402:1423–1433. [https://doi.org/10.1016/S0140-6736\(23\)01245-X](https://doi.org/10.1016/S0140-6736(23)01245-X)
- Sabbaghi, M., G. Gil-Gómez, C. Guardia, S. Servitja, O. Arpi, S. García-Alonso, S. Menendez, M. Arumi-Uria, L. Serrano, M. Salido, et al. 2017. Defective cyclin B1 induction in Trastuzumab-emtansine (T-DM1) acquired resistance in HER2-positive breast cancer. *Clin. Cancer Res.* 23:7006–7019. <https://doi.org/10.1158/1078-0432.CCR-17-0696>
- Sands, J., M.-J. Ahn, A. Lisberg, B.C. Cho, G. Blumenschein, E. Shum, E. Pons Tostivint, Y. Goto, K. Yoh, R. Heist, et al. 2025. Datopotamab deruxtecan in advanced or metastatic non-small cell lung cancer with actionable genomic alterations: Results from the phase II TROPION-Lung05 study. *J. Clin. Oncol.* 43:1254–1265. <https://doi.org/10.1200/JCO-24-01349>
- Schlam, I., P. Tarantino, and S.M. Tolaney. 2023. Managing adverse events of sacituzumab govitecan. *Expert Opin. Biol. Ther.* 23:1103–1111. <https://doi.org/10.1080/14712598.2023.2267975>
- Schmid, P., P.J. Wysocki, C.X. Ma, Y.H. Park, R. Fernandes, S. Lord, R.D. Baird, C. Prady, K.H. Jung, J. Asselah, et al. 2023. 379MO Datopotamab deruxtecan (Dato-DXd) + durvalumab (D) as first-line (1L) treatment for unresectable locally advanced/metastatic triple-negative breast cancer (a/mTNBC): Updated results from BEGONIA, a phase Ib/II study. *Ann. Oncol.* 34:S337. <https://doi.org/10.1016/j.annonc.2023.09.556>
- Schulte, N., M. Ikuta, A. Mertes, I. Kanchev, T. Deguchi, and G. Polier. 2025. Abstract 2160: Trastuzumab deruxtecan induces immunogenic cell death, immune cell activation and -migration in viable human breast cancer slices. *Cancer Res.* 85:2160. <https://doi.org/10.1158/1538-7445.AM2025-2160>
- Sega, E., S. Kotapati, Y.B. Poudel, Q. Cheng, K. Sadanala, B. Schneider, E.P. Chekler, C. Rao, S. Gangwar, T. Sproul, et al. 2025. Targeted delivery of TLR7 agonists to the tumor microenvironment enhances tumor immunity via activation of tumor-resident myeloid cells. *Bioconjug. Chem.* 36:437–448. <https://doi.org/10.1021/acs.bioconjugchem.4c00534>
- Shah, N.M., and F. Meric-Bernstam. 2025. The present and future of precision oncology and tumor-agnostic therapeutic approaches. *The Oncologist*. 30:oyaf152. <https://doi.org/10.1093/oncolo/oyaf152>
- Shatsky, R.A., M.S. Trivedi, C. Yau, R. Nanda, H.S. Rugo, M. Davidian, B. Tsiatis, A.M. Wallace, A.J. Chien, E. Stringer-Reasor, et al. 2024. Datopotamab–deruxtecan plus durvalumab in early-stage breast cancer: The sequential multiple assignment randomized I-SPY2.2 phase 2 trial. *Nat. Med.* 30:3737–3747. <https://doi.org/10.1038/s41591-024-03267-1>
- Shitara, K., Y.-J. Bang, S. Iwasa, N. Sugimoto, M.-H. Ryu, D. Sakai, H.-C. Chung, H. Kawakami, H. Yabusaki, J. Lee, et al. 2020. Trastuzumab deruxtecan in previously treated HER2-positive gastric cancer. *N. Engl. J. Med.* 382:2419–2430. <https://doi.org/10.1056/NEJMoa2004413>
- Shitara, K., Y.-J. Bang, S. Iwasa, N. Sugimoto, M.-H. Ryu, D. Sakai, H.C. Chung, H. Kawakami, H. Yabusaki, Y. Sakamoto, et al. 2024. Trastuzumab deruxtecan in HER2-positive advanced gastric cancer: Exploratory biomarker analysis of the randomized, phase 2 DESTINY-Gastric01 trial. *Nat. Med.* 30:1933–1942. <https://doi.org/10.1038/s41591-024-02992-x>
- Slamon, D.J., G.M. Clark, S.G. Wong, W.J. Levin, A. Ullrich, and W.L. McGuire. 1987. Human breast cancer: Correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science*. 235:177–182. <https://doi.org/10.1126/science.3798106>
- Sledge, G., J. Xiu, R.L. Mahtani, A.C. Sandoval Leon, M.J. Oberley, M. Rado-vich, and D. Spetzler. 2025. Abstract PS13-09: Mechanisms of resistance to trastuzumab deruxtecan in breast cancer elucidated by multi-omic molecular profiling. *Clin. Cancer Res.* 31:PS13–PS19. <https://doi.org/10.1158/1557-3265.SABCS24-PS13-09>
- Smit, E.F., E. Felip, D. Uprety, M. Nagasaka, K. Nakagawa, L.P.-A. Rodríguez, J.M. Pacheco, B.T. Li, D. Planchard, C. Baik, et al. 2024. Trastuzumab deruxtecan in patients with metastatic non-small-cell lung cancer (DESTINY-Lung01): Primary results of the HER2-overexpressing cohorts from a single-arm, phase 2 trial. *Lancet Oncol.* 25:439–454. [https://doi.org/10.1016/S1470-2045\(24\)00064-0](https://doi.org/10.1016/S1470-2045(24)00064-0)
- Smith, L.M., E. Willmore, C.A. Austin, and N.J. Curtin. 2005. The novel Poly-(ADP-Ribose) Polymerase inhibitor, AG14361, sensitizes cells to topoisomerase I poisons by increasing the persistence of DNA strand breaks. *Clin. Cancer Res.* 11:8449–8457. <https://doi.org/10.1158/1078-0432.CCR-05-1224>
- Stepan, L.P., E.S. Trueblood, K. Hale, J. Babcock, L. Borges, and C.L. Sutherland. 2011. Expression of Trop2 cell surface glycoprotein in normal and tumor tissues. *J. Histochem. Cytochem.* 59:701–710. <https://doi.org/10.1369/0022155411410430>
- Storey, C.M., M. Altai, K. Lückerrath, W. Zedan, H. Zhu, L. Breuer, M. Trajkovic-Arsic, J. Park, A. Hasson, J. Sivek, et al. 2025. Development of a leucine-rich repeat-containing protein 15-targeted radio-immunotheranostic approach to deplete pro-tumorigenic mechanisms and immunotherapy resistance. *Signal Transduct. Target. Ther.* 10:319. <https://doi.org/10.1038/s41392-025-02410-9>
- Su, D., and D. Zhang. 2021. Linker design impacts antibody–drug conjugate pharmacokinetics and efficacy via modulating the stability and payload release efficiency. *Front. Pharmacol.* 12:687926. <https://doi.org/10.3389/fphar.2021.687926>
- Su, Z., D. Xiao, F. Xie, L. Liu, Y. Wang, S. Fan, X. Zhou, and S. Li. 2021. Antibody–drug conjugates: Recent advances in linker chemistry. *Acta Pharm. Sin. B*. 11:3889–3907. <https://doi.org/10.1016/j.apsb.2021.03.042>
- Sudhan, D.R., and D.W. Siemann. 2015. Cathepsin L targeting in cancer treatment. *Pharmacol. Ther.* 155:105–116. <https://doi.org/10.1016/j.pharmthera.2015.08.007>
- Tahara, M., T. Inoue, F. Sato, Y. Miyakura, H. Horie, Y. Yasuda, H. Fujii, K. Kotake, and K. Sugano. 2014. The use of olaparib (AZD2281) Potentiates SN-38 cytotoxicity in colon cancer cells by indirect inhibition of Rad51-Mediated repair of DNA Double-strand breaks. *Mol. Cancer Ther.* 13:1170–1180. <https://doi.org/10.1158/1535-7163.MCT-13-0683>
- Thangaretnam, K., M.O. Islam, J. Lv, A. El-Rifai, A. Perloff, H.L. Soutto, D. Peng, and Z. Chen. 2025. WEE1 inhibition in cancer therapy: Mechanisms, synergies, preclinical insights, and clinical trials. *Crit. Rev. Oncol. Hematol.* 211:104710. <https://doi.org/10.1016/j.critrevonc.2025.104710>
- Thomas, P., B.P. Valderrama, S. Gupta, B. Jens, K. Eiji, H.-C. Jean, I. Gopa, V. Christof, S.H. Park, S.S. Joon, et al. 2024. Enfortumab vedotin and pembrolizumab in untreated advanced urothelial cancer. *N. Engl. J. Med.* 390:875–888. <https://doi.org/10.1056/NEJMoa2312117>
- Tolaney, S.M., E. de Azambuja, K. Kalinsky, S. Loi, S.-B. Kim, C. Yam, B.L. Rapoport, S.-A. Im, B. Pistilli, W. McHayleh, et al. 2025a. Sacituzumab govitecan (SG) + pembrolizumab (pembro) vs chemotherapy (chemo) + pembro in previously untreated PD-L1-positive advanced triple-negative breast cancer (TNBC): Primary results from the randomized phase 3 ASCENT-04/KEYNOTE-D19 study. *J. Clin. Oncol.* 43:LBA109. https://doi.org/10.1200/JCO.2025.43.17_suppl.LBA109
- Tolaney, S.M., T.M. Cardillo, C.-C. Chou, C. Dornan, and M. Faris. 2025b. The mode of action and clinical outcomes of sacituzumab govitecan in solid tumors. *Clin. Cancer Res.* 31:1390–1399. <https://doi.org/10.1158/1078-0432.CCR-24-1525>

- Tolaney, S.M., Z. Jiang, Q. Zhang, R. Barroso-Sousa, Y.H. Park, M.F. Rimawi, C. Saura, A. Schneeweiss, M. Toi, Y.S. Chae, et al. 2026. Trastuzumab deruxitecan plus pertuzumab for HER2-positive metastatic breast cancer. *N. Engl. J. Med.* 394:551–562. <https://doi.org/10.1056/NEJMoa2508668>
- Trerotola, M., P. Cantanelli, E. Guerra, R. Tripaldi, A.L. Aloisi, V. Bonasera, R. Lattanzio, R. de Lange, U.H. Weidle, M. Piantelli, and S. Alberti. 2013. Upregulation of Trop-2 quantitatively stimulates human cancer growth. *Oncogene*. 32:222–233. <https://doi.org/10.1038/onc.2012.36>
- Tsao, L.-C., J.S. Wang, X. Ma, S. Sodhi, J.V. Ragusa, B. Liu, J. McBane, T. Wang, J. Wei, C.-X. Liu, et al. 2025. Effective extracellular payload release and immunomodulatory interactions govern the therapeutic effect of trastuzumab deruxitecan (T-DXd). *Nat. Commun.* 16:3167. <https://doi.org/10.1038/s41467-025-58266-8>
- Tu, Y.-P., E. Hanze, F. Zhu, H.M. Lagrassau, C.M. Sloss, M. Method, B. Esteves, E.H. Westin, and A. Berkenblit. 2024. Population pharmacokinetics of mirvetuximab soravtansine in patients with folate receptor- α positive ovarian cancer: The antibody–drug conjugate, payload and metabolite. *Br. J. Clin. Pharmacol.* 90:568–581. <https://doi.org/10.1111/bcp.15937>
- Uba, A.I., C. Bui-Linh, J.M. Thornton, M. Olivieri, and C. Wu. 2023. Computational analysis of drug resistance of taxanes bound to human β -tubulin mutant (D26E). *J. Mol. Graph. Model.* 123:108503. <https://doi.org/10.1016/j.jmgm.2023.108503>
- Uzunparmak, B., C. Haymaker, G. Raso, S. Masciari, L. Wang, H. Lin, A. Gorur, B. Kirby, A.-M. Cimo, A. Kennon, et al. 2023. HER2-low expression in patients with advanced or metastatic solid tumors. *Ann. Oncol.* 34:1035–1046. <https://doi.org/10.1016/j.annonc.2023.08.005>
- Varaganti, P., V. Buddolla, B.A. Lakshmi, and Y.-J. Kim. 2023. Recent advances in using folate receptor 1 (FOLR1) for cancer diagnosis and treatment, with an emphasis on cancers that affect women. *Life Sci.* 326:121802. <https://doi.org/10.1016/j.lfs.2023.121802>
- Veneziani, A.C., S. Sneha, and A.M. Oza. 2024. Antibody–drug conjugates: Advancing from magic bullet to biological missile. *Clin. Cancer Res.* 30:1434–1437. <https://doi.org/10.1158/1078-0432.CCR-23-3414>
- Vergote, I., A. González-Martín, K. Fujiwara, E. Kalbacher, A. Bagaméri, S. Ghamande, J.-Y. Lee, S. Banerjee, F.C. Maluf, D. Lorusso, et al. 2024. Tisotumab vedotin as second- or third-line therapy for recurrent cervical cancer. *N. Engl. J. Med.* 391:44–55. <https://doi.org/10.1056/NEJMoa2313811>
- Verma, S., D. Miles, L. Gianni, I.E. Krop, M. Welslau, J. Baselga, M. Pegram, D.-Y. Oh, V. Diéras, E. Guardino, et al. 2012. Trastuzumab emtansine for HER2-positive advanced breast cancer. *N. Engl. J. Med.* 367:1783–1791. <https://doi.org/10.1056/NEJMoa1209124>
- Vidarsson, G., G. Dekkers, and T. Rispen. 2014. IgG Subclasses and allotypes: From structure to effector functions. *Front. Immunol.* 5:520. <https://doi.org/10.3389/fimmu.2014.00520>
- von Minckwitz, G., C.-S. Huang, M.S. Mano, S. Loibl, E.P. Mamounas, M. Untch, W. Norman, P. Rastogi, A. Schneeweiss, A. Redondo, et al. 2019. Trastuzumab emtansine for residual invasive HER2-positive breast cancer. *N. Engl. J. Med.* 380:617–628. <https://doi.org/10.1056/NEJMoa1814017>
- Vuaroqueaux, V., D. Feger, A.-L. Peille, O. Siedentopf, S. Panzade, S. Ulrich, S. Dempe, H.-H. Fiebig, and J.E. Ehlert. 2022. Abstract 1132: A systems biology approach combining ProLiFiler and cancer data miner for an enhanced preclinical characterization of the WEE-1 inhibitor adavosertib. *Cancer Res.* 82:1132. <https://doi.org/10.1158/1538-7445.AM2022-1132>
- Walsh, S.J., J.D. Bargh, F.M. Dannheim, A.R. Hanby, H. Seki, A.J. Counsell, X. Ou, E. Fowler, N. Ashman, Y. Takada, et al. 2021. Site-selective modification strategies in antibody–drug conjugates. *Chem. Soc. Rev.* 50:1305–1353. <https://doi.org/10.1039/D0CS00310G>
- Wang, J., L. Goetsch, L. Tucker, Q. Zhang, A. Gonzalez, K.S. Vaidya, A. Oleksijew, E. Boghaert, M. Song, I. Sokolova, et al. 2016. Anti-c-Met monoclonal antibody ABT-700 breaks oncogene addiction in tumors with MET amplification. *BMC Cancer*. 16:105. <https://doi.org/10.1186/s12885-016-2138-z>
- Wang, H., W. Wang, Y. Xu, Y. Yang, X. Chen, H. Quan, and L. Lou. 2017a. Aberrant intracellular metabolism of T-DM1 confers T-DM1 resistance in human epidermal growth factor receptor 2-positive gastric cancer cells. *Cancer Sci.* 108:1458–1468. <https://doi.org/10.1111/cas.13253>
- Wang, J., M.G. Anderson, A. Oleksijew, K.S. Vaidya, E.R. Boghaert, L. Tucker, Q. Zhang, E.K. Han, J.P. Palma, L. Naumovski, and E.B. Reilly. 2017b. ABBV-399, a c-met antibody–drug conjugate that targets both MET-amplified and c-met-overexpressing tumors, irrespective of MET pathway dependence. *Clin. Cancer Res.* 23:992–1000. <https://doi.org/10.1158/1078-0432.CCR-16-1568>
- Wang, X., D. Venet, F. Liffrange, D. Larsimont, M. Rediti, L. Stenbeck, F. Dupont, G. Rouas, A.J. Garcia, L. Craciun, et al. 2024. Spatial transcriptomics reveals substantial heterogeneity in triple-negative breast cancer with potential clinical implications. *Nat. Commun.* 15:10232. <https://doi.org/10.1038/s41467-024-54145-w>
- Wang, A., P. Zeng, T. Ma, H. Shi, R. Li, H. Xu, Y. Feng, Q. Liu, M. Wang, T. Chen, et al. 2025a. Establishment of a new method for detection of TROP2-positive circulating tumor cells in breast cancer. *BMC Cancer*. 25:1797. <https://doi.org/10.1186/s12885-025-14184-y>
- Wang, R., B. Hu, Z. Pan, C. Mo, X. Zhao, G. Liu, P. Hou, Q. Cui, Z. Xu, W. Wang, et al. 2025b. Antibody–drug conjugates (ADCs): Current and future biopharmaceuticals. *J. Hematol. Oncol./J Hematol. Oncol.* 18:51. <https://doi.org/10.1186/s13045-025-01704-3>
- Wei, Q., T. Yang, J. Zhu, Z. Zhang, L. Yang, Y. Zhang, C. Hu, J. Chen, J. Wang, X. Tian, et al. 2024. Spatiotemporal quantification of HER2-targeting antibody–drug conjugate bystander activity and enhancement of solid tumor penetration. *Clin. Cancer Res.* 30:984–997. <https://doi.org/10.1158/1078-0432.CCR-23-1725>
- Wen, M., A. Yu, Y. Park, D. Calarese, H.-P. Gerber, and G. Yin. 2025. Homogeneous antibody–drug conjugates with dual payloads: Potential, methods and considerations. *mAbs*. 17:2498162. <https://doi.org/10.1080/19420862.2025.2498162>
- Wu, G., Y. Gao, D. Liu, X. Tan, L. Hu, Z. Qiu, J. Liu, H. He, and Y. Liu. 2019. Study on the heterogeneity of T-DM1 and the analysis of the unconjugated linker structure under a stable conjugation process. *ACS Omega*. 4:8834–8845. <https://doi.org/10.1021/acsomega.9b00430>
- Yadati, T., T. Houben, A. Bitorina, and R. Shiri-Sverdlov. 2020. The ins and outs of cathepsins: Physiological function and role in disease management. *Cells*. 9:1679. <https://doi.org/10.3390/cells9071679>
- Yao, X., Y. Lu, S. Shi, and Y. Wang. 2026. Antibody–drug conjugates (ADCs): “Smart missiles” targeting cancer therapy. *Colloids Surf. B Biointerfaces*. 257:115161. <https://doi.org/10.1016/j.colsurfb.2025.115161>
- Ye, Q., X.-Z. Zhuang, J. Li, and X. Zhou. 2025. Targeting the inhibitors of apoptosis proteins (IAPs) to combat drug resistance in cancers. *Front. Pharmacol.* 16:1562167. <https://doi.org/10.3389/fphar.2025.1562167>
- Yim, W.W.-Y., and N. Mizushima. 2020. Lysosome biology in autophagy. *Cell Discov.* 6:6. <https://doi.org/10.1038/s41421-020-0141-7>
- Yu, J., Y. Song, and W. Tian. 2020. How to select IgG subclasses in developing anti-tumor therapeutic antibodies. *J. Hematol. Oncol.* 13:45. <https://doi.org/10.1186/s13045-020-00876-4>
- Zeng, P., M.-B. Chen, L.-N. Zhou, M. Tang, C.-Y. Liu, and P.-H. Lu. 2016. Impact of TROP2 expression on prognosis in solid tumors: A systematic review and meta-analysis. *Sci. Rep.* 6:33658. <https://doi.org/10.1038/srep33658>
- Zhao, W., H. Zhu, S. Zhang, H. Yong, W. Wang, Y. Zhou, B. Wang, J. Wen, Z. Qiu, G. Ding, et al. 2015. Trop2 is overexpressed in gastric cancer and predicts poor prognosis. *Oncotarget*. 7:6136–6145. <https://doi.org/10.18632/oncotarget.6733>
- Zhao, Y., W. Ye, Y.-D. Wang, and W.-D. Chen. 2022. HGF/c-Met: A key promoter in liver regeneration. *Front. Pharmacol.* 13:808855. <https://doi.org/10.3389/fphar.2022.808855>
- Zhao, M., T.P. DiPeri, M.G. Raso, X. Zheng, Y.Q. Rizvi, K.W. Evans, F. Yang, A. Akcakanat, M. Roberto Estecio, D. Tripathy, et al. 2023. Epigenetically upregulating TROP2 and SLFN11 enhances therapeutic efficacy of TROP2 antibody drug conjugate sacituzumab govitecan. *Npj Breast Cancer*. 9:1–14. <https://doi.org/10.1038/s41523-023-00573-8>
- Zhou, X., and Y.M. Agazie. 2012. The signaling and transformation potency of the overexpressed HER2 protein is dependent on the normally-expressed EGFR. *Cell. Signal.* 24:140–150. <https://doi.org/10.1016/j.cellsig.2011.08.015>
- Zhou, W., Z. Xu, S. Liu, X. Lou, P. Liu, H. Xie, S. Zhang, X. Liu, B. Zhuo, and H. Huang. 2024a. Landscape of clinical drug development of ADCs used for the pharmacotherapy of cancers: An overview of clinical trial registry data from 2002 to 2022. *BMC Cancer*. 24:898. <https://doi.org/10.1186/s12885-024-12652-5>
- Zhou, Z.Z., Y. Si, J. Zhang, K. Chen, A. George, S. Kim, L. Zhou, and L.X. Margaret. 2024b. A Dual-Payload Antibody–Drug Conjugate Targeting CD276/B7-H3 elicits cytotoxicity and immune activation in triple-negative breast cancer. *Cancer Res.* 84:3848–3863. <https://doi.org/10.1158/0008-5472.CAN-23-4099>
- Zidel, A., A. Benton, E. Brown, S. Shmuel, A. Vanover, S.S. Panikar, R. Bose, H. Park, A.A. Davis, and P.M.R. Pereira. 2025. Modulation of HER2 internalization enhances single-dose antibody–drug potency in HER2+ gastric cancer. *Sci. Rep.* 15:16964. <https://doi.org/10.1038/s41598-025-01947-7>
- Znojek, P., E. Willmore, and N.J. Curtin. 2014. Preferential potentiation of topoisomerase I poison cytotoxicity by PARP inhibition in S phase. *Br. J. Cancer* 111:1319–1326. <https://doi.org/10.1038/bjc.2014.378>