

Jones, L.P., S.-k. Wei, and D.T. Yue.

The Journal of General Physiology. Volume 112, No. 2, August 1998. 125–143.

Page 137.

Line five of Figure 9 legend contained an error in which two symbols were incorrectly labelled. The correct legend is shown below.

FIGURE 9. Activation and inactivation properties of a COOH-terminal truncated version of α_{1E} ($\alpha_{1E\Delta}$, amino acids 1–1871) that lacks a putative β binding site. All measurements for both α_{1E} and $\alpha_{1E\Delta}$ are in 10 mM Ba^{2+} . (A) Traces illustrating activation of ionic currents for $\alpha_{1E\Delta}\alpha_2\delta$ (cell 375_19) or $\alpha_{1E\Delta}\beta_{2a}\alpha_2\delta$ (cell 370_6) using the same voltage protocol as in Fig. 2. Data are shown for test pulse potentials of -30 , -20 , -10 , 10 , and 50 mV. (B) Comparison of G-V for $\alpha_{1E\Delta}$ and α_{1E} indicates that modulation of $\alpha_{1E\Delta}$ activation by β subunits is preserved. Symbols correspond to data for $\alpha_{1E\Delta}$ and wild-type α_{1E} ($\alpha_{1E\Delta}\beta_{2a}\alpha_2\delta$, $\bigcirc$, $n = 4$; $\alpha_{1E\Delta}\alpha_2\delta$, $\bullet$, $n = 11$; $\alpha_{1E}\alpha_2\delta$, $\square$, $n = 11$; and $\alpha_{1E}\beta_{2a}$, $\blacksquare$, $n = 8$). Lines are single-Boltzmann fits to the $\alpha_{1E}\alpha_2$ and $\alpha_{1E}\beta_{2a}$ data with fit parameters $z = 2.4$, $V_{1/2} = -10.1$; and $z = 3.4$, $V_{1/2} = -18.2$, respectively. The shift in $V_{1/2}$ values relative to Fig. 6 C corresponds to a surface potential shift between 10 mM (used here) and 2 mM Ba^{2+} (Fig. 6). (C) Steady state inactivation. Traces after a 20-s prepulse (as in Fig. 7) to the indicated potentials are shown for $\alpha_{1E\Delta}$ in combination with $\alpha_2\delta$ (cell 376_9), β_3 (cell 375_4), $\beta_{2a}\alpha_2\delta$ (cell 370_11), and $\beta_3\alpha_2\delta$ (cell 371_10). (D) Comparison of $h(\infty)$ -V relations for $\alpha_{1E\Delta}$ (symbols) and α_{1E} (same data as in Fig. 7, shown as lines connecting mean data values, without explicit reproduction of data points as symbols). Symbols correspond to the following constructs: $\alpha_{1E\Delta}\alpha_2\delta$, $\bullet$; $\alpha_{1E\Delta}\beta_{2a}\alpha_2\delta$, $\bigcirc$; $\alpha_{1E\Delta}\beta_3$, ∇ ; $\alpha_{1E\Delta}\beta_3\alpha_2\delta$, $\times$. Data for α_{1E} is shifted by -7 mV to overlay the $\alpha_{1E\Delta}$ data. Average fit values and cell numbers are summarized in Tables V (G-V) and III [$h(\infty)$ -V] for both α_{1E} and $\alpha_{1E\Delta}$.