

# Role for SUR2A ED Domain in Allosteric Coupling within the $K_{ATP}$ Channel Complex

Amy B. Karger, Sungjo Park, Santiago Reyes, Martin Bienengraeber, Roy B. Dyer, Andre Terzic, and Alexey E. Alekseev

Division of Cardiovascular Diseases, Department of Medicine and Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, MN 55905

Allosteric regulation of heteromultimeric ATP-sensitive potassium ( $K_{ATP}$ ) channels is unique among protein systems as it implies transmission of ligand-induced structural adaptation at the regulatory SUR subunit, a member of ATP-binding cassette ABCC family, to the distinct pore-forming  $K^+$  (Kir6.x) channel module. Cooperative interaction between nucleotide binding domains (NBDs) of SUR is a prerequisite for  $K_{ATP}$  channel gating, yet pathways of allosteric intersubunit communication remain uncertain. Here, we analyzed the role of the ED domain, a stretch of 15 negatively charged aspartate/glutamate amino acid residues (948–962) of the SUR2A isoform, in the regulation of cardiac  $K_{ATP}$  channels. Disruption of the ED domain impeded cooperative NBDs interaction and interrupted the regulation of  $K_{ATP}$  channel complexes by MgADP, potassium channel openers, and sulfonylurea drugs. Thus, the ED domain is a structural component of the allosteric pathway within the  $K_{ATP}$  channel complex integrating transduction of diverse nucleotide-dependent states in the regulatory SUR subunit to the open/closed states of the  $K^+$ -conducting channel pore.

## INTRODUCTION

Adoption of different functional states in response to effector binding at a site distal from the catalytic site within homomeric systems, with one or more axes of molecular symmetry, defines the allosteric regulation of enzymes (Monod et al., 1965; Monod, 1966). This is the case with ATP-binding cassette (ABC) transporters where adenine nucleotide interactions with nucleotide binding domains (NBDs) drive structural rearrangement necessary for cargo translocation across the cell membrane (Linton and Higgins, 2007). Within the ABC superfamily, regulation of the ATP-sensitive  $K^+$  ( $K_{ATP}$ ) channel complex is unique as it relies not only on structural coupling and cooperativity, induced by nucleotide interaction within the ATPase-harboring regulatory sulfonylurea receptors (SUR), but also on transmission of such interaction to the associated pore-forming Kir6.x channel (Alekseev et al., 2005).


The  $K_{ATP}$  channel heteromer is comprised of four SUR modules, members of the ABCC subfamily, and four  $K^+$  conducting pores (Aguilar-Bryan et al., 1995; Inagaki et al., 1995; 1996; Tucker et al., 1997; Lorenz and Terzic, 1999; Nichols, 2006). SUR, like other ABCC proteins, consists of two bundles of six hydrophobic transmembrane-spanning domains (TMDs) fused to hydrophilic nucleotide-binding domains (NBDs) (TMD<sub>1</sub>-NBD1-TMD<sub>2</sub>-NBD2) (Lorkowski and Cullen, 2002). An additional TMD<sub>0</sub> module with five transmembrane domains (TMD<sub>0</sub>-TMD<sub>1</sub>-NBD1-TMD<sub>2</sub>-NBD2) anchors the channel pore to the SUR module (Babenko and Bryan, 2003; Fang

et al., 2006; Hosy et al., 2007). While SUR shares the properties of ATP interaction and hydrolysis with ABC proteins (Bienengraeber et al., 2000; Matsuo et al., 2000), a catalysis-coupled transport function has not been identified for  $K_{ATP}$  channels. Catalysis in SUR, while not required for ion permeation down the electrochemical gradient, has been implicated in allosteric regulation of the heteromeric  $K_{ATP}$  channel complex (Zingman et al., 2001). In this way, the concept of allosteric regulation, developed for homomers, could be expanded to heteromers, where ligand-induced conformation adaptation within the regulatory module is translated to the functional state of the distal subunit.

While the allosteric hierarchy controlling  $K_{ATP}$  channel operation has not been dissected, the proposed model of nucleotide-dependent channel gating (Abraham et al., 2002; Selivanov et al., 2004) implies that four identical binding sites for ATP and ATP/ADP coexist on the pore-forming and regulatory subunits, respectively, within the octameric complex (Clement et al., 1997). Binding of ATP to the Kir6.x pore inhibits channel opening (Tucker et al., 1997; Drain et al., 1998), whereas MgADP at the corresponding regulatory SUR subunit antagonizes ATP-induced pore closure (Shyng et al., 1997; Matsuo et al., 2000). Cooperative rather than individual contribution of SUR NBDs gate channel operation as

Abbreviations used in this paper: ABC, ATP-binding cassette; CL, cytoplasmic loop; HRP, horseradish peroxidase; KCO, potassium channel opener; NBD, nucleotide binding domain; SUR, sulfonylurea receptor; TMD, transmembrane-spanning domain.

Correspondence to Alexey E. Alekseev: aea03@mayo.edu



**Figure 1.** The ED domain, a candidate for allosteric intersubunit communication. (A) The ED domain in SUR, located in the CL6 downstream of NBD1, consists of 15 linear aspartate and glutamate residues, and structurally overlaps with the R domain in CFTR. (B) The ED domain (in red) of SUR isoforms is not conserved in other ABC proteins. ClustalW software ([www.ch.embnet.org](http://www.ch.embnet.org)) was used to align CL6, a region between TMD1 and TMD2 that includes NBD1, for SUR isoforms, i.e., SUR1 (606–1005) and SUR2A (591–987), as well as two representative ABCC family members, MRP1 (613–963) and CFTR (351–854). The SUR ED domains are in red, while the PKA-phosphorylated serines in the R domain of CFTR are shaded in blue. While the SUR ED domains are not conserved among ABCC proteins, they match the location of the serine-rich R domain of CFTR. All sequences are from mouse.

ATP binding to NBD1, supported by hydrolysis at NBD2, is a necessary step in securing structural arrangement of both NBDs (Ueda et al., 1999; Matsuo et al., 2000, 2002; Zingman et al., 2002). The outcome on  $K_{ATP}$  channel pore state is not ultimately determined by the rate of NBD2-mediated catalysis at SUR (Zingman et al., 2001; 2002), but relies on the probability that NBD2 will adopt post- versus prehydrolytic conformations reflecting cooperative NBD interaction (Bienengraeber et al., 2004; Alekseev et al., 2005). Although this model is in agreement with experimentally defined nucleotide-dependent gating (Alekseev et al., 1998; Abraham et al., 2002), it does not elucidate the structural determinants underlying intersubunit communication.

Pharmacological  $K_{ATP}$  channel regulators, potassium channel openers and sulfonylurea drugs, albeit with distinct binding topologies within SUR also rely on nucleotide-dependent NBD states. Nucleotides and drugs induce an identical fingerprint in channel behavior, modulating termination of burst openings while maintaining an intact pattern of interburst kinetics (Alekseev et al., 1998; Bienengraeber et al., 2000). Potassium channel openers require cooperative NBD interaction, since drug effects are dependent on intracellular nucleotides and  $Mg^{2+}$ , intact NBDs, and post-hydrolytic SUR conformations (Gribble et al., 1997; Schwanstecher et al., 1998; Bienengraeber et al., 2000; Ashcroft and Gribble, 2000; Zingman et al., 2001; Moreau et al., 2005). Biochemical studies further demonstrate that binding of sulfonylureas to SUR modulates cooperative interaction between NBDs, inducing ATP dissociation from NBD1 (Ueda et al., 1999). Further, neutralization of positively charged arginine moieties in Kir6.x produces a  $K_{ATP}$  channel phenotype insensitive to  $MgADP$ -induced activation and sulfonylurea-induced inhibition (John et al., 2001). These features of  $K_{ATP}$  channel regulation collectively indicate that conformational rearrangements of NBDs induced by exogenous (openers and sulfonylurea) or endogenous ( $MgADP$ ) ligands are coupled with the open or

closed state of the channel pore through a common allosteric pathway.


To serve as transducer of diverse signals to the channel pore, components of such a pathway should not be conserved among ABC proteins, which, unlike SUR, function as complete enzymatic modules and do not require communication with a functionally distinct protein. This domain should not interfere with critical binding sites for drugs or nucleotides, yet to facilitate readout of nucleotide binding it may be located in the vicinity of NBDs. Lastly, it may comprise a significant negative charge recognized in protein–protein interactions (Shin et al., 2000; Ma et al., 2001).

The present study analyzes an aspartate/glutamate-rich domain as a candidate common transducer securing allosteric regulation in the cardiac  $K_{ATP}$  channel complex, comprised of Kir6.2 and SUR2A channel isoforms. SUR2A contains a stretch of 15 negatively charged aspartate and glutamate residues, designated the ED domain adjacent to NBD1 within the sixth cytoplasmic loop (Fig. 1 A). The ED domain does not overlap with known binding sites for  $K_{ATP}$  channel ligands (Aguilar-Bryan et al., 1998; Uhde et al., 1999; Moreau et al., 2000; Mikhailov et al., 2001; Gribble and Reimann, 2003). Sequence alignment revealed that the ED domain is not conserved among ABCC subfamily members (Fig. 1 B), yet it matches the location of the R domain in ABCC7 encoding the CFTR channel (Fig. 1 A), which when phosphorylated permits nucleotide-dependent gating of  $Cl^{-}$  conduction (Aguilar-Bryan et al., 1998; Ostedgaard et al., 2001). Thus, the ED domain emerges as a candidate allosteric transducer within  $K_{ATP}$  channel complexes.

## MATERIALS AND METHODS

### Recombinant $K_{ATP}$ Channels

SUR2A and Kir6.2 were subcloned into the pcDNA3.1 vector. Overlap PCR mutagenesis was used to create deletions and mutations within the acidic domain (Ho et al., 1989).  $K_{ATP}$  channel



**Figure 2.** ED mutants and ATP sensitivity of  $K_{ATP}$  channels. (A) Amino acid composition of the wild-type (WT) SUR2A ED domain versus ED domain mutant constructs, i.e., entire deletion, first half deletion, second half deletion, and neutrally charged entire Q/N substitution. ED domain SUR2A mutants, when expressed with Kir6.2, generated  $K_{ATP}$  channels with a respective open state probability ( $P_o$ ) similar to the wild-type counterpart, indicating that disruption of ED domain does not affect intrinsic gating of the channel pore. (B) Membrane expression of  $K_{ATP}$  channel complexes measured as chemiluminescence from extracellular HA epitope-labeled WT versus ED domain-mutated SUR2A coexpressed with Kir6.2. For each construct, a p-value is provided indicating no significant reduction in surface expression, except Kir6.2 alone, which was used as a negative control. Sample number is indicated in parentheses. (C) ATP concentration–

response curves for each of the ED mutants measured in the presence of  $Mg^{2+}$  demonstrated a moderate reduction in ATP sensitivity manifested by a rightward shift relative to wild type (WT). Curves are Hill plots constructed based on fitting results revealing the following  $IC_{50}$  values: WT,  $28 \pm 3.3 \mu M$  ( $n = 8$ ); entire deletion,  $51 \pm 4.6 \mu M$  ( $n = 6$ ); first half deletion,  $73 \pm 3.7 \mu M$  ( $n = 3$ ); second half deletion,  $35 \pm 4.1 \mu M$  ( $n = 5$ ); entire Q/N mutant,  $64 \pm 4.9 \mu M$  ( $n = 6$ ). (D) ATP concentration–response curves for WT, entire deletion, and entire Q/N mutant measured in the absence of  $Mg^{2+}$ , and characterized by indistinguishable  $IC_{50}$  values:  $14 \pm 3.7$ ,  $16 \pm 3.3$ , and  $15 \pm 2.6 \mu M$ , respectively;  $n = 3$  for each construct.

subunits were expressed in HEK293 cells, cultured in Dulbecco's Modified Eagle's Medium with 10% FBS and 2 mM glutamine at 5%  $CO_2$ . Subunits were transiently transfected (5  $\mu g$  SUR2A:1  $\mu g$  Kir6.2) with 0.5  $\mu g$  of the reporter gene GFP using FuGene (Roche). Experiments were performed 48–72 h post-transfection.

#### Photoaffinity Labeling

Labeling protocols using 8-azido- $[\gamma\text{-}^{32}P]$ ATP followed established post-incubation procedures (Ueda et al., 1999). Membrane samples (40  $\mu g$ ) isolated from HEK293 cells cotransfected with wild-type or mutated SUR2A and Kir6.2 were incubated with 100  $\mu M$  8-azido- $[\gamma\text{-}^{32}P]$ ATP, 3 mM  $MgCl_2$ , 0.1 mM EGTA, 2 mM ouabain, and 40 mM Tris-HCl pH 7.5 at 37°C for 3 min. Unbound 8-azido- $[\gamma\text{-}^{32}P]$ ATP was removed after centrifugation (13,000  $g$ , for 5 min at 4°C). Pellets were resuspended with the same buffer and then mixed with the indicated concentration of ADP. The mixture solutions were incubated for 15 min at 37°C and then irradiated on ice at maximal output ( $\sim 1 J/cm^2$ ) for 8 min in a UVP CL1000 ultraviolet cross-linker. Samples were electrophoresed on 12% SDS-PAGE and autoradiographed.

#### Membrane Expression of $K_{ATP}$ Channel Constructs

A surface expression of  $K_{ATP}$  channel constructs were determined by using HA tag incorporated in the extracellular loop between transmembrane helices 16 and 17 of wild-type or mutated SUR2A (Schwappach et al., 2000). Tagged SUR2A were cotransfected with Kir6.2. After 48 h, transfected cells were fixed with 4% formaldehyde for 15 min and washed twice with PBS. After 30 min of blocking (1% FBS in PBS), cells were incubated with monoclonal HA antibody 3F10 (Roche) for 1 h, washed, and incubated with goat anti-mouse horseradish peroxidase (HRP)–conjugated secondary antibody (20 min). After washing, power signal ELISA solution (Pierce Chemical Co.) was added

and chemiluminescence read after 15 s in a plate reader (Molecular Devices). Chemiluminescence signal was normalized to the protein content.


#### Immunoblotting

Proteins were separated in 3–8% Tris-Acetate gel (Invitrogen) from membrane samples isolated from HEK293 cells cotransfected with Kir6.2 and wild-type or ED-mutated SUR2A. Resolved proteins were transferred by electrophoresis onto PVDF membrane. The presence of SUR2A was probed by incubation with SUR2A-specific antibody (Santa Cruz Biotechnology, Inc.) counterstained by anti-goat IgG HRP conjugate. Membranes were incubated in Super Signal West Pico Substrate (Pierce Chemical Co.), and signals captured on CL-XPosure film (Pierce Chemical Co.).

#### Electrophysiology

Recombinant  $K_{ATP}$  channel activity was recorded using the inside-out configuration of the patch-clamp methodology at a membrane potential of  $-60$  mV. Patch pipettes contained (in mM) KCl 140,  $CaCl_2$  1,  $MgCl_2$  1, HEPES-KOH 5 (pH 7.3). Cells were bathed in internal solution containing (in mM) KCl 140,  $MgCl_2$  1, EGTA 5, HEPES-KOH 5 (pH 7.3). After patch excision,  $K_{ATP}$  channel activity was measured in the same internal solution, supplemented by drugs and/or nucleotides. Mg-free internal solution was supplemented by 5 mM EDTA with omitted  $MgCl_2$ . Recordings were performed at  $32 \pm 1^\circ C$  using the HI-25x heating element and the TC $^2$  temperature controller (Cell Micro Controls). Determination of the open state probability ( $P_o$ ), through calculation of  $NP_o$  values, was performed in channel recordings when a clear determination of the number of simultaneously operating  $K_{ATP}$  channels ( $n < 10$ ) was possible (Jovanovic et al., 1996).  $P_o$  were calculated with known  $N$  using obtained  $NP_o$  values averaged for a 30–40-s-long duration of channel activity.





**Figure 3.** Intact ED domain required for MgADP-dependent channel activation. (A) Mg-nucleotide-dependent  $K_{ATP}$  channel gating was probed as the channel response to 0.3 mM MgADP in WT and ED mutants inhibited by 0.3 mM ATP. Disruption of the ED domain prevented MgADP-dependent channel activation in inside-out patches. Dotted lines represent zero-current levels. (B) Activation of ED domain mutants by MgADP is significantly blunted when compared with WT ( $P < 0.05$ ) in inside-out patches. Channel activity expressed relative to that measured in the absence of nucleotides. Sample size is indicated in parentheses.

#### Statistics

Data are expressed as mean  $\pm$  SEM; p-values were calculated using unpaired Student's *t* test. A p-value  $< 0.05$  was considered significant.


## RESULTS

### Role of the ED Domain in Nucleotide-dependent Channel Gating

To examine the role of the ED domain in  $K_{ATP}$  channel regulation, deletion and mutations were made within this unique region of SUR2A. Specifically, three deletion mutants were constructed: first half deletion ( $\Delta 948-954$ ), second half deletion ( $\Delta 955-962$ ), and entire deletion ( $\Delta 948-962$ ) (Fig. 2 A). In addition, to ensure that the effects with deletion mutants are not due to nonspecific shortening of the cytoplasmic loop, the entire Q/N mutant was made replacing negatively charged aspartates and glutamates with neutrally charged asparagines and glutamines (Fig. 2 A). Expression of Kir6.2 with the ED domain mutants generated channels that retained basic properties of  $K_{ATP}$  channels. These included an identical single channel conductance ( $\sim 85$  pS at  $32^\circ\text{C}$ ) and intrinsic probability of the open state ( $P_o$ ) measured based on  $NP_o$  values in the absence of drug or nucleotides ( $P_o \approx 0.76-0.8$ ), indicating unimpaired pore gating in ED mutants versus wild-type  $K_{ATP}$  channel constructs (Fig. 2 A). All SUR2A ED domain mutants coexpressed with Kir6.2 produced vigorous  $K_{ATP}$  channel activity, detected via patch-clamp, indicating unaltered surface expression compared with wild-type  $K_{ATP}$  channels, further confirmed by chemiluminescence measured from extracellular HA epitope-labeled channel constructs (Fig. 2 B).

The ED domain mutants exhibited a moderate reduction in ATP sensitivity manifested by a rightward shift of MgATP concentration-dependent curves with the observed  $IC_{50}$  ranging from 35 to 73  $\mu\text{M}$ , relative to 28  $\mu\text{M}$  measured for the wild type (Fig. 2 C). However, in the absence of  $Mg^{2+}$ , the ATP concentration-dependent relationships constructed for wild-type, entire deletion, and entire Q/N mutant  $K_{ATP}$  channels revealed identical  $IC_{50}$  values (Fig. 2 D), indicating that deletion or neutralization of negative charges of the ED domain did not affect apparent ATP binding to the channel pore. Inhibition of wild-type  $K_{ATP}$  channels by 300  $\mu\text{M}$  ATP was readily reversed by 300  $\mu\text{M}$  MgADP (Fig. 3 A). While 300  $\mu\text{M}$  ATP inhibited the ED mutant SUR2A/Kir6.2 channels similarly to the wild type, MgADP was unable to reverse such inhibition in ED mutants (Fig. 3, A and B). Thus, while the ED domain of SUR2A is not directly involved in ATP-induced pore inhibition, it is essential in mediating MgADP-dependent channel activation.

**ED Domain Critical for Cooperative Interaction of NBDs**  
MgADP, through binding or as a product of MgATP hydrolysis at NBD2, stabilizes nucleotide binding to NBD1 of SUR, suggesting that cooperative interaction between NBDs is a critical conformational rearrangement for ADP-induced  $K_{ATP}$  channel activation (Ueda et al., 1999; Matsuo et al., 2000; Zingman et al., 2002). To test the role of the ED domain in cooperative NBD interaction, the effect of ADP on photoaffinity labeling of membrane proteins by 8-azido- $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  in cells coexpressing Kir6.2 with wild-type or ED-mutated SUR2A was assessed using a post-incubation procedure established to evaluate cooperative binding (Ueda et al., 1999). Untransfected cells, used as control, were photolabeled in the



**Figure 4.** ED domain is critical for cooperative NBD interaction. Membrane proteins isolated from HEK293 cells were photoaffinity labeled. (A) Nonspecific photoaffinity labeling was determined in samples from untransfected cells using a range of ADP concentrations. Nonspecific labeling was essentially independent from ADP concentration and used as background to define specific labeling. (B) Effect of ADP on photoaffinity SUR2A labeling was assessed in samples from cells expressing Kir6.2 along with of WT or ED mutant constructs. Untransfected lanes, in the absence of ADP, indicate nonspecific labeling relative to each construct. (C) Control Western blots of membranes purified from cells transfected with Kir6.2/WT SUR2A indicate the expected molecular weight for SUR2A. (D and E) Photoaffinity labeling was quantified based on densitometry expressed in arbitrary units ( $n = 3$ ) with regard to nonspecific labeling detected from membranes of untransfected cells in the absence (D) or presence of ADP (E). Data for nonspecific ATP photolabeling measured from untransfected cells in the range of ADP concentra-

tions are presented in absolute arbitrary units ( $n = 3$ ). In comparison with WT, ED domain mutants exhibited reduced ATP photoaffinity labeling of NBD1 in the ADP range from 0 to 1 mM, and lack of potentiation of labeling by ADP.

absence or presence of MgADP in the range from 0.05 to 1.0 mM. Nonspecific labeling in untransfected samples was negligible and unaffected by ADP titration (Fig. 4 A). Due to inherent catalytic activity centered at NBD2, leading to hydrolysis of the ATP  $\gamma$ -phosphate, 8-azido- $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  labeling reflects the presence of nucleotides at NBD1 that harbors low or no ATPase activity (Ueda et al., 1999; Bienengraeber et al., 2000; Matsuo et al., 2000; Fig. 4 B). Immunoblotting with an SUR2A-specific antibody validated that photolabeled bands in wild-type or ED domain mutants corresponded to SUR2A (Fig. 4 C). In the absence of MgADP, the entire deletion and Q/N mutants demonstrated reduced 8-azido- $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  labeling compared with wild type (Fig. 4, B and D). After titration with MgADP, wild-type constructs exhibited increased 8-azido- $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  labeling, reaching a maximum at 500  $\mu\text{M}$ , consistent with previous findings (Ueda et al., 1999). In contrast, MgADP within the applied concentration range did not affect labeling of the ED-mutated constructs (Fig. 4, B and E). Thus, an intact ED domain is critical for cooperative NBD interaction and may therefore be essential for readout and transmission of nucleotide interactions to the channel pore.


#### Role of ED Domain in Opener-induced $\text{K}_{\text{ATP}}$ Channel Activation

ED disruption blunted the activating effect of the potassium channel opener (KCO) pinacidil (Fig. 5), although the ED domain does not overlap with binding sites for pinacidil and related cyanoguanidines that interact with

the cytoplasmic loop 7 (CL7) between TM helices 13 and 14, as well as TM helices 16 and 17 of SUR2 (Ude et al., 1999; Moreau et al., 2000). Specifically, 50  $\mu\text{M}$  pinacidil antagonized the inhibition of wild-type  $\text{K}_{\text{ATP}}$  channels by 300  $\mu\text{M}$  ATP, reversing channel activity close to the maximum level measured in the absence of nucleotides (Fig. 5). In contrast, channel activity induced by the same concentration of pinacidil, in the presence of 300  $\mu\text{M}$  ATP, was 30–50% lower for the ED domain mutants when compared with the wild-type counterpart (Fig. 5). Thus, pinacidil-induced antagonism of ATP-induced Kir6.2 pore was impaired, albeit not abolished, in ED domain mutants.

#### Role of ED Domain in Sulfonylurea-induced $\text{K}_{\text{ATP}}$ Channel Inhibition

On wild-type  $\text{K}_{\text{ATP}}$  channels, the sulfonylurea glyburide (10  $\mu\text{M}$ ) exhibited a dual time course of inhibition, with fast and slow rates manifested by instantaneous blockade of only a fraction ( $59 \pm 11\%$ ;  $n = 9$ ) of the channel population (Fig. 6, A and B) followed by gradual dissipation of channel activity ( $93 \pm 3\%$ ;  $n = 9$ ) in the remaining subpopulation within 2–3 min (Fig. 6 A). Deletion of the entire ED domain generated a monophasic response to glyburide, with loss of the slow phase of inhibition and a reduced fast phase so that glyburide-induced reduction in overall channel activity was only  $21 \pm 5\%$ ;  $n = 3$  (Fig. 6, A and B). Similarly, first half deletion, second half deletion, and the entire Q/N mutant produced channels with reduced glyburide responsiveness manifested




**Figure 5.** Intact ED domain required for potassium channel opener-induced channel activation. (A) Disruption of ED domain blunts the activating effect of pinacidil (50  $\mu$ M) on recombinant  $K_{ATP}$  channels in the presence of inhibitory concentration of ATP (300  $\mu$ M). (B) In all mutant constructs, the effect of pinacidil was significantly blunted when compared with WT channels ( $P < 0.05$ ). Number of values averaged for each experimental condition is indicated in parentheses. Channel activity expressed relative to that measured in the absence of nucleotides and drugs.


primarily by a modest slow phase of glyburide-induced inhibition compared with wild type (Fig. 6, A and B). Thus, the ED domain contributes to sulfonylurea-induced  $K_{ATP}$  channel inhibition affecting to various degrees both phases of glyburide-mediated inhibition.

To assess the mechanism underlying the dual phase of the  $K_{ATP}$  channel response to glyburide, the effect of the

sulfonylurea was tested on wild-type channels. Removal of glyburide did not reverse current inhibition, which required reapplication of ATP (Fig. 6 A), suggesting that at least the slow phase of inhibition is pertinent to the nucleotide-dependent state of the channel complex and resembles the described phenomenon of  $K_{ATP}$  channel rundown (Trube and Hescheler, 1984; Findlay and



**Figure 6.** Role of ED domain in glyburide-induced  $K_{ATP}$  channel inhibition. (A) All ED domain mutants, in comparison with WT, exhibited an essentially monophasic response toward glyburide (10  $\mu$ M), with little to no slow phase. (B)  $K_{ATP}$  channel activity was measured at the end of the fast phase and at the end of 3 min after glyburide application (slow phase) and expressed relative to the control activity in the absence of drugs and nucleotide. The effect of the drug on ED-mutated constructs was significantly different from WT ( $P < 0.05$ ) except for the fast phase measured in the second half deletion mutant. Number of values averaged for each experimental condition is indicated in parentheses.




**Figure 7.** Biphasic glyburide-induced channel inhibition of wild-type  $K_{ATP}$  channels. (A) In the absence of nucleotide only a fraction of the channel population present in the patch (100–200 channels) was instantaneously blocked by the sulfonylurea glyburide (10  $\mu$ M). The rest manifested gradual and eventually complete dissipation of activity after glyburide application. Reapplication and removal of ATP (300  $\mu$ M) restored  $K_{ATP}$  channel activity. In the presence of 50  $\mu$ M MgATP, glyburide induced only a fast phase of  $K_{ATP}$  channel inhibition. In the presence of 300  $\mu$ M MgADP, glyburide maintained the dual pattern of  $K_{ATP}$  channel blockade as measured without nucleotides, but both phases of inhibition were blunted. In the presence of AMP-PNP, a nonhydrolyzable ATP analogue, the profile of glyburide inhibition was virtually indistinguishable from the effect of glyburide in the absence of nucleotides. (B) Number of values averaged for each experimental condition is indicated in parentheses. Fast and slow phases of glyburide inhibition were quantified as in Fig. 6 B. (C) In the presence of hydrolyzable ATP (50  $\mu$ M), glyburide-induced channel inhibition was reversed after washout of glyburide. (D) AMP-PNP (50  $\mu$ M) did not prevent channel rundown initiated by the sulfonylurea and did not support restoration of channel activity after removal of glyburide.

Dunne, 1986; Nichols and Lederer, 1991; Tung and Kurachi, 1991; Findlay, 1994; Terzic et al., 1994). One of the contributing conditions of  $K_{ATP}$  channel rundown is a fall in  $PIP_2$  levels, which however is a slow process lasting >1 h following nucleotide removal (Ribalet et al., 2000), and therefore per se cannot explain the slow phase of glyburide-induced channel inhibition that occurs with a significantly faster time course. To test whether the slow phase of glyburide-induced channel inhibition could be accounted based on nucleotide-dependent states of SUR2A, glyburide action was further probed in the presence of adenine nucleotides or the nonhydrolyzable AMP-PNP analogue known to induce distinct NBD formations (Gribble et al., 1998b; Matsuo et al., 2000; Zingman et al., 2001; Alekseev et al., 2005; Nichols, 2006).

At a membrane potential of  $-60$  mV and a single channel conductance of 85 pS, the single channel current ampli-

tude was estimated at  $\sim 5$  pA and at a given level of protein expression and open state probability of  $\sim 0.8$ , patches with estimated 100–200 operating recombinant channels were selected to ensure discrimination of the probabilistic properties of the time course of  $K_{ATP}$  channel closure (Fig. 7 A). In the presence of 50  $\mu$ M MgATP, which partially blocked  $K_{ATP}$  channels but was used to prevent channel rundown, glyburide (10  $\mu$ M) induced only the fast phase of  $K_{ATP}$  channel inhibition with an amplitude of inhibition double that obtained in the absence of MgATP (Fig. 7, A and B). Despite a more prominent fast phase, glyburide in the presence of 50  $\mu$ M MgATP did not induce maximal inhibition of channel activity seen with glyburide alone. On average, the steady level of  $K_{ATP}$  channel activity in the presence of 50  $\mu$ M MgATP and glyburide was  $22 \pm 4\%$  ( $n = 5$ ) of that measured before sulfonylurea application (Fig. 7 B). In the presence of 300  $\mu$ M MgADP,





**Figure 8.** Suggested role of the ED domain in  $K_{ATP}$  channel gating. (A) In the presence of nucleotides, the intact SUR2A ED domain contributes to optimal coordination of both NBDs, essential in the antagonism of ATP-induced Kir6.2 inhibition. Cooperative NBD formation, controlled by the ED domain, is also a prerequisite in mediating  $K_{ATP}$  channel activation in response to potassium channel opener (KCO) interaction with SUR2A. (B) Glyburide binding to SUR2A destabilizes cooperative NBDs interaction, with ED domain participation, accomplished by dissociation of ATP from NBD1. Such NBDs/ED domain conformational rearrangement is critical for transduction of inhibitory channel gating originating from SUR2A to pore closure. (C) Disruption of the ED domain stabilizes the NBDs of SUR2A in a suboptimal conformation characterized by residual communication between constitutive channel subunits. Hypothetic occlusion of MgADP within such suboptimal NBD conformation endows moderate antagonism of ATP-induced pore inhibition (right shift of ATP concentration responses), partial channel activation by KCO, and impaired MgADP-induced channel opening. The inflexibility of this NBD formation in the absence of the ED domain precludes glyburide-induced  $K_{ATP}$  channel inhibition, yet prevents the rundown process associated with nucleotide-dependent states of SUR2A.

glyburide maintained the dual pattern of  $K_{ATP}$  channel blockade as measured without nucleotides, but both phases were blunted (Fig. 7 A). Although it partially antagonized sulfonylurea-induced channel inhibition as previously reported (Gribble et al., 1998a; Reimann et al., 2003), MgADP apparently could not equivalently replace MgATP in supporting nucleotide-driven NBD formation. This is further supported by the observation that in the presence of AMP-PNP, a nonhydrolyzable analogue of ATP, the profile of glyburide inhibition was indistinguishable from the effect of glyburide in the absence of nucleotides (Fig. 7, A and B). While in the presence of hydrolyzable ATP, channel inhibition could be reversed (Takano et al., 1990; Furukawa et al., 1994), following washout of glyburide, the nonhydrolyzable ATP analogue did not prevent channel rundown initiated by the sulfonylurea and did not support restoration of channel activity after removal of glyburide (Fig. 7, C and D). Restoration of channel activity after glyburide treatment in the presence of AMP-PNP required MgATP (Fig. 7 D). Taken together, these observations suggest that the effect of the sulfonylurea glyburide relies upon nucleotide-dependent conformations of SUR2A associated with its catalytic function involving cooperative interaction between NBDs (Ueda et al., 1999; Matsuo et al., 2000; Zingman et al., 2001, 2002). Thus, disruption or neutralization of the ED domain interrupted allosteric communication of channel inhibition induced by a sulfonylurea at SUR2A to the  $K_{ATP}$  channel pore.

## DISCUSSION


Gating of the cardiac  $K_{ATP}$  channel complex implies transduction of ligand-induced conformational transitions in the regulatory SUR2A module to the Kir6.2 pore. In the

present study, the negatively charged ED domain was identified as a structural component within SUR2A contributing to allosteric signal communication between constitutive  $K_{ATP}$  channel subunits. Specifically,  $K_{ATP}$  channel regulation by MgADP, potassium channel openers, and sulfonylurea drugs, which relies on nucleotide-dependent NBD states, was dependent on the intactness of the ED domain. Indeed, while the ED domain does not overlap with known ligand binding sites, deletion or charge neutralization of this acidic domain impeded  $K_{ATP}$  channel pore function in response to either activating or inhibitory SUR2A ligands. The strategic location of the ED domain downstream of NBD1, in the cytoplasmic loop 6 (CL6) between TMD1 and TMD2, facilitated readout of cooperative NBD formation communicated to the channel pore. Whether communication of SUR2A and Kir6.2 involves direct electrostatic interaction via the ED domain or implies additional intermediate structural components remains to be determined and will require more detailed elaboration of the  $K_{ATP}$  channel architecture.

## ED Domain Equivalents in ABC Proteins

Although the ED domain is not conserved among other ABCC proteins, structural alignment indicated topological equivalence of the ED domain in SUR with the R domain in CFTR channels, a related family member. In fact, PKA-mediated phosphorylation of serine residues within the R domain negatively charges this region, permitting  $Cl^-$  conductance by the CFTR protein (Rich et al., 1993; Seibert et al., 1999; Ostedgaard et al., 2001) presumably after cooperative NBD interaction (Ostedgaard et al., 1997; Aleksandrov et al., 2001, 2002; Vergani et al., 2003; Howell et al., 2004). This effect is mimicked by mutating serines to negatively charged aspartate residues, allowing CFTR operation in the absence of PKA phosphorylation





**Figure 9.** Model for biphasic glyburide-mediated inhibition of  $K_{ATP}$  channels. (A) The  $K_{ATP}$  channel population exhibiting cooperative formation of NBDs is denoted by filled rectangles and attached diamonds. The rest of the  $K_{ATP}$  channel population is in a noncooperative NBD state, with either or no nucleotide bound, is denoted in half-filled shapes in brackets. In the absence of ATP, glyburide (G) inhibits  $K_{ATP}$  channels by “shaking” the cooperative state, promoting dissociation of ATP from NBD1 (see Ueda et al., 1999). The channel population in noncooperative states contributes only to the rundown state (denoted as open circles) through the slow phase promoted by glyburide. In the absence of ATP the effect of glyburide is irreversible since, following sulfonylurea removal, closed channel may transit exclusively to rundown, which is functionally indistinguishable from the closed state. (B) In contrast, ATP can support nucleotide binding and hydrolysis and thereby the steady state between cooperative and noncooperative states, revealing the fast phase of the glyburide effect as incomplete and reversible.

(Rich et al., 1993; Ostedgaard et al., 2001). While the SUR ED domain, and its R domain analogue in CFTR, may be uniquely required for ion channel gating, the role of equivalent domains may be more broadly applicable to other members of the ABC protein family. Indeed, protein sequence alignment demonstrated that the linker region in p-glycoprotein (P-gp), an ABCB subfamily member, structurally overlaps the ED and R domains and contains several consensus sites for phosphorylation (Orr et al., 1993). Additionally, replacement of the CFTR R domain residues 780–830 with P-gp linker residues 644–689 resulted in stimulated channel activity indistinguishable from that of wild-type CFTR (Vankeerberghen et al., 1999). Thus, the ED domain shares topological and functional properties of domains critical for the regulation of related ABC proteins.

#### ED Domain Controls Cooperative NBD Interaction

Nucleotide-dependent gating of  $K_{ATP}$  channels is coupled to the catalytic function of NBD2 and requires nucleotide coordination at NBD1 (Gribble et al., 1997, 1998b; Zingman et al., 2001, 2002; Matsuo et al., 2002), implying determinate linkage between catalysis-driven SUR conformations and pore gating (Aleksiev et al., 2005). Evidence for conformational association of SUR NBDs stems from the dimeric structures of distinct ABC proteins (Hung et al., 1998; Diederichs et al., 2000; Hopfner et al., 2000; Campbell et al., 2003) and is supported by direct labeling studies (Ueda et al., 1999; Matsuo et al., 2000). Mutations in the ED domain were here found to disrupt cooperative nucleotide binding at NBD. Thus, the ED domain is integral to the allosteric machinery controlling the readout of NBD states essential for optimal channel regulation (Fig. 8).

#### ED Domain Mediates $K_{ATP}$ Channel Activation

Both NBDs as well as the engagement of NBD2 into post-hydrolytic conformations are prerequisites for  $K_{ATP}$  channel activation by MgADP and potassium channel openers (Gribble et al., 1997; Schwanstecher et al., 1998; Bienengraeber et al., 2000; Zingman et al., 2001). Disruption of the ED domain antagonized  $K_{ATP}$  channel opening induced by MgADP and the potassium channel opener pinacidil. This could be explained by destabilization of NBD dimer formation required for channel activation (Gribble et al., 1998b; Zingman et al., 2002; Aleksiev et al., 2005). Moreover, the observed minor rightward shift of MgATP (but not Mg-free ATP)-induced  $K_{ATP}$  channel inhibition in the ED mutants despite loss of MgADP efficacy could result from stabilization of the NBD2 ATPase cycle in a post-hydrolytic conformation (Bienengraeber et al., 2004). Although an alternative pathway for potassium channel openers cannot be excluded (Russ et al., 2003), the residual effectiveness of potassium channel openers measured in the ED domain mutants could also be related to the stabilization of such post-hydrolytic conformation (Schwanstecher et al., 1998; Zingman et al., 2001). Thus, the intact ED domain secures optimal cooperativity of NBDs and proper  $K_{ATP}$  channel activation, whereas disruption in the ED domain locks NBDs in a suboptimal state, allowing only residual communication between SUR2A and the channel pore (Fig. 8).

#### ED Domain and Inhibitory Channel Gating

The reduced sensitivity of ED-mutated  $K_{ATP}$  channel constructs toward the inhibitory effect of glyburide, which interacts by its benzamido moiety with the cytoplasmic loop 3 (CL3) of SUR2 between TMD0 and TMD1 (Mikhailov et al., 2001), may also be explained by a suboptimal NBD

state induced by disruption of the ED domain. The present findings provide a framework for a mechanistic model of channel regulation to interpret the nucleotide dependence of glyburide inhibition of  $K_{ATP}$  channels (Fig. 9). Upon removal of ATP, a population of  $K_{ATP}$  channels remains in a nucleotide-bound state in line with cooperative dimer conformation of SUR2A NBDs. This is in accord with SUR properties to stabilize bound ATP and MgADP at NBD1 and NBD2, respectively (Ueda et al., 1999; Matsuo et al., 2000). When glyburide is added, two phases of inhibition are observed. Binding of the sulfonylurea to dimerized SUR disrupts cooperative interaction of NBDs and promotes ATP dissociation from NBD1 (Ueda et al., 1999) allosterically associated with immediate pore closure (Fig. 9 A). The  $K_{ATP}$  channel subpopulation that at glyburide application is outside of cooperative NBD interaction can resist sulfonylurea-induced channel closure, and contributes only to the slower phase of channel inhibition characterized as sulfonylurea-accelerated rundown. This is supported by glyburide-induced  $K_{ATP}$  channel inhibition in the presence of 50  $\mu$ M MgATP (Fig. 9 B), an ATP concentration that prevents rundown and, although insufficient per se to keep channel closed, can apparently support cooperative interaction of NBDs. Under this condition, virtually the whole  $K_{ATP}$  channel population is primed for sulfonylurea-induced channel block manifested by exclusively fast inhibition (Fig. 9 B). MgADP blunted both phases of glyburide-induced  $K_{ATP}$  channel inhibition in line with the known property of nucleotide diphosphates to prevent rundown, and the inability to mimic ATP-bound SUR conformations that are most sensitive to sulfonylurea (Nichols and Lederer, 1991; Tung and Kurachi, 1991; Alekseev et al., 1998; Reimann et al., 2003). Thus, the attenuated sensitivity of ED domain mutants to glyburide, which revealed essentially a monophasic profile of channel inhibition, could indicate that deletion or neutralization of negative charges in the ED domain stabilizes the NBDs in suboptimal conformation, rendering  $K_{ATP}$  channels less sensitive to rundown. Collectively, sulfonylurea, which binds to SUR2A, induced pore inhibition through a common NBD-dependent mechanism involving the ED domain-mediated pathway (Fig. 8).

In conclusion, transmission of ligand-induced structural adaptation of NBDs at the SUR2A module to the channel pore defines the operation of cardiac  $K_{ATP}$  channels. The present study identifies the acidic ED domain of SUR2A as a structural component of allosteric  $K_{ATP}$  channel gating integrating transduction of diverse nucleotide-dependent states in the regulatory module to the  $K^+$ -conducting channel pore. In this way, the identified role for the ED domain in allosteric coupling within the  $K_{ATP}$  channel complex could provide a foundation for the further understanding of structure–function relationships among ABC proteins.

This work was supported by National Institutes of Health (HL64822).

Lawrence G. Palmer served as editor.

Submitted: 6 July 2007

Accepted: 5 February 2008

## REFERENCES

- Abraham, M.R., V.A. Selivanov, D.M. Hodgson, D. Pucar, L.V. Zingman, B. Wieringa, P.P. Dzeja, A.E. Alekseev, and A. Terzic. 2002. Coupling of cell energetics with membrane metabolic sensing. Integrative signaling through creatine kinase phosphotransfer disrupted by M-CK gene knock-out. *J. Biol. Chem.* 277:24427–24434.
- Aguilar-Bryan, L., C.G. Nichols, S.W. Wechsler, J.P. Clement IV, A.E. Boyd III, G. Gonzalez, H. Herrera-Sosa, K. Nguy, J. Bryan, and D.A. Nelson. 1995. Cloning of the  $\beta$  cell high-affinity sulfonylurea receptor: a regulator of insulin secretion. *Science*. 268:426–429.
- Aguilar-Bryan, L., J.P. Clement IV, G. Gonzalez, K. Kunjilwar, A. Babenko, and J. Bryan. 1998. Toward understanding the assembly and structure of  $K_{ATP}$  channels. *Physiol. Rev.* 78:227–245.
- Aleksandrov, L., A.A. Aleksandrov, X.B. Chang, and J.R. Riordan. 2002. The first nucleotide binding domain of cystic fibrosis transmembrane conductance regulator is a site of stable nucleotide interaction, whereas the second is a site of rapid turnover. *J. Biol. Chem.* 277:15419–15425.
- Aleksandrov, L., A. Mengos, X. Chang, A. Aleksandrov, and J.R. Riordan. 2001. Differential interactions of nucleotides at the two nucleotide binding domains of the cystic fibrosis transmembrane conductance regulator. *J. Biol. Chem.* 276:12918–12923.
- Alekseev, A.E., P.A. Brady, and A. Terzic. 1998. Ligand-insensitive state of cardiac ATP-sensitive  $K^+$  channels. Basis for channel opening. *J. Gen. Physiol.* 111:381–394.
- Alekseev, A.E., D.M. Hodgson, A.B. Karger, S. Park, L.V. Zingman, and A. Terzic. 2005. ATP-sensitive  $K^+$  channel channel/enzyme multimer: metabolic gating in the heart. *J. Mol. Cell. Cardiol.* 38:895–905.
- Ashcroft, F.M., and F.M. Gribble. 2000. New windows on the mechanism of action of  $K_{ATP}$  channel openers. *Trends Pharmacol. Sci.* 21:439–445.
- Babenko, A.P., and J. Bryan. 2003. SUR domains that associate with and gate  $K_{ATP}$  pores define a novel gatekeeper. *J. Biol. Chem.* 278:41577–41580.
- Bienengraeber, M., A.E. Alekseev, M.R. Abraham, A.J. Carrasco, C. Moreau, M. Vivaudou, P.P. Dzeja, and A. Terzic. 2000. ATPase activity of the sulfonylurea receptor: a catalytic function for the  $K_{ATP}$  channel complex. *FASEB J.* 14:1943–1952.
- Bienengraeber, M., T.M. Olson, V.A. Selivanov, E.C. Kathmann, F. O’Coilain, F. Gao, A.B. Karger, J.D. Ballew, D.M. Hodgson, L.V. Zingman, et al. 2004. ABCC9 mutations identified in human dilated cardiomyopathy disrupt catalytic  $K_{ATP}$  channel gating. *Nat. Genet.* 36:382–387.
- Campbell, J.D., M.S. Sansom, and F.M. Ashcroft. 2003. Potassium channel regulation. *EMBO Rep.* 4:1038–1042.
- Clement, J.P., K. Kunjilwar, G. Gonzalez, M. Schwanstecher, U. Panten, L. Aguilar-Bryan, and J. Bryan. 1997. Association and stoichiometry of  $K_{ATP}$  channel subunits. *Neuron*. 18:827–838.
- Diederichs, K., J. Diez, G. Greller, C. Muller, J. Breed, C. Schnell, C. Vornrhein, W. Boos, and W. Welte. 2000. Crystal structure of MalK, the ATPase subunit of the trehalose/maltose ABC transporter of the archaeon *Thermococcus litoralis*. *EMBO J.* 19:5951–5961.
- Drain, P., L. Li, and J. Wang. 1998.  $K_{ATP}$  channel inhibition by ATP requires distinct functional domains of the cytoplasmic C terminus of the pore-forming subunit. *Proc. Natl. Acad. Sci. USA*. 95:13953–13958.
- Fang, K., L. Csanády, and K.W. Chan. 2006. The N-terminal transmembrane domain (TMD0) and a cytosolic linker (L0) of sulfonylurea receptor define the unique intrinsic gating of  $K_{ATP}$  channels. *J. Physiol.* 576:379–389.

- Findlay, I., and M.J. Dunne. 1986. ATP maintains ATP-inhibited K<sup>+</sup> channels in an operational state. *Pflugers Arch.* 407:238–240.
- Findlay, I. 1994. Interactive regulation of the ATP-sensitive potassium channels of cardiac muscle. *J. Cardiovasc. Pharmacol.* 24:S6–S11.
- Furukawa, T., L. Virag, N. Furukawa, T. Sawanobori, and M. Hiraoka. 1994. Mechanism for reactivation of the ATP-sensitive K<sup>+</sup> channel by MgATP complexes in guinea-pig ventricular myocytes. *J. Physiol.* 479:95–107.
- Gribble, F.M., S.J. Tucker, and F.M. Ashcroft. 1997. The essential role of the Walker A motifs of SUR1 in K<sub>ATP</sub> channel activation by Mg-ADP and diazoxide. *EMBO J.* 16:1145–1152.
- Gribble, F.M., S.J. Tucker, S. Seino, and F.M. Ashcroft. 1998a. Tissue specificity of sulfonylureas: studies on cloned cardiac and  $\beta$ -cell K<sub>ATP</sub> channels. *Diabetes.* 47:1412–1418.
- Gribble, F.M., S.J. Tucker, T. Haug, and F.M. Ashcroft. 1998b. MgATP activates the  $\beta$  cell K<sub>ATP</sub> channel by interaction with its SUR1 subunit. *Proc. Natl. Acad. Sci. USA.* 95:7185–7190.
- Gribble, F.M., and F. Reimann. 2003. Sulphonylurea action revisited: the post-cloning era. *Diabetologia.* 46:875–891.
- Ho, S.N., H.D. Hunt, R.M. Horton, J.K. Pullen, and L.R. Pease. 1989. Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene.* 77:51–59.
- Hopfner, K.P., A. Karcher, D.S. Shin, L. Craig, L.M. Arthur, J.P. Carney, and J.A. Tainer. 2000. Structural biology of Rad50 ATPase: ATP-driven conformational control in DNA double-strand break repair and the ABC-ATPase superfamily. *Cell.* 101:789–800.
- Hosy, E., R. Dérand, J. Revilloud, and M. Vivaldoux. 2007. Remodelling of the SUR-Kir6.2 interface of the K<sub>ATP</sub> channel upon ATP binding revealed by the conformational blocker rhodamine 123. *J. Physiol.* 582:27–39.
- Howell, L.D., R. Borchardt, J. Kole, A.M. Kaz, C. Randak, and J.A. Cohn. 2004. Protein kinase A regulates ATP hydrolysis and dimerization by a CFTR (cystic fibrosis transmembrane conductance regulator) domain. *Biochem. J.* 378:151–159.
- Hung, L.W., I.X. Wang, K. Nikaido, P.Q. Liu, G.F. Ames, and S.H. Kim. 1998. Crystal structure of the ATP-binding subunit of an ABC transporter. *Nature.* 396:703–707.
- Inagaki, N., T. Gono, J.P. Clement IV, N. Namba, J. Inazawa, G. Gonzalez, L. Aguilar-Bryan, S. Seino, and J. Bryan. 1995. Reconstitution of I<sub>KATP</sub>: an inward rectifier subunit plus the sulfonylurea receptor. *Science.* 270:1166–1170.
- Inagaki, N., T. Gono, J.P. Clement IV, C.Z. Wang, L. Aguilar-Bryan, J. Bryan, and S. Seino. 1996. A family of sulfonylurea receptors determines the pharmacological properties of ATP-sensitive K<sup>+</sup> channels. *Neuron.* 16:1011–1017.
- John, S.A., J.N. Weiss, and B. Ribalet. 2001. Regulation of cloned ATP-sensitive K<sup>+</sup> channels by adenine nucleotides and sulfonylureas: interactions between SUR1 and positively charged domains on Kir6.2. *J. Gen. Physiol.* 118:391–405.
- Jovanovic, A., A.E. Alekseev, and A. Terzic. 1996. Cardiac ATP-sensitive K<sup>+</sup> channel: a target for diadenosine 5',5''-P<sub>1</sub>,P<sub>5</sub>-pentaphosphate. *Naunyn Schmiedeberg's Arch. Pharmacol.* 353:241–244.
- Linton, K.J., and C.F. Higgins. 2007. Structure and function of ABC transporters: the ATP switch provides flexible control. *Pflugers Arch.* 453:555–567.
- Lorenz, E., and A. Terzic. 1999. Physical association between recombinant cardiac ATP-sensitive K<sup>+</sup> channel subunits Kir6.2 and SUR2A. *J. Mol. Cell. Cardiol.* 31:425–434.
- Lorkowski, S., and P. Cullen. 2002. ABCG subfamily of human ATP-binding cassette proteins. *Pure Appl. Chem.* 74:2057–2081.
- Ma, Z., E. Cshai, K.M. Chow, and L.B. Hersh. 2001. Expression of the acidic stretch of nardilysin as a functional binding domain. *Biochemistry.* 40:9447–9452.
- Matsuo, M., K. Tanabe, N. Kioka, T. Amachi, and K. Ueda. 2000. Different binding properties and affinities for ATP and ADP among sulfonylurea receptor subtypes, SUR1, SUR2A, and SUR2B. *J. Biol. Chem.* 275:28757–28763.
- Matsuo, M., M. Dabrowski, K. Ueda, and F.M. Ashcroft. 2002. Mutations in the linker domain of NBD2 of SUR inhibit transduction but not nucleotide binding. *EMBO J.* 21:4250–4258.
- Mikhailov, M.V., E.A. Mikhailova, and S.J.H. Ashcroft. 2001. Molecular structure of the glibenclamide binding site of the  $\beta$ -cell channel. *FEBS Lett.* 499:154–160.
- Monod, J. 1966. From enzymatic adaptation to allosteric transitions. *Science.* 154:475–483.
- Monod, J., J. Wyman, and J.-P. Changeux. 1965. On the nature of allosteric transitions: a plausible model. *J. Mol. Biol.* 12:88–118.
- Moreau, C., H. Jacquet, A.L. Prost, N. D'Hahan, and M. Vivaldoux. 2000. The molecular basis of the specificity of action of K<sub>ATP</sub> channel openers. *EMBO J.* 19:6644–6651.
- Moreau, C., A.L. Prost, R. Dérand, and M. Vivaldoux. 2005. SUR, ABC proteins targeted by K<sub>ATP</sub> channel openers. *J. Mol. Cell. Cardiol.* 38:951–963.
- Nichols, C.G. 2006. K<sub>ATP</sub> channels as molecular sensors of cellular metabolism. *Nature.* 440:470–476.
- Nichols, C.G., and W.J. Lederer. 1991. Adenosine triphosphate-sensitive potassium channels in the cardiovascular system. *Am. J. Physiol.* 261:H1675–H1686.
- Orr, G.A., E.K.H. Han, P.C. Browne, E. Nieves, B.M. O'Connor, C.P.H. Yang, and S.B. Horwitz. 1993. Identification of the major phosphorylation domain of murine *mdr1b* P-glycoprotein. *J. Biol. Chem.* 268:25054–25062.
- Ostedgaard, L.S., O. Baldursson, and M.J. Welsh. 2001. Regulation of the cystic fibrosis transmembrane conductance regulator Cl<sup>−</sup> channel by its R domain. *J. Biol. Chem.* 276:7689–7692.
- Ostedgaard, L.S., D.P. Rich, L.G. DeBerg, and M.J. Welsh. 1997. Association of domains within the cystic fibrosis transmembrane conductance regulator. *Biochemistry.* 36:1287–1294.
- Reimann, F., M. Dabrowski, P. Jones, F.M. Gribble, and F.M. Ashcroft. 2003. Analysis of the differential modulation of sulphonylurea block of  $\beta$ -cell and cardiac ATP-sensitive K<sup>+</sup> (K<sub>ATP</sub>) channels by Mg-nucleotides. *J. Physiol.* 547:159–168.
- Ribalet, B., S. John, and J. Weiss. 2000. Regulation of cloned ATP-sensitive K channels by phosphorylation, MgADP, and phosphatidylinositol bisphosphate (PIP<sub>2</sub>): a study of channel rundown and reactivation. *J. Gen. Physiol.* 116:391–410.
- Rich, D.P., H.A. Berger, S.H. Cheng, S.M. Travis, M. Saxena, A.E. Smith, and M.J. Welsh. 1993. Regulation of the cystic fibrosis transmembrane conductance regulator Cl<sup>−</sup> channel by negative charge in the R domain. *J. Biol. Chem.* 268:20259–20267.
- Russ, U., U. Lange, C. Löffler-Walz, A. Hambrock, and U. Quast. 2003. Binding and effect of K<sub>ATP</sub> channel openers in the absence of Mg<sup>2+</sup>. *Br. J. Pharmacol.* 139:368–380.
- Schwanstecher, M., C. Sieverding, H. Dörschner, I. Gross, L. Aguilar-Bryan, C. Schwanstecher, and J. Bryan. 1998. Potassium channel openers require ATP to bind to and act through sulfonylurea receptors. *EMBO J.* 17:5529–5535.
- Schwappach, B., N. Zerangue, Y.N. Jan, and L.Y. Jan. 2000. Molecular basis for K<sub>ATP</sub> assembly: transmembrane interactions mediate association of a K<sup>+</sup> channel with an ABC transporter. *Neuron.* 26:155–167.
- Seibert, F.S., X.B. Chang, A.A. Aleksandrov, D.M. Clarke, J.W. Hanrahan, and J.R. Riordan. 1999. Influence of phosphorylation by protein kinase A on CFTR at the cell surface and endoplasmic reticulum. *Biochim. Biophys. Acta.* 1461:275–283.
- Selivanov, V.A., A.E. Alekseev, D.M. Hodgson, P.P. Dzeja, and A. Terzic. 2004. Nucleotide-gated K<sub>ATP</sub> channels integrated with creatine and adenylate kinases: amplification, tuning and sensing of energetic signals in the compartmentalized cellular environment. *Mol. Cell. Biochem.* 256-257:243–256.

- Shin, D.W., J. Ma, and D.H. Kim. 2000. The asp-rich region at the carboxyl-terminus of calsequestrin binds to  $\text{Ca}^{2+}$  and interacts with triadin. *FEBS Lett.* 486:178–182.
- Shyng, S., T. Ferrigni, and C.G. Nichols. 1997. Regulation of  $\text{K}_{\text{ATP}}$  channel activity by diazoxide and MgADP. Distinct functions of the two nucleotide binding folds of the sulfonylurea receptor. *J. Gen. Physiol.* 110:643–654.
- Takano, M., D. Qin, and A. Noma. 1990. ATP-dependent decay and recovery of  $\text{K}^+$  channels in guinea-pig cardiomyocytes. *Am. J. Physiol.* 258:H45–H50.
- Terzic, A., I. Findlay, Y. Hosoya, and Y. Kurachi. 1994. Dualistic behavior of ATP-sensitive  $\text{K}^+$  channels toward intracellular nucleoside diphosphates. *Neuron.* 12:1049–1058.
- Trube, G., and J. Hescheler. 1984. Inward-rectifying channels in isolated patches of the heart cell membrane: ATP-dependence and comparison with cell-attached patches. *Pflugers Arch.* 401:178–184.
- Tucker, S.J., F.M. Gribble, C. Zhao, S. Trapp, and F.M. Ashcroft. 1997. Truncation of Kir6.2 produces ATP-sensitive  $\text{K}^+$  channels in the absence of the sulphonylurea receptor. *Nature.* 387:179–183.
- Tung, R.T., and Y. Kurachi. 1991. On the mechanism of nucleotide diphosphate activation of the ATP-sensitive  $\text{K}^+$  channel in ventricular cell of guinea-pig. *J. Physiol.* 437:239–256.
- Ueda, K., J. Komine, M. Matsuo, S. Seino, and T. Amachi. 1999. Cooperative binding of ATP and MgADP in the sulfonylurea receptor is modulated by glibenclamide. *Proc. Natl. Acad. Sci. USA.* 96:1268–1272.
- Uhde, I., A. Toman, I. Gross, C. Schwanstecher, and M. Schwanstecher. 1999. Identification of the potassium channel opener site on sulfonylurea receptors. *J. Biol. Chem.* 274:28079–28082.
- Vankeerberghen, A., W. Lin, M. Jaspers, H. Cuppens, B. Nilius, and J.J. Cassiman. 1999. Functional characterization of the CFTR R domain using CFTR/MDR1 hybrid and deletion constructs. *Biochemistry.* 38:14988–14998.
- Vergani, P., A.C. Nairn, and D.C. Gadsby. 2003. On the mechanism of MgATP-dependent gating of CFTR  $\text{Cl}^-$  channels. *J. Gen. Physiol.* 121:17–36.
- Zingman, L.V., A.E. Alekseev, M. Bienengraeber, D.M. Hodgson, A.B. Karger, P.P. Dzeja, and A. Terzic. 2001. Signaling in channel/enzyme multimers: ATPase transitions in SUR module gate ATP-sensitive  $\text{K}^+$  conductance. *Neuron.* 31:233–245.
- Zingman, L.V., D.M. Hodgson, M. Bienengraeber, A.B. Karger, E.C. Kathmann, A.E. Alekseev, and A. Terzic. 2002. Tandem function of nucleotide binding domains confers competence to sulfonylurea receptor in gating ATP-sensitive  $\text{K}^+$  channels. *J. Biol. Chem.* 277:14206–14210.