Local elementary purinergic-induced Ca²⁺ transients: from optical mapping of nerve activity to local Ca²⁺ signaling networks

David C. Hill-Eubanks, Matthias E. Werner, and Mark T. Nelson^{1,2}

The autonomic nervous system regulates smooth muscle contractility through both sympathetic and parasympathetic influences. In some tissues, such as the urinary bladder, parasympathetic influences predominate and nerves communicate to detrusor smooth muscle through the release of acetylcholine (ACh). In other tissues, such as the vas deferens and mesenteric arterial circulation, the primary autonomic influence is sympathetic, and norepinephrine (NE) is the predominant neurotransmitter. NE and ACh act on smooth muscle cells through G_q-coupled α-adrenergic and muscarinic receptors, respectively, which signal through PLC to elevate diacylglycerol and inositol trisphosphate (IP₃), which in turn, activate PKC and IP3 receptors (IP3Rs) in the SR. IP₃-mediated Ca²⁺ release from the SR of vascular smooth muscle cells gives rise to Ca²⁺ waves (Iino and Tsukioka, 1994; Jaggar and Nelson, 2000; Wray et al., 2005; Kim et al., 2008), which are propagating elevations in Ca²⁺ that are thought to contribute to vascular smooth muscle contraction (Mauban et al., 2001; Zang et al., 2006). The consequences of G protein-coupled signaling events manifest after a characteristic lag, reflecting the temporal dynamics of multiple sequential and parallel molecular linkages.

The Journal of General Physiology

Although NE and ACh are the prototypical transmitters released by autonomic nerves, it has long been known that ATP is coreleased with NE at sympathetic nerve–muscular junctions and with ACh at parasympathetic nerve–muscular junctions. Coreleased ATP acts on P2X receptor channels in the plasma membrane of smooth muscle cells. Because P2X receptors are ion channels, once activated, their effects are experienced almost immediately by the cell. This rapid time course is in contrast to the more delayed influence of the G protein–coupled adrenergic and muscarinic receptors.

Correspondence to Mark T. Nelson: Mark.Nelson@uvm.edu

Abbreviations used in this paper: α,β -meATP, α,β -methylene ATP; ACh, acetylcholine; CICR, Ca^{2^+} -induced Ca^{2^+} release; EJC, excitatory junction current; EJP, excitatory junction potential; IP $_3$, inositol trisphosphate; IP $_3$ R, IP $_3$ receptor; jCaT, junctional Ca^{2^+} transient; NCT, neuroeffector Ca^{2^+} transient; NE, norepinephrine; NEPCaT, nerve-evoked elementary purinergic Ca^{2^+} transient; P2X $_1$ R-KO, P2X $_1$ receptor knockout; VDCC, voltage-dependent Ca^{2^+} channel.

P2X receptors represent a family of seven receptors (P2X₁₋₇) that belong to the transmitter-gated ion channel superfamily, which also includes nicotinic-like receptors and glutamate-like receptors (for review see Khakh, 2001). Each P2X receptor subunit possesses intracellular N and C termini and two membrane-spanning domains linked by a large extracellular domain (for review see Khakh, 2001; North, 2002). P2X receptors are thought to consist of three subunits (Aschrafi et al., 2004), which is also the simplest stoichiometry among ionotropic receptors. At least three ATP molecules bind to the extracellular domain of P2X channels (Jiang et al., 2003). Upon binding ATP, P2X receptors undergo conformational changes that result in the opening of the pore within milliseconds, although the underlying molecular details have not yet been elucidated.

P2X receptors are nonselective cation channels that exhibit a permeability to Ca2+ approximately equal to that of sodium (Na+) (Schneider et al., 1991). Thus, activation of P2X receptors by ATP released at nerve-muscle junctions causes a rapid local influx of Na⁺ and Ca²⁺ (Lamont and Wier, 2002; Lamont et al., 2006). Although most of the excitatory junction current (EJC) associated with P2X activation is carried by the more abundant (\sim 70-fold) Na⁺ ions, the influx of Ca²⁺ is quite substantial. In fact, the fractional Ca²⁺ currents mediated by the rat $(\sim 12.4\%)$ and human $(\sim 11\%)$ P2X₁ isoforms are not significantly different from that of the NMDA channel $(\sim 14\%)$ (Egan and Khakh, 2004), long considered the gold standard for high-level, ligand-gated Ca2+ entry. The current mediated by Na⁺ and Ca²⁺ influx creates an excitatory junction potential (EJP) that contributes directly to the increase in postjunctional excitability associated with autonomic stimulation.

The P2X₁ receptor is the predominant P2X receptor isoform expressed in smooth muscle. It was originally cloned from the vas deferens (Valera et al., 1994), and immunocytochemical studies in mice have shown that

¹Department of Pharmacology, University of Vermont, Burlington, VT 05405

²Division of Cardiovascular and Endocrine Sciences, University of Manchester, Manchester M13 9PL, England, UK

^{© 2010} Hill-Eubanks et al. This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see http://www.rupress.org/terms). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at http://creativecommons.org/licenses/by-nc-sa/3.0/).

P2X₁ expression in the urinary bladder is restricted to detrusor smooth muscle (Vial and Evans, 2000). The most compelling evidence for the prominence of the P2X₁ isoform in smooth muscle comes from studies using P2X₁ receptor knockout (P2X₁R-KO) mice. These studies have shown that ATP-evoked EJCs and EJPs are absent in the vas deferens from P2X₁R-KO mice (Mulryan et al., 2000). Similarly, these mice lack nerve-evoked purinergic contractile responses in bladders (Vial and Evans, 2000) and mesenteric arteries (Vial and Evans, 2002; Lamont et al., 2006).

Nerve-evoked elementary purinergic Ca²⁺ transients: NCTs, jCaTs, and NEPCaTs

As first demonstrated by Brain et al. (2002), the postjunctional action of ATP can be detected optically in the form of discrete, focal Ca²⁺ increases in smooth muscle cells. Using confocal microscopy and a mouse vas deferens preparation in which both smooth muscle and nerve varicosities are loaded with the Ca²⁺ indicator dye Oregon Green 488 BAPTA-1, these authors found that nerve stimulation evokes intermittent Ca2+ transients at tightly clustered sites immediately adjacent to nerve varicosities. These events, termed neuroeffector Ca2+ transients (NCTs), are temporally linked to the stimulating impulse (average delay, ~6 ms) and are preceded by an increase in Ca²⁺ in the adjacent nerve varicosity. NCTs are abolished by persistent exposure to the P2X agonist/desensitizing agent α,β-methylene ATP (α,β-meATP) and unaffected by inhibition of voltagedependent Ca²⁺ channels (VDCCs), α₁-adrenergic receptors, or IP₃Rs (Brain et al., 2003), establishing their likely identity as Ca²⁺ influx mediated by ATP-activated P2X receptors.

Shortly thereafter, Wier and colleagues reported similar spatially localized Ca²⁺ transients in vascular smooth muscle cells of pressurized mesenteric arteries (Lamont

and Wier, 2002). The authors termed these events junctional Ca²⁺ transients (jCaTs). Using the Ca²⁺-binding dye fluo-4 and fluorescence confocal microscopy, these authors showed that jCaTs are largely unaffected by ryanodine, which abolishes RYR-mediated Ca²⁺ sparks. Instead, they are blocked by the nonselective P2X receptor antagonist suramin, transiently induced by the application of the P2X receptor agonist (desensitizing agent) α,β-meATP, and absent in P2X₁-KO mice, confirming that these events are records of Ca²⁺ influx through postjunctional P2X1 receptors (Lamont and Wier, 2002; Lamont et al., 2006). ¡CaTs induced by electrical field stimulation of associated nerves exhibit a close temporal relationship to the stimulus (latency, generally <3 ms). ¡CaTs also occur spontaneously, reflecting spontaneous neurotransmitter release.

Work in our laboratory has shown that both spontaneous and nerve-evoked elementary purinergic Ca²⁺ transients (NEPCaTs) can also been detected in the urinary bladder. These events are blocked by suramin and desensitization with α,β -meATP in rat urinary bladder smooth muscle, and are absent in P2X₁R-KO mice (Heppner et al., 2009), showing that they reflect Ca²⁺ entry through P2X₁ receptor channels. They are also unaffected by inhibitors of IP₃Rs (2-APB), RYRs (ryanodine), or VDCCs (dihydropyridines) (Heppner et al., 2005), confirming that they are distinct from Ca²⁺ "puffs," Ca²⁺ sparks (Pérez et al., 1999; Jaggar et al., 2000), and the VDCCmediated Ca2+ sparklets described by Santana and Navedo (2009) (see also Table I). In mouse bladders, spontaneous Ca²⁺ transients are coincident with spontaneous EJPs, and their magnitudes are correlated (Young et al., 2008), clearly linking these optical events with long-studied, postjunctional electrical events.

The kinetic properties of purinergic Ca²⁺ transients identified in vas deferens (NCTs), mesenteric arteries (jCaTs), and urinary bladder (NEPCaTs) are similar to

TABLE I

Comparison of local Ca²⁺ transients in smooth muscle

Parameter	Purinergic Ca ²⁺ transients			Ca ²⁺ spark ^{d,e}	Ca ²⁺ puff ^f	Ca ²⁺ sparklet ^g
	Vas deferens (NCT) ^a	Mesenteric artery (jCaT) ^b	Urinary bladder (NEPCaT) ^c			
Duration (t _{1/2}) (ms)	120 ^h ; 280 ⁱ	~145	~112	\sim 56	\sim 375	(τ) 23; 104
Area (μm²)	\sim 12	\sim 25	\sim 14	\sim 13.6	2-4	\sim 0.8
Amplitude (F/F ₀)	n/d	~ 2.8	\sim 2.0	~ 2.0	n/a	n/a
Amplitude (nM)	n/d	n/a	n/a	100-200	50-500	38
Latency (ms)	\sim 6	<3	8-16	n/a	n/a	n/a

^aBrain et al., 2002.

^bLamont and Wier, 2002.

^cHeppner et al., 2005.

dPerez et al., 1999.

^eJaggar et al., 2000.

Ledoux et al., 2008.

gSantana and Naveda, 2009.

^hLine scan.

ⁱxy scan.

one another and are clearly distinct from those of other focal Ca²⁺ transients (Table I). The spatial spread and duration ($t_{1/2}$) of these events in the bladder are 14 µm² and \sim 112 ms, respectively, and the corresponding values for mesenteric artery jCaTs are \sim 20 µm² and 145 ms. Using line scanning to analyze the kinetics of NCTs, Brain et al. (2002) showed that NCTs measured in mouse vas deferens have a spatial spread of \sim 12 µm² and decay with a first-order time constant ($t_{1/2}$) of \sim 120 ms. The decay time constant obtained by xy scanning is much larger (\sim 280 ms), a difference that was attributed to the contribution of cytoplasmic diffusion of Ca²⁺ near the site of entry.

The kinetic properties of spontaneous and evoked purinergic transients are the same, suggesting that these events are caused by the quantal release of ATP. This is consistent with earlier evidence that quantal release of ATP is responsible for EJPs and/or EJCs in femoral and mesenteric arteries, rat tail artery, and vas deferens (for review see Stjärne and Stjärne, 1995). The low probability, highly intermittent quality of the events recorded in these electrophysiological studies conforms well with the predictions of the "intermittent model" developed to describe NE release from sympathetic nerves, which posits that a single vesicle in $\sim 1\%$ of all varicosities releases its entire content in response to a nerve impulse (Stjärne and Stjärne, 1995, and references therein).

Exploiting this logic, Cunnane and colleagues have used NCTs as a means to detect "packeted release" of ATP from nerve terminals (Brain et al., 2002; Young et al., 2007;

Brain, 2009). Their results based on electrophysiological measurements in single smooth muscle cells showed that the amplitude distribution of spontaneous EJPs is skewed, suggesting a broad distribution of spontaneously released neurotransmitter packet size (Young et al., 2007). Although bulk changes in ATP release can be monitored electrophysiologically as EJPs (or EJCs), this approach is less suitable for mapping quantal transmitter release because smooth muscle cells are large and electrically coupled, making it difficult to determine whether the recorded event originates in the cell being recorded (and if so, where), or is caused by release events that occur at some distance removed from the recording site. Optically measuring Ca2+, released focally by ATP-activated P2X₁Rs, overcomes these limitations, allowing more accurate spatial mapping of ATP release sites. And because ATP-induced, P2X receptor-mediated Ca²⁺ influx is very rapid, optical mapping also provides fine temporal resolution of the underlying transmitter release events. jCaTs in mesenteric arteries (Lamont and Wier, 2002) and NEPCaTs in urinary bladder (Heppner et al., 2005) have also been used to optically map ATP release by sympathetic and parasympathetic nerves, respectively.

In addition to spatial and temporal mapping of ATP release events, NCTs can provide information about coreleased transmitters and their potential local modulation of transmitter release probability. One example of this is using NCTs to monitor the prejunctional autoinhibitory effects of neurally released transmitters. In this context, the frequency of nerve-evoked NCTs was




Figure 1. Local elementary purinergic-induced Ca2+ transients and possible local Ca2+ signaling networks. ATP released from a nerve varicosity activates smooth muscle P2X₁Rs, which then allow influx of Na+ and Ca2+ ions. Ca2+ influx can induce CICR from RYRs (Brain et al., 2003) and, in theory, also from IP₃Rs. Local influx of Ca²⁺ may also lead to activation of NFAT (via calcineurin) or Ca2+dependent $K^{\scriptscriptstyle +}$ (K_{Ca}) channels. Finally, membrane depolarization caused by Na⁺ and Ca²⁺ influx through P2X1Rs would also activate voltage-dependent ion channels, such as VDCCs or K+ (K_V) channels.

shown to increase in the presence of the α_2 -adrenoceptor inhibitor yohimbine (Brain et al., 2002), providing evidence that coreleased NE acts through prejunctional α_2 -adrenoceptors to reduce nerve terminal Ca^{2+} concentration and decrease the probability of exocytosis (Brain et al., 2002; Brain, 2009). In a similar vein, potential off-target effects of pharmacological agents on prejunctional targets can be inferred from changes in the frequency of purinergic Ca^{2+} transients upon the application of such agents, a strategy we have used in studies on the urinary bladder and mesenteric arteries (unpublished data).

To the extent that release of different transmitters is coupled (i.e., not differentially regulated), detection of local purinergic Ca²⁺ transients could provide the means to optically map nerve activity generally. Whether ATP and NE in sympathetic nerve terminals are stored and/or released together has been extensively studied by Stjärne and colleagues. This seemingly straightforward question is deceptively difficult to answer, especially given the available experimental tools. Early reports from this group based on electrochemical and electrophysiological studies in rat tail arteries suggested that ATP and NE are indeed released in parallel by nerve stimulation, with apparent deviations from this conclusion likely reflecting differences in clearance rates (for review see Stjärne and Stjärne, 1995). The use of a paired-pulse stimulus paradigm provided support for this interpretation, showing that the dramatic depression of paired-pulse transmitter release caused by K⁺ channel block has similar effects on NE oxidation currents and ATP-mediated EJCs, and purinergic and adrenergic contractile responses (Msghina et al., 1998). However, more recent work by these researchers presents a more nuanced picture. The results of these studies suggest that ATP and NE are stored in separate small vesicles that are released in parallel upon low frequency stimulation (<2 Hz), but above this frequency show increasingly nonparallel release (Stjärne, 2001).

Implications of local Ca²⁺ signaling:

Ca²⁺-signaling networks

Individual purinergic Ca²⁺ transients have the potential to signal locally to modulate Ca²⁺-sensitive processes (Fig. 1). Although this is largely unexplored territory, some features of such local signaling networks can be discerned from published reports, and it is possible to speculate about others.

Action potential trigger or current injection. A single purinergic Ca²⁺ transient represents the activation of a cluster of P2X₁Rs by local ATP from a nerve varicosity. This local injection of current could conceivably trigger an action potential. Indeed, Young et al. (2008) recently demonstrated that in intact UBSM strips from mice, single NCTs cause spontaneous depolarizations ("sDeps")

and can trigger "spontaneous" action potentials. These action potentials cause phasic contractions, which contribute to muscle tone during bladder filling; an increase in their activity is a hallmark of unstable detrusor and urinary bladder dysfunction. The observation that purinergic Ca²⁺ transients can trigger action potentials suggests that bladder filling is also under local neurogenic control.

In vas deferens, NCTs do not map to action potentials in a simple one-to-one relationship (Brain et al., 2002). Although not all NCTs elicit an action potential, $\sim 20\%$ of NCTs are rapidly (<0.5 s) followed by an action potential. Consistent with this, purinergic Ca²⁺ transients in bladder co-occur with large increases in global Ca²⁺ termed "flashes" (Heppner et al., 2005). Both purinergic Ca²⁺ transients and Ca²⁺ flashes occur spontaneously, and the frequency of both types of events is increased by nerve stimulation. Ca²⁺ flashes are associated with tissue contraction and are eliminated by dihydropyridines, indicating that they are caused by Ca2+ influx through VDCCs during an action potential. Purinergic Ca²⁺ transients are unaffected by inhibition of VDCCs, but inhibition of P2X receptors abrogates Ca²⁺ flashes, implying that the cationic flux registered by the optical NCT/ jCaT/NEPCaT event lies upstream of the action potential and is responsible for triggering it. If sufficiently large, the current and associated depolarization associated with a single purinergic Ca²⁺ transient is capable of triggering an action potential. Presumably, the associated membrane potential depolarization responsible for activating VDCCs is attributable to the much larger influx of Na⁺ rather than the optically registered influx of Ca²⁺, although this has not been directly tested.

Ca²⁺-induced Ca²⁺ release (CICR): NCT/jCaT/NEPCaT to RYR/IP₃R communication. Ca²⁺ influx through a cluster of P2X1Rs may activate nearby RYRs in the SR, which should contribute to the purinergic Ca²⁺ transient (Fig. 1). In vas deferens, unlike mesenteric artery and urinary bladder smooth muscle, inhibition of RYRs with ryanodine substantially reduces the amplitude of NCTs (\sim 45%), and activation of RYRs with caffeine (3 mM) induces a dramatic (16-fold) increase in the frequency of NCTs (Brain et al., 2003). In addition, the inhibition of SR Ca²⁺ uptake by Ca²⁺ SR/ER-ATPase with cyclopiazonic acid increases the half-life of these events. These results suggest a functional unit in which Ca²⁺ influx mediated by P2X1 stimulates CICR from RYRs, which augments the local P2X₁-mediated Ca²⁺ signal. According to the model proposed by Brain et al. (2003), the duration of the Ca²⁺ signal is governed by the summation of these two Ca2+ release events as well as the rate at which released Ca²⁺ is sequestered by the SR through Ca²⁺ SR/ ER-ATPase pump activity. Such a mechanism would be consistent with the larger spread and longer half-life of NCT/jCaT/NEPCaTs compared with sparks. Ryanodine reduces jCaT amplitude by a much more modest, but

significant, \sim 13% in mesenteric arteries (Lamont and Wier, 2002), and does not appear to affect purinergic Ca²⁺ transients in the urinary bladder (Heppner et al., 2005), suggesting that the effect of RYR inhibition is at least quantitatively different among these different smooth muscle tissues. Consistent with the lack of CICR in the regulation of spontaneous phasic contractions, the inhibition of RYRs does not decrease, but instead enhances, the frequency of phasic contractions in the urinary bladder smooth muscle from the guinea pig (Herrera et al., 2000). Depending on the relative speed of IP₃ production by concurrent activation of adrenergic or muscarinic receptors, it is also possible that Ca²⁺ influx through P2X₁Rs could amplify local IP₃R activation by IP₃ (Fig. 1). This remains to be explored.

NCT/jCaT/NEPCaT to K_{Ca} channel communication. The timing, nature, and proximity of the actions of coreleased ATP and NE/ACh suggest that local Ca²⁺ influx through P2X₁ receptors might modulate the subsequent effects of nerve-evoked NE/ACh release on smooth muscle. Nerve-released ATP acts rapidly on postjunctional P2XRs to cause a rapid influx of Ca²⁺ (and Na⁺), and the coreleased NE and ACh act more slowly on their respective G_q-coupled receptors. One intriguing possibility that has not yet been experimentally tested is that the influx of Ca²⁺ associated with a purinergic Ca²⁺ transient might activate Ca²⁺-sensitive K⁺ (K_{Ca}) channels, such as small-conductance SK or large-conductance BK channels (Fig. 1). Such a mechanism might conceivably account for our observation that the cholinergic component of parasympathetic nerve-evoked action potentials in mouse urinary bladder is augmented by inhibition of P2X receptors (with suramin or α,β -meATP, or by genetic ablation of P2X₁Rs) (Heppner et al., 2009). These results imply that ATP-mediated P2X₁R activity normally exerts an inhibitory influence on the subsequent ACh-muscarinic receptor signaling pathway. If K_{Ca} channels are, in fact, activated by purinergic Ca²⁺ transients, their activity would be predicted to dampen cholinergic signaling and limit the duration of the cholinergic action potential through their membrane hyperpolarizing action.

Interestingly, purinergic signaling may have the opposite effect on NE signaling in mesenteric arteries, at least in fully pressurized arteries (90 mmHg). Here, the purinergic component of sympathetic nerve-evoked vasoconstriction is similar in the presence and absence of the α_1 -adrenergic receptor antagonist prazosin, but the adrenergic component is substantially higher in the presence of functional P2X receptors than it is with P2X receptors blocked with suramin (Rummery et al., 2007). Thus, it appears that purinergic activity may exert a potentiating effect on adrenergic signaling in this setting, consistent with a possible postjunctional influence of smooth muscle P2X receptors.

NCT/jCaT/NEPCaT to transcription factor communication. Recent evidence from vascular smooth muscle indicates that VDCCs are associated with a macromolecular complex containing PKC and AKAP150, as well as calcineurin and the Ca²⁺-dependent transcription factor NFAT (Navedo et al., 2008, 2010). Using this parallel, it is possible to speculate that complexes of P2X₁Rs with kinases and phosphatases, including those that regulate NFAT activation, might also be present in postjunctional smooth muscle cell membranes (Fig. 1). Colocalization of transmitter receptors (adrenergic and cholinergic) and ion channels (e.g., VDCCs and K_{Ca} channels) in membrane microdomains might add a further level of regulation to such Ca²⁺-dependent signaling.

Closing thoughts. Purinergic Ca²⁺ transients, by whatever name, are likely a common feature of nerve-smooth muscle junctions, where they may feed into tissue-specific local Ca²⁺ signaling networks and potentially modulate a myriad of Ca²⁺-dependent processes. Most of the potential connections between NCT/jCaT/NEPCaT-like events and intracellular signal transduction have not yet been experimentally tested and remain a matter of conjecture. Regardless of the functional roles that these events prove to play in postsynaptic smooth muscle cells, they should provide a convenient and sensitive optical readout of neurally released ATP specifically and, to an asyet-undetermined extent, of neural activity generally. This Perspectives series includes articles by Gordon, Parker and Smith, Xie et al., Prosser et al., and Santana and Navedo.

This work was supported by National Institutes of Health (grants R37DK 053832, RO1 DK065947, RO1 HL44455, PO1 HL077378, P20 R016435, and RO1 HL098243); Totman Trust for Medical Research, Research into Ageing (grant P332); The Royal Society (grant RG080197); and the British Heart Foundation (grant PG/07/115).

REFERENCES

Aschrafi, A., S. Sadtler, C. Niculescu, J. Rettinger, and G. Schmalzing. 2004. Trimeric architecture of homomeric P2X2 and heteromeric P2X1+2 receptor subtypes. *J. Mol. Biol.* 342:333–343. doi:10.1016/j.jmb.2004.06.092

Brain, K.L. 2009. Neuroeffector Ca2+ transients for the direct measurement of purine release and indirect measurement of cotransmitters in rodents. *Exp. Physiol.* 94:25–30. doi:10.1113/expphysiol.2008.043679

Brain, K.L., V.M. Jackson, S.J. Trout, and T.C. Cunnane. 2002. Intermittent ATP release from nerve terminals elicits focal smooth muscle Ca2+ transients in mouse vas deferens. *J. Physiol.* 541:849–862. doi:10.1113/jphysiol.2002.019612

Brain, K.L., A.M. Cuprian, D.J. Williams, and T.C. Cunnane. 2003. The sources and sequestration of Ca(2+) contributing to neuroeffector Ca(2+) transients in the mouse vas deferens. *J. Physiol.* 553:627–635. doi:10.1113/jphysiol.2003.049734

Egan, T.M., and B.S. Khakh. 2004. Contribution of calcium ions to P2X channel responses. *J. Neurosci.* 24:3413–3420. doi:10.1523/JNEUROSCI.5429-03.2004

- Heppner, T.J., A.D. Boney, and M.T. Nelson. 2005. Elementary purinergic Ca2+ transients evoked by nerve stimulation in rat urinary bladder smooth muscle. J. Physiol. 564:201-212. doi:10.1113/jphysiol.2004.077826
- Heppner, T.J., M.E. Werner, B. Nausch, C. Vial, R.J. Evans, and M.T. Nelson. 2009. Nerve-evoked purinergic signalling suppresses action potentials, Ca2+ flashes and contractility evoked by muscarinic receptor activation in mouse urinary bladder smooth muscle. J. Physiol. 587:5275–5288. doi:10.1113/jphysiol.2009.178806
- Herrera, G.M., T.J. Heppner, and M.T. Nelson. 2000. Regulation of urinary bladder smooth muscle contractions by ryanodine receptors and BK and SK channels. Am. J. Physiol. Regul. Integr. Comp. Physiol. 279:R60-R68.
- Iino, M., and M. Tsukioka. 1994. Feedback control of inositol trisphosphate signalling bycalcium. Mol. Cell. Endocrinol. 98:141-146. doi:10.1016/0303-7207(94)90132-5
- Jaggar, J.H., and M.T. Nelson. 2000. Differential regulation of Ca(2+) sparks and Ca(2+) waves by UTP in rat cerebral artery smooth muscle cells. Am. J. Physiol. Cell Physiol. 279:C1528-C1539.
- Jaggar, J.H., V.A. Porter, W.J. Lederer, and M.T. Nelson. 2000. Calcium sparks in smooth muscle. Am. J. Physiol. Cell Physiol. 278:C235-C256.
- Jiang, L.H., M. Kim, V. Spelta, X. Bo, A. Surprenant, and R.A. North. 2003. Subunit arrangement in P2X receptors. J. Neurosci. 23:8903-8910.
- Khakh, B.S. 2001. Molecular physiology of P2X receptors and ATP signalling at synapses. Nat. Rev. Neurosci. 2:165-174. doi:10.1038/ 35058521
- Kim, M., G.W. Hennig, T.K. Smith, and B.A. Perrino. 2008. Phospholamban knockout increases CaM kinase II activity and intracellular Ca2+ wave activity and alters contractile responses of murine gastric antrum. Am. J. Physiol. Cell Physiol. 294:C432-C441. doi:10.1152/ajpcell.00418.2007
- Lamont, C., and W.G. Wier. 2002. Evoked and spontaneous purinergic junctional Ca2+ transients (¡CaTs) in rat small arteries. Circ. Res. 91:454-456. doi:10.1161/01.RES.0000035060.98415.4B
- Lamont, C., C. Vial, R.J. Evans, and W.G. Wier. 2006. P2X1 receptors mediate sympathetic postjunctional Ca2+ transients in mesenteric small arteries. Am. J. Physiol. Heart Circ. Physiol. 291:H3106-H3113. doi:10.1152/ajpheart.00466.2006
- Mauban, J.R., C. Lamont, C.W. Balke, and W.G. Wier. 2001. Adrenergic stimulation of rat resistance arteries affects Ca(2+) sparks, Ca(2+) waves, and Ca(2+) oscillations. Am. J. Physiol. Heart Circ. Physiol. 280:H2399-H2405.
- Msghina, M., F. Gonon, and L. Stjärne. 1998. Paired pulse analysis of ATP and noradrenaline release from sympathetic nerves of rat tail artery and mouse vas deferens: effects of K+ channel blockers. Br. J. Pharmacol. 125:1669–1676. doi:10.1038/sj.bjp.0702246
- Mulryan, K., D.P. Gitterman, C.J. Lewis, C. Vial, B.J. Leckie, A.L. Cobb, J.E. Brown, E.C. Conley, G. Buell, C.A. Pritchard, and R.J. Evans. 2000. Reduced vas deferens contraction and male infertility in mice lacking P2X1 receptors. Nature. 403:86-89. doi:10.1038/47495
- Navedo, M.F., M. Nieves-Cintrón, G.C. Amberg, C. Yuan, V.S. Votaw, W.J. Lederer, G.S. McKnight, and L.F. Santana. 2008. AKAP150 is required for stuttering persistent Ca2+ sparklets

- and angiotensin II-induced hypertension. Circ. Res. 102:e1-e11. doi:10.1161/CIRCRESAHA.107.167809
- Navedo, M.F., Y. Takeda, M. Nieves-Cintrón, J.D. Molkentin, and L.F. Santana. 2010. Elevated Ca2+ sparklet activity during acute hyperglycemia and diabetes in cerebral arterial smooth muscle cells. Am. J. Physiol. Cell Physiol. 298:C211-C220. doi:10.1152/ajpcell.00267.2009
- North, R.A. 2002. Molecular physiology of P2X receptors. Physiol. Rev. 82:1013-1067.
- Pérez, G.J., A.D. Bonev, J.B. Patlak, and M.T. Nelson. 1999. Functional coupling of ryanodine receptors to K_{Ca} channels in smooth muscle cells from rat cerebral arteries. J. Gen. Physiol. 113:229–238. doi:10.1085/jgp.113.2.229
- Rummery, N.M., J.A. Brock, P. Pakdeechote, V. Ralevic, and W.R. Dunn. 2007. ATP is the predominant sympathetic neurotransmitter in rat mesenteric arteries at high pressure. J. Physiol. 582:745-754. doi:10.1113/jphysiol.2007.134825
- Santana, L.F., and M.F. Navedo. 2009. Molecular and biophysical mechanisms of Ca2+ sparklets in smooth muscle. J. Mol. Cell. Cardiol. 47:436-444. doi:10.1016/j.yjmcc.2009.07.008
- Schneider, P., H.H. Hopp, and G. Isenberg. 1991. Ca2+ influx through ATP-gated channels increments [Ca2+]i and inactivates ICa in myocytes from guinea-pig urinary bladder. J. Physiol. 440:479-496.
- Stjärne, L. 2001. Novel dual 'small' vesicle model of ATP- and noradrenaline-mediated sympathetic neuromuscular transmission. Auton. Neurosci. 87:16-36. doi:10.1016/S1566-0702(00)00246-0
- Stjärne, L., and E. Stjärne. 1995. Geometry, kinetics and plasticity of release and clearance of ATP and noradrenaline as sympathetic cotransmitters: roles for the neurogenic contraction. Prog. Neurobiol. 47:45-94. doi:10.1016/0301-0082(95)00018-Q
- Valera, S., N. Hussy, R.J. Evans, N. Adami, R.A. North, A. Surprenant, and G. Buell. 1994. A new class of ligand-gated ion channel defined by P2x receptor for extracellular ATP. Nature. 371:516-519. doi:10.1038/371516a0
- Vial, C., and R.J. Evans. 2000. P2X receptor expression in mouse urinary bladder and the requirement of P2X(1) receptors for functional P2X receptor responses in the mouse urinary bladder smooth muscle. Br. J. Pharmacol. 131:1489-1495. doi:10.1038/ sj.bjp.0703720
- Vial, C., and R.J. Evans. 2002. P2X(1) receptor-deficient mice establish the native P2X receptor and a P2Y6-like receptor in arteries. Mol. Pharmacol. 62:1438-1445. doi:10.1124/mol.62.6.1438
- Wray, S., T. Burdyga, and K. Noble. 2005. Calcium signalling in smooth muscle. Cell Calcium. 38:397–407. doi:10.1016/j.ceca.2005.06.018
- Young, J.S., K.L. Brain, and T.C. Cunnane. 2007. The origin of the skewed amplitude distribution of spontaneous excitatory junction potentials in poorly coupled smooth muscle cells. Neuroscience. 145:153-161. doi:10.1016/j.neuroscience.2006.11.054
- Young, J.S., E. Meng, T.C. Cunnane, and K.L. Brain. 2008. Spontaneous purinergic neurotransmission in the mouse urinary bladder. J. Physiol. 586:5743–5755. doi:10.1113/jphysiol.2008.162040
- Zang, W.J., J. Zacharia, C. Lamont, and W.G. Wier. 2006. Sympathetically evoked Ca2+ signaling in arterial smooth muscle. Acta Pharmacol. Sin. 27:1515-1525. doi:10.1111/j.1745-7254.2006.00465.x

154