

No evidence for inositol 1,4,5-trisphosphate–dependent Ca^{2+} release in isolated fibers of adult mouse skeletal muscle

Bert Blaauw,^{1,2} Paola del Piccolo,¹ Laura Rodriguez,¹ Victor-Hugo Hernandez Gonzalez,¹ Lisa Agatea,¹ Francesca Solagna,¹ Fabio Mammano,^{1,3} Tullio Pozzan,^{1,2,4} and Stefano Schiaffino^{1,4}

¹Venetian Institute of Molecular Medicine, 35129 Padova, Italy

²Department of Biomedical Sciences and ³Department of Physics, University of Padova, 35122 Padova, Italy

⁴Institute of Neuroscience, Consiglio Nazionale delle Ricerche, 56124 Pisa, Italy

The presence and role of functional inositol 1,4,5-trisphosphate (IP_3) receptors (IP_3Rs) in adult skeletal muscle are controversial. The current consensus is that, in adult striated muscle, the relative amount of IP_3Rs is too low and the kinetics of Ca^{2+} release from IP_3R is too slow compared with ryanodine receptors to contribute to the Ca^{2+} transient during excitation–contraction coupling. However, it has been suggested that IP_3 -dependent Ca^{2+} release may be involved in signaling cascades leading to regulation of muscle gene expression. We have reinvestigated IP_3 -dependent Ca^{2+} release in isolated flexor digitorum brevis (FDB) muscle fibers from adult mice. Although Ca^{2+} transients were readily induced in cultured C2C12 muscle cells by (a) UTP stimulation, (b) direct injection of IP_3 , or (c) photolysis of membrane-permeant caged IP_3 , no statistically significant change in calcium signal was detected in adult FDB fibers. We conclude that the IP_3 – IP_3R system does not appear to affect global calcium levels in adult mouse skeletal muscle.

INTRODUCTION

Skeletal muscle cells contain a major SR Ca^{2+} release channel, the RyR, which is responsible for excitation–contraction (EC) coupling. Early reports suggested a role of inositol 1,4,5-trisphosphate (IP_3) signaling in EC coupling in skeletal muscle fibers, but this view was challenged by subsequent studies. It is now generally agreed that in both cardiac and skeletal muscle, the relative amount of IP_3 receptors (IP_3Rs) is too low and the kinetics of Ca^{2+} release from IP_3R is too slow compared with RyRs to contribute to the Ca^{2+} transient during EC coupling (see Kocksämper et al., 2008). However, several controversial issues remain unsolved concerning the role of the IP_3 – IP_3R system in skeletal muscle, including (a) the expression level of the IP_3R , (b) whether IP_3R releases a significant amount of Ca^{2+} , and (c) whether IP_3 signaling has a role in the activity-dependent regulation of muscle gene expression, a process referred to as excitation–transcription coupling. In mammalian skeletal muscle, IP_3 was reported to release Ca^{2+} from isolated SR fractions of rabbit fast-twitch skeletal muscle and to elicit isometric force development in chemically skinned muscle fibers (Volpe et al., 1985). In frog muscles, IP_3 was found to be released by electrical stimulation in intact muscle fibers and to induce contractures of

skinned fibers (Vergara et al., 1985). Subsequent studies reported divergent results (see below); however, the reason for these discrepancies remains largely obscure. It is our biased opinion that the different results may depend on the use of different types of muscle fibers, developmental stage, or species. An additional complication in the study of IP_3Rs is represented by the existence of three isoforms, $\text{IP}_3\text{R1}$, $\text{IP}_3\text{R2}$, and $\text{IP}_3\text{R3}$, derived from three distinct genes in mammals (Iwai et al., 2005), showing both specific and redundant roles in organ development and function (Matsumoto et al., 1996; Futatsugi et al., 2005).

Tissue variations in IP_3R distribution are known to be present in cardiac muscle. IP_3Rs are more abundant in atrial than in ventricular cardiomyocytes (Lipp et al., 2000) and even more abundant in conduction tissue cells (Gorza et al., 1993), with $\text{IP}_3\text{R1}$ being the predominant isoform in Purkinje fibers (Gorza et al., 1993) and $\text{IP}_3\text{R2}$ being predominant in sinoatrial node and atrial tissue (Ju et al., 2011). In adult rabbit ventricular myocytes, IP_3Rs were implicated in the regulation of gene expression by a local Ca^{2+} -dependent pathway at the nuclear envelope, based on the finding that the endothelin 1-induced mobilization of Ca^{2+} from the nuclear envelope was blocked by the IP_3R inhibitor 2-aminoethoxydiphenyl borate (2-APB) (Wu et al., 2006).

Correspondence to Stefano Schiaffino: stefano.schiaffino@unipd.it

V.-H.H. Gonzalez's present address is InnerEarLab, Dept. of Otolaryngology, University of Göttingen, 37073 Göttingen, Germany.

Abbreviations used in this paper: 2-APB, 2-aminoethoxydiphenyl borate; EC, excitation–contraction; FDB, flexor digitorum brevis; IP_3 , inositol 1,4,5-trisphosphate; IP_3R , IP_3 receptor; NMJ, neuromuscular junction.

The role of IP₃Rs in skeletal muscle cells is more controversial. Most available data support the existence of a functional IP₃–IP₃R system in cultured skeletal muscle cells, and it has been suggested that IP₃Rs regulate Ca²⁺-dependent gene transcription in these cells (Powell et al., 2001; Stüber et al., 2005). In cultured mouse muscle cells, high potassium-induced depolarization was reported to induce, in addition to the fast Ca²⁺ transients associated with EC coupling, a slower calcium wave, mostly confined to the nuclear and perinuclear regions

of the myotubes, which was inhibited by 2-APB (Powell et al., 2001; Cárdenas et al., 2005). The depolarization-induced phosphorylation of the transcription factor CREB (Powell et al., 2001) and the activation of the early genes *c-fos* and *c-jun* (Carrasco et al., 2003) was also inhibited by 2-APB in skeletal muscle cells. However, another study reported a differential effect of IP₃ signaling according to the degree of muscle cell differentiation, as Ca²⁺ released via IP₃R promoted nuclear entry of the transcription factor NFAT in myoblasts but




Figure 1. UTP addition induces calcium release from cultured muscle cells but not from adult muscle fibers. Here, individual traces represent the $[Ca^{2+}]_i$ responses measured by Fura-2 video microscopy in single cells from the same field in a representative experiment. Similar results were obtained in at least three different experiments. Variations in $[Ca^{2+}]_i$ over time are represented by the ratio between the fluorescence intensities at 340- and 380-nm excitation wavelengths. (A) Changes in the 340/380 ratio of Fura-2 in C2C12 myotubes after UTP stimulation (100 μ M). Top panels show the amount of calcium released at various time points (as indicated by a–d), with pseudocolors indicating the relative intensities. Each trace represents the calcium changes in an individual myotube (four different myotubes are shown, outlined in red, green, blue, and yellow). (B) Preincubation with 70 μ M 2-APB inhibits $[Ca^{2+}]_i$ elicited by 100 μ M UTP in C2C12 myotubes. The addition of ionomycin leads to a significant increase in intracellular calcium. (C) Response of adult fibers to the addition of 100 μ M UTP. Traces of two individual fibers are shown. (D) Quantification of the response of C2C12 myotubes and adult single fibers ($n = 25$ for C2C12 and $n = 10$ for adult fibers). (E) Increase in $[Ca^{2+}]_i$ induced in two adult fibers by 10 mM caffeine. (F) The effect of caffeine is significantly reduced by the addition of 20 μ M dantrolene, an inhibitor of RyR.

nuclear exit of NFAT in myotubes (Stiber et al., 2005). Interpretation of these studies is complicated by the use of inhibitors, like 2-APB and heparin, which are not completely specific. For example, 2-APB is a blocker of store-operated Ca^{2+} entry (Bootman et al., 2002).

The role and the very existence of functional IP_3Rs in adult mammalian skeletal muscle are even less clear. Immunodetectable IP_3R proteins and specific $[^3\text{H}]\text{IP}_3$ -binding sites were reported to be preferentially expressed in slow oxidative (type I) and fast oxidative glycolytic (type IIA) fibers, but not in fast glycolytic (type IIB) fibers in rat muscles (Moschella et al., 1995). Accordingly, the application of IP_3 to rat slow- and fast-twitch saponin-skinned fibers induced contractile responses

that were larger in slow compared with fast muscles but decreased during postnatal development (Talon et al., 1999, 2002). On the contrary, muscle fibers isolated from the fast flexor digitorum brevis (FDB) but not from slow soleus muscle of young adult (5–7-wk-old) mice were found to respond to electrical stimulation with a fast calcium signal associated with muscle contraction, followed by a slower signal, which was inhibited by the IP_3R inhibitors xestospongin B or C (Casas et al., 2010). Other immunofluorescence studies reported the presence of IP_3Rs in both mouse and rabbit skeletal muscle (Salanova et al., 2002), and a recent study reported that $\text{IP}_3\text{R}2$ and $\text{IP}_3\text{R}3$ are expressed in all muscle fiber types in adult mouse skeletal muscle, whereas




Figure 2. Injection of IP_3 does not lead to an increase in intracellular calcium in adult skeletal muscle fibers. (A) Microinjection of 50 μM IP_3 mediated by patch clamp triggers a rapid Ca^{2+} response in a C2C12 myotube. Note that an increase in $[\text{Ca}^{2+}]_i$ is seen in the myotube injected with IP_3 (outlined in red) but not in noninjected neighboring myotube (outlined in green). (B) Microinjection of intracellular solution alone does not elicit any $[\text{Ca}^{2+}]_i$ variation in C2C12 myotubes. (C) Microinjection of IP_3 in adult single fibers does not induce any changes in $[\text{Ca}^{2+}]_i$. As a positive control for injection, the membrane-impermeable dye Alexa red was injected simultaneously. Microinjection of ionomycin leads to a marked increase in the Fura-2 signal. Similar results were obtained in three different experiments; a total of 10 fibers were examined. (D) Microinjection of the fluorescent dye calcein was used to monitor the velocity of diffusion in adult single fibers. Four regions of interest are shown in a single fiber (yellow, red, green, and blue squares), with their respective increases in fluorescence in time shown in the graph (color of the line corresponds to the color of the square). The top panels are taken at two time points (indicated by a and b). Note that in all these experiments (A–D), the resting ratios differ from those in the experiments of Fig. 1, because the light source and the permissivity of the filters for both Fura-2 wavelengths were not the same in the two setups.

IP₃R1 was only found in fast type IIX fibers (Casas et al., 2010). IP₃Rs appear to be especially abundant at the neuromuscular junctions (NMJs) (Powell et al., 2003), and recent studies showed that IP₃R1 is detectable by immunofluorescence at mouse NMJs, but apparently not in extrajunctional regions, and is involved in synaptic gene expression, as determined by RNAi-mediated IP₃R1 gene silencing (Zayas et al., 2007; Zhu et al., 2011).

Here, we have reexamined the existence of functional IP₃R signaling triggered by IP₃ in isolated FDB muscle fibers from adult mice by monitoring IP₃-dependent Ca²⁺ release induced by (a) UTP stimulation, a classical way to activate IP₃Rs; (b) direct injection of IP₃; or (c) photolysis of membrane-permeant caged IP₃. As a control, we determined the effect of UTP, IP₃ injection, and activation of caged IP₃ in cultured C2C12 myotubes, which were shown to possess a functional IP₃–IP₃R system. The results of these experiments indicate that, although a functional IP₃–IP₃R system is readily detectable in cultured muscle cells, no significant change in calcium signal was detected in adult muscle fibers after IP₃ microinjection or photolysis of membrane-permeant IP₃. This is to our knowledge the first work in which these direct approaches have been used to address the adult muscle IP₃ controversy.

MATERIALS AND METHODS

Muscle cells and UTP stimulation

Cultured C2C12 myotubes were prepared as described previously (Di Lisi et al., 2007). FDB muscle fibers were isolated from 4-mo-old CD1 mice and 5-wk-old BALB/c mice, and cultured as described previously (Zhao et al., 2007). UTP, ionomycin, caffeine, and dantrolene were from Sigma-Aldrich.

IP₃ injection

Muscle cells loaded with Fura-2 were microinjected by passive dialysis using borosilicate patch pipettes filled with 50 μM IP₃ dissolved in an intracellular solution containing (in mM): 145 potassium aspartate, 3.5 NaCl, 1 MgCl₂, 6.5 NaOH, 10 HEPES, and 5 MgATP, pH 7.2. Pipette resistances were 3–5 MΩ when immersed in the bath. Recordings were made under whole cell voltage-clamp conditions at room temperature using a patch-clamp amplifier (EPC-7; List). The holding potential was set at –60 mV for C2C12 myotubes (see Fioretti et al., 2005) and at –90 mV for adult muscle fibers. The fluorescent dye Alexa Fluor 568 (1 mM) was added to the patch pipette solution to monitor microinjection in adult muscle fibers.

Caged IP₃ photolysis

C2C12 myotubes and myofibers were incubated for 30 min at 37° in DMEM F-12 supplemented with FluoForte AM (16 μM DMSO; Enzo Life Sciences), 5 μM of caged IP₃ AM (Enzo Life Sciences), pluronic F-127 (0.01% wt/vol; Sigma-Aldrich), and 250 μM sulphinpyrazone (Sigma-Aldrich). For photostimulation with caged IP₃, the output of a TTL-controlled semiconductor-laser module (20 mW, 379 nm) was injected into a UV-permissive fiber-optic cable (multimode step index 0.22 N.A., 105 μm core; part no. AFS105/125YCUSTOM; Thorlabs, Inc.). Fiber output was projected onto the specimen plane by an aspheric condenser lens

(20-mm effective focal length; part no. ACL2520; Thorlabs, Inc.), and the recollimated beam was directed onto a dichromatic mirror (400 dclp; Chroma Technology Corp.) placed at 45° just above the objective lens of the microscope. For recording, muscle cells were transferred to the stage of an upright fluorescence microscope (BX51; Olympus) and perfused in EXM (an extracellular medium containing 138 mM NaCl, 5 mM KCl, 200 μM CaCl₂, 0.3 mM NaH₂PO₄, 0.4 mM KH₂PO₄, 10 mM HEPES-NaOH, and 6 mM D-glucose, pH 7.2, 320 mOsm) for 10 min at 2 ml/min to allow for de-esterification. FluoForte fluorescence was excited at 460 nm by an LED and directed onto the sample through a 505-dcxr dichromatic mirror (Chroma Technology Corp.). Fluorescence emission was selected by an HQ520/40-M filter (Chroma Technology Corp.), centered around a 520-nm wavelength, to form fluorescence images on a scientific-grade CCD camera (PCO AG; SensiCam) using a 20× water-immersion objective (NA 0.95; LumPlan FL; Olympus). For uncaging experiments, baseline (pre-stimulus) fluorescence emission (*f*₀) was recorded for 20 s; thereafter, a UV laser pulse of 300 ms was applied to release IP₃, and fluorescence emission was monitored for up to 3 min. Because intense UV illumination can damage cells and trigger intercellular Ca²⁺ signals also in the absence of caged IP₃, we performed a series of preliminary control experiments to find the minimal UV dose sufficient for uncaging IP₃ and determined that, under our experimental conditions, it corresponded to the above-mentioned laser-pulse duration of 300 ms. To establish a safety margin, we additionally verified that no Ca²⁺ signal was evoked by UV pulses of the same intensity and duration up to 2 s if caged IP₃ was omitted from the loading solution. Image sequences were acquired using software developed in the laboratory, stored on disk and processed offline using the Matlab 7.0 software package (The MathWorks, Inc.). In particular, signals were measured as relative changes of fluorescence emission intensity, ($\Delta f/f_0$), where *f* is fluorescence at post-stimulus time *t*, and $\Delta f = f - f_0$. All live cell-imaging experiments were performed at room temperature (24–26°C).

RESULTS AND DISCUSSION

Purinergic receptor agonists are known to induce G protein-coupled generation of IP₃ and IP₃R-dependent mobilization of intracellular Ca²⁺ stores in different cell types (Ralevic and Burnstock, 1998). Thus, we first examined the effect of UTP, which was shown to induce IP₃-dependent Ca²⁺ release in cultured muscle cells (Stüber et al., 2005). As expected, in cultured C2C12 myotubes, UTP induced Ca²⁺ transients (Fig. 1 A) that were blocked by 2-APB (Fig. 1 B). In contrast, only a minimal Ca²⁺ transient was detected in adult muscle fibers after the addition of UTP (Fig. 1, C and D). On the other hand, adult muscle fibers showed clear Ca²⁺ transients in response to caffeine (3.41 ± 0.56 peak Δ ratio; *n* = 8), which was markedly reduced by the RyR channel blocker dantrolene (1.76 ± 0.14 peak Δ ratio; *n* = 6) (Fig. 1, E and F). The finding that adult muscle fibers are unresponsive to UTP may be because of the down-regulation of purinergic receptors observed during rat (Cheung et al., 2003) and chick (Wells et al., 1995) embryonic development, and is consistent with the negligible Ca²⁺ response to ATP of human mature myofibers compared with young myotubes (Cseri et al., 2002). However, extracellular ATP was considered to be an important mediator for

calcium transients evoked by electrical stimulation in adult skeletal fibers, based on the reduced slow Ca^{2+} signal induced by apyrase, an enzyme that metabolizes extracellular ATP to AMP (Buvinic et al., 2009).

To determine directly whether adult muscle fibers respond to IP_3 , we examined the effect of IP_3 injected into cultured muscle cells and adult fibers loaded with Fura-2. As shown in Fig. 2 A, Ca^{2+} transients from Fura-2-loaded C2C12 myotubes were readily detected after injection of IP_3 (1.52 ± 0.16 peak Δ ratio; $n = 8$). Injection of intracellular solution alone did not elicit any variation in intracellular Ca^{2+} in C2C12 myotubes (Fig. 2 B). In contrast, no Ca^{2+} signal was detectable after IP_3 injection in adult muscle fibers (Fig. 2 C). Successful microinjection was documented by coinjection of Alexa red, and the final addition of $1 \mu\text{M}$ ionomycin to the medium led to a rapid increase in Ca^{2+} concentration. To test whether the different response of adult muscle fibers compared with cultured muscle cells may be caused by a slower diffusion, we monitored the velocity of diffusion of intracellular tracers after microinjection in single muscle fibers. The fluorescent dye calcein, which has a size comparable to IP_3 , was injected in adult FDB fibers (Fig. 2 D). The intensity of the dye fluorescence emission in various regions of interest indicates that the diffusion velocity is sufficient for IP_3 to diffuse throughout the fiber in the time frame considered. However, interpretation of these results is complicated by the possibility that microinjected IP_3 is rapidly hydrolyzed by IP_3 -5-phosphatase, the first enzyme responsible for its degradation, and thus may not reach its targets in sufficient concentrations to activate the IP_3 Rs. To rule out this possibility, we incubated cultured muscle cells and adult muscle fibers with a membrane-permeant caged IP_3 , leaving enough time so that the caged compound can diffuse throughout the muscle fibers before flash photolysis. In previous studies using skinned fibers from frog semitendinosus muscle, it was shown that caged IP_3 readily diffuses in adult muscle fibers (Walker et al., 1987). Flash photolysis of caged IP_3 in C2C12 myoblasts coloaded with Fluo-4 leads to a rapid increase in intracellular calcium (Fig. 3, A and C). In contrast, adult muscle fibers showed no statistically significant response to IP_3 uncaging (Fig. 3, B and D), although a minimal increase (2–3%) was observed in 11 out of 35 fibers. This result is in contrast with a recent report of IP_3 -mediated Ca^{2+} release in mouse FDB fibers from 5-wk-old BALB/c mice, as determined by the response to the IP_3 R inhibitor, xestospongin (Casas et al., 2010). To determine whether the discrepancy is caused by the younger age of the mice used in that study, we repeated the experiments using photolysis of membrane-permeant caged IP_3 in single fibers of 5-wk-old BALB/c mice. As shown in Fig. 3 D, these fibers also showed no significant global calcium release after photolysis. Only 4 out of 24 fibers showed a minimal increase (2–3%) in fluorescence signal.

The different conclusion between our study and that of Casas et al. (2010) may be because their conclusion is based on a pharmacological inhibitor, xestospongin C,




Figure 3. Uncaging of IP_3 does not elicit appreciable calcium release in adult single fibers. (A) Pseudocolor images of Fluo-Forte Ca^{2+} -dependent fluorescence from C2C12 myoblasts at various time points in response to photorelease of caged IP_3 . The shown images were obtained at 5.7 and 12.9 s after photo liberating IP_3 from its caged precursor. Traces show the time course of the relative percentage change in fluorescence ($\Delta f/f_0$) caused by intracellular Ca^{2+} increase for four different regions of interest (identified by the colored boxes). (B) Changes in $[\text{Ca}^{2+}]_i$ after photorelease of caged IP_3 in four FDB fibers. A minimal signal was detected in 11 out of 35 fibers. (C) Quantification of the peak relative increase of $[\text{Ca}^{2+}]_i$ after photorelease, as measured by the change in FluoForte intensity, in C2C12 muscle cells ($n = 85$ cells). (D) Same as C for adult muscle fibers isolated from 4-mo-old CD1 mice (left; $n = 35$) and 5-wk-old BALB/c mice (right; $n = 24$). The difference between fibers incubated with caged IP_3 and control fibers is not statistically significant.

whose specificity is not absolute, whereas we examine the direct effect of IP₃.

Collectively, the results presented here indicate that IP₃-mediated Ca²⁺ release, which is clearly present and functional in cultured muscle cells, is unlikely to play a significant functional role in adult muscle fibers from the mouse FDB muscle. The existence of major developmental changes in the mechanisms that regulate Ca²⁺ release from intracellular Ca²⁺ stores of skeletal muscle fibers should be considered when inferring a physiological role of IP₃ in muscle gene regulation. Our findings, although not supporting the notion of global IP₃-dependent Ca²⁺ release in adult mammalian skeletal muscle fibers, do not rule out the possibility that IP₃Rs may be present and functional in specific microdomains, such as the NMJ. A role of IP₃R signaling at the NMJ was suggested by the finding that several minutes of repeated acetylcholine applications directly to the NMJs of isolated adult mouse muscle fibers caused a localized subsynaptic Ca²⁺ increase, which was blocked by xestospongin C and by the phospholipase C blocker, U-73122 (Zayas et al., 2007). Ca²⁺ was found to accumulate much faster around the NMJs of muscle fibers from transgenic mice with specific mutations in acetylcholine receptor, leading to postsynaptic Ca²⁺ accumulation, a model of the slow-channel congenital myasthenic syndrome (Zayas et al., 2007). In adult mouse skeletal muscles, IP₃R1 was specifically localized in the subsynaptic region of NMJs. In addition, IP₃R1 knockdown by siRNA electroporation was found to induce up-regulation of several synaptic genes, including AChR subunits, and HDAC4 up-regulation followed by nuclear accumulation of HDAC4 in junctional nuclei (Zhu et al., 2011). These changes are in part similar to those induced by denervation; however, in contrast to denervation, IP₃R1 silencing had no effect on extrajunctional muscle genes and extrajunctional HDAC4 nuclear translocation, supporting the notion of a selective role of muscle IP₃R1 in compartmentalized Ca²⁺ signals at the NMJ (Zhu et al., 2011).

This work is supported in part by grants from the European Commission (FP7 Integrated Project MYOAGE to S. Schiaffino), the Italian Space Agency (project OSMA to S. Schiaffino), MIUR PRIN (grant 2009CCZSES to T. Pozzan and F. Mammano), and Progetto di Ateneo (to B. Blaauw).

Richard L. Moss served as editor.

Submitted: 28 November 2011

Accepted: 18 June 2012

REFERENCES

Bootman, M.D., T.J. Collins, L. Mackenzie, H.L. Roderick, M.J. Berridge, and C.M. Peppiatt. 2002. 2-aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated Ca²⁺ entry but an inconsistent inhibitor of InsP₃-induced Ca²⁺ release. *FASEB J.* 16:1145–1150. <http://dx.doi.org/10.1096/fj.02-0037rev>

Buvinic, S., G. Almaraz, M. Bustamante, M. Casas, J. López, M. Riquelme, J.C. Sáez, J.P. Huidobro-Toro, and E. Jaimovich. 2009.

ATP released by electrical stimuli elicits calcium transients and gene expression in skeletal muscle. *J. Biol. Chem.* 284:34490–34505. <http://dx.doi.org/10.1074/jbc.M109.057315>

Cárdenas, C., J.L. Liberona, J. Molgó, C. Colasante, G.A. Mignery, and E. Jaimovich. 2005. Nuclear inositol 1,4,5-trisphosphate receptors regulate local Ca²⁺ transients and modulate cAMP response element binding protein phosphorylation. *J. Cell Sci.* 118:3131–3140. <http://dx.doi.org/10.1242/jcs.02446>

Carrasco, M.A., N. Riveros, J. Ríos, M. Müller, F. Torres, J. Pineda, S. Lantadilla, and E. Jaimovich. 2003. Depolarization-induced slow calcium transients activate early genes in skeletal muscle cells. *Am. J. Physiol. Cell Physiol.* 284:C1438–C1447.

Casas, M., R. Figueroa, G. Jorquera, M. Escobar, J. Molgó, and E. Jaimovich. 2010. IP₃-dependent, post-tetanic calcium transients induced by electrostimulation of adult skeletal muscle fibers. *J. Gen. Physiol.* 136:455–467. <http://dx.doi.org/10.1085/jgp.200910397>

Cheung, K.K., M. Ryten, and G. Burnstock. 2003. Abundant and dynamic expression of G protein-coupled P2Y receptors in mammalian development. *Dev. Dyn.* 228:254–266. <http://dx.doi.org/10.1002/dvdy.10378>

Cseri, J., H. Szappanos, G.P. Szigeti, Z. Csernátó, L. Kovács, and L. Csernoch. 2002. A purinergic signal transduction pathway in mammalian skeletal muscle cells in culture. *Pflugers Arch.* 443:731–738. <http://dx.doi.org/10.1007/s00424-001-0757-x>

Di Lisi, R., A. Picard, S. Ausoni, and S. Schiaffino. 2007. GATA elements control repression of cardiac troponin I promoter activity in skeletal muscle cells. *BMC Mol. Biol.* 8:78. <http://dx.doi.org/10.1186/1471-2199-8-78>

Fioretti, B., T. Pietrangelo, L. Catacuzzeno, and F. Franciolini. 2005. Intermediate-conductance Ca²⁺-activated K⁺ channel is expressed in C2C12 myoblasts and is downregulated during myogenesis. *Am. J. Physiol. Cell Physiol.* 289:C89–C96. <http://dx.doi.org/10.1152/ajpcell.00369.2004>

Futatsugi, A., T. Nakamura, M.K. Yamada, E. Ebisui, K. Nakamura, K. Uchida, T. Kitaguchi, H. Takahashi-Iwanaga, T. Noda, J. Aruga, and K. Mikoshiba. 2005. IP₃ receptor types 2 and 3 mediate exocrine secretion underlying energy metabolism. *Science*. 309:2232–2234. <http://dx.doi.org/10.1126/science.1114110>

Gorza, L., S. Schiaffino, and P. Volpe. 1993. Inositol 1,4,5-trisphosphate receptor in heart: evidence for its concentration in Purkinje myocytes of the conduction system. *J. Cell Biol.* 121:345–353. <http://dx.doi.org/10.1083/jcb.121.2.345>

Iwai, M., Y. Tateishi, M. Hattori, A. Mizutani, T. Nakamura, A. Futatsugi, T. Inoue, T. Furuichi, T. Michikawa, and K. Mikoshiba. 2005. Molecular cloning of mouse type 2 and type 3 inositol 1,4,5-trisphosphate receptors and identification of a novel type 2 receptor splice variant. *J. Biol. Chem.* 280:10305–10317. <http://dx.doi.org/10.1074/jbc.M413824200>

Ju, Y.K., J. Liu, B.H. Lee, D. Lai, E.A. Woodcock, M. Lei, M.B. Cannell, and D.G. Allen. 2011. Distribution and functional role of inositol 1,4,5-trisphosphate receptors in mouse sinoatrial node. *Circ. Res.* 109:848–857. <http://dx.doi.org/10.1161/CIRCRESAHA.111.243824>

Kocksämper, J., A.V. Zima, H.L. Roderick, B. Pieske, L.A. Blatter, and M.D. Bootman. 2008. Emerging roles of inositol 1,4,5-trisphosphate signaling in cardiac myocytes. *J. Mol. Cell. Cardiol.* 45:128–147. <http://dx.doi.org/10.1016/j.jmcc.2008.05.014>

Lipp, P., M. Laine, S.C. Tovey, K.M. Burrell, M.J. Berridge, W. Li, and M.D. Bootman. 2000. Functional InsP₃ receptors that may modulate excitation-contraction coupling in the heart. *Curr. Biol.* 10:939–942. [http://dx.doi.org/10.1016/S0960-9822\(00\)00624-2](http://dx.doi.org/10.1016/S0960-9822(00)00624-2)

Matsumoto, M., T. Nakagawa, T. Inoue, E. Nagata, K. Tanaka, H. Takano, O. Minowa, J. Kuno, S. Sakakibara, M. Yamada, et al. 1996. Ataxia and epileptic seizures in mice lacking type 1 inositol 1,4,5-trisphosphate receptor. *Nature*. 379:168–171. <http://dx.doi.org/10.1038/379168a0>

- Moschella, M.C., J. Watras, T. Jayaraman, and A.R. Marks. 1995. Inositol 1,4,5-trisphosphate receptor in skeletal muscle: differential expression in myofibres. *J. Muscle Res. Cell Motil.* 16:390–400. <http://dx.doi.org/10.1007/BF00114504>
- Powell, J.A., M.A. Carrasco, D.S. Adams, B. Drouet, J. Rios, M. Müller, M. Estrada, and E. Jaimovich. 2001. IP(3) receptor function and localization in myotubes: an unexplored Ca(2+) signaling pathway in skeletal muscle. *J. Cell Sci.* 114:3673–3683.
- Powell, J.A., J. Molgó, D.S. Adams, C. Colasante, A. Williams, M. Bohlen, and E. Jaimovich. 2003. IP3 receptors and associated Ca2+ signals localize to satellite cells and to components of the neuromuscular junction in skeletal muscle. *J. Neurosci.* 23:8185–8192.
- Ralevic, V., and G. Burnstock. 1998. Receptors for purines and pyrimidines. *Pharmacol. Rev.* 50:413–492.
- Salanova, M., G. Priori, V. Barone, E. Intravaia, B. Flucher, F. Ciruela, R.A. McIlhinney, J.B. Parys, K. Mikoshiba, and V. Sorrentino. 2002. Homer proteins and InsP(3) receptors co-localise in the longitudinal sarcoplasmic reticulum of skeletal muscle fibres. *Cell Calcium.* 32:193–200. <http://dx.doi.org/10.1016/S0143416002001549>
- Stüber, J.A., N. Tabatabaei, A.F. Hawkins, T. Hawke, P.F. Worley, R.S. Williams, and P. Rosenberg. 2005. Homer modulates NFAT-dependent signaling during muscle differentiation. *Dev. Biol.* 287:213–224. <http://dx.doi.org/10.1016/j.ydbio.2005.06.030>
- Talon, S., C. Huchet-Cadiou, and C. Léoty. 1999. Inositol 1,4,5-trisphosphate-sensitive Ca2+ release in rat fast- and slow-twitch skinned muscle fibres. *Pflugers Arch.* 438:804–816. <http://dx.doi.org/10.1007/s004240051110>
- Talon, S., O. Vallot, C. Huchet-Cadiou, A.M. Lompré, and C. Léoty. 2002. IP(3)-induced tension and IP(3)-receptor expression in rat soleus muscle during postnatal development. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 282:R1164–R1173.
- Vergara, J., R.Y. Tsien, and M. Delay. 1985. Inositol 1,4,5-trisphosphate: a possible chemical link in excitation-contraction coupling in muscle. *Proc. Natl. Acad. Sci. USA.* 82:6352–6356. <http://dx.doi.org/10.1073/pnas.82.18.6352>
- Volpe, P., G. Salviati, F. Di Virgilio, and T. Pozzan. 1985. Inositol 1,4,5-trisphosphate induces calcium release from sarcoplasmic reticulum of skeletal muscle. *Nature.* 316:347–349. <http://dx.doi.org/10.1038/316347a0>
- Walker, J.W., A.V. Somlyo, Y.E. Goldman, A.P. Somlyo, and D.R. Trentham. 1987. Kinetics of smooth and skeletal muscle activation by laser pulse photolysis of caged inositol 1,4,5-trisphosphate. *Nature.* 327:249–252. <http://dx.doi.org/10.1038/327249a0>
- Wells, D.G., M.J. Zawisa, and R.I. Hume. 1995. Changes in responsiveness to extracellular ATP in chick skeletal muscle during development and upon denervation. *Dev. Biol.* 172:585–590. <http://dx.doi.org/10.1006/dbio.1995.8062>
- Wu, X., T. Zhang, J. Bossuyt, X. Li, T.A. McKinsey, J.R. Dedman, E.N. Olson, J. Chen, J.H. Brown, and D.M. Bers. 2006. Local InsP3-dependent perinuclear Ca2+ signaling in cardiac myocyte excitation-transcription coupling. *J. Clin. Invest.* 116:675–682. <http://dx.doi.org/10.1172/JCI27374>
- Zayas, R., J.S. Groshong, and C.M. Gomez. 2007. Inositol-1,4,5-trisphosphate receptors mediate activity-induced synaptic Ca2+ signals in muscle fibers and Ca2+ overload in slow-channel syndrome. *Cell Calcium.* 41:343–352. <http://dx.doi.org/10.1016/j.ceca.2006.07.007>
- Zhao, J., J.J. Brault, A. Schild, P. Cao, M. Sandri, S. Schiaffino, S.H. Lecker, and A.L. Goldberg. 2007. FoxO3 coordinately activates protein degradation by the autophagic/lysosomal and proteasomal pathways in atrophying muscle cells. *Cell Metab.* 6:472–483. <http://dx.doi.org/10.1016/j.cmet.2007.11.004>
- Zhu, H., B.J. Bhattacharyya, H. Lin, and C.M. Gomez. 2011. Skeletal muscle IP3R1 receptors amplify physiological and pathological synaptic calcium signals. *J. Neurosci.* 31:15269–15283. <http://dx.doi.org/10.1523/JNEUROSCI.3766-11.2011>