Skeletal muscle fibers: Inactivated or depleted after long depolarizations?

Werner Melzer

Institute of Applied Physiology, Ulm University, D-89081 Ulm, Germany

Skeletal muscle consists of large multinucleated fibers that rapidly contract in response to action potentials generated in their surface membrane. The fiber's contractile apparatus is activated by Ca²⁺ ions that are released into the cytoplasm from an extensive intracellular storage compartment, the sarcoplasmic reticulum (SR), in response to the action potential. Ca²⁺ concentration changes in the cytoplasm during the excitation—contraction (EC) coupling process have been recorded for several decades, but only in recent years have methods been developed to monitor Ca²⁺ in the SR. In this issue of the Journal, Robin and Allard make use of this approach to study the functional basis of changes in the filling state of the SR during long-lasting membrane depolarizations.

Voltage-controlled Ca²⁺ release in muscle fibers

The Journal of General Physiology

The efflux of Ca²⁺ from the SR is mediated by ryanodine receptors (RyRs), members of a family of giant intracellular channel proteins. The subtype 1 (RyR1) forms regular arrays in specialized zones of the SR membrane that communicate with the transverse tubules (TTs), narrow invaginations of the surface membrane, which conduct the surface action potential into the center of the fiber (Manno et al., 2013). The TT-SR junction is the basis of a unique voltage control mechanism for the rapid opening and closing of the RyR1 in response to action potential depolarization and repolarization (Melzer et al., 1995; Dulhunty, 2006; Baylor and Hollingworth, 2012). Whereas "standard" voltage-activated channels contain both of their essential functional elements, i.e., a voltage sensor and an ion-conducting pore, within a single protein, the RyR1 protein has no comparable voltage-sensitive region. The SR membrane does not seem to develop a substantial voltage, either at rest or during Ca²⁺ release (Fink and Veigel, 1996). Instead, it relies on the voltage of the TT membrane and the voltage sensitivity of a separate protein located in this membrane, the dihydropyridine receptor (DHPR; an L-type Ca²⁺ channel, CaV1.1), which is mechanically connected to the RyR1. The DHPR itself is not a very effective Ca²⁺ channel. Its pore-opening reaction is much slower than the coupling reaction that bridges the junctional gap

and gates the RyR1. Moreover, opening the DHPR pore requires stronger depolarization than does the coupling reaction, and the Ca²⁺ inward current it passes is far smaller than the Ca²⁺ release flux from the SR (Ursu et al., 2005). In fact, in some fish (higher teleosts), the DHPR does not pass Ca²⁺ at all (Schredelseker et al., 2010).

The TT voltage controls the gate of the RyR1 but being generated in a separate membrane—has no direct influence on the driving force for the efflux of Ca²⁺ from the SR, which is provided by the steep transmembrane Ca²⁺ concentration gradient between the SR lumen and the cytoplasmic space. The resting gradient is established by the constant action of adenosine triphosphate (ATP)-driven Ca²⁺ pumps (SERCA) and supported by Ca²⁺ buffers in the SR lumen, of which calsequestrin is the most important (Murphy et al., 2009). Calsequestrin's high capacity (up to 80 Ca²⁺ per molecule) enables storage of large amounts of Ca2+ in the SR and further stabilizes the concentration gradient during Ca²⁺ release. The affinity of calsequestrin for Ca²⁺ appears to decrease when the free Ca²⁺ concentration in the SR starts to fall, and it may even act as a funnel that guides Ca²⁺ ions to the releasing sites (Royer and Ríos, 2009). Moreover, calsequestrin probably also acts as a Ca²⁺ sensor that modulates RyR permeability depending on the Ca²⁺loading state of the SR (Beard et al., 2009).

Voltage-dependent inactivation of EC coupling

Loss of the normal resting membrane potential may occur in extensively stressed muscle fibers. If not counteracted by protective mechanisms, this condition would lead to maintained activation of Ca²⁺ entry from the extracellular space and Ca²⁺ release from the SR, and to dramatically increased ATP consumption. Long-lasting depolarization leads to the spontaneous relaxation of force, a phenomenon first analyzed by Hodgkin and Horowicz (1960) in frog muscle using potassium-induced contractures. They suggested that depolarization "liberates an activator which is used up in generating tension." Later, this spontaneous relaxation process was linked to the immobilization of a transverse tubular charge

Correspondence to Werner Melzer: werner.melzer@uni-ulm.de

movement (Chandler et al., 1976) and attributed to an inactivation process. During strong depolarization the transition from the activatable to the inactivated state takes seconds to develop, and it takes minutes to fully recover from inactivation after repolarization. This inactivation is enhanced by L-type Ca²⁺ channel antagonists (Berwe et al., 1987; Feldmeyer et al., 1990). Indeed, like other voltage-dependent channels, the DHPR enters an inactivated state during long-lasting membrane depolarization (Cota et al., 1984). Inactivation can be studied by recording the L-type Ca²⁺ current or the gating charge movement produced by the DHPR. Like its voltage-dependent activation, inactivation of the DHPR is communicated to the RyR1 and leads to its closure.

Availability curves, i.e., the graded sigmoidal dependence of fractional force or Ca²⁺ release activation on steady-state voltage, belong to the standard repertoire in the characterization of the EC coupling mechanism. It is generally assumed that these curves demonstrate the voltage dependence of the inactivation mechanism of the voltage sensor. Yet, in experiments in which L-type Ca²⁺ current and Ca²⁺ release are measured in parallel, Ca²⁺ release availability reaches its half-maximal value at more negative holding potentials compared with the availability of the Ca²⁺ inward current produced by the DHPR (Andronache et al., 2009). One explanation would be that the voltage control of RyR1 inactivation and the inactivation of the L-type channel permeation pathway are mediated by different states of the DHPR. A comparable situation is found in the activation process in which different states of the DHPR, attained at different voltages, appear to control Ca²⁺ release and Ca²⁺ entry, respectively (García et al., 1994; Dietze et al., 2000). In this issue, Robin and Allard (2013) challenge the view that DHPR inactivation is the dominant cause of the decline in Ca²⁺ release during long-lasting depolarization. Instead, they suggest that Ca²⁺ depletion in the SR lumen shapes the "steady-state availability curve" of Ca²⁺ release.

Monitoring Ca²⁺ in the storage compartment

It is obvious that the large voltage-activated flux of Ca²⁺ from the SR may also appreciably alter its own driving force. However, separating RyR gating from SR depletion is difficult. Until recently, indirect methods were used to estimate the Ca²⁺ content of the SR and its changes during various Ca²⁺ release paradigms (Pape et al., 1995; Posterino and Lamb, 2003). Within the last couple of years, progress has been made in imaging intra-SR calcium using fluorescent probes (Rudolf et al., 2006; Ziman et al., 2010; Sztretye et al., 2011; Robin and Allard, 2012). Two different methods have been established. One uses the permeable acetoxymethyl ester form of a synthetic fluorimetric dye loaded into the SR lumen. The other uses protein-based Ca²⁺ sensors that can specifically be targeted to the SR. The indicators should be of low affinity to avoid saturation by the high resting free Ca²⁺

concentration in the SR (\sim 0.5 mM). The advantages and disadvantages of the two methods have recently been summarized by Manno et al. (2013). Robin and Allard (2013) used the first method in their experiments and the indicator Fluo-5N (Kabbara and Allen, 2001; Ziman et al., 2010) to record intra-SR Ca²⁺ concentration changes. They combined this with an electrophysiological technique termed "silicon clamp" (Pouvreau et al., 2007), which allows single electrode voltage clamping of large muscle cells using conventional whole cell patch clamp circuitry. The notorious problems of space-clamping muscle fibers are avoided by electrically isolating a small region of the cell by covering the rest of it with a silicon grease. The micropipette used for voltage clamping dialyses the intracellular space with an artificial solution to dilute the indicator present in the cytoplasm. In this study, the pipette solution also contained a very high concentration of the Ca²⁺ chelator EGTA (50 mM). This ensures that any residual cytoplasmic Fluo-5N produces negligible fluorescent Ca2+ transients during depolarization and, therefore, permits the specific recording of free Ca²⁺ concentration changes within the SR. Conditions remain remarkably stable for considerable periods of time in this configuration and thus permit the repeated fluorimetric recording of Ca²⁺ concentration changes within the SR during the long-lasting prepulse protocols that are necessary to determine the steadystate voltage dependence of Ca²⁺ release availability. Under comparable conditions, Robin and Allard (2013) also recorded cytoplasmic Ca²⁺ transients using Fura-2. Inspecting the decay of a cytoplasmic Ca²⁺ transient that occurs during maintained depolarization does not allow one to determine whether it results from inactivation of the release pathway or SR Ca²⁺ depletion. When looking at the Ca²⁺ level within the SR, however, the distinction is possible: depletion will cause a decrease, whereas inactivation will lead to the opposite. Exploiting these properties, the study came to the interesting conclusion that the slow decline in Ca2+ transients recorded with cytoplasmic indicators during long-lasting depolarization is largely caused by depletion. Inactivation gets started only at lower (more depolarized) membrane potential and follows with a delay.

These findings are compatible with the discrepancy between steady-state availability curves of Ca^{2+} release and L-type Ca^{2+} current we have observed in our own experiments (Ursu et al., 2004; Andronache et al., 2009). We noticed a markedly elevated free cytoplasmic Ca^{2+} concentration in a voltage range whose center was $\sim\!20$ mV more positive than the threshold for release activation. We viewed this steady increase in basal Ca^{2+} concentration as resulting from a window Ca^{2+} release (Ursu et al., 2004; Andronache et al., 2009) in analogy to the window current known from "conventional" voltage-dependent channels. Window currents arise from the overlap between activation and availability curves at voltages where

there is some degree of activation but incomplete steadystate inactivation. During long depolarizations, even a small fractional activation of the Ca²⁺ release mechanism may shift a substantial amount of Ca2+ from the SR lumen to the cytoplasmic space. When we depolarized the muscle fiber membrane beyond a certain value, the elevated cytoplasmic basal Ca2+ concentration declined to normal values. The fluorescence signal from the intra-SR Ca²⁺ sensor provides a direct demonstration of this effect in Fig. 3 A of Robin and Allard (2013) and in a previous paper by the Allard laboratory (Robin et al., 2012): When the voltage step exceeds a certain depolarization level and duration, the fluorescence signal starts rising instead of further decreasing, showing that Ca²⁺ returns into the release compartment. This is interpreted as occurring because more and more DHPR voltage sensors and consequently RyRs enter inactivated states, and release finally falls to a level that can be overcompensated by the activity of the SERCA pumps. Thus, it is likely that the shape of the steady-state availability curve of Ca²⁺ release assessed by cytoplasmic Ca²⁺ indicators does not simply reflect the voltage dependence of the DHPRmediated inactivation gating of the RyR1 but is mixed with or even dominated by changes of the driving force for Ca²⁺ efflux from the SR as a result of depletion.

There is agreement in general but deviation in detail between the present results and our previous findings. In the present study, the voltage ranges of Ca²⁺ release leading to depletion and inactivation are apparently farther apart from each other than found in our experiments (Andronache et al., 2009). As a consequence, the predicted window range is substantially broader. This mainly seems to result from a considerably more negative voltage threshold for Ca²⁺ release activation. Because the experimental design differs, it is not evident what causes these differences. One possible reason is the buffering to very low cytoplasmic free Ca²⁺ concentrations in the present experiments because the internal solution did not contain any Ca²⁺. On one hand this may impede reloading of the SR by the SERCA pumps and might cause a right shift in the apparent voltage dependence of DHPR inactivation, on the other hand it may interfere with Ca²⁺-dependent inactivation (Schneider and Simon, 1988; Jong et al., 1995) and therefore lower the voltage threshold for activation.

Is SR depletion a player in muscle fatigue?

A further question emerging from these findings is whether SR Ca²⁺ depletion or DHPR inactivation could contribute to skeletal muscle fatigue caused by strenuous exercise. To investigate this point, Robin and Allard (2013) studied intra-SR Ca²⁺ changes during various protocols in which long trains of action potentials or voltage pulses were applied. Action potentials did in fact produce significant SR Ca²⁺ depletion but no DHPR inactivation even if applied in rapid succession. Only repetitions of

rectangular pulses that were considerably longer than an action potential produced clear evidence of inactivation, i.e., a recovery of SR Ca2+ after an initial decline during the course of stimulation. According to current knowledge, it is unlikely that muscle fatigue can be attributed to a single cellular event; instead, it includes contributions from action potential changes, alterations in the contractile apparatus, and Ca²⁺ release (Allen et al., 2008). Therefore, the authors' conclusion that SR depletion may predominantly contribute to fatigue should be considered with a bit of caution. Their method required intracellular conditions that are quite far from physiological. ATP turnover was probably far lower than under normal conditions in working muscle during high frequency stimulation, and the very high concentration of the Ca²⁺ chelator in the cytoplasm likely favored SR depletion. It seems worthwhile to further investigate this question using less invasive methods. Here, protein-based Ca²⁺ indicators that can be used even in intact muscle (Rudolf et al., 2006) could show their full potential.

I thank Dr. Zoita Andronache for helpful discussions.

This work was supported by Deutsche Forschungsgemeinschaft (DFG) grant ME 713/18-2

Edward N. Pugh Jr. served as editor.

REFERENCES

Allen, D.G., G.D. Lamb, and H. Westerblad. 2008. Skeletal muscle fatigue: cellular mechanisms. *Physiol. Rev.* 88:287–332. http://dx.doi.org/10.1152/physrev.00015.2007

Andronache, Z., S.L. Hamilton, R.T. Dirksen, and W. Melzer. 2009. A retrograde signal from RyR1 alters DHP receptor inactivation and limits window Ca2+ release in muscle fibers of Y522S RyR1 knock-in mice. *Proc. Natl. Acad. Sci. USA*. 106:4531–4536. http://dx.doi.org/10.1073/pnas.0812661106

Baylor, S.M., and S. Hollingworth. 2012. Intracellular calcium movements during excitation–contraction coupling in mammalian slow-twitch and fast-twitch muscle fibers. *J. Gen. Physiol.* 139:261–272. http://dx.doi.org/10.1085/jgp.201210773

Beard, N.A., L. Wei, and A.F. Dulhunty. 2009. Ca(2+) signaling in striated muscle: the elusive roles of triadin, junctin, and calsequestrin. *Eur. Biophys. J.* 39:27–36. http://dx.doi.org/10.1007/s00249-009-0449-6

Berwe, D., G. Gottschalk, and H.C. Lüttgau. 1987. Effects of the calcium antagonist gallopamil (D600) upon excitation-contraction coupling in toe muscle fibres of the frog. *J. Physiol.* 385: 693–707.

Chandler, W.K., R.F. Rakowski, and M.F. Schneider. 1976. Effects of glycerol treatment and maintained depolarization on charge movement in skeletal muscle. *J. Physiol.* 254:285–316.

Cota, G., L. Nicola Siri, and E. Stefani. 1984. Calcium channel inactivation in frog (Rana pipiens and Rana moctezuma) skeletal muscle fibres. J. Physiol. 354:99–108.

Dietze, B., J. Henke, H.M. Eichinger, F. Lehmann-Horn, and W. Melzer. 2000. Malignant hyperthermia mutation Arg615Cys in the porcine ryanodine receptor alters voltage dependence of Ca2+ release. *J. Physiol.* 526:507–514. http://dx.doi.org/10.1111/j.1469-7793.2000.t01-1-00507.x

Dulhunty, A.F. 2006. Excitation-contraction coupling from the 1950s into the new millennium. *Clin. Exp. Pharmacol. Physiol.* 33:763–772. http://dx.doi.org/10.1111/j.1440-1681.2006.04441.x

- Feldmeyer, D., W. Melzer, and B. Pohl. 1990. Effects of gallopamil on calcium release and intramembrane charge movements in frog skeletal muscle fibres. *J. Physiol.* 421:343–362.
- Fink, R.H., and C. Veigel. 1996. Calcium uptake and release modulated by counter-ion conductances in the sarcoplasmic reticulum of skeletal muscle. *Acta Physiol. Scand.* 156:387–396. http://dx.doi.org/10.1046/j.1365-201X.1996.212000.x
- García, J., T. Tanabe, and K.G. Beam. 1994. Relationship of calcium transients to calcium currents and charge movements in myotubes expressing skeletal and cardiac dihydropyridine receptors. J. Gen. Physiol. 103:125–147. http://dx.doi.org/10.1085/jgp.103.1.125
- Hodgkin, A.L., and P. Horowicz. 1960. Potassium contractures in single muscle fibres. J. Physiol. 153:386–403.
- Jong, D.S., P.C. Pape, S.M. Baylor, and W.K. Chandler. 1995. Calcium inactivation of calcium release in frog cut muscle fibers that contain millimolar EGTA or Fura-2. J. Gen. Physiol. 106:337– 388. http://dx.doi.org/10.1085/jgp.106.2.337
- Kabbara, A.A., and D.G. Allen. 2001. The use of the indicator fluo-5N to measure sarcoplasmic reticulum calcium in single muscle fibres of the cane toad. *J. Physiol.* 534:87–97. http://dx.doi.org/10.1111/j.1469-7793.2001.00087.x
- Manno, C., L. Figueroa, R. Fitts, and E. Ríos. 2013. Confocal imaging of transmembrane voltage by SEER of di-8-ANEPPS. J. Gen. Physiol. 141:371–387. http://dx.doi.org/10.1085/jgp.201210936
- Melzer, W., A. Herrmann-Frank, and H.C. Lüttgau. 1995. The role of Ca2+ ions in excitation-contraction coupling of skeletal muscle fibres. *Biochim. Biophys. Acta.* 1241:59–116. http://dx.doi.org/10.1016/0304-4157(94)00014-5
- Murphy, R.M., N.T. Larkins, J.P. Mollica, N.A. Beard, and G.D. Lamb. 2009. Calsequestrin content and SERCA determine normal and maximal Ca2+ storage levels in sarcoplasmic reticulum of fast-and slow-twitch fibres of rat. *J. Physiol.* 587:443–460. http://dx.doi.org/10.1113/jphysiol.2008.163162
- Pape, P.C., D.S. Jong, and W.K. Chandler. 1995. Calcium release and its voltage dependence in frog cut muscle fibers equilibrated with 20 mM EGTA. J. Gen. Physiol. 106:259–336. http://dx.doi.org/ 10.1085/jgp.106.2.259
- Posterino, G.S., and G.D. Lamb. 2003. Effect of sarcoplasmic reticulum Ca2+ content on action potential-induced Ca2+ release in rat skeletal muscle fibres. *J. Physiol.* 551:219–237. http://dx.doi.org/10.1113/jphysiol.2003.040022
- Pouvreau, S., C. Collet, B. Allard, and V. Jacquemond. 2007. Wholecell voltage clamp on skeletal muscle fibers with the silicone-clamp

- technique. Methods Mol. Biol. 403:185–194. http://dx.doi.org/10.1007/978-1-59745-529-9_12
- Robin, G., and B. Allard. 2012. Dihydropyridine receptors actively control gating of ryanodine receptors in resting mouse skeletal muscle fibres. *J. Physiol.* 590:6027–6036. http://dx.doi.org/10.1113/jphysiol.2012.237321
- Robin, G., and B. Allard. 2013. Major contribution of sarcoplasmic reticulum Ca²⁺ depletion during long-lasting activation of skeletal muscle. *J. Gen. Physiol.* 141:557–565.
- Robin, G., C. Berthier, and B. Allard. 2012. Sarcoplasmic reticulum Ca²⁺ permeation explored from the lumen side in *mdx* muscle fibers under voltage control. *J. Gen. Physiol.* 139:209–218. http://dx.doi.org/10.1085/jgp.201110738
- Royer, L., and E. Ríos. 2009. Deconstructing calsequestrin. Complex buffering in the calcium store of skeletal muscle. *J. Physiol.* 587: 3101–3111. http://dx.doi.org/10.1113/jphysiol.2009.171934
- Rudolf, R., P.J. Magalhães, and T. Pozzan. 2006. Direct in vivo monitoring of sarcoplasmic reticulum Ca²⁺ and cytosolic cAMP dynamics in mouse skeletal muscle. *J. Cell Biol.* 173:187–193. http://dx.doi.org/10.1083/jcb.200601160
- Schneider, M.F., and B.J. Simon. 1988. Inactivation of calcium release from the sarcoplasmic reticulum in frog skeletal muscle. *J. Physiol.* 405:727–745.
- Schredelseker, J., M. Shrivastav, A. Dayal, and M. Grabner. 2010. Non-Ca2+-conducting Ca2+ channels in fish skeletal muscle excitation-contraction coupling. *Proc. Natl. Acad. Sci. USA*. 107:5658–5663. http://dx.doi.org/10.1073/pnas.0912153107
- Sztretye, M., J. Yi, L. Figueroa, J. Zhou, L. Royer, and E. Ríos. 2011. D4cpv-calsequestrin: A sensitive ratiometric biosensor accurately targeted to the calcium store of skeletal muscle. *J. Gen. Physiol.* 138:211–229. http://dx.doi.org/10.1085/jgp.201010591
- Ursu, D., R.P. Schuhmeier, M. Freichel, V. Flockerzi, and W. Melzer. 2004. Altered inactivation of Ca²⁺ current and Ca²⁺ release in mouse muscle fibers deficient in the DHP receptor γ₁ subunit. *J. Gen. Physiol.* 124:605–618. http://dx.doi.org/10.1085/jgp.200409168
- Ursu, D., R.P. Schuhmeier, and W. Melzer. 2005. Voltage-controlled Ca2+ release and entry flux in isolated adult muscle fibres of the mouse. *J. Physiol.* 562:347–365. http://dx.doi.org/10.1113/jphysiol. 2004.073882
- Ziman, A.P., C.W. Ward, G.G. Rodney, W.J. Lederer, and R.J. Bloch. 2010. Quantitative measurement of Ca²⁺ in the sarcoplasmic reticulum lumen of mammalian skeletal muscle. *Biophys. J.* 99:2705–2714. http://dx.doi.org/10.1016/j.bpj.2010.08.032