Drug-induced ion channel opening tuned by the voltage sensor charge profile

Nina E. Ottosson, Sara I. Liin, and Fredrik Elinder

Department of Clinical and Experimental Medicine, Linköping University, SE-581 83 Linköping, Sweden

Polyunsaturated fatty acids modulate the voltage dependence of several voltage-gated ion channels, thereby being potent modifiers of cellular excitability. Detailed knowledge of this molecular mechanism can be used in designing a new class of small-molecule compounds against hyperexcitability diseases. Here, we show that arginines on one side of the helical K-channel voltage sensor S4 increased the sensitivity to docosahexaenoic acid (DHA), whereas arginines on the opposing side decreased this sensitivity. Glutamates had opposite effects. In addition, a positively charged DHA-like molecule, arachidonyl amine, had opposite effects to the negatively charged DHA. This suggests that S4 rotates to open the channel and that DHA electrostatically affects this rotation. A channel with arginines in positions 356, 359, and 362 was extremely sensitive to DHA: $70\,\mu\text{M}$ DHA at pH 9.0 increased the current >500 times at negative voltages compared with wild type (WT). The small-molecule compound pimaric acid, a novel Shaker channel opener, opened the WT channel. The 356R/359R/362R channel drastically increased this effect, suggesting it to be instrumental in future drug screening.

INTRODUCTION

The Journal of General Physiology

Polyunsaturated fatty acids (PUFAs) are naturally occurring substances with important functions in normal physiology. As a component of the cell membrane, PUFAs and other fatty acids can directly affect the activity of membrane proteins like voltage-gated ion channels (Schmidt et al., 2006; Boland and Drzewiecki, 2008; Börjesson and Elinder, 2008; Xu et al., 2008b). In addition, free PUFAs can affect different ion channels (Boland and Drzewiecki, 2008; Moreno et al., 2012), and their beneficial effects on heart arrhythmias and epilepsy have been known for a while (McLennan et al., 1985; Hock et al., 1990; Billman et al., 1994; Xiao et al., 1995; Vreugdenhil et al., 1996; Xiao and Li, 1999; Spector, 2001). PUFAs have been suggested to regulate neuronal excitability by closing sodium or calcium channels (Vreugdenhil et al., 1996; Tigerholm et al., 2012) and/or by opening potassium (K) channels (Börjesson et al., 2008, 2010; Xu et al., 2008a; Börjesson and Elinder, 2011; Tigerholm et al., 2012).

A voltage-gated ion channel consists of a pore-forming unit surrounded by four voltage sensor domains (VSDs; Long et al., 2007; Börjesson and Elinder, 2008). Each VSD is composed of four transmembrane helices (S1–S4), where S4 contains several regularly spaced positively charged amino acid residues (Fig. 1, A and B). These positive charges respond to alterations in the membrane voltage by sliding along negative countercharges in S1–S3, and this movement regulates whether

the channel is open or closed (Papazian et al., 1995; Keynes and Elinder, 1998; Broomand and Elinder, 2008; DeCaen et al., 2008, 2009; Catterall, 2010; Henrion et al., 2012; Jensen et al., 2012). The VSD of voltage-gated K channels can be in at least four closed/resting configurations (C1 to C4) and one open/activated configuration (O; Delemotte et al., 2011; Henrion et al., 2012). PUFAs open the voltage-gated Shaker K channel by shifting the voltage dependence of the opening toward more negative voltages (Börjesson et al., 2008, 2010; Xu et al., 2008a; Börjesson and Elinder, 2011). The PUFA molecule is suggested to be inserted into the lipid membrane, close to or in direct contact with the ion channel (Fig. 1 B). The negative charge of the fatty acid attracts the positively charged voltage sensor S4, primarily the last molecular conformational step (C1 \rightarrow O) that swings out the top charge (arginine R362 called R1) against the lipid bilayer (Fig. 1, C and D; Börjesson and Elinder, 2011; Henrion et al., 2012). Thus, R1 is suggested to be a key player in determining the sensitivity to PUFA and PUFA-like molecules (Börjesson and Elinder, 2011). From an earlier study, we know that the Shaker ILT mutant is more sensitive to docosahexaenoic acid (DHA) compared with the WT Shaker channel, supporting an effect by DHA on the last opening step (Börjesson and Elinder, 2011).

In the present investigation, we aimed to construct a channel with increased sensitivity to PUFAs. Such a

Correspondence to Fredrik Elinder: fredrik.elinder@liu.se

Abbreviations used in this paper: AA⁺, arachidonyl amine; DHA, docosahexaenoic acid; PiMA, pimaric acid; PUFA, polyunsaturated fatty acid; VSD, voltage sensor domain.

^{© 2014} Ottosson et al. This article is distributed under the terms of an Attribution–Non-commercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see http://www.rupress.org/terms). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at http://creativecommons.org/licenses/bv-nc-sa/3.0/).

channel would (a) gain us more insight into the biophysical mechanism of action of PUFA, (b) explain why some ion channels are more sensitive for PUFAs than others (Boland and Drzewiecki, 2008), and (c) function as an important tool in the search for new substances with lipoelectric properties, acting as drugs against epilepsy, cardiac arrhythmias, and pain. In addition, we also report on electrostatic channel-opening effects of a small-molecule compound.

MATERIALS AND METHODS

Molecular biology and expression of ion channels

Experiments were performed on the Shaker H4 channel (Kamb et al., 1987), made incapable of fast inactivation by the $\Delta(6-46)$ deletion (Hoshi et al., 1990). Mutagenesis, cRNA synthesis, oocyte preparation, cRNA injection, and oocyte storage were performed according to the procedures described previously (Börjesson et al., 2010; Börjesson and Elinder, 2011). Animal experiments were approved by the local Animal Care and Use Committee at Linköping University.

Electrophysiology

Currents were measured with the two-electrode voltage-clamp technique (GeneClamp 500B amplifier, Digidata 1440A digitizer, and pClamp 10 software; Molecular Devices) 1-6 d after injection of RNA. The amplifier's leak and capacitance compensation were used, and currents were low-pass filtered at 5 kHz. All experiments were performed at room temperature (20-23°C). The holding voltage was set to -80 mV (-120 mV for the L361R/R362Q) mutant), and steady-state currents were achieved by stepping to voltages between -80 and 50 mV (adjusted for some of the mutants) for 80 ms in 5-mV increments. The control solution contained (mM): 88 NaCl, 1 KCl, 15 HEPES, 0.4 CaCl₂, and 0.8 MgCl₂. pH was adjusted to 7.4 with NaOH, yielding a final sodium concentration of \sim 100 mM. Pure control solution was added using a gravitydriven perfusion system. DHA was prepared, stored, and applied as previously described (Börjesson et al., 2008). Arachidonyl amine (AA+) was provided by T. Parkkari (University of Eastern Finland, Kuopio, Finland) and prepared, stored, and applied as previously described (Börjesson et al., 2010). For AA⁺ measurements, cells were preincubated in 1 µM indomethacin, and all recording solutions were supplemented with 1 µM indomethacin, as previously described (Börjesson et al., 2010), to prevent COX-induced metabolization of AA+. Pimaric acid (PiMA) was obtained from Alomone Labs and treated as DHA (however, the stock concentration of PiMA was 50 mM). The effective DHA, AA⁺, and PiMA concentrations were assumed to be 70% of the nominal concentration because of binding to the chamber walls (Börjesson et al., 2008). All concentrations given in the main article are the effective concentrations. To improve the washout of DHA and PiMA, albuminsupplemented (100 mg/liter) control solution was added manually to the bath, followed by continuous wash by control solution. For low concentrations of DHA, the recovery was almost complete, but for higher concentrations less complete. For 70 µM DHA, the recovery ranged from 40 to 85% for the different mutants. All chemicals were obtained from Sigma-Aldrich, if not stated otherwise.

Analysis of data

The K conductance $G_K(V)$ was calculated as

$$G_{K}(V) = I_{K} / (V - V_{rev}),$$

where $I_{\rm K}$ is the steady-state current at the end of an 80-ms pulse, $V_{\rm rev}$ is the absolute membrane voltage, and $V_{\rm rev}$ is the reversal potential for the K channel, set to -80 mV. s and $V_{1/2}$ for Shaker mutants was determined by fitting a simple Boltzmann (n=1) curve to the conductance data:

$$G_{K}(V) = A / \left(1 + \exp\left(\left(V_{1/2} - V\right) / s\right)\right)^{n},$$

where A is the amplitude of the curve and $V_{1/2}$ and s are the midpoint and the slope, respectively. The DHA-induced shift of the $G_K(V)$ curve was quantified at the 10% level as previously described (Börjesson et al., 2008). For illustrative reasons, the figures presented in this manuscript were generated by fitting a Boltzmann curve raised to the n^{th} power (i.e., no restriction for n) to the conductance data. When several concentrations (in increasing order) of DHA were applied on the same oocyte, all shifts were calculated compared with the first control curve.

Statistical analysis

Average values are expressed as mean \pm SEM. When comparing DHA-induced shifts of mutants with control (R362Q), one-way ANOVA together with Dunnett's multiple comparison test was used. When comparing groups, one-way ANOVA together with Bonferroni's multiple comparison tests was used. Correlation analysis was performed by Pearson's correlation test and linear regression. P < 0.05 is considered significant for all tests.

Molecular K channel structure

The crystal structure of the Shaker K channel is not determined. Therefore, we used the structure of the Kv1.2/2.1 chimera channel (Long et al., 2007) with Shaker side chains (Henrion et al., 2012) for the structural evaluations. The Kv1.2/2.1 chimera shares high sequence identity with the Shaker K channel and has previously been shown to serve as an accurate Shaker model (Tao et al., 2010). This chimera was also used for generation of models for the closed states as previously described (Henrion et al., 2012).

Online supplemental material

Fig. S1 shows representative G(V) curves and DHA-induced G(V) shifts for arginine mutants. Fig. S2 shows the correlation between s values of G(V) curves and DHA-induced G(V) shifts for arginine mutants. Fig. S3 shows representative data for the AA^+ effect on A359E/R362Q. Online supplemental material is available at http://www.jgp.org/cgi/content/full/jgp.201311087/DC1.

RESULTS

Place dependence of the top arginine for the effect of DHA

To explore the place dependence of the top arginine (R1) in S4 for DHA effects, we expressed the Shaker K channel in *Xenopus laevis* oocytes and measured ion currents with a two-electrode voltage-clamp technique. We studied eight channels in which either the region 356–362 was neutral or in which one charge at a time was introduced in each of the seven positions 356–362 (Fig. 1 E). All mutants expressed well, and the opening kinetics was essentially unaffected by the mutations (opening time constants were 0.5–2.0 times the opening time constant for the WT channel).

By shifting the positive charge along S4, the voltage dependence for the conductance versus voltage, G(V), curve also shifted along the voltage axis (Fig. 1 F); some mutated channels were opened at more positive voltages (green symbols) than the channel with a neutral 356–362 segment (open symbol), and some were opened at more negative voltages (red symbols; for G(V) curves, see Fig. S1). The two groups of residues are positioned on opposite sides of the S4 helix (Fig. 1 G). The simplest explanation for this orientation dependence is that a charged residue dislikes a hydrophobic environment and therefore destabilizes either the open state or closed state depending on in which state a particular residue faces the lipid bilayer (Yang et al., 2011). Side chain interactions within the VSD may also contribute to the voltage dependence of the arginine mutants: The right-shifted G(V) curve for M356R and A359R is probably supported by charge interactions with negative countercharges in S2 and S3 normally interacting with the gating charges in S4 (Yang et al., 2011; Henrion et al., 2012), whereas the open conformation for S357R, L358R, and L361R might be stabilized by interactions with negatively charged

clusters in the S1–S2 loop or the S3–S4 loop (Henrion et al., 2012).

Shifting the top charge along S4 also altered the Shaker K channel sensitivity to DHA. Moving the arginine from position 362 to 361 abolished the effect of DHA (Fig. 2, A and B), whereas moving it two steps further clearly potentiated the effect (Fig. 2 C). A summary of DHA-induced shifts of the G(V) curves for all mutations (Fig. 2 D) shows that three (yellow bars) are not significantly different from the neutral mutant (open bar), two are significantly less affected (red bars), and two are significantly more affected (green bars), with A359R/R362Q as the most potent mutation, nearly doubling the shift from -6.1 ± 0.6 (n = 9) to -11.8 ± 0.8 mV (n = 9; Fig. S1). When coloring the mutated residues in two suggested Ky channel structures (Henrion et al., 2012), C3 and O, based on their impact on DHA sensitivity, a clear pattern emerges (Fig. 2 E). The red residues point toward the lipid bilayer and the expected position of DHA (Börjesson and Elinder, 2011) in the closed C3 state, supporting the lack of channel-opening effects of DHA on these mutations. The green residues point toward




Figure 1. Charge distribution and voltage dependence of opening of WT and mutated Shaker K channels. (A and B) Gating charges R1–R4 in S4 denoted as blue sticks in a VSD structure (side view [A] and top view [B]) of an open K channel (the Kv1.2/2.1 chimera structure with Shaker side chains; Long et al., 2007; Henrion et al., 2012). The approximate interaction site for DHA is marked with the red encircled negative sign in B (Börjesson and Elinder, 2011). (C and D) R1 (R362) is highlighted in five superimposed structures of S4 (side view [C] from residue 356 and top view [D] from residue 361), for four closed (C1–C4) and one open (O) model states (Henrion et al., 2012). Backbones are color coded according to opening level, from light gray (C4) to black (O). In D, the approximate interaction site for DHA is marked with the red encircled negative sign (Börjesson and Elinder, 2011). The arrow in D denotes the movement of R1 when S4 moves from C1 to O, the step most sensitive to DHA (Börjesson and Elinder, 2011). (E) Amino acid sequences of the extracellular end of S4 (Shaker channel) from the eight single-residue mutations investigated. The positively charged arginines (R) are marked in blue. The mutated region is shown in gray. R1 = R362. (F) The voltage required to reach 50% of the maximum conductance ($V_{1/2}$) plotted against the residue number of the positive charge. The dotted line corresponds to $V_{1/2}$ for the channel without charges in the studied region (R362Q). Green symbols denote positive midpoint voltages relative the neutral 356–362 segment (open symbol), and the red symbols denote negative midpoint voltages. (G) Residues 356–362 are denoted as sticks on the open Shaker VSD structure with same color coding as in F.

the lipid bilayer and the expected DHA position in the open state, thus supporting DHA-induced promotion of channel opening. The yellow residues are positioned toward the lipid bilayer in the open state, but at larger distances (vertically or horizontally) from the expected position of DHA in the open state, supporting a lack of electrostatic interaction. There is a significant correlation between the DHA-induced shifts of voltage for activation and the voltage required to reach 50% of the maximum conductance ($V_{1/2}$; Fig. 2 F). This correlation suggests that, at least to some extent, the mechanisms for the induced shifts and the opening of the channels are common. A similar correlation has been found for metal ion effects on different WT as well as mutated K channels and suggested to depend on alterations in fixed surface charges (Elinder et al., 1996; Elinder and Århem, 2003; Broomand et al., 2007). In contrast, there is no correlation between the slope of the fitted Boltzmann curves of the arginine mutants and the DHA-induced shifts, suggesting that the altered sensitivity to DHA does not depend on altered energy barriers between the states with altered state occupancy as consequence (Fig. S2).

Rotation of S4 is required to open the channel: Sidedness of the charge effects

To further test the hypothesis that there is an electrostatic interaction between charges in S4 and the DHA molecule, and not simply that additional charges in S4 dislike the lipid bilayer or hydrophobic pockets of the channel protein, we tested whether negatively charged glutamates at three consecutive positions, 359 to 361, had opposite effects to the arginines. A glutamate in position 359 significantly reduced the DHA-induced G(V) shift from -6.1 ± 0.6 to -3.8 ± 0.4 mV (n = 7; Fig. 3, A and B). This alteration is in the opposite direction of the positively charged arginine. The 359E mutant was further investigated by testing whether AA+, a positively charged PUFA analogue, shows the opposite effect compared with DHA. Although AA+ previously was shown to close the Shaker WT channel by shifting the G(V) curve in a positive direction along the voltage axis (Börjesson et al., 2010), AA+ opened the 359E channel by shifting the G(V) curve by -8.9 ± 1.7 mV (n = 4; Fig. S3), which is significantly more than the shift induced by the negatively charged DHA on the same channel




Figure 2. DHA sensitivity of the single arginine mutants. (A–C) Representative current traces for voltages corresponding to 10% of G_{max} in control solution at pH 7.4. Black traces indicate control, and red traces indicate 70 μM DHA. The increments in current amplitudes are R362R (WT), 2.2 times (A); for L361R/R362Q, 0.9 times (B); and A359R/R362Q, 2.7 times (C). (D) DHA-induced G(V) shifts for the single-charge mutants (70 μM DHA at pH 7.4). The black line equals the DHA-induced shift for R362Q. Mean ± SEM (n = 8, 8, 6, 9, 10, 14, 9, and 9). DHA-induced shifts compared with R362Q (one-way ANOVA together with Dunnett's multiple comparison test: **, P < 0.01; ***, P < 0.001). Green bars denote significantly larger DHA-induced shifts relative to the neutral 356–362 segment (open bar), the red bars denote significantly smaller shifts, and the yellow bars denote no significant differences. (E) Mutated residues are marked on one VSD of the Shaker K channel in states C3 and O. Same color coding as in D. (F) Correlations between the $V_{1/2}$ values and the DHA-induced shifts for the channels described in D. Slope is significantly different from zero (Pearson correlation test, and linear regression: P < 0.01 for both). The symbols denote mean ± SEM, the continuous line is the linear regression, and the dashed lines denote the 95% confidence interval.

 $(-3.8 \pm 0.4 \text{ mV})$. These experiments clearly support an electrostatic mechanism.

In contrast to the experiments for 359 above, swopping the charge at position 361 from positive to negative had the opposite effect; the DHA-induced shift increased from -0.8 to -7.0 mV (Fig. 3, C and D), slightly more than for R362Q. The AA+ effect on 361E was not possible to evaluate because AA+ induced severe cellular toxicity for this mutant (see also Börjesson et al. [2010] for toxicity description) and caused substantial inactivation of 361E. For position 360, there was no difference between the glutamate and the arginine (Fig. 3 E). Calculations of the electrostatic effects are complicated because of several unknown parameters, like the exact positions of the amino acid charges, the dielectric constant, and uncertainty regarding the number of bound DHA molecules. However, the data are consistent with a simple model, locating one DHA molecule in the lipid bilayer just outside of S4 as indicated in Fig. 3 F. The last step in the channel opening, the transition from C1 to O (Henrion et al., 2012), is the step most sensitive to DHA (Börjesson and Elinder, 2011). During this step (Fig. 3 F, arrows), 359 approaches the negatively charged DHA molecule, 361 departs from the DHA molecule, and 360 keeps a relatively constant distance from the DHA molecule (Fig. 3 F). A positive charge at 359 or a negative charge at 361 promotes opening. A negative charge at 359 or a positive charge at 361 prevents opening. Notably, the negative glutamates have less of an effect than the positive arginines, compared with R362Q, probably because the negative charge of the glutamate can partly push away the negative DHA molecule from the channel or deprotonate the DHA molecule. These data strongly suggest that S4 is required to rotate in the last step to open the channel and that DHA electrostatically affects this rotation.

Combining arginine mutations to increase the DHA sensitivity

To search for a channel with even higher sensitivity to DHA than the single mutants, we explored the effect of DHA (70 µM at pH 7.4) on several combinations of positively charged residues in the positions 356, 358, 359, 360, and 362 (Fig. 4). The potentiating effect of A359R was independent (i.e., additive) of single-positive charges in positions 356, 358, or 362, and thus, A359R could easily be combined with these arginines to gain a larger sensitivity to DHA. In contrast, the potentiating effect of I360R was abolished in various combinations with A359R. In combination with A359R alone, I360R even abolished the potentiating effect of an arginine at position 359. Thus, I360R was not a good candidate for the construction of a highly DHA-sensitive channel. The L358R/A359R/R362Q mutant was generated to mimic the charge profile of the Kv2.1 channel because this channel is known to be sensitive to DHA (McKay and

Worley, 2001). Also in this system, the Kv2.1 look-a-like channel was sensitive to DHA, which displayed a relatively large DHA-induced shift of -10.7 ± 1.2 mV (n = 8), but this effect was gained from 359R rather than 358R. The triple-R mutant M356R/A359R/R362R (hereafter referred to as 3R) turned out to be the most sensitive channel and was thus further explored.




Figure 3. Charge-dependent DHA sensitivity. (A) Normalized representative current traces for voltages corresponding to 10% of G_{max} in control solution at pH 7.4. Black and gray traces indicate control solution for A359R/R362Q and A359E/R362Q, respectively (normalized to 1), blue trace denotes 70 µM DHA on A359R/R362Q, and red trace denotes 70 µM DHA on A359E/ R362Q. (B) DHA-induced G(V) shifts for the channels in A. 70 μ M DHA at pH 7.4. R362Q is included for comparison. Mean ± SEM (n = 7, 12, and 9). DHA-induced shifts are compared by one-way ANOVA together with Bonferroni post-hoc test: *, P < 0.05; ***, P < 0.001. (C) As in A. Red trace for L361E/R362Q and blue trace for L361R/R362Q. (D) As in B (n = 9, 12, and 14). (E) As in B (n = 7, 12, and 10; **, P < 0.01). (F) Cartoon to qualitatively explain data in A-E. Figures denote approximate positions of specific residues in the open state. Arrows denote the movements of the respective residues from the C1 state to the O state. The minus sign denotes a DHA position consistent with the experimental data.

The 3R mutant exhibits dramatically increased DHA sensitivity

There are no major differences between the activation kinetics of the 3R mutant and the control channel (i.e., a difference less than a factor of 2; Fig. 5, A and B). For the 3R mutant, 70 μ M DHA at pH 7.4 increased the current 11-fold at 10 mV (Fig. 5 C) and shifted the voltage dependence by -19.6 ± 1.0 mV (n=15; Fig. 5 D), a shift more than three times larger than that for the R362Q and WT channels.

In a previous study, we showed that the DHA-induced G(V) shift of the Shaker WT channel was pH dependent (Börjesson et al., 2008). The pH dependence is explained by incomplete deprotonation of DHA at pH 7.4, as the apparent pKa value of PUFAs in a lipid membrane is \sim 7.5 (Börjesson et al., 2008). Thus, the effect of DHA on the 3R channel was further investigated by altering the pH. At -15 mV and pH 9.0, a DHA concentration ≥ 3 µM clearly increased the current; 7 µM caused a 10-fold increase (orange trace), and 70 µM caused a 40-fold increase (green trace; Fig. 6 A). Increasing concentrations of DHA at pH 9.0 gradually shifted the voltage dependence to more negative voltages (up to -60 mV), with minor effects on the maximal conductance (Fig. 6 B). However, at high concentrations, DHA also induced some inactivation, as previously reported for the WT Shaker channel (Börjesson et al., 2008). The DHA-induced shift caused by the highest DHA concentrations may therefore be the result of two separate mechanisms: lipoelectric opening of the channel and channel inactivation (possibly by DHA-induced conformational changes of the selectivity filter or block of the gate). As inactivation mainly occurs at the more positive voltages, we expect only minor effects of DHA-induced inactivation on the G(V) shifts measured at the 10% level. DHA concentrations >210 µM were not tested because free




Figure 4. Combinations of arginines make channels more PUFA sensitive. DHA-induced shifts for multicharge channels (70 μM DHA at pH 7.4). Mean \pm SEM (n = 9, 12, 10, 15, 11, 8, and 8). Horizontal lines denote shifts of R362Q (black), I360R/R362Q (light blue), and A359R/R362Q (dark blue). DHA-induced shifts compared with R362Q (one-way ANOVA together with Dunnett's multiple comparison test: *, P < 0.05; **, P < 0.01; ***, P < 0.001).

fatty acids form micelles around 100 μ M in physiological solutions (Richieri et al., 1992; Börjesson et al., 2008). The effect of DHA on the control channel versus the 3R channel was measured at different concentrations at pH 7.4 and 9.0 (Fig. 6 C). From these data, it is clear that the DHA sensitivity of the WT channel at pH 9.0 is similar to the DHA sensitivity of the 3R channel in pH 7.4. The increased DHA sensitivity of the 3R channel is likely explained by the additional three positive charges in the 3R channel that all point toward the lipid bilayer and therefore are able to electrostatically interact with DHA (Fig. 6 D).

Thus, although 70 µM DHA at pH 9.0 shifted the G(V) of the WT channel by -18.0 ± 1.4 mV (n=9; Börjesson et al., 2008), it shifted the G(V) of the 3R channel by -47.9 ± 4.2 mV (n=4). At low voltages these shifts can be converted to equivalent increases in current magnitude, $A=\exp(-\Delta V/4.7;$ Börjesson et al., 2010). The amplitude increase from WT to 3R is thus $A_{3R}/A_{WT}=\exp(-\Delta \Delta V/4.7)$. Thus, introducing two extra positive charges at positions 356 and 359 increases the open probability of the Shaker K channel by a factor of 580 at 70 µM DHA at pH 9.0. This large potentiation simplifies the search for other compounds with similar properties.




Figure 5. Characteristics of the 3R channel. (A and B) Current families for R362Q (A) and 3R channel (B) when stepping the membrane voltage from -80 mV to voltages between -80 and 40 mV (-60 and 60 mV for 3R channel) in 5-mV increments in control solution, pH 7.4, at a frequency of 0.2 Hz. (C) Representative current traces at pH 7.4 for 3R in control solution (black) and in 70 µM DHA (red) for a voltage corresponding to 10% of $G_{\rm max}$ in control solution (i.e., 10 mV). The increment in current amplitude is >10-fold. (D) Representative G(V) curves. Same cell as in C (control, black symbols; DHA, red symbols). $\Delta G(V)_{\rm DHA} = -22.1$ mV in this example.

Identifying a novel Shaker channel opener

DHA and other PUFAs display promising anti-excitable effects on the Shaker K channel (Börjesson et al., 2008, 2010; Xu et al., 2008a; Börjesson and Elinder, 2011; Tigerholm et al., 2012). However, they are very promiscuous, a characteristic not appreciated in drug design. Therefore, in an attempt to find other candidate drugs, we have searched for small-molecule compounds with similar effects to DHA. PiMA, an amphipathic resin acid known to open Ca²⁺-activated BK channels (Imaizumi et al., 2002) has, in conformity with the fatty acids, a lipophilic domain and a carboxyl group supposed to be negatively charged at high pH (Fig. 7 A). Therefore, PiMA is also a possible candidate to open Kv channels. For the WT Shaker K channel, 70 µM PiMA at pH 7.4 shifted the G(V) by -4.6 ± 0.7 mV (n = 15; Fig. 7 B). For the 3R channel, the shift was almost doubled to $-8.4 \pm$ 1.2 mV (n = 4; Fig. 7 B), indicating that PiMA also activates the Kv channel by electrostatically affecting the positive charges in the top of S4. Similar to the DHA effect, the PiMA effect was potentiated at increased pH (Fig. 7, B-F). Thus, we have found a small-molecule compound able to shift the G(V) of the 3R channel at pH 9.0 by almost -30 mV.

DISCUSSION

In this study, we have investigated introduced charged amino acid residues in S4, and combinations of them, and how they affect the DHA-induced alteration of the Shaker K channel's voltage dependence. We found that the single mutations A359R and I360R and combinations including A359R, in particular 3R, significantly increased the sensitivity to DHA. We have also identified a small molecule compound, PiMA, with similar effects as DHA on the channel's voltage dependence. Furthermore, we found that residues on the opposite side of S4, S357R and L361R, significantly reduced channel sensitivity to DHA. The place dependence of arginines and glutamates for the DHA sensitivity supports a rotational S4 movement in the last opening step that is promoted by negatively charged lipophilic compounds, like PUFAs and PiMA. This model is supported by data for the positively charged PUFA analogue AA+, which shows opposite the effects to DHA on 359E.

An understanding of the molecular details for the PUFA-channel interactions is important for explaining and predicting differences in PUFA sensitivity between channels. One of the channels constructed in this study mimics the S4 arginine profile of Kv2.1 (L358R/A359R/R362Q). The Kv2.1-mimicking mutant demonstrated increased DHA sensitivity, in line with experimental findings reporting clear shifts in Kv2.1 channel voltage dependence from low micromolar PUFA concentrations (McKay and Worley, 2001). BK channels also display a favorable S4 charge profile for possible PUFA

effects by having a positive charge at the position equivalent to 359 in the Shaker K channel. BK channels are reported to be highly sensitive to PUFAs (Clarke et al., 2003; Sun et al., 2007; Hoshi et al., 2013a,b). However, these large effects on the BK channel could, at least partly, be explained by another mechanism because residues on the intracellular side of the channel are involved in the effect (Hoshi et al., 2013a). Other DHA-promoting or -preventing charge profiles are found in different pseudotetrameric sodium and calcium channels.




Figure 6. pH-dependent DHA sensitivity. (A) Representative current sweeps in control (black), 2.1 μM DHA (red), 7 μM DHA (orange), 21 μM DHA (yellow), 70 μM DHA (green), and 210 μM DHA (blue) at pH 9.0 and -15 mV (the voltage corresponding to 10% of $G_{\rm max}$ in control solution minus 10 mV). Current amplitude is increased >40-fold at 70 μM DHA. (B) Representative G(V) curves. Same cell, color coding, and order as in A. The shifts are -4.9, -15.7, -26.7, -50.0, and -54.6 mV in this example. (C) Dose–response curve for R362Q (gray) and the 3R channel (red) at pH 7.4 (light colored) and at pH 9.0 (dark colored). Error bars indicate SEM (n = 4–15). (D) VSD of the open 3R Shaker structure (top view to the left and side view to the right) with arginines in S4 shown as blue sticks.

However, the physiological relevance is difficult to evaluate because only one out of four VSDs is expected to show increased or decreased PUFA sensitivity.

We have previously reported that part of S4 adapts a 3₁₀-helical structure and that the segment forming a 3₁₀ helix slides along the S4 helix during gating (Henrion et al., 2012). In our previous study, we proposed that the 3₁₀-helical structure is downstream of R3 in the last opening step, the step most sensitive to DHA. Because the electric force varies inversely with the square of the distance between two charges, the electrostatic interaction between DHA and the gating charges in the 3₁₀-helical segment during the opening step will be weak. Hence, the 3₁₀ helix structure and the sliding of that structure will likely be preserved in the presence of DHA.

The main finding in the present work is the 3R mutant, a constructed channel with an increased sensitivity to PUFAs with respect to the channel's voltage sensitivity. 70 µM DHA increased the current of the 3R channel at pH 9.0 and negative voltages >500 times the WT channel. Thus, the 3R channel is a promising tool in the search for pharmacological compounds with beneficial affects against cellular hyperexcitability in diseases

such as cardiac arrhythmia, epilepsy, and pain. As a first example of this, we here demonstrate 3R channel sensitivity to PiMA, a novel Shaker-channel opener.

The AA⁺ was generously provided by Teija Parkkari.

This work was supported by grants from the Swedish Research Council, the Swedish Heart-Lung Foundation, the Swedish Brain Foundation, the County Council of Östergötland, and King Gustaf V and Queen Victoria's Freemasons Foundation.

The authors have no conflicting financial interests.

Kenton J. Swartz served as editor.

Submitted: 16 August 2013 Accepted: 19 December 2013

REFERENCES

Billman, G.E., H. Hallaq, and A. Leaf. 1994. Prevention of ischemia-induced ventricular fibrillation by omega 3 fatty acids. *Proc. Natl. Acad. Sci. USA*. 91:4427–4430. http://dx.doi.org/10.1073/pnas.91.10.4427

Boland, L.M., and M.M. Drzewiecki. 2008. Polyunsaturated fatty acid modulation of voltage-gated ion channels. *Cell Biochem. Biophys.* 52:59–84. http://dx.doi.org/10.1007/s12013-008-9027-2 Börjesson, S.I., and F. Elinder. 2008. Structure, function, and modification of the voltage sensor in voltage-gated ion channels.




Figure 7. Drug sensitivity is increased for 3R. (A) Structure of PiMA, here in the uncharged form with the carboxylic acid group protonated. (B) G(V) shifts induced by 70 µM PiMA at pH 7.4 and 9.0 for WT and 3R. Mean \pm SEM (n = 12, 6, 4, and 6; one-way ANOVA together with Bonferroni's pairwise comparison test: **, P < 0.01; ***, P < 0.001). (C) Representative current traces at pH 9.0 for WT in control solution (black) and in 70 µM PiMA (red) for a voltage corresponding to 10% of G_{max} in control solution (i.e., -35 mV). The current amplitude is increased fourfold. (D) Representative G(V) curves. Same cell as in C (control, black symbols; PiMA, red symbols). $\Delta G(V)_{PiMA} = -9.0 \text{ mV}$ in this example. (E) Representative current traces at pH 9.0 for 3R in control solution (black) and in 70 µM PiMA (red) for a voltage corresponding to 10% of G_{max} in control solution (i.e., 5 mV). The current amplitude is increased >10 times. (F) Representative G(V) curves. Same cell as in F (control, black symbols; PiMA, red symbols). $\Delta G(V)_{PiMA} = -28.5 \text{ mV}$ in this example.

- Cell Biochem. Biophys. 52:149–174. http://dx.doi.org/10.1007/s12013-008-9032-5
- Börjesson, S.I., and F. Elinder. 2011. An electrostatic potassium channel opener targeting the final voltage sensor transition. *J. Gen. Physiol.* 137:563–577. http://dx.doi.org/10.1085/jgp.201110599
- Börjesson, S.I., S. Hammarström, and F. Elinder. 2008. Lipoelectric modification of ion channel voltage gating by polyunsaturated fatty acids. *Biophys. J.* 95:2242–2253. http://dx.doi.org/10.1529/biophysj.108.130757
- Börjesson, S.I., T. Parkkari, S. Hammarström, and F. Elinder. 2010. Electrostatic tuning of cellular excitability. *Biophys. J.* 98:396–403. http://dx.doi.org/10.1016/j.bpj.2009.10.026
- Broomand, A., and F. Elinder. 2008. Large-scale movement within the voltage-sensor paddle of a potassium channel-support for a helical-screw motion. *Neuron.* 59:770–777. http://dx.doi.org/10.1016/j.neuron.2008.07.008
- Broomand, A., F. Osterberg, T. Wardi, and F. Elinder. 2007. Electrostatic domino effect in the Shaker K channel turret. *Biophys. J.* 93:2307–2314. http://dx.doi.org/10.1529/biophysj.107.104349
- Catterall, W.A. 2010. Ion channel voltage sensors: structure, function, and pathophysiology. *Neuron*. 67:915–928. http://dx.doi.org/10.1016/j.neuron.2010.08.021
- Clarke, A.L., S. Petrou, J.V. Walsh Jr., and J.J. Singer. 2003. Site of action of fatty acids and other charged lipids on BKCa channels from arterial smooth muscle cells. Am. J. Physiol. Cell Physiol. 284:C607–C619. http://dx.doi.org/10.1152/ajpcell.00364.2002
- DeCaen, P.G., V. Yarov-Yarovoy, Y. Zhao, T. Scheuer, and W.A. Catterall. 2008. Disulfide locking a sodium channel voltage sensor reveals ion pair formation during activation. *Proc. Natl. Acad. Sci. USA*. 105:15142–15147. http://dx.doi.org/10.1073/pnas.0806486105
- DeCaen, P.G., V. Yarov-Yarovoy, E.M. Sharp, T. Scheuer, and W.A. Catterall. 2009. Sequential formation of ion pairs during activation of a sodium channel voltage sensor. *Proc. Natl. Acad. Sci. USA*. 106:22498–22503. http://dx.doi.org/10.1073/pnas.0912307106
- Delemotte, L., M. Tarek, M.L. Klein, C. Amaral, and W. Treptow. 2011. Intermediate states of the Kv1.2 voltage sensor from atomistic molecular dynamics simulations. *Proc. Natl. Acad. Sci. USA*. 108:6109–6114. http://dx.doi.org/10.1073/pnas.1102724108
- Elinder, F., and P. Århem. 2003. Metal ion effects on ion channel gating. *Q. Rev. Biophys.* 36:373–427. http://dx.doi.org/10.1017/S0033583504003932
- Elinder, F., M. Madeja, and P. Århem. 1996. Surface Charges of K channels. Effects of strontium on five cloned channels expressed in *Xenopus* oocytes. *J. Gen. Physiol.* 108:325–332. http://dx.doi.org/10.1085/jgp.108.4.325
- Henrion, U., J. Renhorn, S.I. Börjesson, E.M. Nelson, C.S. Schwaiger, P. Bjelkmar, B. Wallner, E. Lindahl, and F. Elinder. 2012. Tracking a complete voltage-sensor cycle with metal-ion bridges. *Proc. Natl. Acad. Sci. USA*. 109:8552–8557. http://dx.doi.org/10.1073/ pnas.1116938109
- Hock, C.E., L.D. Beck, R.C. Bodine, and D.K. Reibel. 1990. Influence of dietary n-3 fatty acids on myocardial ischemia and reperfusion. Am. J. Physiol. 259:H1518–H1526.
- Hoshi, T., W.N. Zagotta, and R.W. Aldrich. 1990. Biophysical and molecular mechanisms of Shaker potassium channel inactivation. *Science*. 250:533–538. http://dx.doi.org/10.1126/science.2122519
- Hoshi, T., Y. Tian, R. Xu, S.H. Heinemann, and S. Hou. 2013a. Mechanism of the modulation of BK potassium channel complexes with different auxiliary subunit compositions by the omega-3 fatty acid DHA. *Proc. Natl. Acad. Sci. USA*. 110:4822–4827. http://dx.doi.org/10.1073/pnas.1222003110
- Hoshi, T., B. Wissuwa, Y. Tian, N. Tajima, R. Xu, M. Bauer, S.H. Heinemann, and S. Hou. 2013b. Omega-3 fatty acids lower blood

- pressure by directly activating large-conductance Ca^{2^+} -dependent K^+ channels. *Proc. Natl. Acad. Sci. USA.* 110:4816–4821. http://dx.doi.org/10.1073/pnas.1221997110
- Imaizumi, Y., K. Sakamoto, A. Yamada, A. Hotta, S. Ohya, K. Muraki, M. Uchiyama, and T. Ohwada. 2002. Molecular basis of pimarane compounds as novel activators of large-conductance Ca(2+)-activated K(+) channel alpha-subunit. *Mol. Pharmacol.* 62:836–846. http://dx.doi.org/10.1124/mol.62.4.836
- Jensen, M.Ø., V. Jogini, D.W. Borhani, A.E. Leffler, R.O. Dror, and D.E. Shaw. 2012. Mechanism of voltage gating in potassium channels. *Science*. 336:229–233. http://dx.doi.org/10.1126/science.1216533
- Kamb, A., L.E. Iverson, and M.A. Tanouye. 1987. Molecular characterization of Shaker, a *Drosophila* gene that encodes a potassium channel. *Cell*. 50:405–413. http://dx.doi.org/10.1016/0092-8674(87)90494-6
- Keynes, R.D., and F. Elinder. 1998. Modelling the activation, opening, inactivation and reopening of the voltage-gated sodium channel. *Proc. Biol. Sci.* 265:263–270. http://dx.doi.org/10.1098/rspb.1998.0291
- Long, S.B., X. Tao, E.B. Campbell, and R. MacKinnon. 2007. Atomic structure of a voltage-dependent K+ channel in a lipid membrane-like environment. *Nature*. 450:376–382. http://dx.doi.org/10.1038/nature06265
- McKay, M.C., and J.F. Worley III. 2001. Linoleic acid both enhances activation and blocks Kv1.5 and Kv2.1 channels by two separate mechanisms. *Am. J. Physiol. Cell Physiol.* 281:C1277–C1284.
- McLennan, P.L., M.Y. Abeywardena, and J.S. Charnock. 1985. Influence of dietary lipids on arrhythmias and infarction after coronary artery ligation in rats. *Can. J. Physiol. Pharmacol.* 63:1411– 1417. http://dx.doi.org/10.1139/y85-232
- Moreno, C., A. Macías, A. Prieto, A. de la Cruz, T. González, and C. Valenzuela. 2012. Effects of n-3 Polyunsaturated Fatty Acids on Cardiac Ion Channels. Front Physiol. 3:245. http://dx.doi.org/ 10.3389/fphys.2012.00245
- Papazian, D.M., X.M. Shao, S.A. Seoh, A.F. Mock, Y. Huang, and D.H. Wainstock. 1995. Electrostatic interactions of S4 voltage sensor in Shaker K+ channel. *Neuron.* 14:1293–1301. http://dx.doi.org/10.1016/0896-6273(95)90276-7
- Richieri, G.V., R.T. Ogata, and A.M. Kleinfeld. 1992. A fluorescently labeled intestinal fatty acid binding protein. Interactions with fatty acids and its use in monitoring free fatty acids. *J. Biol. Chem.* 267:23495–23501.
- Schmidt, D., Q.-X. Jiang, and R. MacKinnon. 2006. Phospholipids and the origin of cationic gating charges in voltage sensors. *Nature*. 444:775–779. http://dx.doi.org/10.1038/nature05416
- Spector, A.A. 2001. Plasma free fatty acid and lipoproteins as sources of polyunsaturated fatty acid for the brain. *J. Mol. Neurosci.* 16:159–165. http://dx.doi.org/10.1385/JMN:16:2-3:159
- Sun, X., D. Zhou, P. Zhang, E.G. Moczydlowski, and G.G. Haddad. 2007. Beta-subunit-dependent modulation of hSlo BK current by arachidonic acid. *J. Neurophysiol.* 97:62–69. http://dx.doi.org/10.1152/jn.00700.2006
- Tao, X., A. Lee, W. Limapichat, D.A. Dougherty, and R. MacKinnon. 2010. A gating charge transfer center in voltage sensors. *Science*. 328:67–73. http://dx.doi.org/10.1126/science.1185954
- Tigerholm, J., S.I. Börjesson, L. Lundberg, F. Elinder, and E. Fransén. 2012. Dampening of hyperexcitability in CA1 pyramidal neurons by polyunsaturated fatty acids acting on voltage-gated ion channels. *PLoS ONE.* 7:e44388. http://dx.doi.org/10.1371/journal.pone.0044388
- Vreugdenhil, M., C. Bruehl, R.A. Voskuyl, J.X. Kang, A. Leaf, and W.J. Wadman. 1996. Polyunsaturated fatty acids modulate sodium and calcium currents in CA1 neurons. *Proc. Natl. Acad. Sci. USA*. 93:12559–12563. http://dx.doi.org/10.1073/pnas.93.22.12559
- Xiao, Y., and X. Li. 1999. Polyunsaturated fatty acids modify mouse hippocampal neuronal excitability during excitotoxic

- or convulsant stimulation. Brain Res. 846:112–121. http://dx.doi .org/10.1016/S0006-8993(99)01997-6
- Xiao, Y.F., J.X. Kang, J.P. Morgan, and A. Leaf. 1995. Blocking effects of polyunsaturated fatty acids on Na+ channels of neonatal rat ventricular myocytes. *Proc. Natl. Acad. Sci. USA*. 92:11000–11004. http://dx.doi.org/10.1073/pnas.92.24.11000
- Xu, X.P., D. Erichsen, S.I. Börjesson, M. Dahlin, P. Amark, and F. Elinder. 2008a. Polyunsaturated fatty acids and cerebrospinal fluid from children on the ketogenic diet open a voltage-gated K channel: a putative mechanism of antiseizure
- action. *Epilepsy Res.* 80:57-66. http://dx.doi.org/10.1016/j.eplepsyres.2008.03.013
- Xu, Y., Y. Ramu, and Z. Lu. 2008b. Removal of phospho-head groups of membrane lipids immobilizes voltage sensors of K+ channels. *Nature*. 451:826–829. http://dx.doi.org/10.1038/nature06618
- Yang, Y.-C., S. Lin, P.-C. Chang, H.-C. Lin, and C.-C. Kuo. 2011. Functional extension of amino acid triads from the fourth transmembrane segment (S4) into its external linker in Shaker K(+) channels. *J. Biol. Chem.* 286:37503–37514. http://dx.doi.org/10.1074/jbc.M111.237792