Location, Location: Juxtaposed calcium-signaling microdomains as a novel model of the vascular smooth muscle myogenic response

Nathan R. Tykocki¹ and Mark T. Nelson^{1,2}

Small arteries and arterioles constrict in response to increases in intraluminal pressure through a process called "myogenic tone" or the "myogenic response" (McCarron et al., 1989). Development of myogenic tone has profound physiological implications; it contributes substantially to regulation of systemic blood pressure, maintenance of capillary/tissue perfusion, and autoregulation of blood flow to specific organs, such as kidneys and brain (Davis, 2012). Thus, understanding the mechanisms that govern the myogenic response is germane to our understanding of the cardiovascular system as a whole. The current thinking is that myogenic tone is regulated by a "push/pull" relationship between vasoconstrictor and vasodilator mechanisms, both of which are regulated through focal increases in intracellular Ca²⁺ concentration within cellular microdomains (Hill-Eubanks et al., 2011). The interaction and proximity of these two vasoregulatory pathways are critical to the regulation of myogenic tone and proper vascular function. However, even with the breadth and extent of investigation into the myogenic response, a definitive model of the structure, localization, and composition of these cellular-signaling domains remains elusive.

Karlin (2015) describes a new model of an arterial smooth muscle cell that, unlike previous models (Bennett et al., 2005; Kapela et al., 2008), incorporates five interconnected Ca²⁺ compartments and 37 separate protein components into two distinct microdomains. When tested, the proposed model accurately recapitulated the changes in membrane potential and intracellular Ca²⁺ observed experimentally in response to increases in intraluminal pressure. Moreover, when used to simulate pressure-independent responses to vasoconstrictors (ATP, α-adrenergic agonists) and vasodilators (nitric oxide, epoxyeicosatrienoic acids, β -adrenergic agonists), this model continued to accurately represent the changes in smooth muscle cell membrane potential and intracellular Ca2+ recorded experimentally under the same conditions.

Mechanisms of pressure-induced dilation and constriction Of the mechanisms that regulate the development of myogenic tone in smooth muscle, the best defined is the major vasodilatory pathway that opposes myogenic constriction. This pathway involves large conductance Ca²⁺-activated K⁺ (BK) channels and voltage-dependent Ca²⁺ channels (VDCCs) in the plasma membrane, which are closely aligned with one another and with ryanodine receptors (RyRs) on the sarcoplasmic reticulum (Brayden and Nelson, 1992; Nelson and Quayle, 1995; Nelson et al., 1995; Jaggar et al., 2000). Intravascular pressure-induced membrane depolarization increases the open probability of VDCCs, thereby elevating Ca²⁺ influx and intracellular Ca²⁺ concentration (Nelson et al., 1990; Rubart et al., 1996; Knot and Nelson, 1998). This increase in intracellular Ca2+ leads to an increase in the activity of clusters of RyRs in the adjacent sarcoplasmic reticulum, causing a transient focal increase in Ca²⁺ known as a Ca²⁺ spark (Nelson et al., 1995). Ca²⁺ sparks, in turn, activate nearby BK channels to cause membrane hyperpolarization and closure of VDCCs, which ultimately leads to vasodilation (Jaggar et al., 2000).

The upstream molecular players in the pathway leading to myogenic constriction are less well characterized, but it is generally accepted that increases in intraluminal pressure leads to the activation of cation channels in smooth muscle cells to cause membrane depolarization. Evidence collected to date suggests that the most likely candidates for these cation channels are members of the transient receptor potential (TRP) superfamily (Welsh et al., 2002). One such channel is TRPM4, a member of the melastatin subfamily of TRP channels that primarily conducts Na⁺, and as such causes membrane depolarization and subsequent Ca2+ influx through VDCCs (Earley and Brayden, 2015). However, other mechanosensitive TRP channels present in smooth muscle could also locally increase cytosolic Ca²⁺ concentrations in response to stretch (Bulley et al., 2012). Whether resulting

The Journal of General Physiology

¹Department of Pharmacology, University of Vermont, Burlington, VT 05405

²Institute of Cardiovascular Sciences, University of Manchester, Manchester M13 9PL, England, UK

^{© 2015} Tykocki and Nelson This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see http://www.rupress.org/terms). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at http://creativecommons.org/licenses/by-nc-sa/3.0/).

from the opening of VDCCs or of TRP channels, localized increases in Ca²⁺ serve as part of a feed-forward mechanism that opens adjacent Ca²⁺-activated Cl⁻ channels (CaCCs) to augment membrane depolarization, Ca²⁺ influx, global intracellular Ca²⁺ concentration, and vasoconstriction (Jaggar et al., 2000; Bulley et al., 2012).

Ca²⁺-signaling microdomains and Ca²⁺ sensitivity

Strict regulation of global intracellular Ca²⁺ is of the utmost importance for the controlled development of myogenic tone because a relatively small change in Ca²⁺ concentration (from <100 to ~400 nM global Ca²⁺) is sufficient to span the range from complete dilation to complete constriction (Knot and Nelson, 1998; Hill-Eubanks et al., 2011). As such, complex regulatory mechanisms orchestrate local Ca²⁺ dynamics within the smooth muscle cell to control both contractile and relaxant mechanisms. A logical way to achieve this regulation is to arrange the requisite component proteins into spatially distinct contractile and relaxant microdomains, which confine and insulate the necessary molecular players within an extremely limited space. In this arrangement, localized increases in intracellular Ca²⁺ can be high enough to activate pro-contractile and prorelaxant mechanisms separately, without causing a short circuit between these mechanisms or increasing global cytosolic Ca2+ to an extent that could be deleterious to cell viability (Hill-Eubanks et al., 2011). Spatial constraints imposed within these microdomains also help impart signal fidelity and temporal control over the development of myogenic tone (Parekh, 2008).

However, microdomain proximity and component density are not the sole determinants of myogenic responsiveness, nor are they capable of controlling intracellular Ca²⁺ concentrations over such a narrow range. Rather, differences in the Ca2+ sensitivity of individual microdomain elements combine with microdomain arrangement to impart the resolution and precision necessary to control myogenic tone. The component with the highest sensitivity to Ca²⁺ is the CaCC, which is activated by high nanomolar to low micromolar concentrations of Ca²⁺ and thereby elicits membrane depolarization in some types of smooth muscle (Hartzell et al., 2005). At local Ca²⁺ concentrations of 1–5 μM, RyRs on the adjacent sarcoplasmic reticular membrane open; the resultant Ca²⁺ spark can raise local Ca²⁺ to a concentration high enough (1–100 µM) to activate BK channels at physiological membrane potentials (-40 mV) and hyperpolarize the cell (Nelson et al., 1995; Jaggar et al., 2000; Pérez et al., 2001). The combined proximity and differential Ca²⁺ sensitivity of these individual elements, as well as the separation of contractile and relaxant microdomains, leads to extremely strict regulation of Ca²⁺ influx through VDCCs and subsequent changes in global cytosolic Ca²⁺ (Fig. 1).

Modeling myogenic responses in vascular smooth muscle cells

The model presented by Karlin (2015) is novel in its consideration of the individual Ca²⁺ dynamics within each microdomain, while simultaneously calculating the effects of the interaction between microdomains. This framework accurately modeled the changes in both intracellular Ca²⁺ and membrane potential in response to increases in intraluminal pressure (Knot and Nelson, 1998). It also accurately simulated the effects of BK channel and RyR blockade determined experimentally in pressurized arteries (Knot et al., 1998), suggesting




Figure 1. Arrangement of microdomains within a theoretical smooth muscle cell regulates myogenic tone. As modeled by Karlin (2015), myogenic tone development is regulated by the crosstalk between pro-contractile and pro-relaxant mechanisms that exist in close proximity within the smooth muscle cell. In the correct spatial arrangement, nonselective cation channels in stretchsensing microdomains on the cell wall respond to deformation caused by increases in intraluminal pressure, allowing influx of Na⁺ (NSC_{Na}) and Ca²⁺ (NSC_{Ca}). This leads to CaCC (Cl_{Ca}) opening, depolarization of the cell membrane (+V_M), and opening of VDCCs (Ca_V). In close proximity to stretch-sensing microdomains are pro-relaxant microdomains, formed at the interface between the junctional sarcoplasmic reticulum and the plasma membrane and containing RyRs and BK channels (BK). Increases in cytosolic Ca²⁺ cause RyR-mediated Ca²⁺ sparks that open BK channels to hyperpolarize the cell $(-V_M)$.

that the model provides a reasonable representation of the relative contributions of these two channels. The Karlin model makes some interesting predictions regarding the nature of CaCCs and TRP channels, suggesting that these two entities must be in close proximity to one another to affect membrane potential without inadvertently increasing global cytosolic Ca²⁺ to a level capable of causing maximal contraction. It also suggests that the Na⁺/Ca²⁺ exchanger, regardless of mode or location, has little or no effect on short-term myogenic responses in vascular smooth muscle cells.

As with all models, the information generated from the simulation are only as precise as the data used to generate the simulation itself. Given the large number of components included in this new model (many of which lack experimentally verified biophysical parameters), several assumptions had to be made about the identity and nature of some key components that potentially affect the model's accuracy. This is especially true with regard to the exact identity and regulation of stretch- and receptor-activated nonselective cation channels. Although they are assumed by Karlin to be members of the TRPM and TRPC subfamilies, respectively, concrete experimental evidence as to the exact nature and arrangement of these channels is still lacking (Earley and Brayden, 2015). Because of this gap in knowledge, Karlin astutely models these nonspecific cation currents in the broadest possible sense, in an attempt to maintain the robustness of the simulation while still allowing for future adaptation and clarification as new experimental evidence arises. Nonetheless, this experimental ambiguity is part of the challenge of modeling smooth muscle cell behavior, particularly as it relates to certain situations, such as predicting membrane potential and intracellular Ca²⁺ during VDCC blockade.

Overall, the model presented by Karlin (2015) represents a major step forward, as it is the first to accurately represent myogenic tone development in vascular smooth muscle. Furthermore, this model improves our understanding of intracellular Ca²⁺ microdomain-signaling dynamics as regulators of myogenic and agonist-induced vascular reactivity by enabling "experimental" manipulations that are either currently impossible or cost prohibitive. Given its reliability and robustness, Karlin's model represents a valuable new tool for testing the potential arrangement and composition of Ca²⁺-signaling microdomains in a single smooth muscle cell, helping to direct and focus future physiological investigations into vascular smooth muscle cell function.

This work was supported by grants to M.T. Nelson from the National Institute of Diabetes and Digestive and Kidney Diseases (R37 DK-053832); National Heart, Lung, and Blood Institute (R01 HL121706 and PO1 HL-095488); Fondation Leducq; and the Totman Medical Research Trust. N.R. Tykocki was also

supported by the National Heart, Lung, and Blood Institute (T32 HL007647-24) and the National Institute of Diabetes and Digestive and Kidney Diseases (K01 DK-103840-01A1).

The authors declare no competing financial interests.

Elizabeth M. Adler served as editor.

REFERENCES

- Bennett, M.R., L. Farnell, and W.G. Gibson. 2005. A quantitative description of the contraction of blood vessels following the release of noradrenaline from sympathetic varicosities. *J. Theor. Biol.* 234:107–122. http://dx.doi.org/10.1016/j.jtbi.2004.11.013
- Brayden, J.E., and M.T. Nelson. 1992. Regulation of arterial tone by activation of calcium-dependent potassium channels. *Science*. 256:532–535. http://dx.doi.org/10.1126/science.1373909
- Bulley, S., Z.P. Neeb, S.K. Burris, J.P. Bannister, C.M. Thomas-Gatewood, W. Jangsangthong, and J.H. Jaggar. 2012. TMEM16A/ANO1 channels contribute to the myogenic response in cerebral arteries. *Circ. Res.* 111:1027–1036. http://dx.doi.org/10.1161/CIRCRESAHA.112.277145
- Davis, M.J. 2012. Perspective: Physiological role(s) of the vascular myogenic response. *Microcirculation*. 19:99–114. http://dx.doi.org/10.1111/j.1549-8719.2011.00131.x
- Earley, S., and J.E. Brayden. 2015. Transient receptor potential channels in the vasculature. *Physiol. Rev.* 95:645–690. http://dx.doi.org/10.1152/physrev.00026.2014
- Hartzell, C., I. Putzier, and J. Arreola. 2005. Calcium-activated chloride channels. Annu. Rev. Physiol. 67:719–758. http://dx.doi.org/10.1146/annurev.physiol.67.032003.154341
- Hill-Eubanks, D.C., M.E. Werner, T.J. Heppner, and M.T. Nelson. 2011. Calcium signaling in smooth muscle. *Cold Spring Harb. Perspect. Biol.* 3:a004549. http://dx.doi.org/10.1101/cshperspect a004549
- Jaggar, J.H., V.A. Porter, W.J. Lederer, and M.T. Nelson. 2000. Calcium sparks in smooth muscle. Am. J. Physiol. Cell Physiol. 278:C235–C256.
- Kapela, A., A. Bezerianos, and N.M. Tsoukias. 2008. A mathematical model of Ca²⁺ dynamics in rat mesenteric smooth muscle cell: Agonist and NO stimulation. *J. Theor. Biol.* 253:238–260. http://dx.doi.org/10.1016/j.jtbi.2008.03.004
- Karlin, A. 2015. Membrane potential and Ca²⁺ concentration dependence on pressure and vasoactive agents in arterial smooth muscle: A model. *J. Gen. Physiol.* 146:79–96. http://dx.doi.org/10.1085/jgp.201511380
- Knot, H.J., and M.T. Nelson. 1998. Regulation of arterial diameter and wall [Ca²⁺] in cerebral arteries of rat by membrane potential and intravascular pressure. *J. Physiol.* 508:199–209. http://dx.doi .org/10.1111/j.1469-7793.1998.199br.x
- Knot, H.J., N.B. Standen, and M.T. Nelson. 1998. Ryanodine receptors regulate arterial diameter and wall [Ca²⁺] in cerebral arteries of rat via Ca²⁺-dependent K⁺ channels. *J. Physiol.* 508:211–221. http://dx.doi.org/10.1111/j.1469-7793.1998.211br.x
- McCarron, J.G., G. Osol, and W. Halpern. 1989. Myogenic responses are independent of the endothelium in rat pressurized posterior cerebral arteries. *Blood Vessels*. 26:315–319.
- Nelson, M.T., and J.M. Quayle. 1995. Physiological roles and properties of potassium channels in arterial smooth muscle. Am. J. Physiol. 268:C799–C822.
- Nelson, M.T., J.B. Patlak, J.F. Worley, and N.B. Standen. 1990. Calcium channels, potassium channels, and voltage dependence of arterial smooth muscle tone. Am. J. Physiol. 259:C3–C18.
- Nelson, M.T., H. Cheng, M. Rubart, L.F. Santana, A.D. Bonev, H.J. Knot, and W.J. Lederer. 1995. Relaxation of arterial smooth

- muscle by calcium sparks. Science. 270:633–637. http://dx.doi .org/10.1126/science.270.5236.633
- Parekh, A.B. 2008. Ca²⁺ microdomains near plasma membrane Ca²⁺ channels: impact on cell function. *J. Physiol.* 586:3043–3054. http://dx.doi.org/10.1113/jphysiol.2008.153460
- Pérez, G.J., A.D. Bonev, and M.T. Nelson. 2001. Micromolar Ca²⁺ from sparks activates Ca²⁺-sensitive K⁺ channels in rat cerebral artery smooth muscle. *Am. J. Physiol. Cell Physiol.* 281:C1769–C1775.
- Rubart, M., J.B. Patlak, and M.T. Nelson. 1996. Ca²⁺ currents in cerebral artery smooth muscle cells of rat at physiological Ca²⁺ concentrations. *J. Gen. Physiol.* 107:459–472. http://dx.doi.org/10.1085/jgp.107.4.459
- Welsh, D.G., A.D. Morielli, M.T. Nelson, and J.E. Brayden. 2002. Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circ. Res.* 90:248–250. http://dx.doi.org/10.1161/hh0302.105662