

The structural basis of ryanodine receptor ion channel function

Gerhard Meissner

Department of Biochemistry and Biophysics, School of Medicine, University of North Carolina, Chapel Hill, NC

Large-conductance Ca^{2+} release channels known as ryanodine receptors (RyRs) mediate the release of Ca^{2+} from an intracellular membrane compartment, the endo/sarcoplasmic reticulum. There are three mammalian RyR isoforms: RyR1 is present in skeletal muscle; RyR2 is in heart muscle; and RyR3 is expressed at low levels in many tissues including brain, smooth muscle, and slow-twitch skeletal muscle. RyRs form large protein complexes comprising four 560-kD RyR subunits, four ~ 12 -kD FK506-binding proteins, and various accessory proteins including calmodulin, protein kinases, and protein phosphatases. RyRs share $\sim 70\%$ sequence identity, with the greatest sequence similarity in the C-terminal region that forms the transmembrane, ion-conducting domain comprising ~ 500 amino acids. The remaining $\sim 4,500$ amino acids form the large regulatory cytoplasmic “foot” structure. Experimental evidence for Ca^{2+} , ATP, phosphorylation, and redox-sensitive sites in the cytoplasmic structure have been described. Exogenous effectors include the two Ca^{2+} releasing agents caffeine and ryanodine. Recent work describing the near atomic structures of mammalian skeletal and cardiac muscle RyRs provides a structural basis for the regulation of the RyRs by their multiple effectors.

Introduction

The release of Ca^{2+} ions from intracellular membrane-bound stores is a key step in a wide variety of biological functions. The release of Ca^{2+} is predominantly mediated by two related Ca^{2+} release channel families: the ryanodine receptors (RyRs; Meissner, 1994; Franzini-Armstrong and Protasi, 1997; Hamilton and Serysheva, 2009; Clarke and Hendrickson, 2016) and inositol 1,4,5-trisphosphate receptors (Patterson et al., 2004; Seo et al., 2015; Baker et al., 2017). Both types of Ca^{2+} release channels are localized in the ER and in muscle in a specialized subcompartment, the SR. Other intracellular organelles that store and release Ca^{2+} include mitochondria and acidic endosomal structures. One distinct feature of the RyRs is their modification by the plant alkaloid ryanodine, hence their name, which distinguishes them from other Ca^{2+} channels. The RyRs are comprised of four polypeptides each of $\sim 5,000$ amino acids and four FK506-binding proteins (FKBPs) each of ~ 110 amino acids. RyRs are high-conductance, monovalent- and divalent-conducting channels regulated by multiple factors that include Ca^{2+} , Mg^{2+} , ATP, calmodulin (CaM), protein kinases and phosphatases, and redox active species. Recent determination of the high-resolution cryo-electron microscopy (cryo-EM) structure of the skeletal muscle RyR1 and cardiac muscle RyR2 isoforms has provided insight into the complex interacting regulatory mechanisms of SR Ca^{2+} release

(Efremov et al., 2015; Yan et al., 2015; Zalk et al., 2015; Bai et al., 2016; des Georges et al., 2016; Peng et al., 2016; Wei et al., 2016). RyR-like structures were identified in colony-forming choanoflagellates that evolved more than 600 million years ago (Mackrill, 2012). Alterations in the activity of the RyRs have been implicated in a number of muscle diseases. More than 100 RyR1 mutations are potentially associated in skeletal muscle with malignant hyperthermia (MH), central core disease (CCD), and multi-minicore disease (MmD; Robinson et al., 2006; Treves et al., 2008), and more than 150 RyR2 mutations are potentially linked to catecholaminergic polymorphic ventricular tachycardia (CPVT), arrhythmogenic right ventricular dysplasia type 2 (ARVD2), and idiopathic ventricular fibrillation in cardiac muscle (George et al., 2007; Leenhardt et al., 2012). Here, the experimental and recent structural studies that led to our current understanding of RyR regulatory mechanisms and their structural basis are reviewed.

Mechanisms of SR Ca^{2+} release in mammalian skeletal and cardiac muscle

Rapid release of Ca^{2+} in striated muscle is triggered by an action potential that spreads rapidly over the surface membrane into tubular infoldings (T-tubule) of the surface membrane. A distinguishing feature is that cardiac muscle excitation–contraction (E-C) coupling depends on extracellular Ca^{2+} , whereas skeletal muscle E-C coupling does not. This finding led to the formulation of two principal mechanisms for E-C coupling: the Ca^{2+} -induced Ca^{2+} release (CICR) hypothesis in heart (Fabiato, 1983) and the mechanical coupling hypothesis

Correspondence to Gerhard Meissner: meissner@med.unc.edu

Abbreviations used: ARVD, arrhythmogenic right ventricular dysplasia; Bsol, bridging solenoid; CaM, calmodulin; CaMK, Ca^{2+} /CaM-dependent protein kinase; CCD, central core disease; CHAPS, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate; CPVT, catecholaminergic polymorphic ventricular tachycardia; cryo-EM, cryo-electron microscopy; Csol, central solenoid; CSQ, calsequestrin; DHPR, dihydropyridine receptor; E-C, excitation–contraction; FKBP, FK506-binding protein; FRET, fluorescence resonance energy transfer; GSH, glutathione; GSSG, GSH disulfide; Jsol, junctional solenoid; MH, malignant hyperthermia; MmD, multi-minicore disease; RyR, ryanodine receptor.

© 2017 Meissner This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms/>). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 4.0 International license, as described at <https://creativecommons.org/licenses/by-nc-sa/4.0/>).



in vertebrate skeletal muscle (Schneider and Chandler, 1973; Ríos and Pizarro, 1991). Voltage- and dihydropyridine-sensitive (L-type) Ca^{2+} channels (dihydropyridine receptors [DHPRs], $\text{Ca}_v1.2$ s) located in the surface membrane and T-tubule in cardiac muscle open closely apposed SR Ca^{2+} release channels (RyR2s) that result in the release of massive amounts of Ca^{2+} . In contrast, in mammalian skeletal muscle, voltage-sensing L-type Ca^{2+} channels (DHPRs, $\text{Ca}_v1.1$ s) in the T-tubule open closely apposed SR Ca^{2+} release channels (RyR1s) through direct protein–protein interactions.

Structural arrangement of RyRs in skeletal and cardiac muscle

Skeletal and cardiac muscle SRs have longitudinal (free) and junctional SRs that are structurally and functionally distinct membrane regions. Longitudinal SR is a sarcotubular network that surrounds the myofibrils and contains a Ca^{2+} pump responsible for Ca^{2+} uptake by SR. The junctional SR is mainly involved in Ca^{2+} release and houses RyR ion channels near L-type Ca^{2+} channels in the plasmalemma (RyR2) and T-tubule (RyR1 and RyR2; Franzini-Armstrong and Protasi, 1997). Groupings of L-type Ca^{2+} channels and RyRs in so-called Ca^{2+} release units or couplons comprise 10–100 RyR1 in fast- and slow-twitch rat, guinea pig, frog, and toadfish skeletal muscle (Franzini-Armstrong et al., 1999). Large protein structures termed “feet” represent the large cytoplasmic domains of the RyR1s that span the narrow gap separating the two membranes. Morphological evidence suggests a well-defined interaction between T-tubular $\text{Ca}_v1.1$ and SR RyR1 channels in skeletal muscle. Clusters of four particles, tetrads representing four $\text{Ca}_v1.1$ s, are located opposite four subunits of every other RyR1 (Block et al., 1988). The formation of $\text{Ca}_v1.1$ tetrads depends on the presence of apposing RyR1s (Takekura et al., 1995; Protasi et al., 1998). However, surface membrane–SR junctions were observed in mouse skeletal muscle primary cultures lacking RyR1. This suggests that other proteins are involved in the close linkage of the two membrane systems. SR junctional proteins that assist in the formation of the triad junction include triadin (Caswell et al., 1991; Guo and Campbell, 1995), junctin (Jones et al., 1995), mitsugumin 29 (Takeshima et al., 1998), and STAC3 (Campiglio and Flucher, 2017).

Two populations of RyR1s in skeletal muscle, one coupled to tetrads and one not, raised the question of how unlinked RyRs are activated. One suggestion was that Ca^{2+} released by $\text{Ca}_v1.1$ -linked RyR1s activates $\text{Ca}_v1.1$ -unlinked RyR1s by a Ca^{2+} -induced Ca^{2+} release mechanism seen in cardiac muscle (Ríos and Pizarro, 1991). However, CICR is slow in the presence of ATP and Mg^{2+} compared with physiological Ca^{2+} release (review by Endo, 2009). Ca^{2+} did not trigger the release of Ca^{2+} in mouse permeabilized skeletal muscle in the presence

of physiological concentrations of ATP and Mg^{2+} , which also suggested that Ca^{2+} is not the principal activator of RyR1 (Figuerola et al., 2012). On the other hand, it was proposed that Ca^{2+} may terminate the release of Ca^{2+} by binding to Ca^{2+} inactivation sites. Posterino and Lamb (2003) observed an increase in SR Ca^{2+} release from skinned rat muscle fibers in the presence of $\sim 400 \mu\text{M}$ 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid tetrakis(acetoxymethyl ester) (BAPTA), a fast Ca^{2+} -chelating agent. Sztretye et al. (2011) noted that BAPTA changed the amplitude and kinetics of Ca^{2+} release in mammalian fibers. An alternative mechanism is that neighboring RyR1s are physically and functionally linked, which leads in lipid bilayers to the simultaneous opening and closing of two or more channels, termed coupled gating (Marx et al., 1998).

Clusters of tetrads of $\text{Ca}_v1.1$ s in skeletal muscle located opposite the four subunits of every other RyR1 resulted in a $\text{Ca}_v1.1/\text{RyR1}$ ratio of 1.4:1.9 (Bers and Stiffel, 1993). A ratio of less than two suggested that not all tetrads were complete, as observed in freeze-fracture studies (Protasi et al., 1997).

The location and arrangement of DHPRs and RyRs in cardiac muscle is more diverse. The $\text{Ca}_v1.2/\text{RyR2}$ ratio is lower, ranging from 0.27 in rabbit to 0.1 in ferret myocytes (Bers and Stiffel, 1993). SR junctional domains containing closely packed groupings of RyR2s that range in size from a few to several 100 RyR2s form discrete junctional domains with the surface membrane (peripheral couplings) and T-tubule (diads). In cardiac cells lacking T-tubules (e.g., conduction cells, avian hearts), junctional SR is near the surface membrane and in extended junctional SR away from the surface membrane (Sommer and Johnson, 1979). Frog ventricle contains few if any RyRs, which suggests a reliance on calcium entry through surface membrane Ca^{2+} channels for muscle contraction (Tijssens et al., 2003).

The third mammalian RyR isoform RyR3 is expressed in diaphragm and slow-twitch skeletal muscle (Conti et al., 1996), brain and smooth muscle (Furuiuchi et al., 1994; Ledbetter et al., 1994; Giannini et al., 1995), and fast-twitch skeletal muscle of young but not adult mice (Flucher et al., 1999). Two immunologically distinct high molecular weight RyR protein bands corresponding to mammalian RyR1 and RyR3 were identified in skeletal muscles from chicken (Airey et al., 1990) and frog (Oyamada et al., 1994). RyR3s in frog skeletal muscle and toadfish swim bladder are segregated in a parajunctional position adjacent to the junctional SR, where they are activated by calcium released from RyR1 channels (Felder and Franzini-Armstrong, 2002). Consistent with a low RyR3/RyR1 ratio, parajunctional feet are less frequently seen in mouse skeletal muscle and diaphragm. The appearance of the RyR1 isoform alone in some fast-contracting muscle of fish suggests that RyR1 is selectively expressed when

rapid contractions are required in muscle (O'Brien et al., 1993). Spontaneous or evoked activation of a few RyRs results in spatially limited Ca^{2+} release events, known as Ca^{2+} sparks (Cheng et al., 1993). Ca^{2+} sparks are observed in skeletal muscle that express RyR3, and in cardiac muscle.

Preparations for studying RyR function

RyR function has been studied using intact and skinned fibers, fragmented SR, and purified RyR preparations. There are two ways of accessing the intracellular compartment in muscle cells without disrupting the SR membrane (review by Stephenson, 1981). One is to mechanically remove the surface membrane from intact skeletal muscle cells. This procedure consists of physically rolling back a portion of the sarcolemma of a single fiber (Natori, 1954). T-tubules will reseal in mechanically skinned fiber preparations. SR Ca^{2+} release and contraction can therefore be elicited by depolarizing T-tubule by ion substitution (Donaldson, 1985; review by Lamb, 2002). Fabiato and Fabiato (1975) studied the mechanism of CICR in cardiac muscle by removing the surface membrane from fragments of single cardiac cells by microdissection. To study SR Ca^{2+} release in the absence of T-tubule depolarization, muscle cells were treated with low concentrations of saponin, a detergent that disrupts cholesterol-containing membranes (surface membrane and T-tubule) but not membranes that lack cholesterol (SR). Skinned fiber studies were performed using mammalian skeletal muscle and amphibian skeletal muscle preparations that contain the RyR3 isoform, in addition to RyR1. These studies identified three principal properties of the RyRs: SR Ca^{2+} release is stimulated by Ca^{2+} and adenine nucleotides and inhibited by Mg^{2+} .

Fragmentation of SR during homogenization of rabbit skeletal muscle and subsequent fractionation by differential and density centrifugation yielded "light" and "heavy" SR vesicles according to sedimentation properties (Meissner, 1975). Light vesicles correspond to the longitudinal SR and contain the SR Ca^{2+} pump as the major protein. Heavy SR vesicles derived from the junctional region of the SR contain " Ca^{2+} release channels" that are activated by Ca^{2+} and adenine nucleotides and inhibited by Mg^{2+} (Meissner, 1984; Meissner et al., 1986). Another important advance was the isolation of membrane fractions enriched in T-tubule segments sandwiched between two junctional SR compartments (Mitchell et al., 1983). These junctional complexes are known as triads. Microsomal membrane fractions enriched in ryanodine-sensitive Ca^{2+} release channels were also isolated from other excitable tissues, such as cardiac muscle, smooth muscle, and brain. In these cases, however, the membrane fractions were typically of a lower purity than those from rabbit skeletal muscle.

Purification and identification of RyRs

Isolation of the SR Ca^{2+} release channel is facilitated by the identification of ryanodine as a channel-specific ligand (Fleischer et al., 1985; Pessah et al., 1985; Sutko et al., 1985; Meissner, 1986). Ryanodine is toxic and has different pharmacological effects depending on muscle type and activity. Ryanodine can increase or decrease the contractile force (Jenden and Fairhurst, 1969). Because the drug binds with high specificity and dissociates slowly from the high-affinity site in the receptor (either membrane-bound or detergent-solubilized), [^3H]ryanodine is an ideal probe to monitor the isolation of RyRs from a variety of tissues and species.

RyRs were initially isolated from rabbit skeletal muscle SR membrane preparations as a spanning protein complex comprised of 300-kD subunits (Kawamoto et al., 1986) and multiprotein ryanodine-binding complexes that displayed a pharmacology (Pessah et al., 1986, 1987; Inui et al., 1987b; Lai et al., 1987) and morphology characteristic of the feet structures in junctional SR (Inui et al., 1987b; Lai et al., 1988a). Insertion of rabbit skeletal muscle (Smith et al., 1985, 1986) and canine cardiac muscle (Rousseau et al., 1986) SR vesicles and purified [^3H]ryanodine-binding protein complexes (Imagawa et al., 1987; Lai et al., 1988b; Smith et al., 1988; Anderson et al., 1989) in lipid bilayers showed that RyR1 and RyR2 are Ca^{2+} - and ATP-activated, monovalent and divalent ion-conducting channels composed of four ~565-kD polypeptides. The high molecular weight RyR polypeptide was associated with isoform-specific immunophilins (FKBPs) that migrated on SDS acrylamide gels as 12-kD proteins (Collins, 1991; Timerman et al., 1996).

Cloning and sequencing rabbit, human, and pig RyR1 (Takeshima et al., 1989; Zorzato et al., 1990; Fujii et al., 1991), rabbit and human cardiac RyR2 (Nakai et al., 1990; Otsu et al., 1990; Tunwell et al., 1996), and rabbit RyR3 (Hakamata et al., 1992) revealed that the mammalian RyRs are a family of closely related Ca^{2+} -conducting channels composed of four polypeptides of ~5,000 amino acids each.

Structure of mammalian RyRs

The morphology of purified RyRs was examined initially by negative-staining electron microscopy (Kawamoto et al., 1986; Inui et al., 1987a,b; Lai et al., 1988b; Saito et al., 1988; Anderson et al., 1989). Mammalian RyR1 and RyR2 have large square, four-leaf clover structures that resemble the feet that span the T-tubule-SR gap. A more detailed understanding of the structure of the intact RyRs was revealed by cryo-EM, a method well suited to resolve the structure of macromolecular complexes in the near-native conformation. Three-dimensional reconstruction from electron micrograph data of frozen-hydrated RyR specimens at a resolution of ~30 Å revealed a $29 \times 29 \times 12$ -nm large cytoplasmic assembly with a 7-nm-length and 8-nm-diameter transmem-




Figure 1. **Open RyR1 channel structure.** The structure (PDB code 5TAL) reveals the major domains and the location of Ca^{2+} , ATP-, and caffeine-binding sites.

brane domain (Wagenknecht et al., 1989; Radermacher et al., 1994; Serysheva et al., 1995). Frozen hydrated specimens at $\sim 10\text{-}\text{\AA}$ resolution indicated that the closed (nonconducting; Ludtke et al., 2005; Samsó et al., 2005) and open (conducting; Samsó et al., 2009) RyR1 has a pore structure characteristic of the voltage-gated ion channel family. Along with revealing the global structure, cryo-EM in combination with fluorescence resonance energy transfer (FRET) measurements identified regulatory sites on RyR1 and RyR2 such as the FKBP site (Wagenknecht et al., 1996; Sharma et al., 2006; Gengenrath et al., 2013) and sites of the Ca^{2+} -free and Ca^{2+} -bound forms of CaM (Wagenknecht et al., 1994; Samsó and Wagenknecht, 2002; Huang et al., 2012).

Recent cryo-EM studies provide more detailed insights into the structure of intact RyRs. The closed and open pore structures of RyR1 and RyR2 were determined at 3.6- to $\sim 6\text{-}\text{\AA}$ resolution (Efremov et al., 2015; Yan et al.,

2015; Zalk et al., 2015; Bai et al., 2016; des Georges et al., 2016; Peng et al., 2016; Wei et al., 2016). More than two thirds of the molecular mass of RyRs was resolved. SR membranes from rabbit skeletal muscle (Efremov et al., 2015; Yan et al., 2015; Zalk et al., 2015; Bai et al., 2016; des Georges et al., 2016; Wei et al., 2016) and porcine cardiac muscle (Peng et al., 2016) were solubilized using the zwitterionic detergent 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS). RyRs were purified on sucrose gradients or by affinity chromatography using glutathione-transferase-tagged FKBP12 and FKBP12.6. Closed RyR2 channels were applied to grids in buffers containing 5 mM EDTA and the nonionic detergent digitonin (Peng et al., 2016). To obtain open channels, RyR2 was transferred to a buffer containing 20 μM Ca^{2+} and 20 μM PCB95, a RyR activator. Closed RyR1 structures were determined in buffers containing EGTA and Tween 20 (Yan et al., 2015) or CHAPS and a

Table 1. Cryo-EM domain nomenclature of RyR1

| Domain name | Alternative names | Color ^a | Sequence ^b |
|-------------------------|-------------------|--------------------|-----------------------|
| N-terminal domain | | Yellow | 1–627 |
| SPRY1 | | Light orange | 628–849 |
| RY1 and 2 | P1 | Light magenta | 850–1,054 |
| SPRY2 and 3 | | Light orange | 1,055–1,656 |
| Jsol | Handle | Green | 1,657–2,144 |
| Bsol | HD1&2 | Cyan | 2,145–3,613 |
| Includes RY3 and 4 | P2 | Orange | 2,735–2,938 |
| Csol | Central domain | Tomato | 3,667–4,174 |
| Includes EF hand motifs | | | 4,060–4,134 |
| TaF | | | 4,175–4,253 |
| Transmembrane domain | | Blue | 4,541–4,956 |
| C-terminal domain | | Brown | 4,957–5,037 |

^aDomain color in Fig. 1.^bdes Georges et al. (2016).

low concentration of lipid (Zalk et al., 2015; des Georges et al., 2016). To obtain closed and open RyR1s, Bai et al. (2016) applied the purified RyR1 channel complex to grids in a buffer that contained Tween 20 or CHAPS, respectively. Wei et al. (2016) determined the structure of the closed RyR1 channel in a buffer containing amphipol A8-35, an amphipathic surfactant that stabilizes and maintains solubilized membrane proteins in detergent-free solutions. The open RyR1 channel was maintained in the presence of 100 μ M Ca^{2+} and 10 μ M ruthenium red, an RyR channel blocker. With some exceptions, retention of function of the purified RyRs was not determined. Four different channel states were obtained according to single-channel measurements that kept RyR1 in a closed state (nominal 0 μ M Ca^{2+}), partially open state (30 μ M Ca^{2+} ; 0 μ M Ca^{2+} , 2 mM ATP, and 5 mM caffeine), or fully open state (30 μ M Ca^{2+} , 2 mM ATP, and 5 mM caffeine; Fig. 1). Addition of 10 μ M ryanodine converted channels to an open subconducting state (des Georges et al., 2016).

Determination of the high-resolution closed- and open-channel structures provides a better understanding of the mechanisms of channel gating and ion permeation and how they are altered in RyR-associated diseases. Each RyR subunit has been subdivided into ~10 domains (Fig. 1 and Table 1). These include (1) the N-terminal domain that is a hotspot of RyR1 and RyR2 disease mutations, (2) the SPRY1 domain that is part of the binding site for FKBP12, (3) the junctional solenoid (Jsol) and bridging solenoid (Bsol), where the latter contains PKA and Ca^{2+} /CaM-dependent protein kinase II (CaMKII) phosphorylation sites, (4) the central solenoid (Csol) domain that contains two EF hand Ca^{2+} binding motifs, (5) the transmembrane domain that forms the conductance pathway for Ca^{2+} , and (6) the C-terminal domain (CTD) that forms part of the Ca^{2+} , ATP, and caffeine activation sites. Crystal structures of smaller RyR fragments at resolutions of <2 Å and covering N-terminal, SPRY1, SPRY2, RY1 and 2, and RY3 and 4 domains have clarified the functional

role of ligands and disease mutations (Yuchi and Van Petegem, 2016). A comprehensive description of the RyR1 domain structure is provided by des Georges et al. (2016) and Samsó (2017).

Ca_v1.1–RyR1 interactions

A unique property of the mammalian skeletal muscle RyR1 is its direct linkage to the skeletal muscle Ca_v1.1 channel isoform, which acts in E-C coupling as a voltage-sensing molecule rather than as a Ca^{2+} channel. Ca_v1.1 is comprised of a pore-forming α_1 subunit and $\alpha_2\delta$, β , and γ subunits. Early death of dysgenic mice lacking the α_1 subunit (Beam et al., 1986) and β -null mutant mice (Gregg et al., 1996) suggested that both subunits are required for skeletal muscle E-C coupling. Several regions of Ca_v1.1 α_1 and β subunits interacted with RyR1 and vice versa (review by Bannister, 2016). Detailed structural information on the interaction between the intact Ca_v1.1 and RyR1 is lacking, in part because a Ca_v1.1–RyR1 complex has not been isolated. However, obtaining this structural information should be facilitated by the availability of the recently determined near-atomic resolution structure of the intact RyR1 (Yan et al., 2015; Zalk et al., 2015; Bai et al., 2016; des Georges et al., 2016; Wei et al., 2016) and Ca_v1.1 (Wu et al., 2016) complexes.

RyRs are high-conductance, cation-selective channels

Fusion of heavy rabbit skeletal muscle vesicles with planar lipid bilayers provided the initial evidence that the SR membrane contains a calcium-conducting channel of 125 pS (Smith et al., 1985). Channel activities recorded in the presence of Ca^{2+} , ATP, and ruthenium red were compared with results obtained in vesicle flux measurements according to a rapid quench protocol. Results indicated that the 125-pS channel mediates rapid Ca^{2+} - and ATP-gated release of Ca^{2+} from skeletal muscle SR vesicles. Subsequent lipid bilayer studies with canine cardiac muscle SR vesicles (Rousseau et al., 1986) and purified [³H]ryanodine-binding protein complexes

(Imagawa et al., 1987; Lai et al., 1988b; Smith et al., 1988; Anderson et al., 1989) established that RyR1 and RyR2 are Ca^{2+} - and ATP-activated, Mg^{2+} -inhibited, monovalent and divalent ion-conducting channels with a $\text{P}_{\text{Ca}}/\text{P}_{\text{K}}$ ratio of ~ 7 – 8 .

Sequence comparisons and mutagenesis studies provided evidence that the RyRs have a pore structure similar to K^+ channels (Balshaw et al., 1999; Zhao et al., 1999; Gao et al., 2000; Du et al., 2001). RyR amino acid residues between the two C-terminal S5 and S6 membrane-spanning segments were suggested to be lumenally located, with a pore helix and an amino acid motif (GGGIG) similar to the selectivity filter motif (TV/IGYG) of K^+ channels. However, there are important differences among the RyRs relative to K^+ channels. One is that RyRs conduct both monovalent and divalent cations with the following relative specificity: $\text{K}^+ > \text{Rb}^+ > \text{Na}^+ \approx \text{Cs}^+ > \text{Li}^+$, and $\text{Ba}^{2+} > \text{Sr}^{2+} > \text{Ca}^{2+} > \text{Mg}^{2+}$ (Lindsay et al., 1991; Tinker and Williams, 1992). A second difference is that single-channel measurements with permeant and impermeant organic cations suggest a wider minimum radius of 3.3 – 3.5 Å for RyR2 (Tinker and Williams, 1993) than the experimentally determined minimum radius of 1.5 Å for K^+ channels (Hille, 1973). Mutagenesis studies showed that two negatively charged amino acid residues (RyR1-D4899 and E4900) immediately after the selectivity filter motif in the luminal vestibule and two negatively charged residues (RyR1-D4938 and D4945) in the cytosolic vestibule have a critical role in RyR1 ion permeation and selectivity (Wang et al., 2005; Xu et al., 2006).

Construction of a RyR1- $\Delta 183$ – 4006 deletion mutant and C-terminal construct of 1,377 amino acid residues showed that the RyR1 C-terminal portion forms a monovalent conducting channel activated by Ca^{2+} and modified by ryanodine (Bhat et al., 1997). The two C-terminal transmembrane segments that included the pore helix and connecting loops formed a homotetrameric assembly that conducted K^+ and Ca^{2+} and also Cl, which suggests the loss of a cation-specific pathway (Eudén et al., 2013b).

Cryo-EM data support the view that RyR1 has a pore structure characteristic of the voltage-gated channel family. Samsó et al. (2005) and Ludtke et al. (2005) determined the pore structure of the closed (nonconducting) RyR1 at a resolution of ~ 10 Å. The pore-forming region as visualized by Ludtke et al. (2005) consisted of a long inner helix comprised of 31 amino acid residues and a pore helix of 15 residues. An increased resolution of ~ 4 Å showed that Ile4937 in the S6 pore lining helix forms the hydrophobic constriction site in the closed RyR1 with a pore radius of less than 1 Å, rendering the channel impermeable to Ca^{2+} (Yan et al., 2015; Zalk et al., 2015). In the open RyR2 state, the pore constriction site was shifted to Gln4864 (corresponding to Gln4933 in RyR1), widening the RyR2 minimal pore radius to

~ 2 Å (Peng et al., 2016). Ion-pulling simulations generated an open-channel conformation of RyR1 (Mowrey et al., 2017) from the 3.8 -Å closed state of RyR1 (Yan et al., 2015). The pore constriction site at RyR1-Gln4933 had a minimum open-channel pore radius of ~ 2.8 Å (Mowrey et al., 2017), in reasonable agreement with the experimentally derived minimum radius of 3.3 – 3.5 Å (Tinker and Williams, 1993). The cryo-EM structure and model data suggest that channel opening involves a rotation of the upper portion of the pore-lining S6 helix away from the fourfold channel axis. Wei et al. (2016) used the RyR channel blocker ruthenium red to obtain an open channel. At variance with the aforementioned studies, Ile4937 in the closed RyR1 conferred a pore radius of 2.5 Å, which increased to 4.9 Å in the “open ruthenium-red-stabilized” channel, allowing the passage of hydrated Ca^{2+} ions.

The RyR1 S6 pore lining helix has two conserved hinge glycines (Gly4934 and Gly4941) associated with channel opening. Determination of the pore structure of the closed (nonconducting) RyR1 at a resolution of ~ 10 Å suggested that the inner S6 helix has a bend at RyR1-Gly4934 (Ludtke et al., 2005). Samsó et al. (2009) determined the cryo-EM structure of both the open and closed RyR1. As observed for the high-resolution structures of K^+ channels (Jiang et al., 2002), the inner helix had an outward bend on a glycine (Gly4934 or Gly4941) in the open but not closed channel. This change along with other cytoplasmic structural changes widened the radius of a cytoplasmic ion gate from ~ 4 to ~ 6 Å, allowing the flow of ions through the channel. Because cryo-EM had a resolution of only ~ 10 Å, the positions of pore residue side chains and the structure of loops connecting the helices remained unknown (Ludtke et al., 2005; Samsó et al., 2009).

The two conserved glycines were replaced with uncharged amino acid residues of an increased side chain volume. Substitution of RyR2-Gly4864 (corresponding to RyR1-Gly4934) with alanine resulted in no significant change of RyR2 function, whereas replacement with valine and proline profoundly altered channel gating and ion permeation (Eudén et al., 2013a). Replacement of RyR1-Gly4934 and Gly4941 with alanine altered RyR1 channel function in single-channel measurements (Mei et al., 2015). Mutations further increasing the side chain volume at these positions (RyR1-G4934V and G4941I) resulted in loss of function. Molecular modeling suggested that the two glycines facilitate RyR channel function by providing flexibility and minimizing amino acid clashes.

A model of RyR2 ion permeation based on Eyring rate theory postulated four potential barriers within the pore corresponding to three putative binding sites (Tinker et al., 1992). Without knowledge of the structure of the pore, the model quantitatively reproduced the conductance data of mono- and divalent cations. Cryo-EM has tentatively assigned two barriers corresponding to




Figure 2. Ca^{2+} dependence of RyR1. Data obtained using the planar lipid bilayer and [³H]ryanodine binding methods. With modifications from Heiny and Meissner (2012).

the selectivity filter and open-channel RyR2-Gln4864 residue. A Poisson–Nernst–Planck density functional theory model accurately modeled the role of negatively charged residues in the luminal (RyR1-Asp4899 and Glu4900; Wang et al., 2005) and cytosolic (RyR1-Asp4938 and Asp4945; Xu et al., 2006) vestibules of RyR1 in generating the high ion conductances of RyRs (Gillespie et al., 2005). Selectivity was attributed to charge-space competition, as Ca^{2+} could accommodate the most charge in the least space compared with K^+ (Gillespie et al., 2005; Gillespie, 2008). Although the large dehydration energy but high conductance of Mg^{2+} suggest that ion dehydration does not play a major role in RyR ion permeation (Gillespie et al., 2014), the exact hydration states of Ca^{2+} ions remain to be determined as they pass through the pore. Gillespie and Fill (2008) predicted that RyR ion channels must mediate their own K^+ countercurrents to minimize the formation of a

membrane potential during SR Ca^{2+} release that would otherwise rapidly impede further release of Ca^{2+} .

Regulation of RyRs by Ca^{2+}

The mechanism of skeletal muscle and cardiac SR Ca^{2+} release has been extensively studied using isolated SR membrane and purified RyR preparations by applying three complementary methods. Macroscopic ion fluxes were measured from actively or passively loaded SR vesicles using rapid flow, quench, and filtration protocols. Microscopic monovalent ion and Ca^{2+} currents were recorded through single Ca^{2+} release channels incorporated into lipid bilayers (Fig. 2). In a widely used but less direct method, RyR channel activity was probed using the specific plant alkaloid ryanodine (Sutko et al., 1997; Fig. 2). All three methods showed that RyRs are activated by micromolar Ca^{2+} and millimolar ATP and are inhibited by millimolar Mg^{2+} and Ca^{2+} .




Figure 3. Ca^{2+} dependence of single purified RyR1, RyR2, and RyR3. Shown are channel P_o values as a function of cytosolic free Ca^{2+} in 250 mM K^+ , pH 7.4 medium. Redrawn from Xu et al. (1998); (RyR1 and RyR2) and Chen et al. (1997); (RyR3).

Evidence for cytosolic RyR sites involved in Ca^{2+} activation. RyRs are activated by micromolar cytosolic Ca^{2+} and inhibited by millimolar cytosolic Ca^{2+} in the absence of other channel effectors. A biphasic behavior implies there are at least two classes of Ca^{2+} -binding sites: a high-affinity activation site and a low-affinity inactivation site. However, differences in Ca^{2+} regulation of the mammalian RyR isoforms have been observed. In single-channel measurements, RyR2 and RyR3 isoforms are activated to a greater extent by cytosolic Ca^{2+} than RyR1 and require higher cytosolic Ca^{2+} concentrations to be inhibited when Ca^{2+} is the sole activating ligand (Fig. 3). Similarly, in avian and amphibian skeletal muscle that express both mammalian RyR1 and RyR3 homologues, RyR3 is activated by Ca^{2+} to a greater extent than RyR1 (Percival et al., 1994; Murayama and Ogawa, 2004).

Comparison of cryo-EM maps prepared in the absence or presence of 30 μM Ca^{2+} revealed a Ca^{2+} -binding site at the interdomain interfaces of the RyR1 Csol and C-terminal domains (des Georges et al., 2016). The site encompasses three amino acids conserved among the RyR and IP_3R families directly interacting with Ca^{2+} , RyR1-Glu3893, and Glu3967 of the Csol and Thr5001 of the C-terminal domain (Fig. 4 A). Two amino acids, Gln3970 and His3895, indirectly interact with Ca^{2+} . The channel apparently remains closed with 30 μM Ca^{2+} as the sole activating ligand, even though single-channel measurements showed a channel open probability (P_o) of 0.22. Similarly, opposite results were obtained using ATP and caffeine alone, whereas the three ligands Ca^{2+} , ATP, and caffeine opened the channel in the structural and single-channel studies. des Georges et al. (2016)

proposed that Ca^{2+} or ATP/caffeine alone primed the channel for opening, but all three ligands were required to overcome a barrier for opening. An alternative explanation is that the number of open channels under the conditions of Ca^{2+} or ATP/caffeine was too low to be scored as a separate class.

Mutagenesis has indicated that additional regions are involved in the regulation of RyRs by Ca^{2+} . In one early study, RyR1- $\Delta 183$ -4006 deletion mutant was activated by micromolar Ca^{2+} , which suggests a Ca^{2+} -binding site different from the one identified by cryo-EM (Bhat et al., 1997). Substitution of RyR1-Glu4032 by alanine and the corresponding amino acid residue in the RyR3 isoform reduced Ca^{2+} -gated channel activity (Chen et al., 1998; Du and MacLennan, 1998; Fessenden et al., 2001). However, RyR1-Glu4032 was not a part of a Ca^{2+} -binding site, but rather stabilized the interaction between two RyR1 regions (des Georges et al., 2016). Other examples include CCD mutations located in the pore-forming region of RyR1 in skeletal muscle (McCarthy et al., 2000; Dirksen and Avila, 2002; Xu et al., 2008), sites in S2 (Gomez and Yamaguchi, 2014), S4-S5 linker (Murayama et al., 2011; Ramachandran et al., 2013), N-terminal region of RyR2 (Liu et al., 2015), and S6 of RyR2 (Sun et al., 2016). Taken together, both structural and functional studies indicate that the transition of the closed to Ca^{2+} -activated open channel involves the coordinated motion of multiple regions outside the Ca^{2+} -binding site.

Evidence for cytosolic RyR sites involved in Ca^{2+} inactivation. Efremov et al. (2015) compared two RyR1 cryo-EM structures in the absence of Ca^{2+} and at 10 mM Ca^{2+} at a




Figure 4. **Structure of regulatory sites in RyR1.** Ca^{2+} (A)-, ATP (B)-, and caffeine (C)-binding sites of open RyR1 (PDB code 5TAL).

resolution of 6.1 Å. Differences in the Ca^{2+} -free and Ca^{2+} -bound maps were interpreted to indicate that binding of Ca^{2+} to a two-EF hand motif in the Csol domain (Fig. 1 and Table 1) increased the mean gate diameter, which indicated formation of an open channel. However, functional studies were not presented. Moreover, the addition of 10 mM Ca^{2+} results in a nearly fully closed and not open RyR1 channel (Fig. 3). Therefore, it seems that Ca^{2+} ions bound to sites involved in Ca^{2+} inactivation and not activation of RyR1.

Earlier [^3H]ryanodine binding studies with RyR1/RyR2 chimeras suggested that Ca^{2+} inactivation sites are located in the C-terminal quarter of RyR1 (Du and MacLennan, 1999). In this region, two EF1 and EF2 hand motifs are located in a CaM-like domain of RyR1 (amino acids 4,064–4,210; Xiong et al., 2006). A peptide corresponding to the CaM-like domain bound Ca^{2+} with an apparent affinity of 60 μM (Xiong et al., 2006). Two smaller recombinant proteins from rabbit skeletal muscle RyR1 (amino acids 4,069–4,139), cardiac muscle RyR2 (amino acids 4,025–4,095), and lobster skeletal muscle containing the two EF hand motifs bound two Ca^{2+} with millimolar affinities, implying that sites in these regions have an inhibitory role (Xiong et al., 1998). Deletion and mutations of the two-EF hand motif in the full-length RyR2 suggested that the EF-hand domain has a role in RyR2 regulation by luminal Ca^{2+} but is not required for activation by cytosolic Ca^{2+} (Guo et al., 2016). Fessenden et al. (2004) assessed the functional significance of five potential Ca^{2+} regulatory sites in RyR1, taking into account that, in addition to the two-EF hand motifs, there are three negatively charged EF-hand-like sequences encompassing amino acid residues 4,253–4,264, 4,407–4,416, and 4,489–4,499. Mutagenesis of the five motifs in RyR1 did not reveal major functional differences in response to depolarization or caffeine compared with wild-type RyR1-expressing myotubes (Fessenden et al., 2004). However, in a binding assay using the RyR specific ligand [^3H]ryanodine,

a mutant with a scrambled EF1 hand motif exhibited altered Ca^{2+} activation and inactivation. In a scrambled EF2 hand mutant, [^3H]ryanodine binding was lost, but Ca^{2+} -dependent activity was maintained in single-channel recordings. Studying a large number of RyR1/RyR2 chimeras, Gomez and Yamaguchi (2014) found that two RyR regions are involved in Ca^{2+} -dependent inactivation, one region containing the two-EF hand motifs and a second that included the S2 transmembrane segment. Among the two-EF hand motifs, EF1 more strongly controlled the isoform-specific inactivation of the RyRs by Ca^{2+} .

Evidence for luminal RyR sites involved in regulation by luminal Ca^{2+} . Both cytosolic and luminal Ca^{2+} regulate RyRs. Accordingly, dependence of channel activity on Ca^{2+} has suggested the presence of at least three types of Ca^{2+} -binding sites, high-affinity (activating) and low-affinity (inactivating) sites accessible from the cytosol in large cytosolic foot region of the RyRs, and luminal Ca^{2+} -binding sites, whose occupancy in skeletal and cardiac muscle depends on SR Ca^{2+} load (Ford and Podolsky, 1972; Fabiato and Fabiato, 1979; Shannon et al., 2000; Györke et al., 2002).

Single-channel measurements suggest two mechanisms to explain the regulation of RyRs by luminal Ca^{2+} . Luminal Ca^{2+} ions access luminal Ca^{2+} -sensing channel sites of RyR1 (Sitsapesan and Williams, 1995) and RyR2 (Sitsapesan and Williams, 1994; Györke and Györke, 1998; Gaburjakova and Gaburjakova, 2006, 2016; Diaz-Sylvester et al., 2011) or bind to cytosolic sites during or after their passage through the open channel (see next section below). Evidence for the presence of luminal Ca^{2+} regulatory sites was obtained in single-channel recordings performed in the presence of cytosolic agonists ATP, caffeine, or sulmazole. Channels recorded in the presence of sulmazole or ATP were activated to a greater extent by luminal Ca^{2+} under voltage conditions that suggested a luminal-to-cytosolic flow through the

channel (Sitsapesan and Williams, 1994, 1995). Differences in the number of channel events and the duration of open and closed events suggested that binding of ATP or sulmazole to cytosolic channel sites uncovered luminal Ca^{2+} -binding sites. Diaz-Sylvester et al. (2011) compared the regulation of RyR2 by luminal Mg^{2+} , Ca^{2+} , Sr^{2+} , and Ba^{2+} . RyR2 activation depended on cytosolic Ca^{2+} and caffeine concentrations. Although a greater activation by divalent cations compared with Cs^+ was observed in the absence of caffeine at 100 μM cytosolic Ca^{2+} , caffeine was required at 100 nM cytosolic Ca^{2+} for luminal Ca^{2+} and Sr^{2+} (but not Mg^{2+} and Ba^{2+}) to activate RyR2 (Diaz-Sylvester et al., 2011). This suggested the presence of two luminal activating cation-binding sites, one specific for Ca^{2+} and another that bound Ca^{2+} and Mg^{2+} . Gaburjakova and Gaburjakova (2016) reported that at 100 nM cytosolic Ca^{2+} and millimolar caffeine, RyR2 was fully activated by luminal Ca^{2+} and Sr^{2+} . Higher caffeine concentrations were required for full activation by luminal Mg^{2+} and Ba^{2+} . A luminal S1-S2 EF hand-related loop region was suggested to bind with an affinity of $\text{Ca}^{2+} > \text{Sr}^{2+} > \text{Mg}^{2+} \approx \text{Ba}^{2+}$.

A “store overload-induced Ca^{2+} release” (SOICR) in cardiac muscle can generate Ca^{2+} waves and Ca^{2+} -triggered arrhythmias. Loss of luminal Ca^{2+} activation suggested the presence of a luminal regulatory site (Jones et al., 2017). A point mutation in RyR2 (RyR2-E4872A corresponding to RyR1-E4942A) eliminated Ca^{2+} regulation by luminal Ca^{2+} but not by cytosolic Ca^{2+} . In mouse hearts, the RyR2-E4872 mutation suppressed SOICR and Ca^{2+} triggered ventricular tachycardia (Chen et al., 2014).

Evidence for cytosolic RyR sites involved in regulation by luminal Ca^{2+} . An alternative suggestion is that luminal Ca^{2+} regulates RyRs by binding to cytosolic regulatory sites after their passage through the open channel (also referred to as feed-through; Herrmann-Frank and Lehmann-Horn, 1996; Tripathy and Meissner, 1996; Xu and Meissner, 1998; Uehara et al., 2017). To distinguish luminal and cytosolic Ca^{2+} regulatory sites, voltage- and Ca^{2+} -dependent regulation of single RyR1 channels was reported (Tripathy and Meissner, 1996). Single channels were partially opened by cytosolic ATP in the presence of low cytosolic Ca^{2+} . As luminal Ca^{2+} was increased to micromolar concentrations, increasing channel activity was initially only seen at negative holding potentials that supported a luminal-to-cytosolic flow through the channels. A further increase in luminal Ca^{2+} to millimolar levels resulted in a decrease in channel activity. The results suggested that luminal Ca^{2+} flowing through the skeletal muscle Ca^{2+} release channel regulates channel activity by accessing cytosolic Ca^{2+} activation and inactivation sites.

A possible resolution of the different mechanisms proposed for the luminal regulation of RyRs is that it involves Ca^{2+} sensing sites on both the luminal and cyto-

solic channel sites (Laver and Honen, 2008). Modeling of single ATP-activated RyR2 channel activities indicated four Ca^{2+} regulatory sites: luminal (L) activation site, cytosolic activation (A) site, and cytosolic high-affinity (I_2) and low-affinity (I_1) inactivation sites. Luminal Mg^{2+} inhibited RyR2 by competing for Ca^{2+} at the L and A sites and binding to the I_1 site. As discussed later, luminal Ca^{2+} also regulates RyRs by binding calsequestrin (CSQ), a low-affinity, high-capacity luminal Ca^{2+} -binding protein.

Regulation by Mg^{2+}

Mg^{2+} is thought to inhibit RyRs by (a) competing with Ca^{2+} for a high-affinity cytosolic Ca^{2+} activation site, (b) binding with similar affinity as Ca^{2+} to a low-affinity cytosolic Ca^{2+} inhibitory site, (c) binding to luminal regulatory sites, and (d) reducing the flow of Ca^{2+} through the channel (Meissner et al., 1986, 1997; Smith et al., 1986; Gusev and Niggli, 2008; Laver and Honen, 2008). Mg^{2+} also has an “activating” effect. Suppression of rat RyR2 and recombinant rabbit RyR2 activities at 10–100 μM Ca^{2+} was relieved by Mg^{2+} in [^3H]ryanodine binding assays (Chugun et al., 2007; Gomez and Yamaguchi, 2014). As discussed later, Mg^{2+} forms a MgATP complex that modulates the regulation of the RyRs by Ca^{2+} .

Regulation by other divalent and trivalent cations

High-resolution cryo-EM identified a C_2H_2 -type Zn^{2+} finger motif formed by two cysteine (RyR1-Cys4958, Cys4961) and two histidine (RyR1-His4978, His4983) residues in the C-terminal domain of RyR1 and RyR2 (Yan et al., 2015; Peng et al., 2016). Although total intracellular Zn^{2+} concentrations are relatively high, the presence of a large number of cellular Zn^{2+} -binding proteins yields a free Zn^{2+} concentration in the picomolar to nanomolar range (Colvin et al., 2010). Binding of Zn^{2+} affects RyR2 activity in the absence or presence of Ca^{2+} (Woodier et al., 2015). In single-channel measurements, picomolar to micromolar Zn^{2+} increased RyR2 channel open probability in a Ca^{2+} -dependent manner, whereas millimolar Zn^{2+} inhibited RyR2. Mutagenesis studies showed that two cysteines in the Zn^{2+} finger motif are critical in maintaining a stable RyR1 channel conformation (Hurne et al., 2005). Expression of RyR1-C4958S or C4961S mutants restored $\text{Ca}_v1.1$ currents in dyspedic myotubes, which indicated restoration of bidirectional coupling between RyR1 and $\text{Ca}_v1.1$ (Sheridan et al., 2006). The two mutants did not restore E-C coupling or the response to RyR1 agonists, but supported RyR1 conformation-sensitive excitation-coupled Ca^{2+} entry (ECCE) that upon depolarization was mediated by three interacting Ca^{2+} channels, $\text{Ca}_v1.1$, RyR1, and a SOCE-like channel. Mutagenesis of RyR2-Cys4888, Cys4891, and His4908 eliminated caffeine activation of RyR2 (Peng et al., 2016).

Among nonphysiological ions tested, Sr^{2+} was optimal in activating RyR2 at 1 mM, whereas Ba^{2+} did not activate RyR2 (Liu et al., 1998; Diaz-Sylvester et al., 2011). Both inhibited RyR1 and RyR2 at high millimolar concentrations (Meissner et al., 1997; Diaz-Sylvester et al., 2011). Fe^{2+} but not Fe^{3+} inhibited [^3H]ryanodine binding by competing with Ca^{2+} for the RyR2 Ca^{2+} activation site (Kim et al., 1995). Lanthanides (Tb^{3+} , Eu^{3+} , Sm^{3+}) are nonconducting ions that activated or inactivated RyR1 at submicromolar to micromolar levels by binding to cytosolic high-affinity Ca^{2+} activating and low-affinity Ca^{2+} inactivating sites (Hadad et al., 1994; Sárközi et al., 2017). Voltage-dependent regulation by cytosolic Eu^{3+} suggested that lanthanides bind to cytosolic activating Ca^{2+} -binding sites in the electrical field of the channel, whereas the inactivating sites are located outside the electrical field (Sárközi et al., 2017).

Regulation by monovalent cations and anions

Nonphysiological assay media have been used to study RyR function. These include 50 mM CaHEPES/TrisHEPES or 0.25 M cesium methanesulfonate in single-channel measurements with SR membrane preparations to suppress SR K^+ and Cl^- channel activities, or 0.25 M KCl in [^3H]ryanodine binding studies or single-channel measurements with purified RyR preparations. However, RyR activities are greatly affected by the ionic composition and strength of the assay medium. An increase in KCl or NaCl stimulated Ca^{2+} release from SR vesicles and increased [^3H]ryanodine binding and single-channel P_o (Michalak et al., 1988; Chu et al., 1990; Ogawa and Harafuji, 1990; Meissner et al., 1997; Liu et al., 1998). Using the [^3H]ryanodine ligand binding assay, RyR1 and RyR2 Ca^{2+} -dependent channel activities were oppositely affected by (a) the competitive binding of monovalent cations to high-affinity Ca^{2+} activating binding sites, and (b) binding of Cl^- to unknown anion activating sites (Meissner et al., 1997; Liu et al., 1998).

Regulation by adenine nucleotides

Comparison of cryo-EM maps prepared in the absence or presence ATP and caffeine revealed an ATP-binding site at the cytoplasmic extension of S6 and elements of the C-terminal domain (des Georges et al., 2016). The site encompasses eight RyR1 amino acids. The adenine base of ATP contacts RyR1-Met4954, Phe4959, Thr4979, and Leu4985; the ribose ring of ATP likely interacts with Glu4955; and the triphosphate tail of ATP with positively charged Lys4211, Lys4214, and Arg4215 (Fig. 4B). Channels apparently remained closed in the presence of ATP and caffeine as the sole activating ligands, even though single-channel measurements showed a channel P_o of 0.13. des Georges et al. (2016) proposed that, similar to Ca^{2+} alone, ATP/caffeine alone primed the channel for opening, and that in addition, Ca^{2+} was required to open RyR1. Cross-linking studies using 2-azidoadenosine

5'-triphosphate 2'-3'-biotin-long-chain-hydrazone identified four glycine-rich consensus motifs in the N-terminal 95-kD fragment of RyR1, which suggests there may be additional nucleotide-binding sites (Popova et al., 2012).

Ca^{2+} and Mg^{2+} greatly affect the activation of RyRs by ATP. In single-channel measurements in the absence of cytosolic Ca^{2+} and Mg^{2+} , millimolar ATP weakly activated the essentially closed RyR1 (Tripathy and Meissner, 1996) and RyR2 (Kermode et al., 1998). On the other hand, RyR1 and RyR2 were nearly fully activated by ATP in the presence of micromolar Ca^{2+} . Other adenine nucleotides such as ADP, AMP, adenosine, or adenosine enhanced the release of Ca^{2+} from skeletal muscle SR vesicles (Morii and Tonomura, 1983; Meissner, 1984) with a decreased potency compared with AMP-PCP (Meissner, 1984). In single channel measurements, ADP was less effective in activating RyR2 than ATP (Kermode et al., 1998). Other nucleotides such as CTP, GTP, ITP, or UTP had no substantial effect on the release of Ca^{2+} (Morii and Tonomura, 1983; Meissner, 1984).

Most cellular ATP is present as MgATP complex that represents the predominant biologically active form of ATP in cells. An increase in Mg^{2+} that increases MgATP and decreases uncomplexed ATP decreased RyR activity (Meissner et al., 1986; Xu et al., 1996; Walweel et al., 2014). However, whether MgATP or ATP is a physiological regulator of RyRs is unclear. In support of an interaction with "free" ATP is that ATP activates RyRs in the absence of Mg^{2+} , and three positively charged amino acid residues in the ATP-binding site complement the negatively charged triphosphate tail of ATP. Experimental evidence whether ATP or MgATP is the preferred physiological regulator of RyR was not obtained by des Georges et al. (2016) because the cryo-EM studies were done in the absence of Mg^{2+} .

Regulation of RyRs by posttranslational modifications

In addition to the regulation by Ca^{2+} and other small molecules, RyRs are regulated by posttranslational modifications involving phosphorylation and redox modification of sites in the large cytosolic domain of the RyRs.

RyR cysteine redox modifications. RyRs contain a large number of amino acid residues that are potential targets for reactive oxygen and nitrogen molecules generated in working muscle under physiological or pathological conditions. These include free thiols, amines, and tyrosines. Most attention has been directed toward the role of cysteines and their modification by redox active molecules. RyRs contain a large number of cysteines: 100 in RyR1 and 89 in RyR2 subunit, and 1 in FKBP12 and 2 in FKBP12.6. It is therefore not surprising that cysteines are involved in the regulation of RyR activity.

RyRs have been studied extensively in ambient O_2 tension ($p\text{O}_2 \sim 150$ mm Hg), often in the absence of a re-

ducing agent. However, muscle pO_2 is ~ 10 mm Hg, and muscle cells have a high ratio of reduced to oxidized glutathione (GSH). Use of the lipophilic thiol-specific probe monobromobimane indicated that nearly half of the 100 cysteines in the RyR1 subunit were free in the presence of 5 mM reduced GSH at $pO_2 \sim 10$ mm Hg (Eu et al., 2000). An increase in oxygen tension from ~ 10 mm Hg to ambient air ($pO_2 \sim 150$ mm Hg) in the presence of 5 mM GSH resulted in loss of 6–8 free thiols/RyR1 subunit. Substitution of GSH with oxidized GSH (GSH disulfide [GSSG]) at $pO_2 \sim 10$ mm Hg or ambient air reduced the number of free thiols/RyR1 subunit to ~ 26 and increased RyR1 activity (Sun et al., 2001b). The number of free thiols in RyR2-enriched fractions in the presence of 5 mM GSH was 58 cysteines/mg protein at $pO_2 \sim 10$ mm Hg and 50 cysteines/mg protein at ~ 150 mm Hg (Sun et al., 2008). In the presence of 5 mM GSSG, a lower number of free cysteines was measured. However, because RyR2 was only partially purified, the exact number of GSH- and pO_2 -sensitive thiols in native RyR2 remains to be determined. The results suggest that RyR1 and RyR2 have functional thiols that respond to cellular pO_2 and the GSH/GSSG redox state.

Treatment of SR vesicles with reduced GSH at $pO_2 \sim 150$ mm Hg and mass spectrometric analysis of tryptic fragments showed that monobromobimane reacted with 40 cysteines in the majority of experiments, with 20 additional monobromobimane reactive cysteines less often detected (Petrotchenko et al., 2011). Seven RyR1 cysteines (Cys1040, 1303, 2436, 2565, 2606, 2611, and 3635) were selectively labeled by 7-diethylamino-3-(4'-maleimidylphenyl)-4-methylcoumarin under conditions that led to loss of RyR1 redox sensitivity (Voss et al., 2004). Nine RyR1 cysteines (Cys36, 315, 811, 906, 1591, 2326, 2363, 3193, and 3635) were endogenously modified, and another three cysteines (Cys253, 1040, and 1303) were modified by exogenous reactive oxygen and nitrogen molecules (Aracena-Parks et al., 2006). Sun et al. (2013) identified 13 oxygen tension-sensitive cysteines (Cys36, 566, 762, 845/854, 1674, 2305/2310, 2555, 2606, 2611, 2704, and 4238). Another eight cysteines (Cys120, 253, 305, 490, 1686, 2021, 3892, and 4663) were oxidized by an NADH oxidase at high pO_2 .

Mutagenesis studies have shown that cysteines located in the Jsol and Bsol domains and a poorly resolved region of RyR1 (des Georges et al., 2016) may serve a redox regulatory function. Substitution of cysteines with serine or alanine identified three cysteines (Cys1781, 2436, and 2606) that responded to a change in GSH redox potential (Petrotchenko et al., 2011). At physiological O_2 concentrations, nitric oxide (NO) specifically S-nitrosylated Cys3635 (Sun et al., 2001a). Cys3635 is part of the high-affinity CaM-binding domain of RyR1, which provides an explanation for the observation that NO was effective only in the presence of CaM. In addition, mutagenesis of Cys4958 and Cys4961 in C₂H₂-type

Zn²⁺ finger motif of RyR1 resulted in an inactive channel, which supported a depolarization-dependent Ca²⁺ entry mechanism (Hurne et al., 2005). Mutagenesis of RyR2-Cys4888 and Cys4891 eliminated caffeine activation of RyR2 (Peng et al., 2016).

RyR phosphorylation. The large cytoplasmic foot region of RyRs has many potential phosphorylation sites (Takeshima et al., 1989; Zorzato et al., 1990) that may be targeted by protein kinases and phosphatases including PKA, CaMKII, and phosphatases 2A (PP2A), PP1, and PP2B (Hohenegger and Suko, 1993; Marx et al., 2000, 2001; Dulhunty et al., 2001; Shin et al., 2002). RyR1 has one and RyR2 has three well-established phosphorylation sites per RyR subunit. Early studies indicated that canine RyR2-Ser2809 is phosphorylated by CaMKII and by PKA to a lesser extent (Witcher et al., 1991). Subsequent studies showed that cAMP-activated kinase (PKA) phosphorylates Ser2843 in rabbit RyR1 (Suko et al., 1993), mouse RyR2 at Ser2030 (Xiao et al., 2005), and human RyR2 at Ser2808 (Marx et al., 2000). CaMKII uniquely phosphorylated recombinant RyR2 at Ser2815 (Wehrens et al., 2004). RyR1 was phosphorylated at Ser2843 (corresponding to Ser2809 in RyR2) by cAMP-, cGMP-, and CaMKs (Suko et al., 1993). Phosphorylation of threonine (Suko et al., 1993) and different functional effects of CaMKII, PKA, and cGMP-dependent protein kinase (Takasago et al., 1991; Hain et al., 1995) suggested the presence of additional phosphorylation sites.

Phosphorylation of rabbit RyR2-Ser2809 in the P2 domain induced a more flexible conformation that favored the transition from the closed to open channel states (Dhindwal et al., 2017). Crystal structures of the P2 domain of RyR1 (amino acids 2,734–2,940; Sharma et al., 2012; Yuchi et al., 2012) and the corresponding P2 domains of RyR2 and RyR3 (Yuchi et al., 2012) have been reported. In vitro phosphorylation showed that PKA phosphorylated four residues corresponding to human residues Ser2808, Thr2810, Ser2811, and Ser2814 in the RyR2 P2 domain peptide (Yuchi et al., 2012). CaMKII also phosphorylated four residues (but Thr2876 instead of Thr2810).

RyR phosphorylation has been linked to the impairment of SR function leading to heart failure. Increased sensitivity to luminal Ca²⁺ of phospho-mimicking mutant RyR2-S2030D suggested that phosphorylation of Ser2030 is involved in stress-induced cardiac arrhythmias (Xiao et al., 2007). Creation of genetically modified mouse models indicated that CaMKII phosphorylation of RyR2 at Ser2814 increases SR Ca²⁺ leak, ventricular arrhythmias, and heart failure (van Oort et al., 2010; Respress et al., 2012; Li et al., 2014). PKA-mediated phosphorylation of RyR1 (Reiken et al., 2003) and RyR2 (Marx et al., 2000; Wehrens et al., 2006) led to leaky channels by dissociating the small FKBP subunit from the channel

complexes, which resulted in defective RyR2 regulation. However, other studies have challenged these findings (Stange et al., 2003; Xiao et al., 2005; Alvarado et al., 2017). A recent study indicates complex PKA phosphorylation-mediated regulation of RyR2-Ser2809. Both the complete loss of phosphorylation or maximal phosphorylation of Ser2809 increased the SR leak in rabbit ventricular myocytes (Bovo et al., 2017).

Modulation of RyR activity by accessory proteins

RyR activity is affected by a large number of proteins that interact with the RyRs, such as the FKBP, CaM, S1001, triadin, junctin, and anchoring proteins for kinases and phosphatases.

FKBPs. The small 12- and 12.6-kD FKBPs are predominantly associated with RyR1 and RyR2, respectively (Lam et al., 1995) and are generally considered part of the massive RyR complexes. FKBPs belong to the family of immunophilins and exhibit cis/trans isomerase activity, and their pharmacological removal using rapamycin or FK506 functionally uncouples groups of channels and increases channel activity (Brillantes et al., 1994). Dissociation of FKBPs induced the formation of substates in single-channel measurements (Brillantes et al., 1994; Ma et al., 1995; Ahern et al., 1997), whereas in other studies, full conductance was maintained (Barg et al., 1997; Mei et al., 2013). Cryo-EM studies located FKBP12 (fused to GSH S-transferase) to a relatively large area of RyR1 (Wagenknecht et al., 1996), which in subsequent studies was narrowed to a site at the interface of three peripheral domains corresponding to the SPRY1, SPRY3, and Jsol domains (Samsó et al., 2006; Sharma et al., 2006). Crystal structure analysis of the RyR2-SPRY1 domain along with FRET and mutagenesis studies and docking to cryo-EM maps suggested FKBP binding to a hydrophobic cluster within SPRY1 (Yuchi et al., 2015).

CaM. CaM is a 16.7-kD cytosolic protein that regulates SR Ca^{2+} release by direct binding to RyRs and through other proteins that interact with RyRs and bind CaM. Other major contributors are the voltage-regulated surface membrane Ca^{2+} channels (Ca_v 1s), CaM-dependent protein kinase (CaMKII), and CaM-stimulated protein phosphatase (calcineurin). CaM inhibits all three mammalian RyR isoforms at free $[\text{Ca}^{2+}] > 1 \mu\text{M}$. At low free Ca^{2+} concentrations ($< 1 \mu\text{M}$), CaM activates RyR1 and RyR3 channel activity, whereas RyR2 is inhibited by CaM (Ikemoto et al., 1998; Fruen et al., 2000; Balshaw et al., 2001; Yamaguchi et al., 2005). RyR1 and RyR2 bind with nanomolar affinity 1 apoCaM (Ca^{2+} -free form of CaM) or 1 CaCaM (Ca^{2+} -bound form of CaM) per RyR subunit (Balshaw et al., 2001; Yamaguchi et al., 2005).

Trypsin digestion and site-directed mutagenesis demonstrated that RyRs have a single conserved high-affinity

CaM-binding domain per RyR subunit (RyR1 amino acids 3,614–3,643, RyR2 amino acids 3,581–3,610, and RyR3 amino acids 3,467–3,498) that interacts with Ca^{2+} -free or Ca^{2+} -bound forms of CaM (Moore et al., 1999; Rodney et al., 2001; Yamaguchi et al., 2001, 2003, 2005). Crystal structure, nuclear magnetic resonance, and FRET data showed that the CaM-binding domain binds both CaM lobes in a complex formed by CaCaM and peptide corresponding to the CaM-binding domain of RyR1 (Maximciuc et al., 2006). Alternatively, CaCaM bound only the C-lobe, with the N-lobe potentially binding to another RyR1 region. This may explain why several RyR1 CaCaM- and apoCaM-binding sites were observed, using synthetic peptides and fusion proteins (Chen and MacLennan, 1994; Guerrini et al., 1995; Lau et al., 2014). Studies with RyR1/RyR2 chimeras and mutants indicated that N-terminal sites and two predicted Ca^{2+} binding motifs (EF1 RyR1-4081-4092 and EF2 RyR1-4116-4127) are involved in the isoform-specific regulation by CaM at less than $1 \mu\text{M}$ Ca^{2+} (Xu et al., 2017). Determination of temperature dependence of CaM binding showed major differences in the energetics of CaM binding to and CaM dissociation from RyR1 and RyR2 (Meissner et al., 2009).

Cryo-EM studies with RyR1 suggest that the CaM-binding site is at least 10 nm away from the transmembrane channel of the receptor and that Ca^{2+} binding to CaM (and RyR1) causes an $\sim 33\text{-\AA}$ shift of the binding site (Wagenknecht et al., 1994; Samsó and Wagenknecht, 2002). CaCaM is located in clefts formed by structural domains corresponding to the Jsol (amino acids 1,657–2,144) and Bsol (amino acids 2,145–3,613) domains, with the latter domain located close to the high-affinity CaM-binding site (Wagenknecht et al., 1997). The sequence corresponding to the high-affinity CaM-binding domain was not sufficiently resolved in high-resolution cryo-EM studies to reveal its structure in intact RyRs (des Georges et al., 2016; Peng et al., 2016).

Studies with RyR1 CaM-deficient mutants suggest a modest role for CaM regulation of RyR1 in skeletal muscle (Yamaguchi et al., 2011). In contrast, dysfunctional regulation of RyR2 by CaM alters cardiac function. Mutations in CaM are associated with RyR2-mediated cardiac arrhythmias (Nomikos et al., 2014; Søndergaard et al., 2017). Substitution of three amino acids in the CaM-binding domain (RyR2-W3587A/L3591D/F3603A) impaired CaM inhibition of RyR2 at diastolic and systolic Ca^{2+} concentrations. In mice, the mutations caused severe cardiac hypertrophy and death 2–3 wks after birth (Yamaguchi et al., 2007). In contrast, mice with a single amino acid mutation in the CaM-binding domain (RyR2-L3591D) showed loss of CaM inhibition of RyR2 at diastolic but not systolic Ca^{2+} , had a normal lifespan, and had only modest changes in heart size and function (Yamaguchi et al., 2013). This was an unexpected finding because a leaky RyR2 at diastolic Ca^{2+} is

commonly implicated in cardiomyopathies. In support of a role of CaM in terminating SR Ca^{2+} release, CaM prolonged the closed durations of RyR2 in the presence of ATP and Mg^{2+} , allowing Ca^{2+} to diffuse away from the release sites and eliminating reopening of these channels by Ca^{2+} (Xu and Meissner, 2004).

S100A1. S100A1 is a small Ca^{2+} -binding protein that modulates the activity of multiple Ca^{2+} -handling proteins in skeletal and cardiac muscle. S100A1 modulated striated muscle function by binding to the CaM-binding domain of RyR1 (Prosser et al., 2008; Wright et al., 2008). S100A1 inhibited RyR2 but not the single-site mutant RyR2-L3591D of the RyR2 CaM-binding domain in single-channel measurements (Yamaguchi et al., 2013). A more recent FRET study suggested that CaM and S100A1 can concurrently modulate RyR1 and RyR2 function without S100A1 competing for CaM at the RyR CaM-binding site (Rebbeck et al., 2016).

Other RyR accessory proteins. RyRs interact with a large number of additional proteins such as triadin, junctin, CSQ, anchoring proteins for kinases and phosphatases, and homer proteins. RyR interacting sites for these proteins have been described. The RyRs form a junctional quaternary complex with triadin, junctin, and CSQ. Triadin and junctin have a single transmembrane-spanning domain and interact directly with the RyRs. Redox-sensitive cysteines on RyR1 and triadin have a role in regulating RyR1 function (Liu and Pessah, 1994). Three positively charged lysines in full-length 95-kD triadin bind to three negatively charged RyR1 residues located in S5 pore helix (RyR1-Asp4878) and S6 pore helix (Asp4907, Glu4908) linkers (Lee et al., 2004; Goonasekera et al., 2007). Junctin-binding sites are less well defined. Different junctin domains may bind to cytosolic and luminal sites (Altschafel et al., 2011; Li et al., 2015).

CSQ is a low-affinity, high-capacity Ca^{2+} storage protein (MacLennan and Wong, 1971) concentrated in the heavy vesicle fraction derived from the SR terminal cisternae of skeletal muscle (Meissner, 1975). Cloning studies indicated the presence of two CSQ isoforms (Zorzato et al., 1994). CSQ1 is present in fast-twitch skeletal muscle and CSQ2 in cardiac muscle. Both CSQ isoforms are expressed in slow-twitch skeletal muscle (Arai et al., 1991; Biral et al., 1992; Murphy et al., 2009). The two isoforms contain a large number of negatively charged aspartic and glutamic acid residues, with CSQ2 having an extended negatively charged C terminus. Association with the RyR complex involved the binding of triadin and junctin KEKE motifs (Guo and Campbell, 1995; Zhang et al., 1997; Kobayashi et al., 2000) to asparagine-rich regions in the C terminus of CSQ (Shin et al., 2000; Beard and Dulhunty, 2015). CSQs formed linear polymers (Park et al., 2004) and dissociated (Zhang

et al., 1997) from their binding proteins as the Ca^{2+} concentration increased.

In addition to increasing Ca^{2+} store size, CSQ modulates RyR channel activity. A deficiency in SR luminal cardiac CSQ in humans (Postma et al., 2002; Lahat et al., 2004), mice (Knollmann et al., 2006), and cardiomyocytes (Terentyev et al., 2003) resulted in an imbalance of Ca^{2+} handling and CPVT. Overexpression of CSQ2 suppressed SR Ca^{2+} transients and led to severe cardiac hypertrophy in mice (Jones et al., 1998; Sato et al., 1998; Terentyev et al., 2003). Abnormal Ca^{2+} handling in association with cardiac myopathies was also observed in mice and cardiomyocytes that lacked or overexpressed the RyR2-accessory proteins triadin and junctin (Gergs et al., 2007; Kirchhof et al., 2007; Yuan et al., 2007; Chopra et al., 2009).

Marx et al. (2001) identified several anchoring proteins that mediate an interaction between RyR2 and accessory kinases and phosphatases through conserved leucine/isoleucine motifs. Spinophilin targeted PP1 to the N-terminal domain, PR130 directed PP2A to the SPRY 3 domain, PR130 directed PP2A to a region overlapping with SPRY3 domain, and A-kinase anchoring protein targeted PKA to Bsol domain. Homer proteins modulate RyR activity by binding putative proline-containing motifs in the RyR SPRY1 and Jsol domains (Feng et al., 2002; Pouliquin and Dulhunty, 2009).

Effects of ryanodine and caffeine

Among the large number of drugs that affect SR Ca^{2+} release (Zucchi and Ronca-Testoni, 1997; Xu et al., 1998; Thomas and Williams, 2012), ryanodine and caffeine have been extensively used in the assessment of SR-controlling cytoplasmic Ca^{2+} concentrations. Molecular dynamic simulations indicate that ryanodine interacts with residues in the cytosolic cavity of the closed RyR1 (Ngo et al., 2017). Single-channel measurements with purified RyRs provide the most direct information on the action of ryanodine. The results show that ryanodine modifies the gating and conductance of RyRs in a characteristic manner (Fig. 5). After the addition of micromolar ryanodine, a single RyR1 channel entered a subconductance state with a channel P_o close to unity. Upon the subsequent addition of millimolar ryanodine, the channel converted to a fully closed state. Because ryanodine binds very slowly to RyRs, ryanodine concentrations in the micromolar to millimolar rather than nanomolar to micromolar range were used. A characteristic property of ryanodine-modified channel states is insensitivity to regulation by Ca^{2+} , Mg^{2+} , and ATP, all of which greatly affect the gating behavior of unmodified channels. [^3H]Ryanodine binding studies have suggested that binding to a single high-affinity site locks RyRs into an open state, and binding of one or more ryanodine to low-affinity sites closes channels




Figure 5. **Effect of ryanodine on single purified RyR1.** Single-channel recordings of K^+ current of purified RyR1 incorporated in a planar lipid bilayer. The top trace shows the appearance of an subconductance state with $P_o \sim 1$, several minutes after the addition of 30 μM ryanodine to the cis cytosolic side of the bilayer. The bottom trace illustrates the transition from the subconductance state to a fully closed state within 1 min after the addition of 2 mM ryanodine. Bars on left indicate the open (o) and closed (c) channel. From Lai et al. (1989).

(Lai et al., 1989; Pessah and Zimanyi, 1991; reviewed by Sutko et al., 1997).

Caffeine binds to a site encompassing the S2-S3 linker and CTD, contacting residues RyR1-Trp4716 and RyR1-Ile4996, respectively (des Georges et al., 2016; Fig. 4 C). In contrast to ryanodine, caffeine (1,3,7-trimethylxanthine) activated purified RyR1 by increasing channel open probability without significantly affecting single-channel conductance (Rousseau et al., 1988). Another difference is that caffeine activates the Ca^{2+} release channel without loss of sensitivity to regulation by Ca^{2+} , Mg^{2+} , and ATP. Among 30 xanthines tested, 1,7-dimethylxanthine and 1-hexyl-3,7-dimethylxanthine (pentifylline) were most effective in activating RyR1 in a [3H]ryanodine binding assay (Liu and Meissner, 1997).

RyRs and muscle disorders

Naturally occurring mutations in RyR1 give rise to a variety of muscle diseases that include MH, CCD, and MmD (Treves et al., 2008). A list of mutations potentially associated with RyR1 is available at http://www.dmd.nl/nmdb2/variants.php?select_db=RYR1&action=search_all&order. MH is an inherited disease that causes a rapid rise in body temperature and inappropriate muscle contraction when the affected persons receive general anesthesia and MH-linked RyR1 mutations release Ca^{2+} ions from the SR. MH-linked RyR1 mutations were initially mapped to the N-terminal and central domains of RyR1; however, more recently identified MH mutations are distributed throughout the RyR1 coding sequence (Robinson et al., 2006; Stowell, 2008; Treves et al., 2008). MH is treated with dantrolene, a muscle relaxant that reduces the rate of SR Ca^{2+} release (Zucchi and Ronca-Testoni, 1997). Recent studies have shown that two RyR1 ligands, CaM (Oo et al., 2015) and an increase in metabolite Mg^{2+} from MgATP hydrolysis

during increased muscle activity (Choi et al., 2017), facilitate dantrolene inhibition. CCD and MmD are congenital diseases associated with a dysfunctional RyR1 (Treves et al., 2008). In many cases, dominant RyR1 mutations linked to CCD localize to the C-terminal domain of the channel. RyR1 with homozygous mutations associated with core myopathies often do not conduct Ca^{2+} , whereas heterozygous RyR1 channels composed of wild-type and mutant subunits conduct variable amounts of Ca^{2+} (Xu et al., 2008). More than 150 RyR2 mutations have been potentially linked to cardiac diseases such as CPVT and ARVD2 (George et al., 2007; Leenhardt et al., 2012). CPVT is an inherited cardiac disorder characterized by life-threatening arrhythmias elicited by stress and emotional disturbances. ARVD is an inherited RyR2-linked disorder that results in arrhythmias in the right ventricle.

Conclusion

Determination of RyR structure at near atomic resolution has opened the way for systematic studies on the molecular properties of a key component of skeletal and cardiac muscle function, as well as in other tissues that express RyRs such as brain and smooth muscle. Understanding the structure–function relationships of RyRs will help to explain the molecular mechanisms associated with myocardial myopathies. The combined results of structural, functional, and computational approaches have contributed basic biochemical, pharmacological, and electrophysiological information on RyR function that has set the stage for future studies of the complex regulatory mechanisms of RyRs by Ca^{2+} and other small molecules, posttranslational modifications, and accessory proteins. However, the recent identification of RyR-binding sites may not sufficiently explain their regulation by Ca^{2+} and ATP, with additional activa-

tion sites possible for these ligands. The use of lanthanides may continue to identify Ca^{2+} -activating sites and the location of Ca^{2+} - and Mg^{2+} -inactivating sites. In skeletal muscle E-C coupling, $\text{Ca}_v1.1$ activation may render a recently identified RyR1 activation site accessible to SR luminal Ca^{2+} , resulting in Ca^{2+} -induced Ca^{2+} release. Further studies are needed to address the mechanism of SR Ca^{2+} release and determine how SR Ca^{2+} release is terminated in cardiac muscle. Accessory proteins such as CaM and posttranslational modifications may affect RyR structure and activity. Structural information may facilitate genetic correction of disease-associated mutations and the development of new small molecules to inhibit disease progression. Studies on the RyRs will continue to contribute to understanding the structural framework that underlies RyR function.

ACKNOWLEDGMENTS

I wish to thank Naohiro Yamaguchi for reading the manuscript and David Mowrey and Jhuma Das for preparing the RyR structures in Figs. 1 and 4, respectively.

Supported by US Public Health Service grant AR018687.

The author declares no competing financial interests.

Lesley C. Anson served as editor.

Submitted: 7 August 2017

Accepted: 12 October 2017

REFERENCES

- Ahern, G.P., P.R. Junankar, and A.F. Dulhunty. 1997. Subconductance states in single-channel activity of skeletal muscle ryanodine receptors after removal of FKBP12. *Biophys. J.* 72:146–162. [https://doi.org/10.1016/S0006-3495\(97\)78654-5](https://doi.org/10.1016/S0006-3495(97)78654-5)
- Airey, J.A., C.F. Beck, K. Murakami, S.J. Tanksley, T.J. Deerinck, M.H. Ellisman, and J.L. Sutko. 1990. Identification and localization of two triad junctional foot protein isoforms in mature avian fast twitch skeletal muscle. *J. Biol. Chem.* 265:14187–14194.
- Altschafli, B.A., D.A. Arvanitis, O. Fuentes, Q. Yuan, E.G. Kranias, and H.H. Valdivia. 2011. Dual role of junctin in the regulation of ryanodine receptors and calcium release in cardiac ventricular myocytes. *J. Physiol.* 589:6063–6080. <https://doi.org/10.1113/jphysiol.2011.215988>
- Alvarado, F.J., X. Chen, and H.H. Valdivia. 2017. Ablation of the cardiac ryanodine receptor phospho-site Ser2808 does not alter the adrenergic response or the progression to heart failure in mice. Elimination of the genetic background as critical variable. *J. Mol. Cell. Cardiol.* 103:40–47. <https://doi.org/10.1016/j.jmcc.2017.01.001>
- Anderson, K., F.A. Lai, Q.Y. Liu, E. Rousseau, H.P. Erickson, and G. Meissner. 1989. Structural and functional characterization of the purified cardiac ryanodine receptor- Ca^{2+} release channel complex. *J. Biol. Chem.* 264:1329–1335.
- Aracena-Parks, P., S.A. Goonasekera, C.P. Gilman, R.T. Dirksen, C. Hidalgo, and S.L. Hamilton. 2006. Identification of cysteines involved in S-nitrosylation, S-glutathionylation, and oxidation to disulfides in ryanodine receptor type 1. *J. Biol. Chem.* 281:40354–40368. <https://doi.org/10.1074/jbc.M600876200>
- Arai, M., N.R. Alpert, and M. Periasamy. 1991. Cloning and characterization of the gene encoding rabbit cardiac calsequestrin. *Gene.* 109:275–279. [https://doi.org/10.1016/0378-1119\(91\)90621-H](https://doi.org/10.1016/0378-1119(91)90621-H)
- Bai, X.C., Z. Yan, J. Wu, Z. Li, and N. Yan. 2016. The central domain of RyR1 is the transducer for long-range allosteric gating of channel opening. *Cell Res.* 26:995–1006. <https://doi.org/10.1038/cr.2016.89>
- Baker, M.R., G. Fan, and I.I. Serysheva. 2017. Structure of IP₃R channel: High-resolution insights from cryo-EM. *Curr. Opin. Struct. Biol.* 46:38–47. <https://doi.org/10.1016/j.sbi.2017.05.014>
- Balshaw, D., L. Gao, and G. Meissner. 1999. Luminal loop of the ryanodine receptor: A pore-forming segment? *Proc. Natl. Acad. Sci. USA.* 96:3345–3347. <https://doi.org/10.1073/pnas.96.7.3345>
- Balshaw, D.M., L. Xu, N. Yamaguchi, D.A. Pasek, and G. Meissner. 2001. Calmodulin binding and inhibition of cardiac muscle calcium release channel (ryanodine receptor). *J. Biol. Chem.* 276:20144–20153. <https://doi.org/10.1074/jbc.M010771200>
- Bannister, R.A. 2016. Bridging the myoplasmic gap II: More recent advances in skeletal muscle excitation-contraction coupling. *J. Exp. Biol.* 219:175–182. <https://doi.org/10.1242/jeb.124123>
- Barg, S., J.A. Copello, and S. Fleischer. 1997. Different interactions of cardiac and skeletal muscle ryanodine receptors with FK-506 binding protein isoforms. *Am. J. Physiol.* 272:C1726–C1733.
- Beam, K.G., C.M. Knudson, and J.A. Powell. 1986. A lethal mutation in mice eliminates the slow calcium current in skeletal muscle cells. *Nature.* 320:168–170. <https://doi.org/10.1038/320168a0>
- Beard, N.A., and A.F. Dulhunty. 2015. C-terminal residues of skeletal muscle calsequestrin are essential for calcium binding and for skeletal ryanodine receptor inhibition. *Skelet. Muscle.* 5:6. <https://doi.org/10.1186/s13395-015-0029-7>
- Bers, D.M., and V.M. Stiffel. 1993. Ratio of ryanodine to dihydropyridine receptors in cardiac and skeletal muscle and implications for E-C coupling. *Am. J. Physiol.* 264:C1587–C1593.
- Bhat, M.B., J. Zhao, H. Takeshima, and J. Ma. 1997. Functional calcium release channel formed by the carboxyl-terminal portion of ryanodine receptor. *Biophys. J.* 73:1329–1336. [https://doi.org/10.1016/S0006-3495\(97\)78166-9](https://doi.org/10.1016/S0006-3495(97)78166-9)
- Biral, D., P. Volpe, E. Damiani, and A. Margreth. 1992. Coexistence of two calsequestrin isoforms in rabbit slow-twitch skeletal muscle fibers. *FEBS Lett.* 299:175–178. [https://doi.org/10.1016/0014-5793\(92\)80241-8](https://doi.org/10.1016/0014-5793(92)80241-8)
- Block, B.A., T. Imagawa, K.P. Campbell, and C. Franzini-Armstrong. 1988. Structural evidence for direct interaction between the molecular components of the transverse tubule/sarcoplasmic reticulum junction in skeletal muscle. *J. Cell Biol.* 107:2587–2600. <https://doi.org/10.1083/jcb.107.6.2587>
- Bovo, E., S. Huke, L.A. Blatter, and A.V. Zima. 2017. The effect of PKA-mediated phosphorylation of ryanodine receptor on SR Ca^{2+} leak in ventricular myocytes. *J. Mol. Cell. Cardiol.* 104:9–16. <https://doi.org/10.1016/j.jmcc.2017.01.015>
- Brillantes, A.B., K. Ondrias, A. Scott, E. Kobrinsky, E. Ondriasová, M.C. Moschella, T. Jayaraman, M. Landers, B.E. Ehrlich, and A.R. Marks. 1994. Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell.* 77:513–523. [https://doi.org/10.1016/0092-8674\(94\)90214-3](https://doi.org/10.1016/0092-8674(94)90214-3)
- Campiglio, M., and B.E. Flucher. 2017. STAC3 stably interacts through its C1 domain with CaV1.1 in skeletal muscle triads. *Sci. Rep.* 7:41003. <https://doi.org/10.1038/srep41003>
- Caswell, A.H., N.R. Brandt, J.P. Brunschwig, and S. Purkerson. 1991. Localization and partial characterization of the oligomeric disulfide-linked molecular weight 95,000 protein (triadin) which binds the ryanodine and dihydropyridine receptors in skeletal muscle triadic vesicles. *Biochemistry.* 30:7507–7513. <https://doi.org/10.1021/bi00244a020>
- Chen, S.R., and D.H. MacLennan. 1994. Identification of calmodulin-, Ca^{2+} -, and ruthenium red-binding domains in the Ca^{2+}

- release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 269:22698–22704.
- Chen, S.R., X. Li, K. Ebisawa, and L. Zhang. 1997. Functional characterization of the recombinant type 3 Ca^{2+} release channel (ryanodine receptor) expressed in HEK293 cells. *J. Biol. Chem.* 272:24234–24246. <https://doi.org/10.1074/jbc.272.39.24234>
- Chen, S.R., K. Ebisawa, X. Li, and L. Zhang. 1998. Molecular identification of the ryanodine receptor Ca^{2+} sensor. *J. Biol. Chem.* 273:14675–14678. <https://doi.org/10.1074/jbc.273.24.14675>
- Chen, W., R. Wang, B. Chen, X. Zhong, H. Kong, Y. Bai, Q. Zhou, C. Xie, J. Zhang, A. Guo, et al. 2014. The ryanodine receptor store-sensing gate controls Ca^{2+} waves and Ca^{2+} -triggered arrhythmias. *Nat. Med.* 20:184–192. <https://doi.org/10.1038/nm.3440>
- Cheng, H., W.J. Lederer, and M.B. Cannell. 1993. Calcium sparks: Elementary events underlying excitation-contraction coupling in heart muscle. *Science*. 262:740–744. <https://doi.org/10.1126/science.8235594>
- Choi, R.H., X. Koenig, and B.S. Launikonis. 2017. Dantrolene requires Mg^{2+} to arrest malignant hyperthermia. *Proc. Natl. Acad. Sci. USA*. 114:4811–4815. <https://doi.org/10.1073/pnas.1619835114>
- Chopra, N., T. Yang, P. Asghari, E.D. Moore, S. Huke, B. Akin, R.A. Cattolica, C.F. Perez, T. Hlaing, B.E. Knollmann-Ritschel, et al. 2009. Ablation of triadin causes loss of cardiac Ca^{2+} release units, impaired excitation-contraction coupling, and cardiac arrhythmias. *Proc. Natl. Acad. Sci. USA*. 106:7636–7641. <https://doi.org/10.1073/pnas.0902919106>
- Chu, A., M. Díaz-Muñoz, M.J. Hawkes, K. Brush, and S.L. Hamilton. 1990. Ryanodine as a probe for the functional state of the skeletal muscle sarcoplasmic reticulum calcium release channel. *Mol. Pharmacol.* 37:735–741.
- Chugun, A., O. Sato, H. Takeshima, and Y. Ogawa. 2007. Mg^{2+} activates the ryanodine receptor type 2 (RyR2) at intermediate Ca^{2+} concentrations. *Am. J. Physiol. Cell Physiol.* 292:C535–C544. <https://doi.org/10.1152/ajpcell.00275.2006>
- Clarke, O.B., and W.A. Hendrickson. 2016. Structures of the colossal RyR1 calcium release channel. *Curr. Opin. Struct. Biol.* 39:144–152. <https://doi.org/10.1016/j.sbi.2016.09.002>
- Collins, J.H. 1991. Sequence analysis of the ryanodine receptor: Possible association with a 12K, FK506-binding immunophilin/protein kinase C inhibitor. *Biochem. Biophys. Res. Commun.* 178:1288–1290. [https://doi.org/10.1016/0006-291X\(91\)91033-9](https://doi.org/10.1016/0006-291X(91)91033-9)
- Colvin, R.A., W.R. Holmes, C.P. Fontaine, and W. Maret. 2010. Cytosolic zinc buffering and muffling: Their role in intracellular zinc homeostasis. *Metallomics*. 2:306–317. <https://doi.org/10.1039/b926662c>
- Conti, A., L. Gorza, and V. Sorrentino. 1996. Differential distribution of ryanodine receptor type 3 (RyR3) gene product in mammalian skeletal muscles. *Biochem. J.* 316:19–23. <https://doi.org/10.1042/bj3160019>
- des Georges, A., O.B. Clarke, R. Zalk, Q. Yuan, K.J. Condon, R.A. Grassucci, W.A. Hendrickson, A.R. Marks, and J. Frank. 2016. Structural basis for gating and activation of RyR1. *Cell*. 167:145–157.e17. <https://doi.org/10.1016/j.cell.2016.08.075>
- Dhindwal, S., J. Lobo, V. Cabra, D.J. Santiago, A.R. Nayak, K. Dryden, and M. Samsó. 2017. A cryo-EM-based model of phosphorylation- and FKBP12.6-mediated allosterism of the cardiac ryanodine receptor. *Sci. Signal*. 10:eaai8842. <https://doi.org/10.1126/scisignal.aai8842>
- Diaz-Sylvestre, P.L., M. Porta, and J.A. Copello. 2011. Modulation of cardiac ryanodine receptor channels by alkaline earth cations. *PLoS One*. 6:e26693. <https://doi.org/10.1371/journal.pone.0026693>
- Dirksen, R.T., and G. Avila. 2002. Altered ryanodine receptor function in central core disease: Leaky or uncoupled Ca^{2+} release channels? *Trends Cardiovasc. Med.* 12:189–197. [https://doi.org/10.1016/S1050-1738\(02\)00163-9](https://doi.org/10.1016/S1050-1738(02)00163-9)
- Donaldson, S.K. 1985. Peeled mammalian skeletal muscle fibers. Possible stimulation of Ca^{2+} release via a transverse tubule-sarcoplasmic reticulum mechanism. *J. Gen. Physiol.* 86:501–525. <https://doi.org/10.1085/jgp.86.4.501>
- Du, G.G., and D.H. MacLennan. 1998. Functional consequences of mutations of conserved, polar amino acids in transmembrane sequences of the Ca^{2+} release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 273:31867–31872. <https://doi.org/10.1074/jbc.273.48.31867>
- Du, G.G., and D.H. MacLennan. 1999. Ca^{2+} inactivation sites are located in the COOH-terminal quarter of recombinant rabbit skeletal muscle Ca^{2+} release channels (ryanodine receptors). *J. Biol. Chem.* 274:26120–26126. <https://doi.org/10.1074/jbc.274.37.26120>
- Du, G.G., X. Guo, V.K. Khanna, and D.H. MacLennan. 2001. Functional characterization of mutants in the predicted pore region of the rabbit cardiac muscle Ca^{2+} release channel (ryanodine receptor isoform 2). *J. Biol. Chem.* 276:31760–31771. <https://doi.org/10.1074/jbc.M102751200>
- Dulhunty, A.F., D. Laver, S.M. Curtis, S. Pace, C. Haarmann, and E.M. Gallant. 2001. Characteristics of irreversible ATP activation suggest that native skeletal ryanodine receptors can be phosphorylated via an endogenous CaMKII. *Biophys. J.* 81:3240–3252. [https://doi.org/10.1016/S0006-3495\(01\)75959-0](https://doi.org/10.1016/S0006-3495(01)75959-0)
- Efremov, R.G., A. Leitner, R. Aebersold, and S. Raunser. 2015. Architecture and conformational switch mechanism of the ryanodine receptor. *Nature*. 517:39–43. <https://doi.org/10.1038/nature13916>
- Endo, M. 2009. Calcium-induced calcium release in skeletal muscle. *Physiol. Rev.* 89:1153–1176. <https://doi.org/10.1152/physrev.00040.2008>
- Eu, J.P., J. Sun, L. Xu, J.S. Stamler, and G. Meissner. 2000. The skeletal muscle calcium release channel: Coupled O_2 sensor and NO signaling functions. *Cell*. 102:499–509. [https://doi.org/10.1016/S0092-8674\(00\)00054-4](https://doi.org/10.1016/S0092-8674(00)00054-4)
- Euden, J., S.A. Mason, C. Viero, N.L. Thomas, and A.J. Williams. 2013a. Investigations of the contribution of a putative glycine hinge to ryanodine receptor channel gating. *J. Biol. Chem.* 288:16671–16679. <https://doi.org/10.1074/jbc.M113.465310>
- Euden, J., S.A. Mason, and A.J. Williams. 2013b. Functional characterization of the cardiac ryanodine receptor pore-forming region. *PLoS One*. 8:e66542. <https://doi.org/10.1371/journal.pone.0066542>
- Fabiato, A. 1983. Calcium-induced release of calcium from the cardiac sarcoplasmic reticulum. *Am. J. Physiol.* 245:C1–C14.
- Fabiato, A., and F. Fabiato. 1975. Contractions induced by a calcium-triggered release of calcium from the sarcoplasmic reticulum of single skinned cardiac cells. *J. Physiol.* 249:469–495. <https://doi.org/10.1113/jphysiol.1975.sp011026>
- Fabiato, A., and F. Fabiato. 1979. Calcium and cardiac excitation-contraction coupling. *Annu. Rev. Physiol.* 41:473–484. <https://doi.org/10.1146/annurev.ph.41.030179.002353>
- Felder, E., and C. Franzini-Armstrong. 2002. Type 3 ryanodine receptors of skeletal muscle are segregated in a parajunctional position. *Proc. Natl. Acad. Sci. USA*. 99:1695–1700. <https://doi.org/10.1073/pnas.032657599>
- Feng, W., J. Tu, T. Yang, P.S. Vernon, P.D. Allen, P.F. Worley, and I.N. Pessah. 2002. Homer regulates gain of ryanodine receptor type 1 channel complex. *J. Biol. Chem.* 277:44722–44730. <https://doi.org/10.1074/jbc.M207675200>

- Fessenden, J.D., L. Chen, Y. Wang, C. Paolini, C. Franzini-Armstrong, P.D. Allen, and I.N. Pessah. 2001. Ryanodine receptor point mutant E4032A reveals an allosteric interaction with ryanodine. *Proc. Natl. Acad. Sci. USA*. 98:2865–2870. <https://doi.org/10.1073/pnas.041608898>
- Fessenden, J.D., W. Feng, I.N. Pessah, and P.D. Allen. 2004. Mutational analysis of putative calcium binding motifs within the skeletal ryanodine receptor isoform, RyR1. *J. Biol. Chem.* 279:53028–53035. <https://doi.org/10.1074/jbc.M411136200>
- Figuerola, L., V.M. Shkryl, J. Zhou, C. Manno, A. Momotake, G. Brum, L.A. Blatter, G.C. Ellis-Davies, and E. Ríos. 2012. Synthetic localized calcium transients directly probe signalling mechanisms in skeletal muscle. *J. Physiol.* 590:1389–1411. <https://doi.org/10.1113/jphysiol.2011.225854>
- Fleischer, S., E.M. Ogunbunmi, M.C. Dixon, and E.A. Fleer. 1985. Localization of Ca^{2+} release channels with ryanodine in junctional terminal cisternae of sarcoplasmic reticulum of fast skeletal muscle. *Proc. Natl. Acad. Sci. USA*. 82:7256–7259. <https://doi.org/10.1073/pnas.82.21.7256>
- Flucher, B.E., A. Conti, H. Takeshima, and V. Sorrentino. 1999. Type 3 and type 1 ryanodine receptors are localized in triads of the same mammalian skeletal muscle fibers. *J. Cell Biol.* 146:621–630. <https://doi.org/10.1083/jcb.146.3.621>
- Ford, L.E., and R.J. Podolsky. 1972. Calcium uptake and force development by skinned muscle fibres in EGTA buffered solutions. *J. Physiol.* 223:1–19. <https://doi.org/10.1113/jphysiol.1972.sp009830>
- Franzini-Armstrong, C., and F. Protasi. 1997. Ryanodine receptors of striated muscles: A complex channel capable of multiple interactions. *Physiol. Rev.* 77:699–729.
- Franzini-Armstrong, C., F. Protasi, and V. Ramesh. 1999. Shape, size, and distribution of Ca^{2+} release units and couplons in skeletal and cardiac muscles. *Biophys. J.* 77:1528–1539. [https://doi.org/10.1016/S0006-3495\(99\)77000-1](https://doi.org/10.1016/S0006-3495(99)77000-1)
- Fruen, B.R., J.M. Bardy, T.M. Byrem, G.M. Strasburg, and C.F. Louis. 2000. Differential Ca^{2+} sensitivity of skeletal and cardiac muscle ryanodine receptors in the presence of calmodulin. *Am. J. Physiol. Cell Physiol.* 279:C724–C733.
- Fujii, J., K. Otsu, F. Zorzato, S. de Leon, V.K. Khanna, J.E. Weiler, P.J. O'Brien, and D.H. MacLennan. 1991. Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science*. 253:448–451. <https://doi.org/10.1126/science.1862346>
- Furuichi, T., D. Furutama, Y. Hakamata, J. Nakai, H. Takeshima, and K. Mikoshiba. 1994. Multiple types of ryanodine receptor/ Ca^{2+} release channels are differentially expressed in rabbit brain. *J. Neurosci.* 14:4794–4805.
- Gaburjakova, J., and M. Gaburjakova. 2006. Comparison of the effects exerted by luminal Ca^{2+} on the sensitivity of the cardiac ryanodine receptor to caffeine and cytosolic Ca^{2+} . *J. Membr. Biol.* 212:17–28. <https://doi.org/10.1007/s00232-006-7018-z>
- Gaburjakova, J., and M. Gaburjakova. 2016. Cardiac ryanodine receptor: Selectivity for alkaline earth metal cations points to the EF-hand nature of luminal binding sites. *Bioelectrochemistry*. 109:49–56. <https://doi.org/10.1016/j.bioelechem.2016.01.002>
- Gao, L., D. Balshaw, L. Xu, A. Tripathy, C. Xin, and G. Meissner. 2000. Evidence for a role of the luminal M3-M4 loop in skeletal muscle Ca^{2+} release channel (ryanodine receptor) activity and conductance. *Biophys. J.* 79:828–840. [https://doi.org/10.1016/S0006-3495\(00\)76339-9](https://doi.org/10.1016/S0006-3495(00)76339-9)
- George, C.H., H. Jundi, N.L. Thomas, D.L. Fry, and F.A. Lai. 2007. Ryanodine receptors and ventricular arrhythmias: emerging trends in mutations, mechanisms and therapies. *J. Mol. Cell. Cardiol.* 42:34–50. <https://doi.org/10.1016/j.yjmcc.2006.08.115>
- Gergs, U., T. Berndt, J. Buskase, L.R. Jones, U. Kirchhefer, F.U. Müller, K.D. Schlüter, W. Schmitz, and J. Neumann. 2007. On the role of junctin in cardiac Ca^{2+} handling, contractility, and heart failure. *Am. J. Physiol. Heart Circ. Physiol.* 293:H728–H734. <https://doi.org/10.1152/ajpheart.01187.2006>
- Giannini, G., A. Conti, S. Mammarella, M. Scrobogna, and V. Sorrentino. 1995. The ryanodine receptor/calcium channel genes are widely and differentially expressed in murine brain and peripheral tissues. *J. Cell Biol.* 128:893–904. <https://doi.org/10.1083/jcb.128.5.893>
- Gillespie, D. 2008. Energetics of divalent selectivity in a calcium channel: The ryanodine receptor case study. *Biophys. J.* 94:1169–1184. <https://doi.org/10.1529/biophysj.107.116798>
- Gillespie, D., and M. Fill. 2008. Intracellular calcium release channels mediate their own countercurrent: The ryanodine receptor case study. *Biophys. J.* 95:3706–3714. <https://doi.org/10.1529/biophysj.108.131987>
- Gillespie, D., L. Xu, Y. Wang, and G. Meissner. 2005. (De) constructing the ryanodine receptor: Modeling ion permeation and selectivity of the calcium release channel. *J. Phys. Chem. B*. 109:15598–15610. <https://doi.org/10.1021/jp052471j>
- Gillespie, D., L. Xu, and G. Meissner. 2014. Selecting ions by size in a calcium channel: The ryanodine receptor case study. *Biophys. J.* 107:2263–2273. <https://doi.org/10.1016/j.bpj.2014.09.031>
- Girgenrath, T., M. Mahalingam, B. Svensson, F.R. Nitu, R.L. Cornea, and J.D. Fessenden. 2013. N-terminal and central segments of the type 1 ryanodine receptor mediate its interaction with FK506-binding proteins. *J. Biol. Chem.* 288:16073–16084. <https://doi.org/10.1074/jbc.M113.463299>
- Gomez, A.C., and N. Yamaguchi. 2014. Two regions of the ryanodine receptor calcium channel are involved in Ca^{2+} -dependent inactivation. *Biochemistry*. 53:1373–1379. <https://doi.org/10.1021/bi401586h>
- Goonasekera, S.A., N.A. Beard, L. Groom, T. Kimura, A.D. Lyfenko, A. Rosenfeld, I. Marty, A.F. Dulhunty, and R.T. Dirksen. 2007. Triadin binding to the C-terminal luminal loop of the ryanodine receptor is important for skeletal muscle excitation contraction coupling. *J. Gen. Physiol.* 130:365–378. <https://doi.org/10.1085/jgp.200709790>
- Gregg, R.G., A. Messing, C. Strube, M. Beurg, R. Moss, M. Behan, M. Sukhareva, S. Haynes, J.A. Powell, R. Coronado, and P.A. Powers. 1996. Absence of the beta subunit (cchb1) of the skeletal muscle dihydropyridine receptor alters expression of the alpha 1 subunit and eliminates excitation-contraction coupling. *Proc. Natl. Acad. Sci. USA*. 93:13961–13966. <https://doi.org/10.1073/pnas.93.24.13961>
- Guerrini, R., P. Menegazzi, R. Anacardio, M. Marastoni, R. Tomatis, F. Zorzato, and S. Treves. 1995. Calmodulin binding sites of the skeletal, cardiac, and brain ryanodine receptor Ca^{2+} channels: Modulation by the catalytic subunit of cAMP-dependent protein kinase? *Biochemistry*. 34:5120–5129. <https://doi.org/10.1021/bi00015a024>
- Guo, W., and K.P. Campbell. 1995. Association of triadin with the ryanodine receptor and calsequestrin in the lumen of the sarcoplasmic reticulum. *J. Biol. Chem.* 270:9027–9030. <https://doi.org/10.1074/jbc.270.16.9027>
- Guo, W., B. Sun, Z. Xiao, Y. Liu, Y. Wang, L. Zhang, R. Wang, and S.R. Chen. 2016. The EF-hand Ca^{2+} binding domain is not required for cytosolic Ca^{2+} activation of the cardiac ryanodine receptor. *J. Biol. Chem.* 291:2150–2160. <https://doi.org/10.1074/jbc.M115.693325>

- Gusev, K., and E. Niggli. 2008. Modulation of the local SR Ca^{2+} release by intracellular Mg^{2+} in cardiac myocytes. *J. Gen. Physiol.* 132:721–730. <https://doi.org/10.1085/jgp.200810119>
- Györke, I., and S. Györke. 1998. Regulation of the cardiac ryanodine receptor channel by luminal Ca^{2+} involves luminal Ca^{2+} sensing sites. *Biophys. J.* 75:2801–2810. [https://doi.org/10.1016/S0006-3495\(98\)77723-9](https://doi.org/10.1016/S0006-3495(98)77723-9)
- Györke, S., I. Györke, V. Lukyanenko, D. Terentyev, S. Viatchenko-Karpinski, and T.F. Wiesner. 2002. Regulation of sarcoplasmic reticulum calcium release by luminal calcium in cardiac muscle. *Front. Biosci.* 7:d1454–d1463. <https://doi.org/10.2741/A852>
- Hadad, N., A.C. Zable, J.J. Abramson, and V. Shoshan-Barmatz. 1994. Ca^{2+} binding sites of the ryanodine receptor/ Ca^{2+} release channel of sarcoplasmic reticulum. Low affinity binding site(s) as probed by terbium fluorescence. *J. Biol. Chem.* 269:24864–24869.
- Hain, J., H. Onoue, M. Mayrleitner, S. Fleischer, and H. Schindler. 1995. Phosphorylation modulates the function of the calcium release channel of sarcoplasmic reticulum from cardiac muscle. *J. Biol. Chem.* 270:2074–2081. <https://doi.org/10.1074/jbc.270.5.2074>
- Hakamata, Y., J. Nakai, H. Takeshima, and K. Imoto. 1992. Primary structure and distribution of a novel ryanodine receptor/calcium release channel from rabbit brain. *FEBS Lett.* 312:229–235. [https://doi.org/10.1016/0014-5793\(92\)80941-9](https://doi.org/10.1016/0014-5793(92)80941-9)
- Hamilton, S.L., and I.I. Serysheva. 2009. Ryanodine receptor structure: progress and challenges. *J. Biol. Chem.* 284:4047–4051. <https://doi.org/10.1074/jbc.R800054200>
- Heiny, J.A., and G. Meissner. 2012. Excitation-contraction coupling in skeletal muscle. In *Cell Physiology Sourcebook*. Fourth edition. N. Sperelakis, editor. Academic Press, Cambridge, MA. 783–800.
- Herrmann-Frank, A., and F. Lehmann-Horn. 1996. Regulation of the purified Ca^{2+} release channel/ryanodine receptor complex of skeletal muscle sarcoplasmic reticulum by luminal calcium. *Pflugers Arch.* 432:155–157. <https://doi.org/10.1007/s004240050117>
- Hille, B. 1973. Potassium channels in myelinated nerve. Selective permeability to small cations. *J. Gen. Physiol.* 61:669–686. <https://doi.org/10.1085/jgp.61.6.669>
- Hohenegger, M., and J. Suko. 1993. Phosphorylation of the purified cardiac ryanodine receptor by exogenous and endogenous protein kinases. *Biochem. J.* 296:303–308. <https://doi.org/10.1042/bj2960303>
- Huang, X., B. Fruen, D.T. Farrington, T. Wagenknecht, and Z. Liu. 2012. Calmodulin-binding locations on the skeletal and cardiac ryanodine receptors. *J. Biol. Chem.* 287:30328–30335. <https://doi.org/10.1074/jbc.M112.383109>
- Hurne, A.M., J.J. O'Brien, D. Wingrove, G. Cherednichenko, P.D. Allen, K.G. Beam, and I.N. Pessah. 2005. Ryanodine receptor type 1 (RyR1) mutations C4958S and C4961S reveal excitation-coupled calcium entry (ECCE) is independent of sarcoplasmic reticulum store depletion. *J. Biol. Chem.* 280:36994–37004. <https://doi.org/10.1074/jbc.M506441200>
- Ikemoto, T., H. Takeshima, M. Iino, and M. Endo. 1998. Effect of calmodulin on Ca^{2+} -induced Ca^{2+} release of skeletal muscle from mutant mice expressing either ryanodine receptor type 1 or type 3. *Pflugers Arch.* 437:43–48. <https://doi.org/10.1007/s004240050744>
- Imagawa, T., J.S. Smith, R. Coronado, and K.P. Campbell. 1987. Purified ryanodine receptor from skeletal muscle sarcoplasmic reticulum is the Ca^{2+} -permeable pore of the calcium release channel. *J. Biol. Chem.* 262:16636–16643.
- Inui, M., A. Saito, and S. Fleischer. 1987a. Isolation of the ryanodine receptor from cardiac sarcoplasmic reticulum and identity with the feet structures. *J. Biol. Chem.* 262:15637–15642.
- Inui, M., A. Saito, and S. Fleischer. 1987b. Purification of the ryanodine receptor and identity with feet structures of junctional terminal cisternae of sarcoplasmic reticulum from fast skeletal muscle. *J. Biol. Chem.* 262:1740–1747.
- Jenden, D.J., and A.S. Fairhurst. 1969. The pharmacology of ryanodine. *Pharmacol. Rev.* 21:1–25.
- Jiang, Y., A. Lee, J. Chen, M. Cadene, B.T. Chait, and R. MacKinnon. 2002. The open pore conformation of potassium channels. *Nature.* 417:523–526. <https://doi.org/10.1038/417523a>
- Jones, L.R., L. Zhang, K. Sanborn, A.O. Jorgensen, and J. Kelley. 1995. Purification, primary structure, and immunological characterization of the 26-kDa calsequestrin binding protein (junctin) from cardiac junctional sarcoplasmic reticulum. *J. Biol. Chem.* 270:30787–30796. <https://doi.org/10.1074/jbc.270.51.30787>
- Jones, L.R., Y.J. Suzuki, W. Wang, Y.M. Kobayashi, V. Ramesh, C. Franzini-Armstrong, L. Cleemann, and M. Morad. 1998. Regulation of Ca^{2+} signaling in transgenic mouse cardiac myocytes overexpressing calsequestrin. *J. Clin. Invest.* 101:1385–1393. <https://doi.org/10.1172/JCI1362>
- Jones, P.P., W. Guo, and S.R.W. Chen. 2017. Control of cardiac ryanodine receptor by sarcoplasmic reticulum luminal Ca^{2+} . *J. Gen. Physiol.* 149:867–875. <https://doi.org/10.1085/jgp.201711805>
- Kawamoto, R.M., J.-P. Brunschwig, K.-C. Kim, and A.H. Caswell. 1986. Isolation, characterization, and localization of the spanning protein from skeletal muscle triads. *J. Cell Biol.* 103:1405–1414. <https://doi.org/10.1083/jcb.103.4.1405>
- Kermode, H., A.J. Williams, and R. Sitsapesan. 1998. The interactions of ATP, ADP, and inorganic phosphate with the sheep cardiac ryanodine receptor. *Biophys. J.* 74:1296–1304. [https://doi.org/10.1016/S0006-3495\(98\)77843-9](https://doi.org/10.1016/S0006-3495(98)77843-9)
- Kim, E., S.N. Giri, and I.N. Pessah. 1995. Iron(II) is a modulator of ryanodine-sensitive calcium channels of cardiac muscle sarcoplasmic reticulum. *Toxicol. Appl. Pharmacol.* 130:57–66. <https://doi.org/10.1006/taap.1995.1008>
- Kirchhof, P., J. Klimas, L. Fabritz, M. Zwiener, L.R. Jones, M. Schäfers, S. Hermann, P. Boknik, W. Schmitz, G. Breithardt, et al. 2007. Stress and high heart rate provoke ventricular tachycardia in mice expressing triadin. *J. Mol. Cell. Cardiol.* 42:962–971. <https://doi.org/10.1016/j.yjmcc.2007.02.012>
- Knollmann, B.C., N. Chopra, T. Hlaing, B. Akin, T. Yang, K. Ettensohn, B.E. Knollmann, K.D. Horton, N.J. Weissman, I. Holinstat, et al. 2006. *Casq2* deletion causes sarcoplasmic reticulum volume increase, premature Ca^{2+} release, and catecholaminergic polymorphic ventricular tachycardia. *J. Clin. Invest.* 116:2510–2520.
- Kobayashi, Y.M., B.A. Alseikhan, and L.R. Jones. 2000. Localization and characterization of the calsequestrin-binding domain of triadin 1. Evidence for a charged beta-strand in mediating the protein-protein interaction. *J. Biol. Chem.* 275:17639–17646. <https://doi.org/10.1074/jbc.M002091200>
- Lahat, H., E. Pras, and M. Eldar. 2004. A missense mutation in *CASQ2* is associated with autosomal recessive catecholamine-induced polymorphic ventricular tachycardia in Bedouin families from Israel. *Ann. Med.* 36(suppl 1):87–91. <https://doi.org/10.1080/17431380410032517>
- Lai, F.A., H. Erickson, B.A. Block, and G. Meissner. 1987. Evidence for a junctional feet-ryanodine receptor complex from sarcoplasmic reticulum. *Biochem. Biophys. Res. Commun.* 143:704–709. [https://doi.org/10.1016/0006-291X\(87\)91411-2](https://doi.org/10.1016/0006-291X(87)91411-2)
- Lai, F.A., K. Anderson, E. Rousseau, Q.Y. Liu, and G. Meissner. 1988a. Evidence for a Ca^{2+} channel within the ryanodine receptor complex from cardiac sarcoplasmic reticulum. *Biochem. Biophys.*

- Res. Commun.* 151:441–449. [https://doi.org/10.1016/0006-291X\(88\)90613-4](https://doi.org/10.1016/0006-291X(88)90613-4)
- Lai, F.A., H.P. Erickson, E. Rousseau, Q.Y. Liu, and G. Meissner. 1988b. Purification and reconstitution of the calcium release channel from skeletal muscle. *Nature*. 331:315–319. <https://doi.org/10.1038/331315a0>
- Lai, F.A., M. Misra, L. Xu, H.A. Smith, and G. Meissner. 1989. The ryanodine receptor-Ca²⁺ release channel complex of skeletal muscle sarcoplasmic reticulum. Evidence for a cooperatively coupled, negatively charged homotetramer. *J. Biol. Chem.* 264:16776–16785.
- Lam, E., M.M. Martin, A.P. Timmerman, C. Sabers, S. Fleischer, T. Lukas, R.T. Abraham, S.J. O'Keefe, E.A. O'Neill, and G.J. Wiederrecht. 1995. A novel FK506 binding protein can mediate the immunosuppressive effects of FK506 and is associated with the cardiac ryanodine receptor. *J. Biol. Chem.* 270:26511–26522. <https://doi.org/10.1074/jbc.270.44.26511>
- Lamb, G.D. 2002. Voltage-sensor control of Ca²⁺ release in skeletal muscle: insights from skinned fibers. *Front. Biosci.* 7:d834–d842.
- Lau, K., M.M. Chan, and F. Van Petegem. 2014. Lobe-specific calmodulin binding to different ryanodine receptor isoforms. *Biochemistry*. 53:932–946. <https://doi.org/10.1021/bi401502x>
- Laver, D.R., and B.N. Honen. 2008. Luminal Mg²⁺, a key factor controlling RYR2-mediated Ca²⁺ release: Cytoplasmic and luminal regulation modeled in a tetrameric channel. *J. Gen. Physiol.* 132:429–446. <https://doi.org/10.1085/jgp.200810001>
- Ledbetter, M.W., J.K. Preiner, C.F. Louis, and J.R. Mickelson. 1994. Tissue distribution of ryanodine receptor isoforms and alleles determined by reverse transcription polymerase chain reaction. *J. Biol. Chem.* 269:31544–31551.
- Lee, J.M., S.H. Rho, D.W. Shin, C. Cho, W.J. Park, S.H. Eom, J. Ma, and D.H. Kim. 2004. Negatively charged amino acids within the intraluminal loop of ryanodine receptor are involved in the interaction with triadin. *J. Biol. Chem.* 279:6994–7000. <https://doi.org/10.1074/jbc.M312446200>
- Leenhardt, A., I. Denjoy, and P. Guicheney. 2012. Catecholaminergic polymorphic ventricular tachycardia. *Circ Arrhythm Electrophysiol.* 5:1044–1052. <https://doi.org/10.1161/CIRCEP.111.962027>
- Li, L., S. Mirza, S.J. Richardson, E.M. Gallant, C. Thekkedam, S.M. Pace, F. Zorzato, D. Liu, N.A. Beard, and A.F. Dulhunty. 2015. A new cytoplasmic interaction between junctin and ryanodine receptor Ca²⁺ release channels. *J. Cell Sci.* 128:951–963. <https://doi.org/10.1242/jcs.160689>
- Li, N., D.Y. Chiang, S. Wang, Q. Wang, L. Sun, N. Voigt, J.L. Respress, S. Ather, D.G. Skapura, V.K. Jordan, et al. 2014. Ryanodine receptor-mediated calcium leak drives progressive development of an atrial fibrillation substrate in a transgenic mouse model. *Circulation*. 129:1276–1285. <https://doi.org/10.1161/CIRCULATIONAHA.113.006611>
- Lindsay, A.R., S.D. Manning, and A.J. Williams. 1991. Monovalent cation conductance in the ryanodine receptor-channel of sheep cardiac muscle sarcoplasmic reticulum. *J. Physiol.* 439:463–480. <https://doi.org/10.1113/jphysiol.1991.sp018676>
- Liu, G., and I.N. Pessah. 1994. Molecular interaction between ryanodine receptor and glycoprotein triadin involves redox cycling of functionally important hyperreactive thiols. *J. Biol. Chem.* 269:33028–33034.
- Liu, W., and G. Meissner. 1997. Structure-activity relationship of xanthines and skeletal muscle ryanodine receptor/Ca²⁺ release channel. *Pharmacology*. 54:135–143. <https://doi.org/10.1159/000139480>
- Liu, W., D.A. Pasek, and G. Meissner. 1998. Modulation of Ca²⁺-gated cardiac muscle Ca²⁺-release channel (ryanodine receptor) by mono- and divalent ions. *Am. J. Physiol.* 274:C120–C128.
- Liu, Y., B. Sun, Z. Xiao, R. Wang, W. Guo, J.Z. Zhang, T. Mi, Y. Wang, P.P. Jones, F. Van Petegem, and S.R. Chen. 2015. Roles of the NH₂-terminal domains of cardiac ryanodine receptor in Ca²⁺ release activation and termination. *J. Biol. Chem.* 290:7736–7746. <https://doi.org/10.1074/jbc.M114.618827>
- Ludtke, S.J., I.I. Serysheva, S.L. Hamilton, and W. Chiu. 2005. The pore structure of the closed RyR1 channel. *Structure*. 13:1203–1211. <https://doi.org/10.1016/j.str.2005.06.005>
- Ma, J., M.B. Bhat, and J. Zhao. 1995. Rectification of skeletal muscle ryanodine receptor mediated by FK506 binding protein. *Biophys. J.* 69:2398–2404. [https://doi.org/10.1016/S0006-3495\(95\)80109-8](https://doi.org/10.1016/S0006-3495(95)80109-8)
- Mackrill, J.J. 2012. Ryanodine receptor calcium release channels: An evolutionary perspective. *Adv. Exp. Med. Biol.* 740:159–182. https://doi.org/10.1007/978-94-007-2888-2_7
- MacLennan, D.H., and P.T.S. Wong. 1971. Isolation of a calcium-sequestering protein from sarcoplasmic reticulum. *Proc. Natl. Acad. Sci. USA*. 68:1231–1235. <https://doi.org/10.1073/pnas.68.6.1231>
- Marx, S.O., K. Ondrias, and A.R. Marks. 1998. Coupled gating between individual skeletal muscle Ca²⁺ release channels (ryanodine receptors). *Science*. 281:818–821. <https://doi.org/10.1126/science.281.5378.818>
- Marx, S.O., S. Reiken, Y. Hisamatsu, T. Jayaraman, D. Burkhoff, N. Roseblit, and A.R. Marks. 2000. PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): Defective regulation in failing hearts. *Cell*. 101:365–376. [https://doi.org/10.1016/S0092-8674\(00\)80847-8](https://doi.org/10.1016/S0092-8674(00)80847-8)
- Marx, S.O., S. Reiken, Y. Hisamatsu, M. Gaburjakova, J. Gaburjakova, Y.M. Yang, N. Roseblit, and A.R. Marks. 2001. Phosphorylation-dependent regulation of ryanodine receptors: A novel role for leucine/isoleucine zippers. *J. Cell Biol.* 153:699–708. <https://doi.org/10.1083/jcb.153.4.699>
- Maximciuc, A.A., J.A. Putkey, Y. Shamoo, and K.R. Mackenzie. 2006. Complex of calmodulin with a ryanodine receptor target reveals a novel, flexible binding mode. *Structure*. 14:1547–1556. <https://doi.org/10.1016/j.str.2006.08.011>
- McCarthy, T.V., K.A. Quane, and P.J. Lynch. 2000. Ryanodine receptor mutations in malignant hyperthermia and central core disease. *Hum. Mutat.* 15:410–417. [https://doi.org/10.1002/\(SICI\)1098-1004\(200005\)15:5<410::AID-HUMU2>3.0.CO;2-D](https://doi.org/10.1002/(SICI)1098-1004(200005)15:5<410::AID-HUMU2>3.0.CO;2-D)
- Mei, Y., L. Xu, H.F. Kramer, G.H. Tomberlin, C. Townsend, and G. Meissner. 2013. Stabilization of the skeletal muscle ryanodine receptor ion channel-FKBP12 complex by the 1,4-benzothiazepine derivative S107. *PLoS One*. 8:e54208. <https://doi.org/10.1371/journal.pone.0054208>
- Mei, Y., L. Xu, D.D. Mowrey, R. Mendez Giraldez, Y. Wang, D.A. Pasek, N.V. Dokholyan, and G. Meissner. 2015. Channel gating dependence on pore lining helix glycine residues in skeletal muscle ryanodine receptor. *J. Biol. Chem.* 290:17535–17545. <https://doi.org/10.1074/jbc.M115.659672>
- Meissner, G. 1975. Isolation and characterization of two types of sarcoplasmic reticulum vesicles. *Biochim. Biophys. Acta*. 389:51–68. [https://doi.org/10.1016/0005-2736\(75\)90385-5](https://doi.org/10.1016/0005-2736(75)90385-5)
- Meissner, G. 1984. Adenine nucleotide stimulation of Ca²⁺-induced Ca²⁺ release in sarcoplasmic reticulum. *J. Biol. Chem.* 259:2365–2374.
- Meissner, G. 1986. Ryanodine activation and inhibition of the Ca²⁺ release channel of sarcoplasmic reticulum. *J. Biol. Chem.* 261:6300–6306.
- Meissner, G. 1994. Ryanodine receptor/Ca²⁺ release channels and their regulation by endogenous effectors. *Annu. Rev. Physiol.* 56:485–508. <https://doi.org/10.1146/annurev.ph.56.030194.002413>
- Meissner, G., E. Darling, and J. Eveleth. 1986. Kinetics of rapid Ca²⁺ release by sarcoplasmic reticulum. Effects of Ca²⁺, Mg²⁺, and

- adenine nucleotides. *Biochemistry*. 25:236–244. <https://doi.org/10.1021/bi00349a033>
- Meissner, G., E. Rios, A. Tripathy, and D.A. Pasek. 1997. Regulation of skeletal muscle Ca^{2+} release channel (ryanodine receptor) by Ca^{2+} and monovalent cations and anions. *J. Biol. Chem.* 272:1628–1638. <https://doi.org/10.1074/jbc.272.3.1628>
- Meissner, G., D.A. Pasek, N. Yamaguchi, S. Ramachandran, N.V. Dokholyan, and A. Tripathy. 2009. Thermodynamics of calmodulin binding to cardiac and skeletal muscle ryanodine receptor ion channels. *Proteins*. 74:207–211. <https://doi.org/10.1002/prot.22148>
- Michalak, M., P. Dupraz, and V. Shoshan-Barmatz. 1988. Ryanodine binding to sarcoplasmic reticulum membrane; comparison between cardiac and skeletal muscle. *Biochim. Biophys. Acta*. 939:587–594. [https://doi.org/10.1016/0005-2736\(88\)90106-X](https://doi.org/10.1016/0005-2736(88)90106-X)
- Mitchell, R.D., A. Saito, P. Palade, and S. Fleischer. 1983. Morphology of isolated triads. *J. Cell Biol.* 96:1017–1029. <https://doi.org/10.1083/jcb.96.4.1017>
- Moore, C.P., G. Rodney, J.Z. Zhang, L. Santacruz-Toloz, G. Strasburg, and S.L. Hamilton. 1999. Apocalmodulin and Ca^{2+} calmodulin bind to the same region on the skeletal muscle Ca^{2+} release channel. *Biochemistry*. 38:8532–8537. <https://doi.org/10.1021/bi9907431>
- Morii, H., and Y. Tonomura. 1983. The gating behavior of a channel for Ca^{2+} -induced Ca^{2+} release in fragmented sarcoplasmic reticulum. *J. Biochem.* 93:1271–1285. <https://doi.org/10.1093/oxfordjournals.jbchem.a134261>
- Mowrey, D.D., L. Xu, Y. Mei, D.A. Pasek, G. Meissner, and N.V. Dokholyan. 2017. Ion-pulling simulations provide insights into the mechanisms of channel opening of the skeletal muscle ryanodine receptor. *J. Biol. Chem.* 292:12947–12958. <https://doi.org/10.1074/jbc.M116.760199>
- Murayama, T., and Y. Ogawa. 2004. RyR1 exhibits lower gain of CICR activity than RyR3 in the SR: Evidence for selective stabilization of RyR1 channel. *Am. J. Physiol. Cell Physiol.* 287:C36–C45. <https://doi.org/10.1152/ajpcell.00395.2003>
- Murayama, T., N. Kurebayashi, T. Oba, H. Oyama, K. Oguchi, T. Sakurai, and Y. Ogawa. 2011. Role of amino-terminal half of the S4-S5 linker in type 1 ryanodine receptor (RyR1) channel gating. *J. Biol. Chem.* 286:35571–35577. <https://doi.org/10.1074/jbc.M111.255240>
- Murphy, R.M., N.T. Larkins, J.P. Mollica, N.A. Beard, and G.D. Lamb. 2009. Calsequestrin content and SERCA determine normal and maximal Ca^{2+} storage levels in sarcoplasmic reticulum of fast- and slow-twitch fibres of rat. *J. Physiol.* 587:443–460. <https://doi.org/10.1113/jphysiol.2008.163162>
- Nakai, J., T. Imagawa, Y. Hakamat, M. Shigekawa, H. Takeshima, and S. Numa. 1990. Primary structure and functional expression from cDNA of the cardiac ryanodine receptor/calcium release channel. *FEBS Lett.* 271:169–177. [https://doi.org/10.1016/0014-5793\(90\)80399-4](https://doi.org/10.1016/0014-5793(90)80399-4)
- Natori, R. 1954. The property and contraction process of isolated myofibrils. *Jikeikai Med. J.* 1:119–126.
- Ngo, V.A., L.L. Perissinotti, W. Miranda, S.R.W. Chen, and S.Y. Noskov. 2017. Mapping ryanodine binding sites in the pore cavity of ryanodine receptors. *Biophys. J.* 112:1645–1653. <https://doi.org/10.1016/j.bpj.2017.03.014>
- Nomikos, M., A. Thanassoulas, K. Beck, V. Vassilakopoulou, H. Hu, B.L. Calver, M. Theodoridou, J. Kashir, L. Blayney, E. Livanou, et al. 2014. Altered RyR2 regulation by the calmodulin F90L mutation associated with idiopathic ventricular fibrillation and early sudden cardiac death. *FEBS Lett.* 588:2898–2902. <https://doi.org/10.1016/j.febslet.2014.07.007>
- O'Brien, J., G. Meissner, and B.A. Block. 1993. The fastest contracting muscles of nonmammalian vertebrates express only one isoform of the ryanodine receptor. *Biophys. J.* 65:2418–2427. [https://doi.org/10.1016/S0006-3495\(93\)81303-1](https://doi.org/10.1016/S0006-3495(93)81303-1)
- Ogawa, Y., and H. Harafuji. 1990. Osmolarity-dependent characteristics of [^3H]ryanodine binding to sarcoplasmic reticulum. *J. Biochem.* 107:894–898. <https://doi.org/10.1093/oxfordjournals.jbchem.a123144>
- Oo, Y.W., N. Gomez-Hurtado, K. Walweel, D.F. van Helden, M.S. Imtiaz, B.C. Knollmann, and D.R. Laver. 2015. Essential role of calmodulin in RyR inhibition by dantrolene. *Mol. Pharmacol.* 88:57–63. <https://doi.org/10.1124/mol.115.097691>
- Otsu, K., H.F. Willard, V.K. Khanna, F. Zorzato, N.M. Green, and D.H. MacLennan. 1990. Molecular cloning of cDNA encoding the Ca^{2+} release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic reticulum. *J. Biol. Chem.* 265:13472–13483.
- Oyamada, H., T. Murayama, T. Takagi, M. Iino, N. Iwabe, T. Miyata, Y. Ogawa, and M. Endo. 1994. Primary structure and distribution of ryanodine-binding protein isoforms of the bullfrog skeletal muscle. *J. Biol. Chem.* 269:17206–17214.
- Park, H., I.Y. Park, E. Kim, B. Youn, K. Fields, A.K. Dunker, and C. Kang. 2004. Comparing skeletal and cardiac calsequestrin structures and their calcium binding: A proposed mechanism for coupled calcium binding and protein polymerization. *J. Biol. Chem.* 279:18026–18033. <https://doi.org/10.1074/jbc.M311553200>
- Patterson, R.L., D. Boehning, and S.H. Snyder. 2004. Inositol 1,4,5-trisphosphate receptors as signal integrators. *Annu. Rev. Biochem.* 73:437–465. <https://doi.org/10.1146/annurev.biochem.73.071403.161303>
- Peng, W., H. Shen, J. Wu, W. Guo, X. Pan, R. Wang, S.R. Chen, and N. Yan. 2016. Structural basis for the gating mechanism of the type 2 ryanodine receptor RyR2. *Science*. 354:1–10. <https://doi.org/10.1126/science.aah5324>
- Percival, A.L., A.J. Williams, J.L. Kenyon, M.M. Grinsell, J.A. Airey, and J.L. Sutko. 1994. Chicken skeletal muscle ryanodine receptor isoforms: Ion channel properties. *Biophys. J.* 67:1834–1850. [https://doi.org/10.1016/S0006-3495\(94\)80665-4](https://doi.org/10.1016/S0006-3495(94)80665-4)
- Pessah, I.N., and I. Zimanyi. 1991. Characterization of multiple [^3H]ryanodine binding sites on the Ca^{2+} release channel of sarcoplasmic reticulum from skeletal and cardiac muscle: Evidence for a sequential mechanism in ryanodine action. *Mol. Pharmacol.* 39:679–689.
- Pessah, I.N., A.L. Waterhouse, and J.E. Casida. 1985. The calcium-ryanodine receptor complex of skeletal and cardiac muscle. *Biochem. Biophys. Res. Commun.* 128:449–456. [https://doi.org/10.1016/0006-291X\(85\)91699-7](https://doi.org/10.1016/0006-291X(85)91699-7)
- Pessah, I.N., A.O. Francini, D.J. Scales, A.L. Waterhouse, and J.E. Casida. 1986. Calcium-ryanodine receptor complex. Solubilization and partial characterization from skeletal muscle junctional sarcoplasmic reticulum vesicles. *J. Biol. Chem.* 261:8643–8648.
- Pessah, I.N., R.A. Stambuk, and J.E. Casida. 1987. Ca^{2+} -activated ryanodine binding: mechanisms of sensitivity and intensity modulation by Mg^{2+} , caffeine, and adenine nucleotides. *Mol. Pharmacol.* 31:232–238.
- Petrotschenko, E.V., N. Yamaguchi, D.A. Pasek, C.H. Borchers, and G. Meissner. 2011. Mass spectrometric analysis and mutagenesis predict involvement of multiple cysteines in redox regulation of the skeletal muscle ryanodine receptor ion channel complex. *Res. Rep. Biol.* 2011:13–21. <https://doi.org/10.2147/RRB.S15776>
- Popova, O.B., M.R. Baker, T.P. Tran, T. Le, and I.I. Serysheva. 2012. Identification of ATP-binding regions in the RyR1 Ca^{2+} release channel. *PLoS One*. 7:e48725. <https://doi.org/10.1371/journal.pone.0048725>
- Posterino, G.S., and G.D. Lamb. 2003. Effect of sarcoplasmic reticulum Ca^{2+} content on action potential-induced Ca^{2+} release

- in rat skeletal muscle fibres. *J. Physiol.* 551:219–237. <https://doi.org/10.1113/jphysiol.2003.040022>
- Postma, A.V., I. Denjoy, T.M. Hoornijde, J.M. Lupoglazoff, A. Da Costa, P. Sebillon, M.M. Mannens, A.A. Wilde, and P. Guicheney. 2002. Absence of calsequestrin 2 causes severe forms of catecholaminergic polymorphic ventricular tachycardia. *Circ. Res.* 91:e21–e26. <https://doi.org/10.1161/01.RES.0000038886.18992.6B>
- Pouliquin, P., and A.F. Dulhunty. 2009. Homer and the ryanodine receptor. *Eur. Biophys. J.* 39:91–102. <https://doi.org/10.1007/s00249-009-0494-1>
- Prosser, B.L., N.T. Wright, E.O. Hernández-Ochoa, K.M. Varney, Y. Liu, R.O. Olojo, D.B. Zimmer, D.J. Weber, and M.F. Schneider. 2008. S100A1 binds to the calmodulin-binding site of ryanodine receptor and modulates skeletal muscle excitation-contraction coupling. *J. Biol. Chem.* 283:5046–5057. <https://doi.org/10.1074/jbc.M709231200>
- Protasi, F., C. Franzini-Armstrong, and B.E. Flucher. 1997. Coordinated incorporation of skeletal muscle dihydropyridine receptors and ryanodine receptors in peripheral couplings of BC3H1 cells. *J. Cell Biol.* 137:859–870. <https://doi.org/10.1083/jcb.137.4.859>
- Protasi, F., C. Franzini-Armstrong, and P.D. Allen. 1998. Role of ryanodine receptors in the assembly of calcium release units in skeletal muscle. *J. Cell Biol.* 140:831–842. <https://doi.org/10.1083/jcb.140.4.831>
- Radermacher, M., V. Rao, R. Grassucci, J. Frank, A.P. Timerman, S. Fleischer, and T. Wagenknecht. 1994. Cryo-electron microscopy and three-dimensional reconstruction of the calcium release channel/ryanodine receptor from skeletal muscle. *J. Cell Biol.* 127:411–423. <https://doi.org/10.1083/jcb.127.2.411>
- Ramachandran, S., A. Chakraborty, L. Xu, Y. Mei, M. Samsó, N.V. Dokholyan, and G. Meissner. 2013. Structural determinants of skeletal muscle ryanodine receptor gating. *J. Biol. Chem.* 288:6154–6165. <https://doi.org/10.1074/jbc.M112.433789>
- Rebeck, R.T., F.R. Nitu, D. Rohde, P. Most, D.M. Bers, D.D. Thomas, and R.L. Cornea. 2016. S100A1 protein does not compete with calmodulin for ryanodine receptor binding but structurally alters the ryanodine receptor-calmodulin complex. *J. Biol. Chem.* 291:15896–15907. <https://doi.org/10.1074/jbc.M115.713107>
- Reiken, S., M. Gaburjakova, S. Guatimosim, A.M. Gomez, J. D'Armiento, D. Burkhoff, J. Wang, G. Vassort, W.J. Lederer, and A.R. Marks. 2003. Protein kinase A phosphorylation of the cardiac calcium release channel (ryanodine receptor) in normal and failing hearts. Role of phosphatases and response to isoproterenol. *J. Biol. Chem.* 278:444–453. <https://doi.org/10.1074/jbc.M207028200>
- Respress, J.L., R.J. van Oort, N. Li, N. Rolim, S.S. Dixit, A. deAlmeida, N. Voigt, W.S. Lawrence, D.G. Skapura, K. Skårdal, et al. 2012. Role of RyR2 phosphorylation at S2814 during heart failure progression. *Circ. Res.* 110:1474–1483. <https://doi.org/10.1161/CIRCRESAHA.112.268094>
- Ríos, E., and G. Pizarro. 1991. Voltage sensor of excitation-contraction coupling in skeletal muscle. *Physiol. Rev.* 71:849–908.
- Robinson, R., D. Carpenter, M.A. Shaw, J. Halsall, and P. Hopkins. 2006. Mutations in RYR1 in malignant hyperthermia and central core disease. *Hum. Mutat.* 27:977–989. <https://doi.org/10.1002/humu.20356>
- Rodney, G.G., C.P. Moore, B.Y. Williams, J.Z. Zhang, J. Krol, S.E. Pedersen, and S.L. Hamilton. 2001. Calcium binding to calmodulin leads to an N-terminal shift in its binding site on the ryanodine receptor. *J. Biol. Chem.* 276:2069–2074. <https://doi.org/10.1074/jbc.M008891200>
- Rousseau, E., J.S. Smith, J.S. Henderson, and G. Meissner. 1986. Single channel and 45Ca^{2+} flux measurements of the cardiac sarcoplasmic reticulum calcium channel. *Biophys. J.* 50:1009–1014. [https://doi.org/10.1016/S0006-3495\(86\)83543-3](https://doi.org/10.1016/S0006-3495(86)83543-3)
- Rousseau, E., J. Ladine, Q.Y. Liu, and G. Meissner. 1988. Activation of the Ca^{2+} release channel of skeletal muscle sarcoplasmic reticulum by caffeine and related compounds. *Arch. Biochem. Biophys.* 267:75–86. [https://doi.org/10.1016/0003-9861\(88\)90010-0](https://doi.org/10.1016/0003-9861(88)90010-0)
- Saito, A., M. Intui, M. Radermacher, J. Frank, and S. Fleischer. 1988. Ultrastructure of the calcium release channel of sarcoplasmic reticulum. *J. Cell Biol.* 107:211–219. <https://doi.org/10.1083/jcb.107.1.211>
- Samsó, M. 2017. A guide to the 3D structure of the ryanodine receptor type 1 by cryoEM. *Protein Sci.* 26:52–68. <https://doi.org/10.1002/pro.3052>
- Samsó, M., and T. Wagenknecht. 2002. Apocalmodulin and Ca^{2+} -calmodulin bind to neighboring locations on the ryanodine receptor. *J. Biol. Chem.* 277:1349–1353. <https://doi.org/10.1074/jbc.M109196200>
- Samsó, M., T. Wagenknecht, and P.D. Allen. 2005. Internal structure and visualization of transmembrane domains of the RyR1 calcium release channel by cryo-EM. *Nat. Struct. Mol. Biol.* 12:539–544. <https://doi.org/10.1038/nsmb938>
- Samsó, M., X. Shen, and P.D. Allen. 2006. Structural characterization of the RyR1-FKBP12 interaction. *J. Mol. Biol.* 356:917–927. <https://doi.org/10.1016/j.jmb.2005.12.023>
- Samsó, M., W. Feng, I.N. Pessah, and P.D. Allen. 2009. Coordinated movement of cytoplasmic and transmembrane domains of RyR1 upon gating. *PLoS Biol.* 7:e85. <https://doi.org/10.1371/journal.pbio.1000085>
- Sárközi, S., I. Komáromi, I. Jóna, and J. Almássy. 2017. Lanthanides report calcium sensor in the vestibule of ryanodine receptor. *Biophys. J.* 112:2127–2137. <https://doi.org/10.1016/j.bpj.2017.03.023>
- Sato, Y., D.G. Ferguson, H. Sako, G.W. Dorn II, V.J. Kadambi, A. Yatani, B.D. Hoit, R.A. Walsh, and E.G. Kranias. 1998. Cardiac-specific overexpression of mouse cardiac calsequestrin is associated with depressed cardiovascular function and hypertrophy in transgenic mice. *J. Biol. Chem.* 273:28470–28477. <https://doi.org/10.1074/jbc.273.43.28470>
- Schneider, M.F., and W.K. Chandler. 1973. Voltage dependent charge movement of skeletal muscle: A possible step in excitation-contraction coupling. *Nature.* 242:244–246. <https://doi.org/10.1038/242244a0>
- Seo, M.D., M. Enomoto, N. Ishiyama, P.B. Stathopoulos, and M. Ikura. 2015. Structural insights into endoplasmic reticulum stored calcium regulation by inositol 1,4,5-trisphosphate and ryanodine receptors. *Biochim. Biophys. Acta.* 1853:1980–1991. <https://doi.org/10.1016/j.bbamcr.2014.11.023>
- Serysheva, I.I., E.V. Orlova, W. Chiu, M.B. Sherman, S.L. Hamilton, and M. van Heel. 1995. Electron cryomicroscopy and angular reconstitution used to visualize the skeletal muscle calcium release channel. *Nat. Struct. Biol.* 2:18–24. <https://doi.org/10.1038/nsb0195-18>
- Shannon, T.R., K.S. Ginsburg, and D.M. Bers. 2000. Potentiation of fractional sarcoplasmic reticulum calcium release by total and free intra-sarcoplasmic reticulum calcium concentration. *Biophys. J.* 78:334–343. [https://doi.org/10.1016/S0006-3495\(00\)76596-9](https://doi.org/10.1016/S0006-3495(00)76596-9)
- Sharma, M.R., L.H. Jeyakumar, S. Fleischer, and T. Wagenknecht. 2006. Three-dimensional visualization of FKBP12.6 binding to an open conformation of cardiac ryanodine receptor. *Biophys. J.* 90:164–172. <https://doi.org/10.1529/biophysj.105.063503>
- Sharma, P., N. Ishiyama, U. Nair, W. Li, A. Dong, T. Miyake, A. Wilson, T. Ryan, D.H. MacLennan, T. Kislinger, et al. 2012. Structural determination of the phosphorylation domain of the ryanodine receptor. *FEBS J.* 279:3952–3964. <https://doi.org/10.1111/j.1742-4658.2012.08755.x>

- Sheridan, D.C., H. Takekura, C. Franzini-Armstrong, K.G. Beam, P.D. Allen, and C.F. Perez. 2006. Bidirectional signaling between calcium channels of skeletal muscle requires multiple direct and indirect interactions. *Proc. Natl. Acad. Sci. USA*. 103:19760–19765. <https://doi.org/10.1073/pnas.0609473103>
- Shin, D.W., J. Ma, and D.H. Kim. 2000. The asp-rich region at the carboxyl-terminus of calsequestrin binds to Ca^{2+} and interacts with triadin. *FEBS Lett.* 486:178–182. [https://doi.org/10.1016/S0014-5793\(00\)02246-8](https://doi.org/10.1016/S0014-5793(00)02246-8)
- Shin, D.W., Z. Pan, A. Bandyopadhyay, M.B. Bhat, D.H. Kim, and J. Ma. 2002. Ca^{2+} -dependent interaction between FKBP12 and calcineurin regulates activity of the Ca^{2+} release channel in skeletal muscle. *Biophys. J.* 83:2539–2549. [https://doi.org/10.1016/S0006-3495\(02\)75265-X](https://doi.org/10.1016/S0006-3495(02)75265-X)
- Sitsapesan, R., and A.J. Williams. 1994. Regulation of the gating of the sheep cardiac sarcoplasmic reticulum Ca^{2+} -release channel by luminal Ca^{2+} . *J. Membr. Biol.* 137:215–226. <https://doi.org/10.1007/BF00232590>
- Sitsapesan, R., and A.J. Williams. 1995. The gating of the sheep skeletal sarcoplasmic reticulum Ca^{2+} -release channel is regulated by luminal Ca^{2+} . *J. Membr. Biol.* 146:133–144. <https://doi.org/10.1007/BF00238004>
- Smith, J.S., R. Coronado, and G. Meissner. 1985. Sarcoplasmic reticulum contains adenine nucleotide-activated calcium channels. *Nature*. 316:446–449. <https://doi.org/10.1038/316446a0>
- Smith, J.S., R. Coronado, and G. Meissner. 1986. Single channel measurements of the calcium release channel from skeletal muscle sarcoplasmic reticulum. Activation by Ca^{2+} and ATP and modulation by Mg^{2+} . *J. Gen. Physiol.* 88:573–588. <https://doi.org/10.1085/jgp.88.5.573>
- Smith, J.S., T. Imagawa, J. Ma, M. Fill, K.P. Campbell, and R. Coronado. 1988. Purified ryanodine receptor from rabbit skeletal muscle is the calcium-release channel of sarcoplasmic reticulum. *J. Gen. Physiol.* 92:1–26. <https://doi.org/10.1085/jgp.92.1.1>
- Sommer, J.R., and E.A. Johnson. 1979. Ultrastructure of cardiac muscle. In *Handbook of Physiology*. R.M. Berne, editor. American Physiological Society, Bethesda, MD. Sec. 2, 113–186.
- Søndergaard, M.T., Y. Liu, K.T. Larsen, A. Nani, X. Tian, C. Holt, R. Wang, R. Wimmer, F. Van Petegem, M. Fill, et al. 2017. The arrhythmogenic calmodulin p.Phe142Leu mutation impairs C-domain Ca^{2+} binding but not calmodulin-dependent inhibition of the cardiac ryanodine receptor. *J. Biol. Chem.* 292:1385–1395. <https://doi.org/10.1074/jbc.M116.766253>
- Stange, M., L. Xu, D. Balshaw, N. Yamaguchi, and G. Meissner. 2003. Characterization of recombinant skeletal muscle (Ser-2843) and cardiac muscle (Ser-2809) ryanodine receptor phosphorylation mutants. *J. Biol. Chem.* 278:51693–51702. <https://doi.org/10.1074/jbc.M310406200>
- Stephenson, E.W. 1981. Activation of fast skeletal muscle: Contributions of studies on skinned fibers. *Am. J. Physiol.* 240:C1–C19.
- Stowell, K.M. 2008. Malignant hyperthermia: A pharmacogenetic disorder. *Pharmacogenomics*. 9:1657–1672. <https://doi.org/10.2217/14622416.9.11.1657>
- Suko, J., I. Maurer-Fogy, B. Plank, O. Bertel, W. Wyskovsky, M. Hohenegger, and G. Hellmann. 1993. Phosphorylation of serine 2843 in ryanodine receptor-calcium release channel of skeletal muscle by cAMP-, cGMP- and CaM-dependent protein kinase. *Biochim. Biophys. Acta*. 1175:193–206. [https://doi.org/10.1016/0167-4889\(93\)90023-I](https://doi.org/10.1016/0167-4889(93)90023-I)
- Sun, B., W. Guo, X. Tian, J. Yao, L. Zhang, R. Wang, and S.R. Chen. 2016. The cytoplasmic region of inner helix S6 is an important determinant of cardiac ryanodine receptor channel gating. *J. Biol. Chem.* 291:26024–26034. <https://doi.org/10.1074/jbc.M116.758821>
- Sun, J., C. Xin, J.P. Eu, J.S. Stamler, and G. Meissner. 2001a. Cysteine-3635 is responsible for skeletal muscle ryanodine receptor modulation by NO. *Proc. Natl. Acad. Sci. USA*. 98:11158–11162. <https://doi.org/10.1073/pnas.201289098>
- Sun, J., L. Xu, J.P. Eu, J.S. Stamler, and G. Meissner. 2001b. Classes of thiols that influence the activity of the skeletal muscle calcium release channel. *J. Biol. Chem.* 276:15625–15630. <https://doi.org/10.1074/jbc.M100083200>
- Sun, J., N. Yamaguchi, L. Xu, J.P. Eu, J.S. Stamler, and G. Meissner. 2008. Regulation of the cardiac muscle ryanodine receptor by O(2) tension and S-nitrosoglutathione. *Biochemistry*. 47:13985–13990. <https://doi.org/10.1021/bi8012627>
- Sun, Q.A., B. Wang, M. Miyagi, D.T. Hess, and J.S. Stamler. 2013. Oxygen-coupled redox regulation of the skeletal muscle ryanodine receptor/ Ca^{2+} release channel (RyR1): Sites and nature of oxidative modification. *J. Biol. Chem.* 288:22961–22971. <https://doi.org/10.1074/jbc.M113.480228>
- Sutko, J.L., K. Ito, and J.L. Kenyon. 1985. Ryanodine: A modifier of sarcoplasmic reticulum calcium release in striated muscle. *Fed. Proc.* 44:2984–2988.
- Sutko, J.L., J.A. Airey, W. Welch, and L. Ruest. 1997. The pharmacology of ryanodine and related compounds. *Pharmacol. Rev.* 49:53–98.
- Sztretye, M., J. Yi, L. Figueroa, J. Zhou, L. Royer, P. Allen, G. Brum, and E. Ríos. 2011. Measurement of RyR permeability reveals a role of calsequestrin in termination of SR Ca^{2+} release in skeletal muscle. *J. Gen. Physiol.* 138:231–247. <https://doi.org/10.1085/jgp.201010592>
- Takasago, T., T. Imagawa, K. Furukawa, T. Ogurusu, and M. Shigekawa. 1991. Regulation of the cardiac ryanodine receptor by protein kinase-dependent phosphorylation. *J. Biochem.* 109:163–170. <https://doi.org/10.1093/oxfordjournals.jbchem.a123339>
- Takekura, H., M. Nishi, T. Noda, H. Takeshima, and C. Franzini-Armstrong. 1995. Abnormal junctions between surface membrane and sarcoplasmic reticulum in skeletal muscle with a mutation targeted to the ryanodine receptor. *Proc. Natl. Acad. Sci. USA*. 92:3381–3385. <https://doi.org/10.1073/pnas.92.8.3381>
- Takeshima, H., S. Nishimura, T. Matsumoto, H. Ishida, K. Kangawa, N. Minamino, H. Matsuo, M. Ueda, M. Hanaoka, T. Hirose, and S. Numa. 1989. Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature*. 339:439–445. <https://doi.org/10.1038/339439a0>
- Takeshima, H., M. Shimuta, S. Komazaki, K. Ohmi, M. Nishi, M. Iino, A. Miyata, and K. Kangawa. 1998. Mitsugumin29, a novel synaptophysin family member from the triad junction in skeletal muscle. *Biochem. J.* 331:317–322. <https://doi.org/10.1042/bj3310317>
- Terentyev, D., S. Viatchenko-Karpinski, I. Györke, P. Volpe, S.C. Williams, and S. Györke. 2003. Calsequestrin determines the functional size and stability of cardiac intracellular calcium stores: Mechanism for hereditary arrhythmia. *Proc. Natl. Acad. Sci. USA*. 100:11759–11764. <https://doi.org/10.1073/pnas.1932318100>
- Thomas, N.L., and A.J. Williams. 2012. Pharmacology of ryanodine receptors and Ca^{2+} -induced Ca^{2+} release. *WIREs Membr. Transp. Signal.* 1:383–397. <https://doi.org/10.1002/wmts.34>
- Tijsskens, P., G. Meissner, and C. Franzini-Armstrong. 2003. Location of ryanodine and dihydropyridine receptors in frog myocardium. *Biophys. J.* 84:1079–1092. [https://doi.org/10.1016/S0006-3495\(03\)74924-8](https://doi.org/10.1016/S0006-3495(03)74924-8)
- Timerman, A.P., H. Onoue, H.B. Xin, S. Barg, J. Copello, G. Wiederrecht, and S. Fleischer. 1996. Selective binding of FKBP12.6 by the cardiac ryanodine receptor. *J. Biol. Chem.* 271:20385–20391. <https://doi.org/10.1074/jbc.271.34.20385>

- Tinker, A., and A.J. Williams. 1992. Divalent cation conduction in the ryanodine receptor channel of sheep cardiac muscle sarcoplasmic reticulum. *J. Gen. Physiol.* 100:479–493. <https://doi.org/10.1085/jgp.100.3.479>
- Tinker, A., and A.J. Williams. 1993. Probing the structure of the conduction pathway of the sheep cardiac sarcoplasmic reticulum calcium-release channel with permeant and impermeant organic cations. *J. Gen. Physiol.* 102:1107–1129. <https://doi.org/10.1085/jgp.102.6.1107>
- Tinker, A., A.R. Lindsay, and A.J. Williams. 1992. A model for ionic conduction in the ryanodine receptor channel of sheep cardiac muscle sarcoplasmic reticulum. *J. Gen. Physiol.* 100:495–517. <https://doi.org/10.1085/jgp.100.3.495>
- Treves, S., H. Jungbluth, F. Muntoni, and F. Zorzato. 2008. Congenital muscle disorders with cores: The ryanodine receptor calcium channel paradigm. *Curr. Opin. Pharmacol.* 8:319–326. <https://doi.org/10.1016/j.coph.2008.01.005>
- Tripathy, A., and G. Meissner. 1996. Sarcoplasmic reticulum luminal Ca^{2+} has access to cytosolic activation and inactivation sites of skeletal muscle Ca^{2+} release channel. *Biophys. J.* 70:2600–2615. [https://doi.org/10.1016/S0006-3495\(96\)79831-4](https://doi.org/10.1016/S0006-3495(96)79831-4)
- Tunwell, R.E., C. Wickenden, B.M. Bertrand, V.I. Shevchenko, M.B. Walsh, P.D. Allen, and F.A. Lai. 1996. The human cardiac muscle ryanodine receptor-calcium release channel: Identification, primary structure and topological analysis. *Biochem. J.* 318:477–487. <https://doi.org/10.1042/bj3180477>
- Uehara, A., T. Murayama, M. Yasukochi, M. Fill, M. Horie, T. Okamoto, Y. Matsuura, K. Uehara, T. Fujimoto, T. Sakurai, and N. Kurebayashi. 2017. Extensive Ca^{2+} leak through K4750Q cardiac ryanodine receptors caused by cytosolic and luminal Ca^{2+} hypersensitivity. *J. Gen. Physiol.* 149:199–218. <https://doi.org/10.1085/jgp.201611624>
- van Oort, R.J., M.D. McCauley, S.S. Dixit, L. Pereira, Y. Yang, J.L. Respress, Q. Wang, A.C. De Almeida, D.G. Skapura, M.E. Anderson, et al. 2010. Ryanodine receptor phosphorylation by calcium/calmodulin-dependent protein kinase II promotes life-threatening ventricular arrhythmias in mice with heart failure. *Circulation.* 122:2669–2679. <https://doi.org/10.1161/CIRCULATIONAHA.110.982298>
- Voss, A.A., J. Lango, M. Ernst-Russell, D. Morin, and I.N. Pessah. 2004. Identification of hyperreactive cysteines within ryanodine receptor type 1 by mass spectrometry. *J. Biol. Chem.* 279:34514–34520. <https://doi.org/10.1074/jbc.M404290200>
- Wagenknecht, T., R. Grassucci, J. Frank, A. Saito, M. Inui, and S. Fleischer. 1989. Three-dimensional architecture of the calcium channel/foot structure of sarcoplasmic reticulum. *Nature.* 338:167–170. <https://doi.org/10.1038/338167a0>
- Wagenknecht, T., J. Berkowitz, R. Grassucci, A.P. Timerman, and S. Fleischer. 1994. Localization of calmodulin binding sites on the ryanodine receptor from skeletal muscle by electron microscopy. *Biophys. J.* 67:2286–2295. [https://doi.org/10.1016/S0006-3495\(94\)80714-3](https://doi.org/10.1016/S0006-3495(94)80714-3)
- Wagenknecht, T., R. Grassucci, J. Berkowitz, G.J. Wiederrecht, H.B. Xin, and S. Fleischer. 1996. Cryoelectron microscopy resolves FK506-binding protein sites on the skeletal muscle ryanodine receptor. *Biophys. J.* 70:1709–1715. [https://doi.org/10.1016/S0006-3495\(96\)79733-3](https://doi.org/10.1016/S0006-3495(96)79733-3)
- Wagenknecht, T., M. Radermacher, R. Grassucci, J. Berkowitz, H.B. Xin, and S. Fleischer. 1997. Locations of calmodulin and FK506-binding protein on the three-dimensional architecture of the skeletal muscle ryanodine receptor. *J. Biol. Chem.* 272:32463–32471. <https://doi.org/10.1074/jbc.272.51.32463>
- Walweel, K., J. Li, P. Molenaar, M.S. Imtiaz, A. Quail, C.G. dos Remedios, N.A. Beard, A.F. Dulhunty, D.F. van Helden, and D.R. Laver. 2014. Differences in the regulation of RyR2 from human, sheep, and rat by Ca^{2+} and Mg^{2+} in the cytoplasm and in the lumen of the sarcoplasmic reticulum. *J. Gen. Physiol.* 144:263–271. <https://doi.org/10.1085/jgp.201311157>
- Wang, Y., L. Xu, D.A. Pasek, D. Gillespie, and G. Meissner. 2005. Probing the role of negatively charged amino acid residues in ion permeation of skeletal muscle ryanodine receptor. *Biophys. J.* 89:256–265. <https://doi.org/10.1529/biophysj.104.056002>
- Wehrens, X.H., S.E. Lehnart, S.R. Reiken, and A.R. Marks. 2004. Ca^{2+} /calmodulin-dependent protein kinase II phosphorylation regulates the cardiac ryanodine receptor. *Circ. Res.* 94:e61–e70. <https://doi.org/10.1161/01.RES.0000125626.33738.E2>
- Wehrens, X.H., S.E. Lehnart, S. Reiken, J.A. Vest, A. Wronska, and A.R. Marks. 2006. Ryanodine receptor/calcium release channel PKA phosphorylation: A critical mediator of heart failure progression. *Proc. Natl. Acad. Sci. USA.* 103:511–518. <https://doi.org/10.1073/pnas.0510113103>
- Wei, R., X. Wang, Y. Zhang, S. Mukherjee, L. Zhang, Q. Chen, X. Huang, S. Jing, C. Liu, S. Li, et al. 2016. Structural insights into Ca^{2+} -activated long-range allosteric channel gating of RyR1. *Cell Res.* 26:977–994. <https://doi.org/10.1038/cr.2016.99>
- Witcher, D.R., R.J. Kovacs, H. Schulman, D.C. Cefali, and L.R. Jones. 1991. Unique phosphorylation site on the cardiac ryanodine receptor regulates calcium channel activity. *J. Biol. Chem.* 266:11144–11152.
- Woodier, J., R.D. Rainbow, A.J. Stewart, and S.J. Pitt. 2015. Intracellular zinc modulates cardiac ryanodine receptor-mediated calcium release. *J. Biol. Chem.* 290:17599–17610. <https://doi.org/10.1074/jbc.M115.661280>
- Wright, N.T., B.L. Prosser, K.M. Varney, D.B. Zimmer, M.F. Schneider, and D.J. Weber. 2008. S100A1 and calmodulin compete for the same binding site on ryanodine receptor. *J. Biol. Chem.* 283:26676–26683. <https://doi.org/10.1074/jbc.M804432200>
- Wu, J., Z. Yan, Z. Li, X. Qian, S. Lu, M. Dong, Q. Zhou, and N. Yan. 2016. Structure of the voltage-gated calcium channel $\text{Ca}_v1.1$ at 3.6 Å resolution. *Nature.* 537:191–196. <https://doi.org/10.1038/nature19321>
- Xiao, B., M.T. Jiang, M. Zhao, D. Yang, C. Sutherland, F.A. Lai, M.P. Walsh, D.C. Warltier, H. Cheng, and S.R.W. Chen. 2005. Characterization of a novel PKA phosphorylation site, serine-2030, reveals no PKA hyperphosphorylation of the cardiac ryanodine receptor in canine heart failure. *Circ. Res.* 96:847–855. <https://doi.org/10.1161/01.RES.0000163276.26083.e8>
- Xiao, B., X. Tian, W. Xie, P.P. Jones, S. Cai, X. Wang, D. Jiang, H. Kong, L. Zhang, K. Chen, et al. 2007. Functional consequence of protein kinase A-dependent phosphorylation of the cardiac ryanodine receptor: Sensitization of store overload-induced Ca^{2+} release. *J. Biol. Chem.* 282:30256–30264. <https://doi.org/10.1074/jbc.M703510200>
- Xiong, H., X. Feng, L. Gao, L. Xu, D.A. Pasek, J.H. Seok, and G. Meissner. 1998. Identification of a two EF-hand Ca^{2+} binding domain in lobster skeletal muscle ryanodine receptor/ Ca^{2+} release channel. *Biochemistry.* 37:4804–4814. <https://doi.org/10.1021/bi971198b>
- Xiong, L., J.Z. Zhang, R. He, and S.L. Hamilton. 2006. A Ca^{2+} -binding domain in RyR1 that interacts with the calmodulin binding site and modulates channel activity. *Biophys. J.* 90:173–182. <https://doi.org/10.1529/biophysj.105.066092>
- Xu, L., and G. Meissner. 1998. Regulation of cardiac muscle Ca^{2+} release channel by sarcoplasmic reticulum luminal Ca^{2+} . *Biophys. J.* 75:2302–2312. [https://doi.org/10.1016/S0006-3495\(98\)77674-X](https://doi.org/10.1016/S0006-3495(98)77674-X)
- Xu, L., and G. Meissner. 2004. Mechanism of calmodulin inhibition of cardiac sarcoplasmic reticulum Ca^{2+} release channel (ryanodine receptor). *Biophys. J.* 86:797–804. [https://doi.org/10.1016/S0006-3495\(04\)74155-7](https://doi.org/10.1016/S0006-3495(04)74155-7)

- Xu, L., G. Mann, and G. Meissner. 1996. Regulation of cardiac Ca^{2+} release channel (ryanodine receptor) by Ca^{2+} , H^+ , Mg^{2+} , and adenine nucleotides under normal and simulated ischemic conditions. *Circ. Res.* 79:1100–1109. <https://doi.org/10.1161/01.RES.79.6.1100>
- Xu, L., A. Tripathy, D.A. Pasek, and G. Meissner. 1998. Potential for pharmacology of ryanodine receptor/calcium release channels. *Ann. N. Y. Acad. Sci.* 853(1 CARDIAC SARCO):130–148. <https://doi.org/10.1111/j.1749-6632.1998.tb08262.x>
- Xu, L., Y. Wang, D. Gillespie, and G. Meissner. 2006. Two rings of negative charges in the cytosolic vestibule of type-1 ryanodine receptor modulate ion fluxes. *Biophys. J.* 90:443–453. <https://doi.org/10.1529/biophysj.105.072538>
- Xu, L., Y. Wang, N. Yamaguchi, D.A. Pasek, and G. Meissner. 2008. Single channel properties of heterotetrameric mutant RyR1 ion channels linked to core myopathies. *J. Biol. Chem.* 283:6321–6329. <https://doi.org/10.1074/jbc.M707353200>
- Xu, L., A.C. Gomez, D.A. Pasek, G. Meissner, and N. Yamaguchi. 2017. Two EF-hand motifs in ryanodine receptor calcium release channels contribute to isoform-specific regulation by calmodulin. *Cell Calcium.* 66:62–70. <https://doi.org/10.1016/j.ceca.2017.05.013>
- Yamaguchi, N., C. Xin, and G. Meissner. 2001. Identification of apocalmodulin and Ca^{2+} -calmodulin regulatory domain in skeletal muscle Ca^{2+} release channel, ryanodine receptor. *J. Biol. Chem.* 276:22579–22585. <https://doi.org/10.1074/jbc.M102729200>
- Yamaguchi, N., L. Xu, D.A. Pasek, K.E. Evans, and G. Meissner. 2003. Molecular basis of calmodulin binding to cardiac muscle Ca^{2+} release channel (ryanodine receptor). *J. Biol. Chem.* 278:23480–23486. <https://doi.org/10.1074/jbc.M301125200>
- Yamaguchi, N., L. Xu, D.A. Pasek, K.E. Evans, S.R. Chen, and G. Meissner. 2005. Calmodulin regulation and identification of calmodulin binding region of type-3 ryanodine receptor calcium release channel. *Biochemistry.* 44:15074–15081. <https://doi.org/10.1021/bi051251t>
- Yamaguchi, N., N. Takahashi, L. Xu, O. Smithies, and G. Meissner. 2007. Early cardiac hypertrophy in mice with impaired calmodulin regulation of cardiac muscle Ca release channel. *J. Clin. Invest.* 117:1344–1353. <https://doi.org/10.1172/JCI29515>
- Yamaguchi, N., B.L. Prosser, F. Ghassemi, L. Xu, D.A. Pasek, J.P. Eu, E.O. Hernández-Ochoa, B.R. Cannon, P.T. Wilder, R.M. Lovering, et al. 2011. Modulation of sarcoplasmic reticulum Ca^{2+} release in skeletal muscle expressing ryanodine receptor impaired in regulation by calmodulin and S100A1. *Am. J. Physiol. Cell Physiol.* 300:C998–C1012. <https://doi.org/10.1152/ajpcell.00370.2010>
- Yamaguchi, N., A. Chakraborty, T.Q. Huang, L. Xu, A.C. Gomez, D.A. Pasek, and G. Meissner. 2013. Cardiac hypertrophy associated with impaired regulation of cardiac ryanodine receptor by calmodulin and S100A1. *Am. J. Physiol. Heart Circ. Physiol.* 305:H86–H94. <https://doi.org/10.1152/ajpheart.00144.2013>
- Yan, Z., X. Bai, C. Yan, J. Wu, Z. Li, T. Xie, W. Peng, C. Yin, X. Li, S.H.W. Scheres, et al. 2015. Structure of the rabbit ryanodine receptor RyR1 at near-atomic resolution. *Nature.* 517:50–55. <https://doi.org/10.1038/nature14063>
- Yuan, Q., G.C. Fan, M. Dong, B. Altschaffl, A. Diwan, X. Ren, H.H. Hahn, W. Zhao, J.R. Waggoner, L.R. Jones, et al. 2007. Sarcoplasmic reticulum calcium overloading in junctin deficiency enhances cardiac contractility but increases ventricular automaticity. *Circulation.* 115:300–309. <https://doi.org/10.1161/CIRCULATIONAHA.106.654699>
- Yuchi, Z., and F. Van Petegem. 2016. Ryanodine receptors under the magnifying lens: Insights and limitations of cryo-electron microscopy and X-ray crystallography studies. *Cell Calcium.* 59:209–227. <https://doi.org/10.1016/j.ceca.2016.04.003>
- Yuchi, Z., K. Lau, and F. Van Petegem. 2012. Disease mutations in the ryanodine receptor central region: Crystal structures of a phosphorylation hot spot domain. *Structure.* 20:1201–1211. <https://doi.org/10.1016/j.str.2012.04.015>
- Yuchi, Z., S.M. Yuen, K. Lau, A.Q. Underhill, R.L. Cornea, J.D. Fessenden, and F. Van Petegem. 2015. Crystal structures of ryanodine receptor SPRY1 and tandem-repeat domains reveal a critical FKBP12 binding determinant. *Nat. Commun.* 6:7947. <https://doi.org/10.1038/ncomms8947>
- Zalk, R., O.B. Clarke, A. des Georges, R.A. Grassucci, S. Reiken, F. Mancía, W.A. Hendrickson, J. Frank, and A.R. Marks. 2015. Structure of a mammalian ryanodine receptor. *Nature.* 517:44–49. <https://doi.org/10.1038/nature13950>
- Zhang, L., J. Kelley, G. Schmeisser, Y.M. Kobayashi, and L.R. Jones. 1997. Complex formation between junctin, triadin, calsequestrin, and the ryanodine receptor. Proteins of the cardiac junctional sarcoplasmic reticulum membrane. *J. Biol. Chem.* 272:23389–23397. <https://doi.org/10.1074/jbc.272.37.23389>
- Zhao, M., P. Li, X. Li, L. Zhang, R.J. Winkfein, and S.R. Chen. 1999. Molecular identification of the ryanodine receptor pore-forming segment. *J. Biol. Chem.* 274:25971–25974. <https://doi.org/10.1074/jbc.274.37.25971>
- Zorzato, F., J. Fujii, K. Otsu, M. Phillips, N.M. Green, F.A. Lai, G. Meissner, and D.H. MacLennan. 1990. Molecular cloning of cDNA encoding human and rabbit forms of the Ca^{2+} release channel (ryanodine receptor) of skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 265:2244–2256.
- Zorzato, F., R. Sacchetto, and A. Margreth. 1994. Identification of two ryanodine receptor transcripts in neonatal, slow-, and fast-twitch rabbit skeletal muscles. *Biochem. Biophys. Res. Commun.* 203:1725–1730. <https://doi.org/10.1006/bbrc.1994.2385>
- Zucchi, R., and S. Ronca-Testoni. 1997. The sarcoplasmic reticulum Ca^{2+} channel/ryanodine receptor: Modulation by endogenous effectors, drugs and disease states. *Pharmacol. Rev.* 49:1–51.