

**COMMENTARY**

# The taming of a scramblase

 Jarred M. Whitlock 

Eukaryotic cells are separated from their environment by a 30-Å-thick delimitation of lipid and protein—the plasma membrane (PM). Although a greater measure of affection has historically been garnered by their protein machines, cells have devoted ~5% of their genes to building lipids. In this pursuit of lipidic diversity, cells have developed repertoires of thousands of lipid species (van Meer et al., 2008), many of which fall into the category of phospholipids (PLs), which are largely responsible for the PM's bilayer arrangement. The PLs within the PM bilayer are actively arranged in an asymmetric orientation rendering the exofacial surface largely composed of cylindrical species (e.g., phosphatidylcholine), while the cytofacial leaflet retains most conical and charged species (e.g., phosphatidylethanolamine and phosphatidylserine [PS]). Despite this organization, many of our cells maintain the ability to rapidly abolish PM lipid asymmetry, exposing cytofacial PLs to the exofacial surface as a means of coordinating intercellular processes (Bevers and Williamson, 2016). Bevers et al. (1982) first described this PL “scrambling” in platelet activation, where PS exposure on the exofacial surface propagates coagulation. A few years after Bevers' description, several others noted that platelet activation was accompanied by a rapid alkalization of intracellular pH ( $pH_i$ ), and that blocking this alkalization inhibited PS exposure (Bucki et al., 2006; Zavoico et al., 1986). In an earlier issue of the *Journal of General Physiology*, Liang and Yang (2021) show us why. In fact, these platelets, and likely other cells that use nonapoptotic scrambling as a signaling module, have tamed their scramblases to fit a distinct biological niche via  $pH_i$ .

In an earlier issue of the *Journal of General Physiology*, Liang and Yang (2021) offer an insightful peek into the mechanism by which the TMEM16F scramblase is regulated by  $pH_i$ . Using simultaneous monitoring of fluorescent PS probe binding and electrophysiological recordings, Liang and Yang (2021) assess how alterations in  $pH_i$  impact both TMEM16F-dependent lipid rearrangements and the ionic current associated with PM scrambling. Under physiological  $Ca^{2+}$  concentrations, Liang and Yang (2021) find that TMEM16F is bidirectionally “tamed” by  $pH_i$ , both entirely inhibited by cytosolic acidification and potentiated by alkalization. However, the inhibition of TMEM16F via low  $pH_i$  can be overcome with super-physiological  $Ca^{2+}$

concentrations.  $pH_i$  appears to exert its effect on TMEM16F by altering the protonation state of the scramblase's  $Ca^{2+}$ -binding sites. To this point, reducing TMEM16F  $Ca^{2+}$  affinity also proportionately reduces the sensitivity of TMEM16F to  $pH_i$ . Either “reined” in by an acidic cytosol or spurred into a “gallop” through alkaline potentiation, these data demonstrate that  $pH_i$  bidirectionally tames the TMEM16 scramblase.

The mechanism described by Liang and Yang (2021) finally illuminates a likely mechanism by which the  $Na^+/H^+$  antiporter NHE1 regulates the activation of platelets. Inhibition of NHE1 function acidifies platelet  $pH_i$  and inhibits the PS exposure and microvesicle release observed in response to  $Ca^{2+}$ , as alluded to above; however, the mechanism by which NHE1 regulates these platelet functions has remained a mystery for decades (Bucki et al., 2006). A vital tool used to study PS exposure and microvesicle release in platelets has historically been the use of cells from Scott's syndrome patients, which lack both of these activities in platelets and red cells. Previous work has demonstrated that the loss of these platelet/red cell activities stems from a loss of their scramblase, TMEM16F (Suzuki et al., 2010). Based on the work of Liang and Yang (2021), it would appear that physiological alkalization of the platelet cytosol may very well be a mechanism platelets employ to potentiate their scramblase and achieve rapid PS exposure and microvesicle release in response to injury. Moreover, the loss of platelet activation in response to NHE1 inhibition likely stems from the acidic environment of the cytosol reigning in the platelet's scramblase, TMEM16F.

For the last 10 yr, we have appreciated that much of the nonapoptotic scrambling that cells employ in intercellular signaling is performed by the TMEM16 family of  $Ca^{2+}$ -activated phospholipid scramblases (Ca $^{2+}$ -PLSases). The most abundant PLSase in mammals is TMEM16F (reviewed in Whitlock and Hartzell, 2017). Of note, this family also includes two dearly loved ion channel members that do not scramble PLs but pass ions through the PM (i.e., TMEM16s A and B). Thanks to the work of many, we now know a great deal about which TMEM16s are scramblases, how they ferry lipids between PM leaflets, how they are activated by  $Ca^{2+}$ , and their structures. Moreover, we also appreciate that these scramblases are nonselective in

Section on Membrane Biology, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD.

Correspondence to Jarred M. Whitlock: [Jarred.Whitlock@nih.gov](mailto:Jarred.Whitlock@nih.gov).


© 2021 Whitlock. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms/>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 4.0 International license, as described at <https://creativecommons.org/licenses/by-nc-sa/4.0/>).

respect to which lipids they move between leaflets, and that some TMEM16s even scramble lipids in membranes other than the PM (Tsuji et al., 2019). What we are just beginning to appreciate are the mechanisms that cells use to manage these potent remodelers of the PM beyond  $\text{Ca}^{2+}$  activation.

In an earlier issue of the *Journal of General Physiology*, Liang and Yang (2021) identify a novel mechanism by which cells can tame TMEM16F activities to fit their biological purpose using  $\text{pH}_i$ . This  $\text{pH}_i$  regulatory mechanism is one of several newly discovered ways that cells can tame TMEM16 scramblases to fit a particular biological system. Moving forward, it will be vital to better elucidate how  $\text{pH}_i$  and other means of taming TMEM16 scramblases fit into the ever-growing list of biological processes that are activated by scrambling.

#### A TMEM16 scramblase module is coming into focus


TMEM16s are unequivocally activated by intracellular  $\text{Ca}^{2+}$ , but how these  $\text{Ca}^{2+}$ -activated machines are adapted to fit distinct biological contexts moves beyond cytosolic  $\text{Ca}^{2+}$ . Because their ion channel cousins were discovered first, our ideas of how TMEM16 scramblases are regulated has largely been colored by the electrophysiological characterization of the  $\text{Cl}^-$  channel TMEM16A. TMEM16A is activated by  $\text{Ca}^{2+}$ , but this activity is regulated by membrane voltage,  $\text{PIP}_2$ , cholesterol, extracellular pH, and  $\text{pH}_i$  (Yu et al., 2019; Le et al., 2019; De Jesús-Pérez et al., 2018; Xiao et al., 2011; Cruz-Rangel et al., 2017; Chun et al., 2015). Although biological functions differ dramatically between TMEM16 ion channels and scramblases, it appears that many routes of regulation are conserved family traits. In particular, TMEM16F is also regulated by membrane voltage,  $\text{PIP}_2$ , and, as we learn from Liang and Yang,  $\text{pH}_i$  (Scudieri et al., 2015; Ye et al., 2018; Liang and Yang, 2021). Moreover, the mechanism by which Liang and Yang (2021) propose  $\text{pH}_i$  regulates TMEM16F (through the protonation of its  $\text{Ca}^{2+}$  binding sites) is shared with TMEM16A (Chun et al., 2015). Together, this shared  $\text{pH}_i$  regulatory mechanism, observed in TMEM16s of differing activities, suggests that most TMEM16s may be regulated by  $\text{pH}_i$ . However, is it reasonable to assume that TMEM16 sensitivity to  $\text{pH}_i$  is similar between family members? All TMEM16s are activated by  $\text{Ca}^{2+}$  binding, and the residues coordinating  $\text{Ca}^{2+}$  in TMEM16s are highly conserved across the family. However, despite this conservation, TMEM16s differ drastically in their sensitivity to  $\text{Ca}^{2+}$  (e.g.,  $\text{Ca}^{2+}$  sensitivity differs ~1,000-fold between TMEM16s A and F). If the mechanism by which  $\text{pH}_i$  regulates TMEM16s involves these same  $\text{Ca}^{2+}$ -coordinating residues, how can we trust that  $\text{pH}_i$  sensitivity would be the same between relatives when  $\text{Ca}^{2+}$  sensitivity is not? Moreover, Liang and Yang (2021) demonstrate that increased  $\text{Ca}^{2+}$  can overcome  $\text{pH}_i$  inhibition. If some TMEM16s are ~1,000-fold more sensitive to  $\text{Ca}^{2+}$ , is it reasonable to expect that all would escape  $\text{pH}_i$  regulation at the same  $\text{Ca}^{2+}$  concentration? Determining the  $\text{pH}_i$  sensitivity of different TMEM16s may help further resolve how the same cell can regulate the activities of several TMEM16s at once. Perhaps evolution's differential tuning of these shared TMEM16 regulatory mechanisms has ensured that some TMEM16s never "see" the activating signals of their relatives, even though they are within the same membrane of the same cell.



**Figure 1. The TMEM16 scramblase module.** A vastly oversimplified cartoon illustrating how disparate regulators of TMEM16 scramblases come together as a "module" employed in many cells to individually tune the "brightness" of the scrambling. This tuning would undoubtedly alter the speed at which a cell promotes lipid redistribution in response to  $\text{Ca}^{2+}$ , as well as the extent that scrambling alters the exposure of cytofacial lipid species and the physical characteristics of the PM during scrambling.

Although TMEM16F and other TMEM16 scramblases are expressed in a diverse array of cell types and tissues, we have now begun to appreciate several unique mechanisms that regulate the location, magnitude, and duration of their activities. Together, these disparate regulatory factors come together to create a "TMEM16 scramblase module" employed by many cells and used to uniquely tame, or tune, TMEM16-dependent scrambling to fit a cell's individual biological niche (Fig. 1).

It should be noted that there is an independent PL scrambling pathway associated with apoptotic caspase activity and the Xk-related protein family (Suzuki et al., 2014). Interestingly, this apoptotic scrambling is also modulated by  $\text{pH}_i$  but in the opposite direction described above, with lower  $\text{pH}_i$  promoting exofacial PS exposure. Importantly, this response is observed in cells from healthy donors and those from Scott's syndrome patients lacking TMEM16F, demonstrating that this is indeed an alternative scrambling response to that described by Liang and Yang (Stout et al., 1997). Caspase- and  $\text{Ca}^{2+}$ -activated scrambling pathways are parallel yet independent processes and differ substantially in a number of ways, including the amount of time between activation of the pathway and PS exposure (hours versus minutes), the distribution of the PS each expose in native cells (whole membrane versus patches), and likely the reversibility of PS exposure (caspases cleave PS transporters while  $\text{Ca}^{2+}$  reversibly inhibits them), and here Liang and Yang (2021) demonstrate that  $\text{Ca}^{2+}$ -activated PM scrambling differs in its response to  $\text{pH}_i$  compared with caspase-activated PM scrambling. Moving forward, it will be important to consider the differences between these scramblase pathways that both result in the exposure of cytofacial lipids and altered PM physical properties.



**Figure 2. The TMEM16F scramblase function in cellular processes.** An illustration highlighting how the regulated activity of TMEM16 scramblases permeates biological processes. TMEM16 scramblases are activated by increases in intracellular calcium ( $\text{Ca}^{2+}_i$ ); however, scramblase activity is regulated by a host of other cellular factors including membrane voltage (mV), phosphatidyl-inositol 4,5-bisphosphate ( $\text{PIP}_2$ ), intracellular pH ( $\text{pH}_i$ ), and likely others (?; see roots). Together, these regulatory factors modulate the exofacial exposure of cytofacial lipids (trunk) that coordinate a variety cellular processes in diverse systems including cell-cell fusion, platelet facilitated blood clotting, membrane patch repair, and enveloped viral entry (see branches). To appreciate how these scramblase regulators fit into distinct, biologically relevant processes, we must begin watching the branches as we tickle the roots.

Downloaded from [http://jupress.org/jgp/article-pdf/115/4/e202012831/1807978/jgp\\_202012831.pdf](http://jupress.org/jgp/article-pdf/115/4/e202012831/1807978/jgp_202012831.pdf) by guest on 09 February 2026

And now for the most important question: how do cells, tissues, and systems employ TMEM16 scramblase regulation for biological purposes (Fig. 2)?

#### How does the taming of a scramblase affect biological processes?

As the list of methods available to cells for taming TMEM16 scramblase function grow, we are left with a major question: how are these regulatory mechanisms employed in the local domestication of scramblases in biologically relevant contexts? In light of this latest contribution by Liang and Yang (2021), a few biological processes come to mind.

In addition to platelet activation, nonapoptotic PS exposure is a reoccurring signal employed in the coordination of cell-cell fusion processes (Whitlock and Chernomordik, 2021). The role of TMEM16 scramblases has been linked to the fusion of gametes, skeletal muscle, osteoclasts, placental trophoblasts, and virus-cell fusion. These could be ideal systems to dissect the mechanistic contribution of  $\text{pH}_i$  in the exposure of PS required

for these fusion events. In particular, during capacitation and/or the acrosome reaction, PS is exposed on the heads of sperm and is required for fertilization (Rival et al., 2019). Interestingly, during capacitation and/or the acrosome reaction, sperm both alkalinize their cytosol and increase its  $\text{Ca}^{2+}$  concentration simultaneously (Chávez et al., 2018). TMEM16E, a close scramblase relative of TMEM16F, is highly expressed in sperm, and its loss perturbs murine fertility. Characterization of  $\text{pH}_i$  in fertilization and its impact on TMEM16E function may offer insight into how mammalian gametes prepare for fusion and better inform our understanding of the regulatory processes surrounding human fertility.

Platelets, red cells, and sperm each exhibit high steady-state levels of a particular TMEM16 scramblase, but many cell types exhibit abundant levels of several TMEM16 scramblases simultaneously. For example, we previously demonstrated that skeletal muscle exhibits high levels of both TMEM16s E and F (Whitlock et al., 2018). What advantage would one cell gain by maintaining multiple scramblases that are each activated by a

single, shared signal? Upon increases in cytosolic  $\text{Ca}^{2+}$ , each of these TMEM16 scramblases should be activated simultaneously. However, we found that loss of TMEM16E perturbed  $\text{Ca}^{2+}$ -activated PS exposure in primary muscle cells despite high levels of TMEM16F, suggesting that the two play distinct roles in the same cell type. At the time, we assumed TMEM16s E and F were functionally separated in muscle by differences in  $\text{Ca}^{2+}$  sensitivity and/or trafficking, but could the function of these scramblase cousins be separated by  $\text{pH}_i$ ?  $\text{pH}_i$  is rather distinct in skeletal muscle as compared with other, nonexcitable cell types with a  $\text{pH}_i$  of ~6 at resting membrane potential (Carter et al., 1967). Depending on the  $\text{Ca}^{2+}$  concentration, the observations of Liang and Yang (2021) would suggest that TMEM16F could be entirely inhibited in resting skeletal muscle fibers. This idea is of course speculative but serves to illustrate how nodes within the TMEM16 scramblase module (Fig. 1) may serve to functionally separate TMEM16s in the same cell. One can easily assert that because the  $\text{Ca}^{2+}$  binding sites of TMEM16s F and E are so highly conserved, their  $\text{pH}_i$  sensitivity should be similar. After all, these are the same residues where  $\text{pH}_i$  likely exerts its regulatory effect in both. However, we found that TMEM16E appears to be much more sensitive to  $\text{Ca}^{2+}$  than TMEM16F (~50% of cells expressing TMEM16E scramble in response to 1  $\mu\text{M}$   $\text{Ca}^{2+}$ , while >100  $\mu\text{M}$   $\text{Ca}^{2+}$  was required to achieve a similar response in TMEM16F expressing cells; Whitlock et al., 2018). If these relatives, with highly conserved  $\text{Ca}^{2+}$  binding sites, can differ dramatically in their sensitivity to  $\text{Ca}^{2+}$ , how can we assume they will not differ in their sensitivity to  $\text{pH}_i$ ? By moving past  $\text{Ca}^{2+}$ , TMEM16 machines may have afforded themselves greater utility and promoted the diversity of their family in many cells.

Apart from healthy, physiological processes, pathogens may also use  $\text{pH}_i$  to tame TMEM16 scramblase function. PD-1 is an immunoregulatory protein that, when exposed on the surface of a variety of different cell types, dampens T cell activation and prevents self-reactive, pathogenic T cells. Perturbed, excessive exposure of PD-1 at the cell surface is a major contributor to T cell exhaustion in chronic HIV-1 infection (Porichis and Kaufmann, 2012). Interestingly, mice lacking TMEM16F also experience T cell exhaustion, and TMEM16F activity appears to play an important role in regulating the trafficking of PD-1 to the PM (Hu et al., 2016; Bricogne et al., 2019). Our laboratory has demonstrated that HIV activates TMEM16F and utilizes exposed PS as a cofactor in the fusion of the viral envelope with the PM (Zaitseva et al., 2017). After HIV gains entry into the cell, the HIV protein Vpr specifically down-regulates the expression of NHE1 and acidifies the cytosol of infected cells (Janket et al., 2007). If the reader will permit some additional speculation, acidification may be a mechanism by which HIV inhibits TMEM16F in infected cells by downregulating NHE1 and acidifying the cytosol. Inhibiting TMEM16F would likely contribute to T cell exhaustion, just as observed in the TMEM16F knockout model, and facilitate the ability of HIV-infected cells to avoid detection by the immune system, likely through increased trafficking of PD-1 to the PM. Perhaps HIV is a master scramblase tamer, both activating TMEM16F to gain cell entry and silencing the scramblase once in the cell?

In this reader's view, Liang and Yang's observations in the earlier issue of the *Journal of General Physiology* are tightly

focused, but the consequences of these observations are broadly reaching (Liang and Yang, 2021). For what purpose do we tame something if not to leverage it for our goals? Over the last decade, we have learned much about the business of taming a scramblase. In the next, let us resolve what that taming is for (Fig. 2).

## Acknowledgments

Crina M. Nimigean served as editor.

The author thanks the work of dozens of laboratories whose contributions were not appropriately acknowledged here due to length restrictions. In spite of the many open questions facing our field, we have entered a golden age of the scramblase. Bethany G. Whitlock contributed mightily to the construction of the figures herein. Moreover, Dr. Criss Hartzell offered much appreciated critical feedback that greatly improved the final manuscript.

The research of J.M. Whitlock is supported by the Intramural Research Program of the Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health.

The author declares no competing financial interests.

## References

Bevers, E.M., P. Comfurius, J.L. van Rijn, H.C. Hemker, and R.F. Zwaal. 1982. Generation of prothrombin-converting activity and the exposure of phosphatidylserine at the outer surface of platelets. *Eur. J. Biochem.* 122: 429–436. <https://doi.org/10.1111/j.1432-1033.1982.tb05898.x>

Bevers, E.M., and P.L. Williamson. 2016. Getting to the Outer Leaflet: Physiology of Phosphatidylserine Exposure at the Plasma Membrane. *Physiol. Rev.* 96:605–645. <https://doi.org/10.1152/physrev.00020.2015>

Bricogne, C., M. Fine, P.M. Pereira, J. Sung, M. Tijani, Y. Wang, R. Henriques, M.K. Collins, and D.W. Hilgemann. 2019. TMEM16F activation by  $\text{Ca}^{2+}$  triggers plasma membrane expansion and directs PD-1 trafficking. *Sci. Rep.* 9:619. <https://doi.org/10.1038/s41598-018-37056-x>

Bucki, R., J.J. Pastore, F. Giraud, P.A. Jamme, and J.C. Sulpice. 2006. Involvement of the  $\text{Na}^+/\text{H}^+$  exchanger in membrane phosphatidylserine exposure during human platelet activation. *Biochim. Biophys. Acta.* 1761: 195–204. <https://doi.org/10.1016/j.bbapap.2005.12.008>

Carter, N.W., F.C. Rector Jr., D.S. Campion, and D.W. Seldin. 1967. Measurement of intracellular pH of skeletal muscle with pH-sensitive glass microelectrodes. *J. Clin. Invest.* 46:920–933. <https://doi.org/10.1172/JCII05598>

Chávez, J.C., J.L. De la Vega-Beltrán, O. José, P. Torres, T. Nishigaki, C.L. Treviño, and A. Darszon. 2018. Acrosomal alkalization triggers  $\text{Ca}^{2+}$  release and acrosome reaction in mammalian spermatozoa. *J. Cell. Physiol.* 233:4735–4747. <https://doi.org/10.1002/jcp.26262>

Chun, H., H. Cho, J. Choi, J. Lee, S.M. Kim, H. Kim, and U. Oh. 2015. Protons inhibit anoctamin 1 by competing with calcium. *Cell Calcium.* 58: 431–441. <https://doi.org/10.1016/j.ceca.2015.06.011>

Cruz-Rangel, S., J.J. De Jesús-Pérez, I.A. Aréchiga-Figueroa, A.A. Rodríguez-Menchaca, P. Pérez-Cornejo, H.C. Hartzell, and J. Arreola. 2017. Extracellular protons enable activation of the calcium-dependent chloride channel TMEM16A. *J. Physiol.* 595:1515–1531. <https://doi.org/10.1113/JP273111>

De Jesús-Pérez, J.J., S. Cruz-Rangel, A.E. Espino-Saldaña, A. Martínez-Torres, Z. Qu, H.C. Hartzell, N.E. Corral-Fernandez, P. Pérez-Cornejo, and J. Arreola. 2018. Phosphatidylinositol 4,5-bisphosphate, cholesterol, and fatty acids modulate the calcium-activated chloride channel TMEM16A (ANO1). *Biochim. Biophys. Acta Mol. Cell Biol. Lipids.* 1863:299–312. <https://doi.org/10.1016/j.bbapap.2017.12.009>

Hu, Y., J.H. Kim, K. He, Q. Wan, J. Kim, M. Flach, T. Kirchhausen, A. Vortkamp, and F. Winau. 2016. Scramblase TMEM16F terminates T cell receptor signaling to restrict T cell exhaustion. *J. Exp. Med.* 213: 2759–2772. <https://doi.org/10.1084/jem.20160612>

Janket, M.L., J.S. DeRicco, L. Borowski, and V. Ayyavoo. 2007. Human immunodeficiency virus (HIV-1) Vpr induced downregulation of NHE1 induces alteration in intracellular pH and loss of ERM complex in target cells. *Virus Res.* 126:76–85. <https://doi.org/10.1016/j.virusres.2007.01.019>

Le, S.C., Z. Jia, J. Chen, and H. Yang. 2019. Molecular basis of PIP<sub>2</sub>-dependent regulation of the Ca<sup>2+</sup>-activated chloride channel TMEM16A. *Nat. Commun.* 10:3769. <https://doi.org/10.1038/s41467-019-11784-8>

Liang, P., and H. Yang. 2021. Molecular underpinning of intracellular pH regulation on TMEM16F. *J. Gen. Physiol.* 153:e202012704. <https://doi.org/10.1085/jgp.202012704>

Porichis, F., and D.E. Kaufmann. 2012. Role of PD-1 in HIV pathogenesis and as target for therapy. *Curr. HIV/AIDS Rep.* 9:81–90. <https://doi.org/10.1007/s11904-011-0106-4>

Rival, C.M., W. Xu, L.S. Shankman, S. Morioka, S. Arandjelovic, C.S. Lee, K.M. Wheeler, R.P. Smith, L.B. Haney, B.E. Isakson, et al. 2019. Phosphatidylserine on viable sperm and phagocytic machinery in oocytes regulate mammalian fertilization. *Nat. Commun.* 10:4456. <https://doi.org/10.1038/s41467-019-12406-z>

Scudieri, P., E. Caci, A. Venturini, E. Sondo, G. Pianigiani, C. Marchetti, R. Ravazzolo, F. Pagani, and L.J. Galieta. 2015. Ion channel and lipid scramblase activity associated with expression of TMEM16F/ANO6 isoforms. *J. Physiol.* 593:3829–3848. <https://doi.org/10.1113/P270691>

Stout, J.G., F. Bassé, R.A. Luhm, H.J. Weiss, T. Wiedmer, and P.J. Sims. 1997. Scott syndrome erythrocytes contain a membrane protein capable of mediating Ca<sup>2+</sup>-dependent transbilayer migration of membrane phospholipids. *J. Clin. Invest.* 99:2232–2238. <https://doi.org/10.1172/JCI19397>

Suzuki, J., E. Imanishi, and S. Nagata. 2014. Exposure of phosphatidylserine by Xk-related protein family members during apoptosis. *J. Biol. Chem.* 289:30257–30267. <https://doi.org/10.1074/jbc.M114.583419>

Suzuki, J., M. Umeda, P.J. Sims, and S. Nagata. 2010. Calcium-dependent phospholipid scrambling by TMEM16F. *Nature.* 468:834–838. <https://doi.org/10.1038/nature09583>

Tsuji, T., J. Cheng, T. Tatematsu, A. Ebata, H. Kamikawa, A. Fujita, S. Gyobu, K. Segawa, H. Arai, T. Taguchi, et al. 2019. Predominant localization of phosphatidylserine at the cytoplasmic leaflet of the ER, and its TMEM16K-dependent redistribution. *Proc. Natl. Acad. Sci. USA.* 116:13368–13373. <https://doi.org/10.1073/pnas.1822025116>

van Meer, G., D.R. Voelker, and G.W. Feigenson. 2008. Membrane lipids: where they are and how they behave. *Nat. Rev. Mol. Cell Biol.* 9:112–124. <https://doi.org/10.1038/nrm2330>

Whitlock, J.M., and L.V. Chernomordik. 2021. Flagging fusion: Phosphatidylserine signaling in cell-cell fusion. *J. Biol. Chem.* In press. <https://doi.org/10.1016/j.jbc.2021.100411>

Whitlock, J.M., and H.C. Hartzell. 2017. Anoctamins/TMEM16 proteins: Chloride channels flirting with lipids and extracellular vesicles. *Annu. Rev. Physiol.* 79:119–143. <https://doi.org/10.1146/annurev-physiol-022516-034031>

Whitlock, J.M., K. Yu, Y.Y. Cui, and H.C. Hartzell. 2018. Anoctamin 5/TMEM16E facilitates muscle precursor cell fusion. *J. Gen. Physiol.* 150:1498–1509. <https://doi.org/10.1085/jgp.201812097>

Xiao, Q., K. Yu, P. Perez-Cornejo, Y. Cui, J. Arreola, and H.C. Hartzell. 2011. Voltage- and calcium-dependent gating of TMEM16A/ANo1 chloride channels are physically coupled by the first intracellular loop. *Proc. Natl. Acad. Sci. USA.* 108:8891–8896. <https://doi.org/10.1073/pnas.1102147108>

Ye, W., T.W. Han, L.M. Nassar, M. Zubia, Y.N. Jan, and L.Y. Jan. 2018. Phosphatidylinositol-(4, 5)-bisphosphate regulates calcium gating of small-conductance cation channel TMEM16F. *Proc. Natl. Acad. Sci. USA.* 115:E1667–E1674. <https://doi.org/10.1073/pnas.1718728115>

Yu, K., T. Jiang, Y. Cui, E. Tajkhorshid, and H.C. Hartzell. 2019. A network of phosphatidylinositol 4,5-bisphosphate binding sites regulates gating of the Ca<sup>2+</sup>-activated Cl<sup>-</sup> channel ANO1 (TMEM16A). *Proc. Natl. Acad. Sci. USA.* 116:19952–19962. <https://doi.org/10.1073/pnas.1904012116>

Zaitseva, E., E. Zaitsev, K. Melikov, A. Arakelyan, M. Marin, R. Villasmil, L.B. Margolis, G.B. Melikyan, and L.V. Chernomordik. 2017. Fusion Stage of HIV-1 Entry Depends on Virus-Induced Cell Surface Exposure of Phosphatidylserine. *Cell Host Microbe.* 22:99–110.e7. <https://doi.org/10.1016/j.chom.2017.06.012>

Zavoico, G.B., E.J. Cragoe Jr., and M.B. Feinstein. 1986. Regulation of intracellular pH in human platelets. Effects of thrombin, A23187, and ionomycin and evidence for activation of Na<sup>+</sup>/H<sup>+</sup> exchange and its inhibition by amiloride analogs. *J. Biol. Chem.* 261:13160–13167. [https://doi.org/10.1016/S0021-9258\(18\)69284-1](https://doi.org/10.1016/S0021-9258(18)69284-1)