

REVIEW

How does mitochondrial Ca²⁺ change during ischemia and reperfusion? Implications for activation of the permeability transition pore

Elizabeth Murphy¹ and David A. Eisner²

Cardiac ischemia followed by reperfusion results in cardiac cell death, which has been attributed to an increase of mitochondrial Ca²⁺ concentration, resulting in activation of the mitochondrial permeability transition pore (PTP). Evaluating this hypothesis requires understanding of the mechanisms responsible for control of mitochondrial Ca²⁺ in physiological conditions and how they are altered during both ischemia and reperfusion. Ca²⁺ influx is thought to occur through the mitochondrial Ca²⁺ uniporter (MCU). However, with deletion of the MCU, an increase in mitochondrial Ca²⁺ still occurs, suggesting an alternative Ca²⁺ influx mechanism during ischemia. There is less certainty about the mechanisms responsible for Ca²⁺ efflux, with contributions from both Ca²⁺/H+ exchange and a Na⁺-dependent Ca²⁺ efflux pathway. The molecular details of both mechanisms are not fully resolved. We discuss this and the contributions of both pathways to the accumulation of mitochondrial Ca²⁺ during ischemia and reperfusion. We further discuss the role of mitochondrial Ca²⁺ in activation of the PTP.

Introduction

There is great interest in the control of the calcium (Ca^{2+}) concentration in the mitochondrial matrix. An increase of mitochondrial Ca²⁺, as often occurs with cellular activity, has been shown to activate enzymes of the tricarboxylic acid cycle and thereby ensure that ATP production is matched to the increased metabolic demand. On the other hand, an excessive increase of mitochondrial Ca2+ has been linked to cell death via activation of the mitochondrial permeability transition pore (PTP) (Murphy and Steenbergen, 2021). The aim of this review is to critically evaluate the hypothesis that an increase in mitochondrial Ca2+ during ischemia and reperfusion is responsible for initiating cell death via activation of the PTP and that inhibition of this increase in mitochondrial Ca2+ might be a cardioprotective strategy. Mitochondrial Ca2+ concentration is determined by a balance between influx and efflux, so the next section will consider the possible transport mechanisms involved.

Regulation of mitochondrial Ca²⁺ Mitochondrial Ca²⁺ uniporter complex

 Ca^{2+} enters the mitochondria via the mitochondrial Ca^{2+} uniporter (MCU) (Baughman et al., 2011; De Stefani et al., 2011). The driving force for Ca^{2+} entry by MCU is provided by the

mitochondrial membrane potential ($\Delta \psi$), with the mitochondrial matrix negative by ~180 mV, which is generated by electron transport. The MCU exists in a complex (MCUC) that contains MCU, essential MCU regulator (EMRE) (Sancak et al., 2013), and three mitochondrial Ca2+ uptake proteins (MICU1, -2, and -3); EMRE is required for MCU activity in metazoans. MICU1 binds directly to MCU and EMRE (Paillard et al., 2018). In addition to binding to MCU and EMRE, MICU1 can either form homodimers with MICU1 or heterodimers with MICU2 or MICU3. The apparent threshold for Ca²⁺ uptake into the mitochondria is set by which MICUs compose the dimer (Csordas et al., 2013; Liu et al., 2016; Kamer et al., 2017). With homodimers of MICU1, the threshold for Ca²⁺ uptake is ~300 nM, whereas the threshold is \sim 600 nM for a complex with MICU1/2 heterodimers (Kamer et al., 2017). The ratio of MICU1 to MCU varies among tissues, and the dimer composition is also tissue dependent (Paillard et al., 2017). The exact mechanisms that regulate the composition of the MCUC are not well established and require further study (Csordas et al., 2013; Plovanich et al., 2013; Patron et al., 2014; Kamer et al., 2017). MCUC and its regulation are discussed in detail elsewhere (Boyman et al., 2021; Murphy and Steenbergen, 2021; Garbincius and Elrod, 2022; Rodríguez-Prados et al., 2023; Vecellio Reane et al., 2024) and are not the focus of this review.

¹Cardiac Physiology Section, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA; ²Unit of Cardiac Physiology, Division of Cardiovascular Sciences, University of Manchester, Manchester, UK.

Correspondence to Elizabeth Murphy: murphy1@nhlbi.nih.gov.

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply. This article is distributed under the terms as described at https://rupress.org/pages/terms102024/.





Efflux mechanisms

At steady state, any Ca²⁺ that enters the mitochondria on MCU must be extruded. Two mitochondrial Ca²⁺ efflux pathways have been described: a Na⁺-independent mechanism generally thought to be a Ca²⁺/H⁺ exchange (CHE) and a Na⁺ dependent Ca²⁺-efflux pathway (Crompton et al., 1977; Nicholls and Crompton, 1980; Brand, 1985). The relative role of each pathway is thought to be tissue specific and the Na⁺-dependent Ca²⁺ efflux mechanism is generally thought to be the primary mitochondrial Ca²⁺ efflux mechanism in cardiac cells (Wei et al., 2011; Garbincius and Elrod, 2022).

CHE

The identity of the Ca2+/H+ exchanger has been debated, and roles for both leucine zipper-EF-hand-containing transmembrane protein 1 (LETM1) and transmembrane BAX inhibitor motif protein 5 (TMBIM5) have been proposed. LETM1 was originally proposed to be a K+/H+ exchanger, although later studies suggested it exchanged Ca²⁺ for H⁺ (Jiang et al., 2009). It was initially reported to be electrogenic, exchanging 1H+ for 1Ca²⁺ (Jiang et al., 2009). This and many studies of pHdependent Ca2+ exchange rely on studies which involve altering pH. One needs to be cautious in interpreting these studies as change in pH affect many transporters as well as the charge state of inorganic phosphate. Previous studies have shown that Ca2+ uptake into the mitochondria is very small in the absence of permeant anions (Wei et al., 2012). Recent studies with purified LETM1 reconstituted into liposomes suggest that LETM1 exchanges 2H⁺ for 1Ca²⁺ and is therefore electroneutral (Tsai et al., 2014).

Recently, three publications have suggested a different molecular identity for mitochondrial CHE (TMBIM5); although they differ in detail regarding the direction of exchange (Austin et al., 2022; Patron et al., 2022; Zhang et al., 2022). Zhang et al. (2022) examined the effect of TMBIM5 overexpression in HEK293 cells. ATP was added to release ER Ca²⁺, and mitochondrial Ca²⁺ was measured using a mitochondrial-targeted genetically encoded Ca²⁺ indicator. Overexpression of TMBIM5 enhanced mitochondria Ca2+ uptake and deletion decreased mitochondrial H+. They also generated a mouse with a mutation in a negatively charged aspartate in the pore domain (D326R), which had been shown previously to block activity. This mutation led to the degradation of the protein. Homozygous mice were born at less than Mendelian ratios and developed a skeletal muscle myopathy. In another study, Patron et al. (2022) reported that TMBIM5 transports protons into the matrix in exchange for Ca2+ efflux, and that with loss of TMBIM5, there is an increase in matrix pH from 7.25 in WT to 7.6, consistent with reduced exchange of matrix Ca²⁺ for cytosolic H⁺. As pointed out below, this direction of net flux would be consistent with an electroneutral exchanger but not with a 1Ca²⁺:1H⁺ exchanger. TMBIM5 was also shown to interact with the m-AAA protease AFG3L2 and inhibit its activity. Austin et al. (2022) also reported data showing that TMBIM5 is a mitochondrial Ca2+/H+ exchanger. In studies in which the Na+-Ca2+-Li+ exchanger (NCLX) was either depleted or inhibited, they showed that compared with WT cells, cells with loss of TMBIM5 had reduced mitochondrial Ca2+ efflux

measured in permeabilized HEK293 cells following the addition of Ru360 to inhibit MCU-mediated Ca²⁺ uptake and thereby reveal mitochondrial Ca²⁺ efflux. They further showed that knocking down LETM1 did not reduce mitochondrial Ca²⁺ efflux and concluded that LETM1 is not responsible for CHE.

Stoichiometry is crucial for the direction of ion flux, and we will illustrate this in detail for CHE. The electron transport chain extrudes protons, thereby generating a large negative $\Delta\psi$ and a pH gradient (Δ pH) with an alkaline matrix pH. A stoichiometry of 1H+ for 1Ca²⁺ means that Ca²⁺ efflux is associated with net movement of positive charge out of the mitochondrion, which will be opposed by the matrix-negative $\Delta\psi$. In contrast, the alkaline matrix pH would increase the driving force for Ca²⁺ transport out of the matrix in exchange for H+ entry. The net direction of flux would therefore depend on the relative magnitudes of the pH gradient and membrane potential. In general, for a transporter that exchanges n protons for one Ca²⁺ ion, net Ca²⁺ efflux will occur when the Ca²⁺ concentration gradient is given by Eq. 1:

$$\frac{\left[\mathsf{Ca}^{2^{+}}\right]_{\mathrm{mito}}}{\left[\mathsf{Ca}^{2^{+}}\right]_{\mathrm{cyto}}} > \left\{ \frac{\left[\mathsf{H}^{+}\right]_{\mathrm{mito}}}{\left[\mathsf{H}^{+}\right]_{\mathrm{cyto}}} \right\}^{n} \cdot e^{\left\{\left[n-2\right] \cdot \frac{\mathsf{VT}}{\mathsf{RT}}\right\}} \tag{1}$$

Therefore, the more acidic the matrix pH, the higher the matrix Ca^{2+} must be to obtain net Ca^{2+} efflux. If matrix pH is 0.5 pH units more alkaline than the cytoplasm, $[H^+]_{mito}/[H^+]_{cyto} = 0.31$. The effects of membrane potential depend on the stoichiometry. If n=1, then the more the negative is $\Delta\psi$, the higher the Ca^{2+} required for net Ca^{2+} efflux. With this stoichiometry, Eq. 1 reduces to:

$$\frac{[Ca^{2+}]_{mito}}{[Ca^{2+}]_{cyto}} > 0.31 \cdot e^{\left\{\frac{-VT}{RT}\right\}}$$
 (2)

Under physiological conditions, with $\Delta\psi$ of about -180 mV and RT/F = 25.2 mV at body temperature, where R, universal gas constant; T, temperature; F, Faraday constant, Eq. 2 predicts that with n = 1, net efflux will only occur if $[Ca^{2+}]_{mito}$ is 392 times greater than $[Ca^{2+}]_{cyto}$. However, under physiological conditions, mitochondrial Ca2+ is 100-200 nM and cytoplasmic Ca^{2+} ranges from about 0.1 μM in diastole to 1 μM in systole, meaning that mitochondrial Ca^{2+} varies from 2 to 0.1 × cytoplasmic, much less than that required for net Ca²⁺ efflux. Peak Ca²⁺ in the space between the sarcoplasmic reticulum and mitochondria is likely much higher (Lu et al., 2013). Therefore with exchange of 1H+ for 1 Ca2+, the expected flux would be net Ca2+ entry and H+ efflux. With loss or reduction in $\Delta \psi$, especially with cytosolic acidification, it has been suggested that LETM1 can function to extrude matrix Ca2+ in exchange for cytosolic H+ (Jiang et al., 2009; Waldeck-Weiermair et al., 2011). It is worth noting that with a 0.5 pH unit gradient and mitochondrial Ca²⁺ twice cytoplasmic, Δψ would have to be reduced to about -47 mV for net Ca efflux to occur. If mitochondrial and cytoplasmic Ca²⁺ are equal, then net efflux will only occur if $\Delta \psi$ is decreased to -30 mV. Finally, if the pH gradient is entirely dissipated, then with mitochondrial and cytoplasmic Ca2+ equal, net efflux can only occur with the unlikely case of a positive $\Delta\psi$.



In contrast, if the exchanger is electroneutral ($2H^+$ for $1Ca^{2+}$), then Eq. 1 reduces to Eq. 3.

$$\frac{[Ca^{2+}]_{\text{mito}}}{[Ca^{2+}]_{\text{cyto}}} > \left\{ \frac{[H^{+}]_{\text{mito}}}{[H^{+}]_{\text{cyto}}} \right\}^{2}$$
(3)

Net Ca^{2+} efflux will occur if $[Ca^{2+}]_{mito}/[Ca^{2+}]_{cyto} > 0.31^2$ (\sim 0.1). In other words, an electroneutral exchanger can pump Ca^{2+} against a 10-fold gradient. It is clear that the stoichiometry will have important implications for Ca^{2+} regulation and it is therefore important to resolve the stoichiometry of the exchanger.

Interestingly, the stoichiometry of TMBIM5-mediated CHE has not been reported. Another question is whether both TMBIM5 and LETM1 exchange Ca²⁺ for H⁺ and if they do so with different stoichiometries? Austin et al. (2022) reported that knocking down LETM1 did not block Na+-independent mitochondrial Ca²⁺ efflux, but further work will be needed to resolve the relative function of TMBIM5 and LETM1. Assuming that LETM1 is electrogenic, in the presence of a large negative $\Delta \psi$, it is likely to function as a Ca2+ influx mechanism and thus Ca2+ efflux might not be observed under these conditions. It is also curious that cardiac cells have high expression of TMBIM5 but very little Na⁺-independent Ca²⁺ efflux. This would be consistent with TMBIM5 acting as a Ca2+ influx pathway. Future studies establishing the stoichiometry will be key to addressing this issue. Alternatively, TMBIM5 might require additional cofactors for activation which are not present in heart under basal conditions, although it is curious that the cardiac cell has such high levels of TMBIM5 at baseline.

Na⁺-dependent Ca²⁺ efflux

Palty et al. (2010) reported that a protein encoded by slc8b1 (subsequently referred to as NCLX) was responsible for Na⁺dependent Ca²⁺ efflux. A Ca²⁺-sensitive mitochondrial-targeted Ca²⁺ indicator was loaded into SHSY-5Y cells and mitochondrial Ca²⁺ efflux was monitored. Mitochondria were loaded with Ca²⁺ using ATP to release it from the ER, and data showed that overexpression of NCLX enhanced and knockdown of NCLX reduced mitochondrial Ca2+ efflux. Luongo et al. (2017) generated a tamoxifen-inducible, cardiac-specific NCLX knockout (KO) mouse. After tamoxifen treatment, 87% of the mice died and this did not occur if the mice were crossed with mice in which cyclophilin D (an activator of the PTP) had been ablated, suggesting that the death was due to Ca2+ overload activation of PTP. In additional studies on digitonin-permeabilized adult cardiomyocytes, mitochondrial Ca²⁺ efflux was reduced by knocking out NCLX (Luongo et al., 2017).

However, the regulation of Na⁺-dependent Ca²⁺ efflux appears to be complicated. Three recent studies have shown that Na⁺-dependent Ca²⁺ efflux is blocked by the deletion of transmembrane protein 65 (TMEM65) (Garbincius et al., 2023, Preprint; Vetralla et al., 2023, Preprint; Zhang et al., 2023, Preprint). Vetralla et al. (2023, Preprint) measured mitochondrial Ca²⁺ efflux in permeabilized HeLa cells in which mitochondria were loaded with an aequorin-based Ca²⁺-sensitive indicator. These measurements were made in the presence and absence of Na⁺ to

follow Na+-dependent and Na+-independent Ca2+ efflux. In the presence of Na⁺, mitochondrial Ca²⁺ efflux was enhanced by overexpression of TMEM65 and reduced by TMEM65 downregulation. In another study, Garbincius et al. (2023, Preprint) identified TMEM65 as interacting with NCLX. They overexpressed TMEM65 in AC16 cells and found an increase in the rate of Na⁺-dependent Ca²⁺ efflux measured using FuraFF to monitor extramitochondrial Ca2+ in digitonin-permeabilized cells. They further showed that CGP37157, an inhibitor of Na+-dependent Ca2+ exchange, blocked this Na+-dependent mitochondrial Ca2+ efflux. Knockdown of TMEM65 in mice using shRNA led to a skeletal muscle myopathy and decreased cardiac contractility. In another study, Zhang et al. (2023, Preprint) generated a global (whole body) TMEM65-KO mouse, which exhibited retarded growth and developed brain seizures leading to death at ~3 wk of age. Similar effects were seen with a neuronal-specific deletion of TMEM65. A skeletal muscle-specific TMEM65-KO exhibited no defects at 2 mo of age but showed growth delay and reduced muscle mass at 3 mo. Mitochondrial Ca2+, measured using Rhod-2, was significantly higher in TMEM65-KO fibers compared with the floxed control. Furthermore, Na+-dependent mitochondrial Ca2+ efflux measured in isolated mitochondria was present in the mitochondria from the floxed control but was absent in TMEM65-KO mitochondria. Interestingly, crossing the global TMEM65-KO mouse with the global MCU-KO mouse rescued the development of seizure and death that occurred in the global TMEM65-KO mouse. Taken together, these three studies agree that Na+-dependent Ca²⁺ efflux requires TMEM65.

The exact details of how TMEM65 and NCLX interact to regulate mitochondrial Na+-dependent Ca2+ efflux is unclear, but perhaps, like the relationship between EMRE and MCU, both are needed for the exchanger to function. It is also possible that TMEM65 and NCLX function independently as Na+-dependent Ca²⁺ exchangers. For this reason, we use NCLX to refer to transport regulated by this protein and Na-dependent Ca²⁺ efflux to refer to overall transport (which may or may not be entirely mediated by NCLX). Another important issue is the stoichiometry. Studies have reported both electrogenic exchange of 3Na+ for 1Ca2+ and electroneutral 2Na+ for 1Ca2+ (Affolter and Carafoli, 1980; Brand, 1985; Jung et al., 1995; Dash and Beard, 2008; Kim and Matsuoka, 2008; Giladi et al., 2022). For the former, net Ca²⁺ efflux is associated with the entry of positive charge and (in contrast to the 1Ca2+:1H+ CHE discussed above) the matrix- negative $\Delta \psi$ will facilitate Ca^{2+} efflux. The electroneutral case will only give net Ca2+ efflux if ([Na+]cvto/ $[Na^+]_{matrix})^2 > ([Ca^{2+}]_{cyto}/[Ca^{2+}]_{matrix})$. Measurement of mitochondrial Na+ would provide some insight into the regulation of mitochondrial Na⁺/Ca²⁺ exchange. There are only a few measurements of mitochondrial Na+, and the studies suggest that basal mitochondrial Na⁺ is ~5 mM, which is lower than cytosolic Na+, and that mitochondrial Na+ increases with addition of cyanide or mitochondrial uncoupler (Donoso et al., 1992). Additional studies are needed to clarify the role of TMEM65 and NCLX in Na+-dependent Ca efflux. Development of mitochondrial targeted Na+ indicators would also help address these questions.



Mitochondrial Ca2+ during ischemia and reperfusion

It has been proposed that an increase in mitochondrial Ca²⁺ during ischemia and reperfusion is responsible for initiating cell death via activation of the mitochondrial PTP and that inhibition of the increase in mitochondrial Ca²⁺ might be cardioprotective (Bauer and Murphy, 2020; Bround et al., 2020; Marta et al., 2021; Bernardi et al., 2023; Murphy and Liu, 2023; Robichaux et al., 2023). Such a strategy would require knowledge of the mechanisms responsible for the increase in mitochondrial Ca²⁺. Furthermore, for inhibition of mitochondrial Ca²⁺ uptake to be protective, ideally the increase in mitochondrial Ca2+ would need to occur on reperfusion so that drugs could be administered before the increase occurred. To address these issues, one needs to have reliable methods to measure mitochondrial Ca2+ in a model that is suitable for inducing ischemia and reperfusion. Ideally, this requires an intact heart. Many studies using isolated myocytes have used "simulated ischemia" by applying solutions to attempt to mimic some of the changes occurring in ischemia and measured mitochondrial-free Ca2+ with a mitochondrialtargeted indicator. However, ischemia is defined as lack of blood flow to an organ which reduces oxygen and nutrient delivery to the organ and reduces washout of end products of metabolism; thus, it is difficult to exactly simulate these conditions with isolated cells. Furthermore, the extracellular volume is frequently very different between an intact heart and isolated cardiomyocytes and this can alter the ion gradients.

In studies carried out >20 years ago, mitochondrial Ca²⁺ was measured in isolated adult rat cardiomyocytes loaded with the Ca²⁺-sensitive indicator indo-1 during hypoxia and reoxygenation (Miyata et al., 1992; Griffiths et al., 1998). Manganese was added to the cells to quench any indo-1 in the cytosol. It should be noted that the addition of manganese can have effects on Ca²⁺ transporters and can slowly enter the mitochondria (Gavin et al., 1999). An increase in mitochondrial Ca²⁺ occurred at the onset of rigor. Approximately 50% of the cardiomyocytes hypercontracted on reoxygenation. At 40 min of hypoxia following the onset of rigor, mitochondrial Ca2+ increased from basal levels (~100 nM) to 280 nM in the cells that subsequently recovered and 743 nM in those that hypercontracted (Griffiths et al., 1998). Studies using clonazepam to inhibit the mitochondrial Na+/Ca2+ exchange showed that none of the cells hypercontracted on reoxygenation. Clonazepam also blocked the increase in mitochondrial Ca²⁺ during the 40 min of hypoxia following rigor. However, on reoxygenation, clonazepam caused an increase in mitochondrial Ca2+, with mitochondrial Ca2+ rising to 465 nM by 15 min of reoxygenation (Griffiths et al., 1998). These data suggest that the increase in mitochondrial Ca2+ during hypoxia is via the reverse mode of Na+/Ca2+ exchange, but that forward Na⁺/Ca²⁺ exchange decreases mitochondrial Ca²⁺ on reoxygenation.

Studies in HL-1 cells (an immortalized cardiomyocyte line) subjected to simulated ischemia and reperfusion reported that addition of 25 μ M ruthenium red (to inhibit MCU) reduced the rise in mitochondrial Ca²⁺ during simulated ischemia; however, this reduction in mitochondrial Ca²⁺ was associated with an increase in lactate dehydrogenase (LDH) release, which was attributed to an increase in cytosolic Ca²⁺ in the presence of

ruthenium red (Ruiz-Meana et al., 2006). Ashok et al. also tested a role for MCU in mitochondrial Ca2+ uptake during 60 min of coverslip-simulated ischemia in neonatal mouse ventricular cardiomyocytes from WT and MCU-KO mice (Ashok et al., 2023). In this study, MCU floxed cells were transfected with a Cre virus to knockdown MCU. Mitochondrial Ca2+ increased during the first 25 min of simulated ischemia. This was, however, followed by a decline in mitochondrial Ca2+ to basal levels occurring at the time of loss of $\Delta \psi$. The mechanism responsible for the decline is unclear. There was no difference in mitochondrial Ca2+ levels during simulated ischemia or reperfusion between the WT and the MCU-KO cardiomyocytes, suggesting that the increase in mitochondrial Ca2+ observed during I/R is not due to Ca²⁺ entry via MCU. Furthermore, there was no difference in LDH release measured following simulated reperfusion, although the levels of LDH released were low. Interestingly, it was observed that addition of CGP37157, an inhibitor of mitochondrial Na+-dependent Ca2+ exchanger, blocked the increase in mitochondrial Ca²⁺ during simulated ischemia, suggesting that Ca²⁺ entry during ischemia is due to mitochondrial Na+/Ca2+ exchange operating in the direction to transport Ca2+ into the mitochondria. This finding is consistent with previous studies mentioned above (Griffiths et al., 1998), which showed in adult cardiomyocytes that inhibition of mitochondrial Na+/Ca2+ exchange blocked the ischemic-induced increase in mitochondrial Ca².

The ability of mitochondrial Na+/Ca2+ exchange to operate in the Ca²⁺ influx mode will depend on the stoichiometry of the exchanger. If Na+/Ca2+ is electroneutral, exchanging 2 Na+ for 1 Ca²⁺, then at equilibrium, the direction of the exchanger would depend only on the gradients for Na⁺ and Ca². With the large rise in cytosolic Ca²⁺ that occurs during ischemia with electroneutral Na+/Ca2+, the exchanger could mediate Ca2+ entry into the mitochondria. However, if the exchanger is electrogenic (3Na+ exchanging with 1 Ca²), then Ca²⁺ entry will be decreased by the negative $\Delta \psi$. Rewriting Eq. 1 for Na⁺, with n = 3 (and assuming that mitochondrial and cytoplasmic Na+ concentrations are equal), we can predict that with a $\Delta \psi$ of -180 mV, net Ca²⁺ uptake (and Na+ efflux) will only occur if cytoplasmic Ca²⁺ is >1,000 times greater than mitochondrial Ca^{2+} . Reducing $\Delta\psi$ to −120 or −60 means that net Ca²⁺ influx requires a concentration gradient of 100 or 10 times, respectively. Thus, it is important to determine both Δψ during ischemia and the stoichiometry of Na⁺/Ca²⁺. As mentioned above, there are reports suggesting both electroneutral and electrogenic mitochondrial Na+/Ca2+ exchange (Affolter and Carafoli, 1980; Brand, 1985; Jung et al., 1995; Dash and Beard, 2008; Kim and Matsuoka, 2008). Interestingly, a recent study utilizing NCLX mutations, the voltage dependence of transport, and molecular dynamic simulation (Giladi et al., 2022) reported that NCLX is electroneutral. Further studies are needed to establish the stoichiometry of the exchanger.

Mastoor et al. (2024) also used neonatal mouse ventricular cardiomyocytes to examine alterations in mitochondria Ca^{2+} during coverslip ischemia and reperfusion. One problem is that simulated ischemia causes a large intracellular acidification, which can alter the K_d for Ca^{2+} binding and fluorescent



properties of Ca²⁺ indicators. To address this concern, a pH-insensitive, genetically encoded, mitochondrial-targeted Ca indicator, TqFLITS, was employed. TqFLITS monitors Ca²⁺ via changes in fluorescent lifetime, which allows measurement of Ca²⁺ that is independent of the indicator concentration, thus allowing one to calculate free Ca²⁺ rather than just changes of fluorescence. In WT cells, after 60 min of simulated ischemia mitochondrial Ca²⁺ increased 2.8-fold, whereas in the MCU-KO the increase was only 1.7-fold, suggesting that a pathway in addition to the MCU contributes to Ca²⁺ entry. Cell death was also monitored during coverslip ischemia and reperfusion using propidium iodide staining. There was no difference in cell death, after 60 min of coverslip ischemia between WT and MCU-KO, again suggesting a role for something other than the MCU.

Given the problems of mimicking ischemia in isolated cells, it is important to measure mitochondrial Ca2+ in an intact beating heart preparation. Miyamae et al. (1996) employed the Ca²⁺ indicator indo-1 to measure mitochondria Ca2+ was not measured during ischemia. We have recently developed a new method to measure mitochondrial Ca2+ transmurally in a beating perfused heart (Petersen et al., 2023). A genetically encoded mitochondrial-targeted Ca2+ indicator, red genetically encoded calcium indicator for optical imaging (R-GECO), is directed to the heart by injecting AAV9-encoding R-GECO into 3-5-day-old mouse pups. At 10-12 wk of age, the heart is perfused in Langendorff mode with a laser inserted into the left ventricle to excite the R-GECO in the heart mitochondria. The heart is placed in an integrating sphere to minimize motion artifacts and the emitted light collected by a spectrometer and analyzed for changes in emission fluorescence. Using this method, a two- to threefold increase in mitochondria Ca2+ was observed during ischemia, followed by a slow recovery of mitochondrial Ca²⁺ toward baseline during reperfusion. The fact that the increase in mitochondrial Ca²⁺ occurs during ischemia limits the ability to target it on reperfusion. However, it is still of interest to understand the mechanisms responsible for the increase in mitochondrial Ca²⁺ during ischemia as this could potentially be useful to develop cardioprotective approaches for use in cardiac surgery.

To test whether MCU is responsible for the increase in mitochondrial Ca^{2+} during I/R, mitochondrial Ca^{2+} was measured in MCU-KO and WT hearts using mitochondrial-targeted R-GECO (Petersen et al., 2023). In WT hearts, mitochondrial Ca^{2+} increased by 2.4-fold during ischemia compared with a 1.7-fold increase in MCU-KO hearts. Thus, the deletion of MCU significantly reduced but did not eliminate the rise in mitochondrial Ca^{2+} during ischemia, suggesting that, as was seen in isolated cardiomyocytes (Mastoor et al., 2024), mechanisms other than MCU contribute to the increase in mitochondrial Ca^{2+} observed during ischemia. It will be interesting to test whether inhibition of pathways such as Na^+/Ca^{2+} can block the rise in mitochondrial Ca^{2+} during ischemia, similar to what has been reported in isolated cardiomyocytes (Griffiths et al., 1998; Ashok et al., 2023).

It is possible that the rise in mitochondrial-free Ca²⁺ may not simply result from altered fluxes across the inner mitochondrial membrane; alterations in Ca²⁺ buffering could also contribute

(Hernansanz-Agustín et al., 2020). Depending on the conditions, it has been reported that >95% of the Ca²⁺ in the mitochondria is bound, much of it in Ca-P granules (reviewed in Eisner et al. [2023]). Acidic conditions, as occur during ischemia, have been shown to promote dissociation of the Ca-P granules (Chalmers and Nicholls, 2003), providing another mechanism to increase mitochondrial-ionized Ca²⁺ that is independent of Ca²⁺ uptake into the mitochondria.

As mentioned previously, it is generally thought that an increase in mitochondrial Ca2+ during ischemia or reperfusion activates the PTP leading to cell death. Studies find that mitochondrial Ca²⁺ rises approximately two- to threefold during simulated ischemia or ischemia (Ashok et al., 2023; Petersen et al., 2023). Basal Ca²⁺ is typically reported to be in the range of ~100 nM; therefore, during ischemia and reperfusion, the rise in mitochondrial Ca^{2+} would be in the range of 200-300 nM. It is important to consider whether this modest increase in mitochondrial Ca²⁺ is sufficient to activate PTP. PTP is typically assayed in isolated mitochondria by measuring the Ca²⁺ retention capacity using an extramitochondrial Ca2+ indicator to determine how much total Ca2+ uptake is needed to trigger PTP opening. When total mitochondrial Ca2+ is low (<10 nmol/mg protein) and phosphate is present, there is a more or less linear relationship between Ca2+ uptake and free mitochondrial Ca2+ levels (Nicholls, 1978). However, as the total mitochondrial Ca²⁺ increases above 10 nmol/mg protein, the added Ca²⁺ is largely buffered and with additional Ca²⁺ uptake, there is little further increase in free mitochondrial Ca²⁺. In fact, it has been reported that activation of PTP is linked to total mitochondrial Ca2+ rather than free Ca²⁺ (Chalmers and Nicholls, 2003; Wei et al., 2012). Thus in isolated mitochondria, under conditions that activate PTP, the free matrix Ca²⁺ is highly buffered. This buffering might explain, in part, why the rise in Ca²⁺ during ischemia is low. Another point to consider is that during ischemia, the matrix would be expected to become more acidic and this would alter the Ca2+ buffering. The studies examining free to bound Ca²⁺ were typically performed at pH 7.2-7.4. The observation that total rather than free mitochondrial Ca2+ activates PTP raises the question of whether inhibition of MCU or Ca2+ efflux mechanisms alters total as well as free mitochondrial Ca2+. It is also important to understand how a change in total mitochondrial Ca²⁺ might lead to activation of PTP. One possibility is that the PTP interacts with the major buffer and therefore senses total Ca2+. Interestingly, polyphosphate has been suggested to regulate PTP and perhaps polyphosphate is involved in sensing total Ca²⁺ (Seidlmayer et al., 2012). Future studies need to examine mechanisms by which total Ca2+ might regulate PTP.

Role of mitochondrial Ca²⁺ transporters in I/R injury

Regardless of whether it is total or free Ca^{2+} that activates PTP and initiates cell death, the question remains as to whether inhibiting mitochondrial Ca^{2+} uptake (which should inhibit the increase in total mitochondrial Ca^{2+}) is cardioprotective. Several studies, using either pharmacological inhibitors or genetic mouse models, have examined the effects of inhibiting the MCU. Studies in perfused rat heart have found that addition of






Figure 1. **Mechanisms regulating mitochondrial Ca²⁺.** The MCU is the primary protein responsible for Ca²⁺ uptake into the mitochondria. The energy for Ca²⁺ uptake is provided by the large negative membrane potential that is generated by the electron transport chain (ETC). The ETC is also responsible for generating an alkaline matrix pH. The inwardly directed proton gradient drives the efflux of a number of ions including Na⁺, K⁺, and Ca²⁺. In addition to CHE, Ca²⁺ efflux from the matrix occurs via a CHE. There are also data suggesting a role for Ca²⁺ entry via an electrogenic Ca²⁺/H⁺ exchanger, although this needs confirmation. Much of the mitochondrial Ca²⁺ is buffered within Ca-P granules. The phosphate enters the mitochondria on the phosphate carrier (PiC) in exchange for proton.

ruthenium red improved after ischemic contractile function (Grover et al., 1990; Benzi and Lerch, 1992; Zhang et al., 2006) and reduced infarct size or enzyme release (Benzi and Lerch, 1992; Zhang et al., 2006). One issue regarding ruthenium red is that it may affect other Ca2+ transporters. Ru360 has been isolated as the active compound in ruthenium red that is responsible for inhibition of MCU, and a study with this more specific compound also found it to improve ischemic contractile function (de Jesus Garcia-Rivas et al., 2005). A potential limitation of Ru360 is that it is reported to poorly permeate the plasma membrane (Hajnoczky et al., 2006). However, arguing against this concern, a number of studies have reported Ru360 inhibition of mitochondrial Ca²⁺ uptake in cardiac cells and tissue (Unitt et al., 1989; Matlib et al., 1998; Kosmach et al., 2021). As discussed, the ability of Ru360 to accumulate in cells at a level sufficient to inhibit MCU may depend on the cell type as Ru360 and MICU1 compete for binding to MCU (Rodríguez-Prados et al., 2023). Therefore, the ability of Ru360 to inhibit MCU is dependent on the tissue levels of MICU1. The heart has a very low ratio of MICU1 to MCU, perhaps explaining why Ru360 is effective in blocking MCU in the heart. A more potent inhibitor of MCU, Ru265, has also been developed (Woods et al., 2019). Interestingly, Cys 97 of MCU appears to be required for inhibition by Ru265, suggesting that Ru265 might inhibit by a different mechanism than Ru360. Experiments on mice with genetic deletion of MCU have yielded mixed results. Perfused hearts from mice with germline deletion of MCU subjected to I/R did

not show protection; infarct size was not different between WT and MCU-KO (Pan et al., 2013). Similarly, perfused hearts from a mouse model in which a dominant negative MCU was overexpressed in cardiac cells using the α -myosin heavy chain promoter were not protected from I/R injury (Rasmussen et al., 2015). In contrast, mice with a tamoxifen-inducible cardiacspecific loss of MCU in adults showed reduced infarct size in an in vivo I/R model (Kwong et al., 2015; Luongo et al., 2015). These differences have typically been attributed to adaptations that occur in mice when MCU is deleted prior to birth. However, recent studies in neonatal mouse ventricular cardiomyocytes, in which MCU is acutely deleted, did not show a reduction in death following simulated ischemia (Ashok et al., 2023; Mastoor et al., 2024). Additional studies will be needed to address these discrepancies.

Very few studies have been performed to test whether the mitochondria Na⁺/Ca²⁺ exchange inhibitor, CGP37157, is cardioprotective when given to hearts before I/R. Griffiths et al. reported that addition of clonazepam blocked the development of hypercontracture on reoxygenation (Griffiths et al., 1998). However, even though Ashok et al. found that addition of CGP37157 reduced the rise in Ca²⁺ during ischemia, there was no reduction in LDH release or alteration in $\Delta\psi$ recovery (Ashok et al., 2023). One issue is that blocking Ca²⁺ efflux at the beginning of reperfusion would be detrimental and it is difficult to remove the inhibitor at the start of reperfusion. Somewhat in contrast to the reduction in mitochondrial Ca²⁺ observed with



addition of CGP37157, when a mouse with cardiac-restricted overexpression of NCLX was subjected to an in vivo model of coronary ligation for 40 min followed by 24 h of reperfusion, the infarct size was less than that in WT (Luongo et al., 2017). Thus, the role of Na⁺-dependent Ca²⁺ exchange in regulating mitochondrial Ca²⁺ during I/R is unclear. There is much we need to better understand about Na⁺-dependent Ca²⁺ exchange, including the stoichiometry and the relationship of TMEM65 and NCLX. We also need a better understanding of the regulation of mitochondrial Na⁺.

Conclusion

Taken together, data in the literature suggest that a rise in mitochondrial Ca2+ occurs during ischemia. However, the mechanisms responsible for this increase are not totally defined. Inhibition of MCU only partially blocks the increase in mitochondrial Ca2+ during ischemia. Studies using inhibitors of mitochondrial Na+-dependent Ca2+ exchange have suggested that during ischemia, Ca²⁺ may enter the mitochondria in exchange for Na⁺. The contribution of CHE to the control of mitochondrial Ca²⁺ is also unclear. As discussed, the roles of these exchangers will depend on their stoichiometry and further work is needed. Release of Ca²⁺ from Ca-P could also provide a source for the rise in mitochondrial Ca²⁺ during ischemia. A major question that needs to be addressed is the mechanism by which increased uptake of Ca²⁺ into the mitochondria actives PTP. It appears that activation of PTP correlates better with an increase in total rather than free Ca²⁺. Despite the fact that mitochondrial Ca²⁺ has been studied for over 60 years, the details of mechanisms regulating mitochondrial Ca²⁺ uptake and release, especially the stoichiometry of the efflux pathways is still unclear (Fig. 1). Furthermore, the role of Ca²⁺ buffering in mitochondrial Ca²⁺ homeostasis and in activation of the PTP also needs additional study.

Acknowledgments

Olaf S. Andersen served as an editor.

E. Murphy was funded by the National Heart, Lung, and Blood Institute intramural program (ZO1HL006059 and ZO1HL002066).

Author contributions: E. Murphy: conceptualization, data curation, funding acquisition, visualization, and writing—original draft, review, and editing. D.A. Eisner: conceptualization, investigation, and writing—original draft, review, and editing.

Disclosures: The authors declare no competing interests exist.

Submitted: 24 September 2024 Revised: 14 November 2024 Accepted: 6 December 2024

References

- Affolter, H., and E. Carafoli. 1980. The Ca²⁺-Na⁺ antiporter of heart mitochondria operates electroneutrally. *Biochem. Biophys. Res. Commun.* 95: 193–196. https://doi.org/10.1016/0006-291X(80)90723-8
- Ashok, D., K. Papanicolaou, A. Sidor, M. Wang, S. Solhjoo, T. Liu, and B. O'Rourke. 2023. Mitochondrial membrane potential instability on

- reperfusion after ischemia does not depend on mitochondrial Ca²⁺ uptake. *J. Biol. Chem.* 299:104708. https://doi.org/10.1016/j.jbc.2023.104708
- Austin, S., R. Mekis, S.E.M. Mohammed, M. Scalise, W.A. Wang, M. Galluccio, C. Pfeiffer, T. Borovec, K. Parapatics, D. Vitko, et al. 2022. TMBIM5 is the Ca²⁺/H⁺ antiporter of mammalian mitochondria. *EMBO Rep.* 23: e54978. https://doi.org/10.15252/embr.202254978
- Bauer, T.M., and E. Murphy. 2020. Role of mitochondrial calcium and the permeability transition pore in regulating cell death. Circ. Res. 126: 280-293. https://doi.org/10.1161/CIRCRESAHA.119.316306
- Baughman, J.M., F. Perocchi, H.S. Girgis, M. Plovanich, C.A. Belcher-Timme, Y. Sancak, X.R. Bao, L. Strittmatter, O. Goldberger, R.L. Bogorad, et al. 2011. Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter. *Nature*. 476:341–345. https://doi.org/10.1038/nature10234
- Benzi, R.H., and R. Lerch. 1992. Dissociation between contractile function and oxidative metabolism in postischemic myocardium. Attenuation by ruthenium red administered during reperfusion. Circ. Res. 71:567–576. https://doi.org/10.1161/01.RES.71.3.567
- Bernardi, P., C. Gerle, A.P. Halestrap, E.A. Jonas, J. Karch, N. Mnatsakanyan, E. Pavlov, S.S. Sheu, and A.A. Soukas. 2023. Identity, structure, and function of the mitochondrial permeability transition pore: Controversies, consensus, recent advances, and future directions. *Cell Death Differ*. 30:1869–1885. https://doi.org/10.1038/s41418-023-01187-0
- Boyman, L., M. Greiser, and W.J. Lederer. 2021. Calcium influx through the mitochondrial calcium uniporter holocomplex, MCU_{cx}. J. Mol. Cell. Cardiol. 151:145–154. https://doi.org/10.1016/j.yjmcc.2020.10.015
- Brand, M.D. 1985. The stoichiometry of the exchange catalysed by the mitochondrial calcium/sodium antiporter. *Biochem. J.* 229:161–166. https://doi.org/10.1042/bj2290161
- Bround, M.J., D.M. Bers, and J.D. Molkentin. 2020. A 20/20 view of ANT function in mitochondrial biology and necrotic cell death. *J. Mol. Cell. Cardiol.* 144:A3–A13. https://doi.org/10.1016/j.yimcc.2020.05.012
- Chalmers, S., and D.G. Nicholls. 2003. The relationship between free and total calcium concentrations in the matrix of liver and brain mitochondria. J. Biol. Chem. 278:19062-19070. https://doi.org/10.1074/jbc.M212661200
- Crompton, M., M. Künzi, and E. Carafoli. 1977. The calcium-induced and sodium-induced effluxes of calcium from heart mitochondria. Evidence for a sodium-calcium carrier. *Eur. J. Biochem.* 79:549–558. https://doi.org/10.1111/j.1432-1033.1977.tb11839.x
- Csordás, G., T. Golenár, E.L. Seifert, K.J. Kamer, Y. Sancak, F. Perocchi, C. Moffat, D. Weaver, S.F. Perez, R. Bogorad, et al. 2013. MICU1 controls both the threshold and cooperative activation of the mitochondrial Ca²⁺ uniporter. *Cell Metab.* 17:976–987. https://doi.org/10.1016/j.cmet.2013
- Dash, R.K., and D.A. Beard. 2008. Analysis of cardiac mitochondrial Na⁺-Ca²⁺ exchanger kinetics with a biophysical model of mitochondrial Ca²⁺ handling suggests a 3:1 stoichiometry. *J. Physiol.* 586:3267–3285. https://doi.org/10.1113/jphysiol.2008.151977
- de Jesús García-Rivas, G., A. Guerrero-Hernández, G. Guerrero-Serna, J.S. Rodríguez-Zavala, and C. Zazueta. 2005. Inhibition of the mitochondrial calcium uniporter by the oxo-bridged dinuclear ruthenium amine complex (Ru360) prevents from irreversible injury in postischemic rat heart. FEBS J. 272:3477-3488. https://doi.org/10.1111/j.1742-4658.2005.04771.x
- De Stefani, D., A. Raffaello, E. Teardo, I. Szabò, and R. Rizzuto. 2011. A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature*. 476:336–340. https://doi.org/10.1038/nature10230
- Donoso, P., J.G. Mill, S.C. O'Neill, and D.A. Eisner. 1992. Fluorescence measurements of cytoplasmic and mitochondrial sodium concentration in rat ventricular myocytes. J. Physiol. 448:493–509. https://doi.org/10.1113/jphysiol.1992.sp019053
- Eisner, D., E. Neher, H. Taschenberger, and G. Smith. 2023. Physiology of intracellular calcium buffering. Physiol. Rev. 103:2767–2845. https://doi. org/10.1152/physrev.00042.2022
- Garbincius, J.F., and J.W. Elrod. 2022. Mitochondrial calcium exchange in physiology and disease. *Physiol. Rev.* 102:893–992. https://doi.org/10 .1152/physrev.00041.2020
- Garbincius, J.F., O. Salik, H.M. Cohen, C. Choya-Foces, A.S. Mangold, A.D. Makhoul, A.E. Schmidt, D.Y. Khalil, J.J. Doolittle, A.S. Wilkinson, et al. 2023. TMEM65 regulates NCLX-dependent mitochondrial calcium efflux. bioRxiv. https://doi.org/10.1101/2023.10.06.561062 (Preprint posted October 09, 2023).
- Gavin, C.E., K.K. Gunter, and T.E. Gunter. 1999. Manganese and calcium transport in mitochondria: Implications for manganese toxicity. Neurotoxicology. 20:445-453.



- Giladi, M., S. Mitra, L. Simhaev, R. Hiller, B. Refaeli, T. Strauss, C.R. Baiz, and D. Khananshvili. 2022. Exploring the Li⁺ transporting mutant of NCX_Mj for assigning ion binding sites of mitochondrial NCLX. Cell Calcium. 107:102651. https://doi.org/10.1016/j.ceca.2022.102651
- Griffiths, E.J., C.J. Ocampo, J.S. Savage, G.A. Rutter, R.G. Hansford, M.D. Stern, and H.S. Silverman. 1998. Mitochondrial calcium transporting pathways during hypoxia and reoxygenation in single rat cardiomyocytes. Cardiovasc. Res. 39:423–433. https://doi.org/10.1016/S0008-6363(98) 00104-7
- Grover, G.J., S. Dzwonczyk, and P.G. Sleph. 1990. Ruthenium red improves postischemic contractile function in isolated rat hearts. *J. Cardiovasc. Pharmacol.* 16:783–789. https://doi.org/10.1097/00005344-199011000-00014
- Hajnóczky, G., G. Csordás, S. Das, C. Garcia-Perez, M. Saotome, S. Sinha Roy, and M. Yi. 2006. Mitochondrial calcium signalling and cell death: Approaches for assessing the role of mitochondrial Ca²⁺ uptake in apoptosis. Cell Calcium. 40:553–560. https://doi.org/10.1016/j.ceca.2006.08
- Hernansanz-Agustín, P., C. Choya-Foces, S. Carregal-Romero, E. Ramos, T. Oliva, T. Villa-Piña, L. Moreno, A. Izquierdo-Álvarez, J.D. Cabrera-García, A. Cortés, et al. 2020. Na⁺ controls hypoxic signalling by the mitochondrial respiratory chain. *Nature*. 586:287-291. https://doi.org/10.1038/s41586-020-2551-y
- Jiang, D., L. Zhao, and D.E. Clapham. 2009. Genome-wide RNAi screen identifies Letm1 as a mitochondrial Ca²⁺/H⁺ antiporter. Science. 326: 144–147. https://doi.org/10.1126/science.1175145
- Jung, D.W., K. Baysal, and G.P. Brierley. 1995. The sodium-calcium antiport of heart mitochondria is not electroneutral. J. Biol. Chem. 270:672–678. https://doi.org/10.1074/jbc.270.2.672
- Kamer, K.J., Z. Grabarek, and V.K. Mootha. 2017. High-affinity cooperative Ca²⁺ binding by MICU1-MICU2 serves as an on-off switch for the uniporter. EMBO Rep. 18:1397–1411. https://doi.org/10.15252/embr .201643748
- Kim, B., and S. Matsuoka. 2008. Cytoplasmic Na⁺-dependent modulation of mitochondrial Ca²⁺ via electrogenic mitochondrial Na⁺-Ca²⁺ exchange. J. Physiol. 586:1683–1697. https://doi.org/10.1113/jphysiol.2007 148726
- Kosmach, A., B. Roman, J. Sun, A. Femnou, F. Zhang, C. Liu, C.A. Combs, R.S. Balaban, and E. Murphy. 2021. Monitoring mitochondrial calcium and metabolism in the beating MCU-KO heart. Cell Rep. 37:109846. https://doi.org/10.1016/j.celrep.2021.109846
- Kwong, J.Q., X. Lu, R.N. Correll, J.A. Schwanekamp, R.J. Vagnozzi, M.A. Sargent, A.J. York, J. Zhang, D.M. Bers, and J.D. Molkentin. 2015. The mitochondrial calcium uniporter selectively matches metabolic output to acute contractile stress in the heart. Cell Rep. 12:15–22. https://doi.org/10.1016/j.celrep.2015.06.002
- Liu, J.C., J. Liu, K.M. Holmström, S. Menazza, R.J. Parks, M.M. Fergusson, Z.X. Yu, D.A. Springer, C. Halsey, C. Liu, et al. 2016. MICU1 serves as a molecular gatekeeper to prevent in vivo mitochondrial calcium overload. Cell Rep. 16:1561–1573. https://doi.org/10.1016/j.celrep.2016.07.011
- Lu, X., K.S. Ginsburg, S. Kettlewell, J. Bossuyt, G.L. Smith, and D.M. Bers. 2013. Measuring local gradients of intramitochondrial [Ca(2+)] in cardiac myocytes during sarcoplasmic reticulum Ca(2+) release. Circ. Res. 112:424-431. https://doi.org/10.1161/CIRCRESAHA.111.300501
- Luongo, T.S., J.P. Lambert, A. Yuan, X. Zhang, P. Gross, J. Song, S. Shanmughapriya, E. Gao, M. Jain, S.R. Houser, et al. 2015. The mitochondrial calcium uniporter matches energetic supply with cardiac workload during stress and modulates permeability transition. *Cell Rep.* 12:23–34. https://doi.org/10.1016/j.celrep.2015.06.017
- Luongo, T.S., J.P. Lambert, P. Gross, M. Nwokedi, A.A. Lombardi, S. Shanmughapriya, A.C. Carpenter, D. Kolmetzky, E. Gao, J.H. van Berlo, et al. 2017. The mitochondrial Na⁺/Ca²⁺ exchanger is essential for Ca²⁺ homeostasis and viability. *Nature*. 545:93–97. https://doi.org/10.1038/nature22082
- Márta, K., P. Hasan, M. Rodríguez-Prados, M. Paillard, and G. Hajnóczky. 2021. Pharmacological inhibition of the mitochondrial Ca²⁺ uniporter: Relevance for pathophysiology and human therapy. J. Mol. Cell. Cardiol. 151:135-144. https://doi.org/10.1016/j.yjmcc.2020.09.014
- Mastoor, Y., M. Harata, K. Silva, C. Liu, C.A. Combs, B. Roman, and E. Murphy. 2024. Monitoring mitochondrial calcium in cardiomyocytes during coverslip hypoxia using a fluorescent lifetime indicator. J. Mol. Cell. Cardiol. Plus. 8:100074. https://doi.org/10.1016/j.jmccpl.2024.100074
- Matlib, M.A., Z. Zhou, S. Knight, S. Ahmed, K.M. Choi, J. Krause-Bauer, R. Phillips, R. Altschuld, Y. Katsube, N. Sperelakis, and D.M. Bers. 1998.

- Oxygen-bridged dinuclear ruthenium amine complex specifically inhibits Ca²⁺ uptake into mitochondria in vitro and in situ in single cardiac myocytes. *J. Biol. Chem.* 273:10223–10231. https://doi.org/10.1074/
- Miyamae, M., S.A. Camacho, M.W. Weiner, and V.M. Figueredo. 1996. Attenuation of postischemic reperfusion injury is related to prevention of [Ca2+]m overload in rat hearts. *Am. J. Physiol.* 271:H2145-H2153. https://doi.org/10.1152/ajpheart.1996.271.5.H2145
- Miyata, H., E.G. Lakatta, M.D. Stern, and H.S. Silverman. 1992. Relation of mitochondrial and cytosolic free calcium to cardiac myocyte recovery after exposure to anoxia. Circ. Res. 71:605–613. https://doi.org/10.1161/01 .RES.71.3.605
- Murphy, E., and J.C. Liu. 2023. Mitochondrial calcium and reactive oxygen species in cardiovascular disease. Cardiovasc. Res. 119:1105–1116. https:// doi.org/10.1093/cvr/cvac134
- Murphy, E., and C. Steenbergen. 2021. Regulation of mitochondrial Ca²⁺ uptake. *Annu. Rev. Physiol.* 83:107–126. https://doi.org/10.1146/annurev-physiol-031920-092419
- Nicholls, D.G. 1978. The regulation of extramitochondrial free calcium ion concentration by rat liver mitochondria. *Biochem. J.* 176:463–474. https://doi.org/10.1042/bj1760463
- Nicholls, D.G., and M. Crompton. 1980. Mitochondrial calcium transport. FEBS Lett. 111:261-268. https://doi.org/10.1016/0014-5793(80) 80806-4
- Paillard, M., G. Csordás, G. Szanda, T. Golenár, V. Debattisti, A. Bartok, N. Wang, C. Moffat, E.L. Seifert, A. Spät, and G. Hajnóczky. 2017. Tissue-specific mitochondrial decoding of cytoplasmic Ca²⁺ signals is controlled by the stoichiometry of MICU1/2 and MCU. Cell Rep. 18: 2291–2300. https://doi.org/10.1016/j.celrep.2017.02.032
- Paillard, M., G. Csordás, K.T. Huang, P. Várnai, S.K. Joseph, and G. Hajnóczky. 2018. MICUI interacts with the D-ring of the MCU pore to control its Ca²⁺ flux and sensitivity to Ru360. Mol. Cell. 72:778–785.e3. https://doi.org/10.1016/j.molcel.2018.09.008
- Palty, R., W.F. Silverman, M. Hershfinkel, T. Caporale, S.L. Sensi, J. Parnis, C. Nolte, D. Fishman, V. Shoshan-Barmatz, S. Herrmann, et al. 2010. NCLX is an essential component of mitochondrial Na⁺/Ca²⁺ exchange. Proc. Natl. Acad. Sci. USA. 107:436–441. https://doi.org/10.1073/pnas.0908099107
- Pan, X., J. Liu, T. Nguyen, C. Liu, J. Sun, Y. Teng, M.M. Fergusson, I.I. Rovira, M. Allen, D.A. Springer, et al. 2013. The physiological role of mitochondrial calcium revealed by mice lacking the mitochondrial calcium uniporter. Nat. Cell Biol. 15:1464–1472. https://doi.org/10.1038/ncb2868
- Patron, M., V. Checchetto, A. Raffaello, E. Teardo, D. Vecellio Reane, M. Mantoan, V. Granatiero, I. Szabò, D. De Stefani, and R. Rizzuto. 2014. MICU1 and MICU2 finely tune the mitochondrial Ca²⁺ uniporter by exerting opposite effects on MCU activity. Mol. Cell. 53:726-737. https://doi.org/10.1016/j.molcel.2014.01.013
- Patron, M., D. Tarasenko, H. Nolte, L. Kroczek, M. Ghosh, Y. Ohba, Y. Lasarzewski, Z.A. Ahmadi, A. Cabrera-Orefice, A. Eyiama, et al. 2022. Regulation of mitochondrial proteostasis by the proton gradient. EMBO J. 41:e110476. https://doi.org/10.15252/embj.2021110476
- Petersen, C.E., J. Sun, K. Silva, A. Kosmach, R.S. Balaban, and E. Murphy. 2023. Increased mitochondrial free Ca²⁺ during ischemia is suppressed, but not eliminated by, germline deletion of the mitochondrial Ca²⁺ uniporter. Cell Rep. 42:112735. https://doi.org/10.1016/j.celrep.2023 .112735
- Plovanich, M., R.L. Bogorad, Y. Sancak, K.J. Kamer, L. Strittmatter, A.A. Li, H.S. Girgis, S. Kuchimanchi, J. De Groot, L. Speciner, et al. 2013. MICU2, a paralog of MICU1, resides within the mitochondrial uniporter complex to regulate calcium handling. PLoS One. 8:e55785. https://doi.org/ 10.1371/journal.pone.0055785
- Rasmussen, T.P., Y. Wu, M.L. Joiner, O.M. Koval, N.R. Wilson, E.D. Luczak, Q. Wang, B. Chen, Z. Gao, Z. Zhu, et al. 2015. Inhibition of MCU forces extramitochondrial adaptations governing physiological and pathological stress responses in heart. Proc. Natl. Acad. Sci. USA. 112:9129–9134. https://doi.org/10.1073/pnas.1504705112
- Robichaux, D.J., M. Harata, E. Murphy, and J. Karch. 2023. Mitochondrial permeability transition pore-dependent necrosis. J. Mol. Cell. Cardiol. 174:47–55. https://doi.org/10.1016/j.yjmcc.2022.11.003
- Rodríguez-Prados, M., E. Berezhnaya, M.T. Castromonte, S.L. Menezes-Filho, M. Paillard, and G. Hajnóczky. 2023. MICU1 occludes the mitochondrial calcium uniporter in divalent-free conditions. Proc. Natl. Acad. Sci. USA. 120:e2218999120. https://doi.org/10.1073/pnas.2218999120
- Ruiz-Meana, M., D. Garcia-Dorado, E. Miró-Casas, A. Abellán, and J. Soler-Soler. 2006. Mitochondrial Ca²⁺ uptake during simulated ischemia does not affect permeability transition pore opening upon simulated



- reperfusion. Cardiovasc. Res. 71:715–724. https://doi.org/10.1016/j.cardiores .2006.06.019
- Sancak, Y., A.L. Markhard, T. Kitami, E. Kovács-Bogdán, K.J. Kamer, N.D. Udeshi, S.A. Carr, D. Chaudhuri, D.E. Clapham, A.A. Li, et al. 2013. EMRE is an essential component of the mitochondrial calcium uniporter complex. Science. 342:1379–1382. https://doi.org/10.1126/science.1242993
- Seidlmayer, L.K., M.R. Gomez-Garcia, L.A. Blatter, E. Pavlov, and E.N. Dedkova. 2012. Inorganic polyphosphate is a potent activator of the mitochondrial permeability transition pore in cardiac myocytes. J. Gen. Physiol. 139:321–331. https://doi.org/10.1085/jgp.201210788
- Tsai, M.F., D. Jiang, L. Zhao, D. Clapham, and C. Miller. 2014. Functional reconstitution of the mitochondrial Ca²⁺/H⁺ antiporter Letml. *J. Gen. Physiol.* 143:67–73. https://doi.org/10.1085/jgp.201311096
- Unitt, J.F., J.G. McCormack, D. Reid, L.K. MacLachlan, and P.J. England. 1989. Direct evidence for a role of intramitochondrial Ca²⁺ in the regulation of oxidative phosphorylation in the stimulated rat heart. Studies using 31P n.m.r. and ruthenium red. Biochem. J. 262:293–301. https://doi.org/10 .1042/bj2620293
- Vecellio Reane, D., J.D.C. Serna, and A. Raffaello. 2024. Unravelling the complexity of the mitochondrial Ca²⁺ uniporter: Regulation, tissue specificity, and physiological implications. *Cell Calcium*. 121:102907. https://doi.org/10.1016/j.ceca.2024.102907
- Vetralla, M., L. Wischhof, V. Cadenelli, E. Scifo, D. Ehninger, R. Rizzuto, D. Bano, and D.D. Stefani. 2023. TMEM65-dependent Ca²⁺ extrusion safeguards mitochondrial homeostasis. *bioRxiv*. https://doi.org/10.1101/2023.10.10.561661 (Preprint posted October 11, 2023).
- Waldeck-Weiermair, M., C. Jean-Quartier, R. Rost, M.J. Khan, N. Vishnu, A.I. Bondarenko, H. Imamura, R. Malli, and W.F. Graier. 2011. Leucine zipper EF hand-containing transmembrane protein 1 (Letml) and uncoupling proteins 2 and 3 (UCP2/3) contribute to two distinct

- mitochondrial Ca²⁺ uptake pathways. *J. Biol. Chem.* 286:28444–28455. https://doi.org/10.1074/jbc.M111.244517
- Wei, A.C., T. Liu, S. Cortassa, R.L. Winslow, and B. O'Rourke. 2011. Mitochondrial Ca²⁺ influx and efflux rates in Guinea pig cardiac mitochondria: Low and high affinity effects of cyclosporine A. Biochim. Biophys. Acta. 1813:1373–1381. https://doi.org/10.1016/j.bbamcr.2011.02.012.
- Wei, A.C., T. Liu, R.L. Winslow, and B. O'Rourke. 2012. Dynamics of matrix-free Ca²⁺ in cardiac mitochondria: Two components of Ca²⁺ uptake and role of phosphate buffering. *J. Gen. Physiol.* 139:465–478. https://doi.org/10.1085/jgp.201210784
- Woods, J.J., N. Nemani, S. Shanmughapriya, A. Kumar, M. Zhang, S.R. Nathan, M. Thomas, E. Carvalho, K. Ramachandran, S. Srikantan, et al. 2019. A selective and cell-permeable mitochondrial calcium uniporter (MCU) inhibitor preserves mitochondrial bioenergetics after hypoxia/reoxygenation injury. ACS Cent. Sci. 5:153–166. https://doi.org/10.1021/acscentsci.8b00773
- Zhang, S.Z., Q. Gao, C.M. Cao, I.C. Bruce, and Q. Xia. 2006. Involvement of the mitochondrial calcium uniporter in cardioprotection by ischemic preconditioning. *Life Sci.* 78:738–745. https://doi.org/10.1016/j.lfs.2005.05 .076
- Zhang, L., F. Dietsche, B. Seitaj, L. Rojas-Charry, N. Latchman, D. Tomar, R.C. Wüst, A. Nickel, K.B. Frauenknecht, B. Schoser, et al. 2022. TMBIM5 loss of function alters mitochondrial matrix ion homeostasis and causes a skeletal myopathy. Life Sci. Alliance. 5:e202201478. https://doi.org/10.26508/lsa.202201478
- Zhang, Y., L. Reyes, J. Sun, C. Liu, D. Springer, A. Noguchi, A.M. Aponte, J. Munasinghe, R. Covian, E. Murphy, and B. Glancy. 2023. Loss of TMEM65 causes mitochondrial disease mediated by mitochondrial calcium. bioRxiv. https://doi.org/10.1101/2022.08.02.502535 (Preprint posted October 11, 2023).