

ARTICLE

Autoantibodies neutralizing type I IFNs in 40% of patients with WNV encephalitis in seven new cohorts

Adrian Gervais^{1,2*}, Francesca Trespidi^{3,4,5*}, Alessandro Ferrari^{6,7*}, Francesca Rovida^{6,8*}, Astrid Marchal^{1,2**}, Stefania Croce^{9***}, Irene Cassaniti^{6,8***}, Mattia Moratti^{10,11}, Jennifer L. Uhrlaub¹², David M. Florian¹³, Karin Stiasny¹³, Elisa Burdino¹⁴, Micol Angelini³, Lucy Bizien^{1,2}, Daniele Lilleri⁶, Veronica Codullo¹⁵, Tal Freund¹⁶, Yael Paran¹⁷, Avi Gadoth¹⁸, Roni Biran¹⁷, Alessandro Mancon¹⁹, Camilla Lucca²⁰, Stefania Vogiatzis²⁰, Monia Pacenti²¹, Mélodie Aubart^{1,2,22}, Marco Zecca²³, Patrizia Comoli⁹, Maria Antonietta Avanzini^{9,23}, Jacques Fellay^{4,24}, Antonio Piralla⁶, Francesca Conti^{10,11}, Alberto Dolci^{19,25}, Luisa Barzon^{20,21}, Valeria Ghisetti¹⁴, Tiziana Lazzarotto^{26,27}, Danilo Cereda²⁸, Alessandro Aiuti^{29,30,31}, Emmanuelle Jouanguy^{1,2,32}, Paul Bastard^{1,2,32,33***}, Margaret R. MacDonald^{34***}, Charles M. Rice^{34***}, Anne Puel^{1,2,32***}, Laurent Abel^{1,2,32***}, Giada Rossini^{26***}, Davide Mileto^{19***}, Yannick Simonin^{35***}, Anna Nagy^{36***}, David Hagin^{16***}, Kristy O. Murray^{37***}, Fausto Baldanti^{6,8***}, Judith H. Aberle^{13***}, Aurélie Cobat^{1,2,32***}, Shen-Ying Zhang^{1,2,32***}, Jean-Laurent Casanova^{1,2,32,38,39****}, and Alessandro Borghesi^{3,4,40,41****}

Mosquito-borne West Nile virus (WNV) infection is a growing global health problem. About 0.5% of infected individuals develop encephalitis. We previously showed that 40% of patients in six cohorts had WNV encephalitis because of circulating autoantibodies (auto-Abs) neutralizing type I IFNs. In seven new cohorts, we found that the prevalence of auto-Abs was highest (40% [17–44%]) in patients with encephalitis and very low in a small sample of individuals with asymptomatic or mild infection. In the 13 European, Middle Eastern, and American cohorts available, odds ratios (OR) for WNV encephalitis in individuals with these auto-Abs relative to those without them in a large sample of the general population untested for WNV infection range from ~20 (OR = 17.7; 95% CI: 13.8–22.8, $P < 10^{-16}$) for auto-Abs neutralizing only 100 pg/ml IFN- α 2 and/or IFN- ω to >2,000 (OR = 2,218.4; 95% CI: 125.1–39,337.7, $P < 10^{-16}$) for auto-Abs neutralizing high concentrations of IFN- α 2 and high or low concentrations of IFN- ω . Preexisting auto-Abs neutralizing type I IFNs are therefore causal for WNV encephalitis in about 40% of patients.

¹Laboratory of Human Genetics of Infectious Diseases, Necker Branch, Institut National de la Santé et de la Recherche Médicale (INSERM) U1163, Necker Hospital for Sick Children, Paris, France; ²Imagine Institute, Paris Cité University, Paris, France; ³Host-Pathogen Interaction Group and Neonatal Intensive Care Unit, San Matteo Research Hospital (Fondazione IRCCS Policlinico San Matteo), Pavia, Italy; ⁴School of Life Sciences, Swiss Federal Institute of Technology, Lausanne, Switzerland; ⁵Medical Genetics, Department of Molecular Medicine, University of Pavia, Pavia, Italy; ⁶Microbiology and Virology Unit, San Matteo Research Hospital (Fondazione IRCCS Policlinico San Matteo), Pavia, Italy; ⁷Department of Public Health, Experimental and Forensic Medicine, University of Pavia, Pavia, Italy; ⁸Department of Clinical, Surgical, Diagnostic and Pediatric Sciences, University of Pavia, Pavia, Italy; ⁹Cell Factory, San Matteo Research Hospital (Fondazione IRCCS Policlinico San Matteo), Pavia, Italy; ¹⁰Pediatric Unit, University Hospital of Bologna, Bologna, Italy; ¹¹Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy; ¹²Department of Immunobiology, The University of Arizona, Tucson, AZ, USA; ¹³Center for Virology, Medical University of Vienna, Vienna, Austria; ¹⁴Laboratory of Microbiology and Virology, Amedeo di Savoia Hospital, Turin, Italy; ¹⁵Rheumatology Unit, San Matteo Research Hospital (Fondazione IRCCS Policlinico San Matteo), Pavia, Italy; ¹⁶Allergy and Clinical Immunology Unit, Department of Medicine, Tel Aviv Sourasky Medical Center and Tel Aviv University, Tel Aviv, Israel; ¹⁷Infection Diseases Department, Tel Aviv Sourasky Medical Center and Tel Aviv University, Tel Aviv, Israel; ¹⁸Department of Neurology, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel; ¹⁹Laboratory of Clinical Microbiology, Virology and Bioemergencies, Luigi Sacco Hospital, Milan, Italy; ²⁰Department of Molecular Medicine, University of Padua, Padua, Italy; ²¹Microbiology and Virology Unit, Padua University Hospital, Padua, Italy; ²²Pediatric Neurology Department, Necker Hospital for Sick Children, Paris, France; ²³Pediatric Hematology-Oncology Unit, San Matteo Research Hospital (Fondazione IRCCS Policlinico San Matteo), Pavia, Italy; ²⁴Precision Medicine Unit, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland; ²⁵Department of Biomedical and Clinical Sciences (DIBIC), University of Milan, Milan, Italy; ²⁶Microbiology Unit, University Hospital of Bologna, Bologna, Italy; ²⁷Section of Microbiology, Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy; ²⁸DG Welfare, Lombardy, Italy; ²⁹Pediatric Immunohematology Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy; ³⁰San Raffaele Telethon Institute for Gene Therapy (SR-TIGET), IRCCS San Raffaele Scientific Institute, Milan, Italy; ³¹Vita-Salute San Raffaele University, Milan, Italy; ³²St. Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, The Rockefeller University, New York, NY, USA; ³³Pediatric Hematology-Immunology and Rheumatology Unit, Necker Hospital for Sick Children, Assistance Publique-Hôpitaux de Paris (AP-HP), Paris, France; ³⁴Laboratory of Virology and Infectious Disease, The Rockefeller University, New York, NY, USA; ³⁵Pathogenesis and Control of Chronic and Emerging Infections, University of Montpellier, INSERM, EFS, Montpellier, France; ³⁶National Reference Laboratory for Viral Zoonoses, National Public Health Center, Budapest, Hungary; ³⁷Department of Pediatrics, Section of Pediatric Tropical Medicine, Center for Human Immunobiology, Baylor College of Medicine and Texas Children’s Hospital, Houston, TX, USA; ³⁸Howard Hughes Medical Institute, New York, NY, USA; ³⁹Department of Pediatrics, Necker Hospital for Sick Children, Paris, France; ⁴⁰Neonatal and Pediatric Intensive Care Unit, Mother, Child and Adolescent Department, Geneva University Hospitals, Geneva, Switzerland; ⁴¹Department of Pediatrics, Gynecology and Obstetrics, University of Geneva, Geneva, Switzerland.

*A. Gervais, F. Trespidi, A. Ferrari, and F. Rovida contributed equally to this paper; **A. Marchal, S. Croce, and I. Cassaniti contributed equally to this paper; ***P. Bastard, M.R. MacDonald, C.M. Rice, A. Puel, L. Abel, G. Rossini, D. Mileto, Y. Simonin, A. Nagy, D. Hagin, K.O. Murray, F. Baldanti, and J.H. Aberle contributed equally to this paper; ****A. Cobat, S.-Y. Zhang, J.-L. Casanova, and A. Borghesi contributed equally to this paper. Correspondence to Alessandro Borghesi: a.borghesi@smatteo.pv.it; Jean-Laurent Casanova: casanova@rockefeller.edu.

Introduction

Mosquito-borne West Nile virus (WNV) is an emerging orthoflavivirus of considerable public health concern due to its dissemination and its role as a major etiology of epidemic encephalitis worldwide (1, 2). In recent years, a wide geographic circulation of WNV across all continents has been observed (3, 4, 5, 6). In 2023, WNV was detected for the first time in 22 new areas of the European Union (EU), corresponding to a 31% increase in the number of affected regions relative to 2022 (1, 7). The 2024 transmission season was marked by the broadest ever geographic distribution of WNV in the EU within a single year, with 212 affected regions across 19 countries, versus only 137 regions in 2023 and 173 regions in 2018 (8). Between June and October 2024, the largest outbreak in more than two decades occurred in Israel, with over 930 reported cases and 73 deaths (9). WNV infections have an enormous impact on healthcare systems, with >90% of reported cases resulting in hospitalization, neurological manifestations occurring in more than two-thirds of cases, and a case fatality rate of ~10% (up to 20% for encephalitis) (8). Tens of thousands of hospitalizations due to WNV infection and thousands of associated deaths have been reported over the last decade in the EU and neighboring countries, in the United States of America (USA), and elsewhere around the world (10, 11, 12). Seroepidemiological studies have shown that reported cases account for only a small proportion of the WNV infections. Indeed, progression to life-threatening WNV disease (WNVD) requiring hospitalization, particularly encephalitis, occurs in <1% of individuals infected with WNV (13). More than 99% of infections are silent or associated with flu-like, self-limiting disease (WNV fever, WNVF) (5, 14, 15, 16, 17, 18).

The root cause of life-threatening WNVD has long remained elusive (19). We recently showed, in six cohorts from the EU and USA, that blood and CSF autoantibodies (auto-Abs) neutralizing IFN- α 2, IFN- β , and/or IFN- ω underlie life-threatening WNVD in ~35% of hospitalized cases (range: 15–50%) (20). This proportion was higher for patients with WNV neuroinvasive disease, ~40% of whom (range: 15–55%) were found to have auto-Abs. auto-Abs neutralizing type I IFNs block the protective antiviral functions of IFN- α 2 and/or IFN- β and/or IFN- ω in Vero E6 cells or ARPE-19 cells infected with WNV *in vitro* (21) and in mice *in vivo* (22). They are present in the general population, with a prevalence of <1% in the under-65s and >4% in the over-70s (23). We have shown that they are not induced by WNV infection and that they precede the appearance of WNV-specific IgM and/or IgG and confer an ~20- to ~600-fold increase in the risk of life-threatening WNVD and neuroinvasive disease, depending on the number and concentration of type I IFNs neutralized (20). These findings, recently replicated in case reports and small case series (24, 25, 26), suggest that auto-Abs neutralizing type I IFNs are strong determinants of WNVD. Moreover, these auto-Abs have been found to underlie tick-borne encephalitis (TBE) in 10% of the most severe cases (27), the most severe cases of the rarer Powassan virus, Usutu virus, and Ross River virus diseases (28), and ~35 and 60% of cases of life-threatening adverse reactions to yellow fever live-attenuated vaccine-17D strain (YFV-17D) (29) and chikungunya virus (CHIKV) live-attenuated

vaccine (VLA1553), respectively (30). Thus, auto-Abs neutralizing type I IFNs are emerging as strong, common, global determinants of a growing range of arboviral diseases, whether caused by orthoflaviviruses or alphaviruses (31). We investigated seven new cohorts of individuals infected with WNV, including six recently recruited cohorts established during the seasonal outbreaks in the summers of 2023 and 2024.

Results

Seven new cohorts of individuals with WNV infection

We studied 225 subjects hospitalized for life-threatening WNVD from seven new cohorts enrolled at three centers in Italy during 2023—Pavia (46 subjects), Milan (13 subjects), and Bologna (11 subjects)—one center in Austria (31 subjects, years 2015–2024), one center in Hungary (68 subjects, during 2024), one center in Israel (32 subjects, during 2024), and one center in Texas, USA (24 subjects, years 2002–2008). Neuroinvasive disease was confirmed in 200/225 (89%) patients—encephalitis (154 cases), meningitis (32 cases), acute flaccid paralysis (7 cases), and unspecified neurological disease (7 cases)—whereas there was no reported clinical evidence of neuroinvasive disease in the remaining 25/225 (11%) hospitalized patients. We also enrolled 61 patients with self-limiting, febrile, and/or flu-like illness (WNVF), managed as outpatients in Italy (11 subjects), Austria (27 subjects), Hungary (6 subjects), Israel (2 subjects), and the USA (15 subjects), and 32 individuals with recent asymptomatic or paucisymptomatic WNV infection (WNV-infected controls, WNVIC) diagnosed on the basis of the detection of WNV RNA in a nucleic acid amplification test on blood performed at the time of blood donation (Fig. 1 A; and Fig. S1, A and B). For all the individuals enrolled, WNV infection was demonstrated by the presence of WNV-specific IgM or seroconversion to IgG, WNV neutralization assays (32), and/or RT-PCR on serum, plasma, or cerebrospinal fluid samples.

Demographics of individuals with WNV infection

The mean age (standard deviation, SD) of patients with WNVD was 68 years (15 years) (range: 10–99 years), higher than that for patients with WNVF (51 years [14], range: 19–86 years). In the WNVIC group, mean age (SD) was 54 years (11) and age ranged from 32 to 78 years (Fig. S1 C). The mean age (SD) of patients with WNVD was higher in the three cohorts enrolled in Italy (Bologna, 77 years [10]; Pavia, 70 years [13]; and Milan, 69 years [11]; overall: 71 years [13]), than that in the cohorts enrolled in Austria (65 years [13], $P = 0.04$), Hungary (63 years [16], $P = 8 \times 10^{-4}$), and the USA (62 years [14], $P = 7 \times 10^{-3}$), but lower than that in the cohort enrolled in Israel (77 years [11], $P = 0.02$) (Fig. S1 D). The proportion of male subjects was 65% (147/225) of all patients with WNVD (130/200 [65%] cases of neuroinvasive disease and 102/154 [66%] cases of encephalitis), 54% (33/61) of patients with WNVF, and 84% (27/32) of those with WNVIC (Fig. S1, E and F). Mortality was assessed from vital status data, which were available for 150/200 (75%) patients with neuroinvasive disease. Mortality was 15% (23/150) for the total population of patients with neuroinvasive disease, 17% (19/114) for those with encephalitis, 33% (2/6) for those with acute flaccid paralysis, and




Figure 1. **Auto-Abs neutralizing type I IFNs in 318 WNV-infected individuals from seven new cohorts. (A)** Age and sex distribution of individuals in the WNVIC, WNVF, and WNV groups. **(B)** Detection of auto-Abs against IFN- α 2, IFN- β , and IFN- ω by ELISA. An OD value >0.5 (dashed line) indicates a sample considered positive based on the signal typically observed for serum/plasma from healthy donors. **(C)** Luciferase-based neutralization assay for the detection of auto-Abs neutralizing 10 ng/ml IFN- α 2, IFN- ω , or IFN- β . Neutralizing activity was defined as a luciferase signal below 15%. **(D)** Luciferase-based neutralization assay for detecting auto-Abs neutralizing 100 pg/ml IFN- α 2 or IFN- ω , or 1 ng/ml IFN- β . Neutralizing activity was defined as a luciferase signal below 15%.

7% (2/27) for those with meningitis; none of the three patients with unspecified neurological disease died, and no deaths were reported among patients hospitalized without evidence of neuroinvasive disease (Table 1).

Auto-Abs neutralizing type I IFNs are rare in asymptomatic WNVIC

We assessed the prevalence of auto-Abs neutralizing type I IFNs in the three groups of individuals with WNV infection (WNVIC,

WNVF, and WNV) from the seven cohorts. We used an enzyme-linked immunosorbent assay (ELISA) to screen serum or plasma samples from all 318 individuals enrolled in this study for circulating IgG auto-Abs against IFN- α 2, IFN- β , or IFN- ω , with positive results defined as an optical density (OD) >0.5 (33, 34). We then assessed the neutralizing activity of these auto-Abs in a well-established luciferase-based assay (20, 23). Blood samples were taken a mean of 18 days (range: 0–341 days) after clinical disease onset or during virological screening before

Table 1. Demographic and clinical characteristics of the study population in the three WNV clinical groups and the two subgroups of WNVD

Characteristic	WNVIC n = 32 ^a	WNVF n = 61 ^a	WNVD n = 225 ^a	Subgroups of WNVD	
				Non-neuroinvasive WNVD n = 25 ^a	Neuroinvasive WNVD n = 200 ^a
Age (years)	54 [11]	51 [14]	68 [15]	68 [12]	68 [15]
Sex					
F	5/32 (16%)	27/61 (44%)	75/225 (33%)	8/25 (32%)	67/200 (34%)
M	27/32 (84%)	33/61 (54%)	147/225 (65%)	17/25 (68%)	130/200 (65%)
Unknown	0/32 (0%)	1/61 (2%)	3/225 (1%)	0/25 (0%)	3/200 (1%)
Mortality	0/32 (0%)	0/61 (0%)	23/225 (10%)	0/25 (0%)	23/200 (12%)
Unknown	0/32 (0%)	0/62 (0%)	57/225 (25%)	7/25 (28%)	50/200 (25%)
Recruitment center					
Italy (PV)	10/32 (31%)	5/61 (8%)	46/225 (20%)	13/25 (52%)	33/200 (17%)
Italy (MI)	2/32 (6%)	6/61 (10%)	13/225 (6%)	1/25 (4%)	12/200 (6%)
Italy (BO)	0/32 (0%)	0/61 (0%)	11/225 (5%)	1/25 (4%)	10/200 (5%)
USA	0/32 (0%)	15/61 (25%)	24/225 (11%)	0/25 (0%)	24/200 (12%)
Austria	17/32 (53%)	27/61 (44%)	31/225 (14%)	5/25 (20%)	26/200 (13%)
Israel	3/32 (9%)	2/61 (3%)	32/225 (14%)	4/25 (16%)	28/200 (14%)
Hungary	0/32 (0%)	6/61 (10%)	68/225 (30%)	1/25 (4%)	67/200 (34%)
Year of recruitment					
Before 2018	5/32 (16%)	10/61 (16%)	27/225 (12%)	0/25 (0%)	27/200 (14%)
2018–2022	7/32 (22%)	14/61 (23%)	11/225 (5%)	3/25 (12%)	8/200 (4%)
2023–2024	20/32 (63%)	28/61 (46%)	187/225 (83%)	22/25 (88%)	165/200 (83%)
Unknown	0/32 (0%)	9/76 (12%)	0/225 (0%)	0/25 (0%)	0/200 (0%)

^aMean [SD] or counts (frequency, %). PV, Pavia; MI, Milan; BO, Bologna.

blood donation for WNVIC. In the WNVIC group, OD values were low for auto-Abs against the three type I IFNs tested in most samples, below the 0.5 OD-unit detection threshold in 25/32 (78%) samples, and slightly above the 0.5 OD-unit detection threshold for at least one type I IFN in 4/32 samples, whereas OD values were very high (>1.0) for auto-Abs against at least one type I IFN in only 3/32 samples (Fig. 1 B). We then tested 1:10 dilutions of the same serum or plasma samples for the neutralization of high (10 ng/ml) or low (100 pg/ml) concentrations of unglycosylated IFN- α 2 and IFN- ω , and high (10 ng/ml) or intermediate (1 ng/ml) concentrations of glycosylated IFN- β . We found no auto-Abs neutralizing high concentrations of IFN- α 2 and/or IFN- β and/or IFN- ω or intermediate concentrations of IFN- β in asymptomatic WNVIC, including samples with high OD values (>1.0) on ELISA. At the more physiological concentration of 100 pg/ml, only 1/32 (3%) individuals carried auto-Abs neutralizing IFN- ω only, corresponding to a sample with an OD <0.5 for auto-Abs against IFN- ω on ELISA (Fig. 1, C and D; Fig. 2, A–D; and Table 2). The correlation between ELISA and neutralization results is illustrated in Fig. S2, A–C. Auto-Abs neutralizing IFN- α or IFN- ω have recently been shown to block the interaction of the corresponding type I IFNs with both type I IFN receptor subunits (IFNAR1/2), whereas high-titer non-neutralizing auto-Abs limit only the interaction of the corresponding type I IFN with a single receptor subunit

(either IFNAR1 or IFNAR2), with the interaction between the IgG and the bound type I IFN being of low avidity (35). This difference may explain the apparent discrepancies between the results obtained by ELISA and those obtained in the neutralization assay in a small number of samples from our cohort.

Auto-Abs neutralizing type I IFNs are rare in patients with WNVF

Similarly, in the WNVF group, OD values were low for auto-Abs against the three type I IFNs tested in most samples, below the 0.5 OD detection threshold in 56/61 (92%) samples and slightly above the 0.5 OD detection threshold for at least one type I IFN in 2/61 (3%) samples, whereas OD values >1.0 for auto-Abs against at least one type I IFN were obtained for 3/61 (5%) samples (Fig. 1 B). In this group, we detected no auto-Abs neutralizing high concentrations of IFN- α 2 and/or IFN- β and/or IFN- ω and/or intermediate concentrations of IFN- β in samples from WNVF patients. We found auto-Abs neutralizing low concentrations of IFN- ω in only 1/61 (2%) samples. The sample testing positive had an OD <0.5 for anti-IFN- ω IgG auto-Abs on ELISA (Fig. 1, C and D; Fig. 2, A–D; Fig. S2, A–C; and Table 2). Thus, the prevalence of auto-Abs neutralizing type I IFNs was similar in the WNVF and WNVIC groups (1/61 [2%] vs. 1/32 [3%], respectively [P = ns]) and overall low (2/93 [2%]) when these two groups were analyzed




Figure 2. **Proportions of individuals with auto-Abs neutralizing type I IFNs in seven new cohorts. (A and B)** Frequency of individuals carrying auto-Abs neutralizing type I IFNs at concentrations of 10 ng/ml (A) or 100 pg/ml (B), as determined in luciferase-based neutralization assays, in the three defined groups

of WNV-infected individuals: WNVIC, WNVF, and WNVD. IFN- α , auto-Abs neutralizing IFN- $\alpha 2$ (regardless of their effects on other IFNs); IFN- ω , auto-Abs neutralizing IFN- ω (regardless of their effects on other IFNs); IFN- $\alpha \pm \omega \pm \beta$, auto-Abs neutralizing IFN- $\alpha 2$ and/or IFN- ω and/or IFN- β ; IFN- $\alpha + \omega$, auto-Abs neutralizing both IFN- $\alpha 2$ and IFN- ω . **(C)** Number of type I IFNs neutralized in the three groups of WNV-infected individuals (WNVIC, WNVF, and WNVD), as determined with the luciferase-based neutralization assay. **(D)** Proportion of type I IFNs neutralized in the three groups of WNV-infected individuals (WNVIC, WNVF, and WNVD) according to the nature and combination of auto-Abs.

together. Importantly, these two groups of individuals carried auto-Abs neutralizing low concentrations of IFN- ω only, suggesting that auto-Abs neutralizing other type I IFNs—IFN- $\alpha 2$ in particular—or more than one type I IFN could be even rarer in groups of WNV-infected individuals selected on the basis of their resistance to severe disease than in the general population (23). These observations also highlight the importance of assessing type I IFN-neutralizing activity (23, 36, 37) in addition to performing assays detecting the binding of type I IFNs to auto-Abs (e.g., Gyros, ELISA, and VIDAS®) (33, 38, 39), when testing samples from individual patients and disease cohorts.

Table 2. Type I IFN-neutralizing auto-Abs in the study population

Characteristic	auto-Ab ⁺ , n = 82 ^a	No auto-Ab, n = 236 ^a
Age (years)	69.52 [15.27]	60.88 [15.32]
≤40	3/29 (10%)	26/29 (90%)
41–65	20/137 (15%)	117/137 (85%)
>65	57/148 (39%)	91/148 (61%)
Unknown	0/4 (0%)	4/4 (100%)
Sex		
F	19/107 (18%)	88/107 (82%)
M	62/207 (30%)	145/207 (70%)
Unknown	1/4 (25%)	3/4 (75%)
WNV groups		
WNVD	81/225 (36%)	144/225 (64%)
WNVF	1/61 (2%)	60/61 (98%)
WNVIC	1/32 (3%)	31/32 (97%)
Subgroups of WNVD		
Non-neuroinvasive WNVD	9/25 (36%)	16/25 (64%)
Neuroinvasive WNVD		
-All	71/200 (36%)	129/200 (64%)
-WNE	59/154 (38%)	95/154 (62%)
-WNM	8/32 (25%)	24/32 (75%)
-AFP	2/7 (29%)	5/7 (71%)
-UNS	2/7 (29%)	5/7 (71%)

^aMean [SD], counts, or frequency (%). WNE, WNV encephalitis; WNM, WNV meningitis; AFP, acute flaccid paralysis; UNS, unspecified neurological syndrome.

The round and square brackets indicate that individuals with ages at the extremes of the range are excluded (40 year-old individuals) or included (65-year-old individuals) in this age group.

Auto-Abs neutralizing IFN- $\alpha 2$, β , or ω in ~35% of patients hospitalized for WNVD from seven new cohorts

In samples from the group of patients with severe WNVD, we found OD values above the 0.5 OD-unit detection threshold for IgG auto-Abs against IFN- $\alpha 2$ and/or IFN- β and/or IFN- ω for 73/225 (32%) samples (Fig. 1 B; and Fig. S3, A and B). We found auto-Abs neutralizing high concentrations of at least one type I IFN in 66/225 (29%) WNVD patients ($P = 8.45 \times 10^{-4}$ and $P = 3.29 \times 10^{-6}$ vs. WNVIC and WNVF, respectively), with 57/225 (25%) neutralizing IFN- $\alpha 2$, 52/225 (23%) neutralizing IFN- ω , and 10/225 (4%) neutralizing IFN- β (Fig. 1 C, Fig. 2, A–D, and Fig. S4, A–D). Specifically, 9/225 (4%) samples neutralized high concentrations of IFN- $\alpha 2$ only, 5/225 (2%) high concentrations of IFN- β only, 4/225 (2%) high concentrations of IFN- ω only, 42/225 (19%) high concentrations of IFN- $\alpha 2$ and IFN- ω , and 6/225 (3%) high concentrations of the three type I IFNs tested; none of the samples neutralized a combination of IFN- $\alpha 2$ and IFN- β only or a combination of IFN- β and IFN- ω only (Fig. 2 D and Table 2). Samples with high OD values usually displayed neutralizing activity (Fig. S2, A–C and Table 2). At more physiological concentrations, 81/225 (36%) samples from patients hospitalized for life-threatening WNVD neutralized at least 100 pg/ml IFN- $\alpha 2$ and/or IFN- ω and/or 1 ng/ml IFN- β , 66/225 (29%) samples neutralized 100 pg/ml of IFN- $\alpha 2$, 65/225 (29%) neutralized 100 pg/ml of IFN- ω , and 19/225 (8%) neutralized 1 ng/ml of IFN- β , with 12/225 (5%) samples neutralizing 100 pg/ml of IFN- $\alpha 2$ only, 11/225 samples (4%) neutralizing 100 pg/ml IFN- ω only, 2/225 (<1%) neutralizing 1 ng/ml IFN- β only, 39/225 (17%) neutralizing 100 pg/ml of both IFN- $\alpha 2$ and IFN- ω , 1/225 (<1%) neutralizing both 100 pg/ml IFN- $\alpha 2$ and 1 ng/ml IFN- β , 2/225 (1%) neutralizing both 100 pg/ml IFN- ω and 1 ng/ml IFN- β , and 14/225 (7%) neutralizing 100 pg/ml IFN- $\alpha 2$ and IFN- ω and 1 ng/ml IFN- β (Fig. 1 D, Fig. 2, B–D, and Fig. S4, A–D). In total, 17/225 (8%) samples neutralized 100 pg/ml IFN- $\alpha 2$ and/or IFN- ω and/or 1 ng/ml IFN- β but did not neutralize high concentrations of any of the three type I IFNs tested.

Auto-Abs neutralizing IFN- $\alpha 2$, β , or ω in ~40% of patients with WNV neuroinvasive disease

Among WNVD patients, the prevalence of auto-Abs was higher in patients with WNV neuroinvasive disease (WNV encephalitis, meningitis, or acute flaccid paralysis), with 64/200 (32%) having an OD >0.5 on ELISA, and 58/200 (29%) displaying the neutralization of high concentrations of at least one of the three type I IFNs tested ($P = 9.75 \times 10^{-4}$ and $P = 4.37 \times 10^{-6}$ vs. WNVIC and WNVF, respectively). In this subgroup of patients, samples from 71/200 (36%) patients contained circulating auto-Abs neutralizing at least 100 pg/ml IFN- $\alpha 2$ and/or IFN- ω and/or 1 ng/ml IFN- β . The prevalence of individuals carrying auto-Abs neutralizing type I IFNs was highest in the subgroup of patients

suffering from encephalitis—the most severe neuroinvasive manifestation of WNV infection—with 52/154 (34%) samples displaying an OD >0.5 on ELISA. In this subgroup of patients, samples from 49/154 (32%) patients neutralized high concentrations of at least one of the three type I IFNs tested. In tests of the neutralization of more physiological concentrations, samples from 59/154 (38%) patients contained auto-Abs neutralizing at least 100 pg/ml IFN- α 2 and/or IFN- ω and/or 1 ng/ml IFN- β , a prevalence higher than that in patients with acute flaccid paralysis (2/7 [29%]), meningitis (8/32 [25%]), or unspecified neurological disease (2/7 [29%]), and hospitalized patients with no documented neuroinvasive disease (9/25 [36%]). Finally, for patients with available mortality data, auto-Abs neutralizing 100 pg/ml IFN- α 2 and/or IFN- ω and/or 1 ng/ml of IFN- β were detected in 43/127 (34%) patients with neuroinvasive disease (36/95 [38%] with encephalitis) who survived, and in 10/23 (44%) patients with neuroinvasive disease (8/19 [42%] with encephalitis) who died from WNV infection. Overall, consistent with our previous report (20), we found auto-Abs neutralizing IFN- α 2 and/or IFN- β and/or IFN- ω in almost 40% of patients with WNV neuroinvasive disease, encephalitis in particular, from seven new cohorts enrolled in five countries, including two new geographic areas.

Auto-Abs neutralizing IFN- α 2, - β , or - ω in 981 individuals with WNV infection from nine centers in five countries

We then performed a combined analysis on the six previously reported cohorts (20) and the seven new cohorts consisting of 981 individuals with WNV infection recruited in nine centers in five countries between 2002 and 2024 (Fig. 3 A). The infecting WNV lineage was determined for 380/981 (39%) individuals (Fig. 3, A and B). There were 666 patients with WNVD (including 548 suffering from neuroinvasive disease, 376 of whom had documented WNV encephalitis), 169 patients managed as outpatients for WNVF, and 146 WNVIC (Fig. S5, A and B). Mean age and sex distribution by disease group, by country, and by recruitment center are shown in Fig. S5, C–F. The overall prevalence of auto-Abs neutralizing at least one type I IFN was 236/666 (35%) in WNVD patients. The highest prevalence was found in WNVD patients from Israel (13/32 [40%]) and Italy (165/423 [39%]), including 7/13 [54%] in Milan, 54/116 [47%] in Pavia, 17/40 [43%] in Turin, 25/58 [38%] in Bologna, 65/196 [33%] in Padua, followed by Austria (11/31 [35%]), Hungary (32/99 [32%]), and the USA (14/81 [17%]) (Fig. 3, C and D). The overall prevalence of auto-Abs neutralizing at least one type I IFN was 206/548 (38%) in patients with WNV neuroinvasive disease (135/376 [36%] in WNV encephalitis patients) and was highest in Israel (12/28 [43%]) and Italy (141/321 [44%]), including 6/12 [50%] in Milan, 47/94 [50%] in Pavia, 16/38 [42%] in Turin, 22/52 [42%] in Bologna, and 50/125 [40%] in Padua, followed by Austria (10/26 [38%]), Hungary (29/92 [32%]), and the USA (14/81 [17%]). The prevalence of auto-Abs neutralizing at least one type I IFN was 16/84 (19%) in WNVD patients recruited between 2002 and 2017, 67/165 (41%) in those recruited in 2018, 5/12 (42%) in 2019, 1/1 (100%) in 2021, 78/217 (36%) in 2022, 34/71 (48%) in 2023, and 35/116 (30%) in 2024 (Fig. 3, E and F). The proportion of individuals with WNVF from the new cohorts carrying auto-Abs

was similar to that in the general population (1/61 [2%]) and lower than that in WNVF patients from the previously reported cohorts (15/108 [14%]) (20). In the WNVF patients of the previously reported cohorts, substantial differences were observed between recruitment centers, possibly reflecting differences in the clinical criteria used to define WNVF and disease severity between centers and across years of recruitment or misclassification due to missing follow-up data and the unreported clinical worsening and hospitalization of some patients initially tested for WNVF.

Auto-Abs neutralizing IFN- α 2, - β , or - ω by infecting WNV lineage

The infecting WNV lineage was determined for 66/146 (45%) WNVIC, 84/169 (50%) patients with WNVF, and 230/666 (35%) WNVD patients (Fig. 3, A and B). The proportions of individuals infected with WNV-1 or WNV-2 were comparable in the three WNV infection groups. WNV-1 was detected in 16/66 (24%) WNVIC, in 21/84 (25%) patients with WNVF, and in 69/230 (30%) of the samples obtained from WNVD patients during the 2022 summer outbreak in Padua (Fig. 3, A and B). WNV-2 was detected in 50/66 (76%) WNVIC, 63/84 (75%) patients with WNVF, and in 161/230 (70%) samples from WNVD patients from Austria, Hungary, Bologna, and Padua obtained in 2018, 2019, 2021, 2022, 2023, and 2024 (Fig. 3, A and B). We found auto-Abs neutralizing low concentrations of IFN- α 2 and/or IFN- ω and/or intermediate concentrations of IFN- β in 23/69 (33%) WNVD patients infected with the WNV-1 lineage, including 19/56 (34%) patients with WNV neuroinvasive disease (14/35 [40%] with meningitis, 1/2 [50%] with acute flaccid paralysis, and 4/19 [21%] with encephalitis), and in 58/161 (36%) WNVD patients infected with the WNV-2 lineage, including 48/104 (46%) patients with WNV neuroinvasive disease (13/31 [42%] with meningitis, none—0/1—with acute flaccid paralysis, 6/19 [32%] with unspecified neurological syndrome, and 29/53 [55%] with encephalitis) (Fig. 3, A and B; and Table S1). Thus, we found no substantial difference in the prevalence of auto-Abs between WNVD patients infected with WNV-1 and WNVD patients infected with WNV-2 viruses (23/69 [33%] vs. 58/161 [37%], $P = 0.81$). The higher prevalence of individuals carrying auto-Abs among those with WNV neuroinvasive disease due to WNV-2 than among those with WNV neuroinvasive disease due to WNV-1 viruses, particularly for patients with encephalitis (29/53 [55%] vs. 4/19 [21%], $P = 0.02$), requires confirmation in future studies and may reflect sample selection at diagnosis for virological studies or, alternatively, differences in the neurotropism of the viral lineages in individuals with impaired type I IFN immunity in the blood or at the blood-brain barrier due to circulating auto-Abs.

Higher prevalence of auto-Abs in male individuals

We previously showed that age and sex influence the prevalence of auto-Abs neutralizing type I IFNs. We therefore analyzed the effects of both variables on the prevalence of auto-Abs neutralizing low concentrations of IFN- α 2 and/or IFN- ω and/or intermediate concentrations of IFN- β in each disease category. For 2/225 WNVD patients, sex and age were not reported; for 2/225




Figure 3. Numbers of patients with auto-Abs neutralizing type I IFNs by center, year of enrollment, and lineage in 13 cohorts from five countries. (A) Distribution by year (x axis), recruitment center (colors), and infecting WNV virus lineage (WNV-1, dashed boxes; WNV-2, dotted boxes) of the individuals with WNVD recruited. Of the 666 patients with WNVD, 423 were recruited in Italy: 116 in Pavia (20 in 2018, 50 in 2022, and 46 in 2023), 196 in Padua (89 in 2018 and 107 in 2022), 58 in Bologna (47 in 2022 and 11 in 2023), 40 in Turin (29 in 2018 and 11 in 2022), and 13 in Milan (13 in 2023); 99 patients were recruited in Budapest, Hungary (20 in 2018, 11 in 2019, and 68 in 2024); 32 were recruited in Tel Aviv, Israel, in 2024; 31 were recruited in Vienna, Austria, between 2015 and 2024; and 81 were recruited in Houston, Texas, USA, between 2002 and 2010. In total, 84/666 (13%) patients were recruited between 2002 and 2017, 165/666 (25%) were recruited in 2018, 12/666 (18%) were in 2019, 1/666 was recruited in 2021, 217/666 (33%) were recruited in 2022, 71/666 (11%) were recruited in 2023, and 116/666 (17%) were recruited in 2024. (B) Number of individuals with auto-Abs neutralizing at least one type I IFN at a concentration of 100 pg/ml (IFN- α 2 and IFN- ω) or 1 ng/ml (IFN- β) in subgroups of WNVD patients: WNE, WNV encephalitis; WNM, WNV meningitis; AFP, acute flaccid paralysis; UNS, unspecified neurological syndrome (subsets of WNND group). (C and D) Number and prevalence of individuals with auto-Abs neutralizing at least one type I IFN at a concentration of 100 pg/ml (IFN- α 2 and IFN- ω) or 1 ng/ml (IFN- β) in the three groups of individuals infected with WNV (WNVIC, WNVF, and WNVD), by enrollment center. (E and F) Number and prevalence of individuals with auto-Abs neutralizing at least one type I IFN at a concentration of 100 pg/ml (IFN- α 2 and IFN- ω) or 1 ng/ml (IFN- β) in the three groups of individuals infected with WNV (WNVIC, WNVF, and WNVD), by enrollment year. WNND, WNV neuroinvasive disease, a subset of WNVD; WNE, WNV encephalitis; WNM, WNV meningitis; AFP, acute flaccid paralysis; UNS, unspecified neurological syndrome (subsets of WNND group).




Figure 4. **Proportions of patients with auto-Abs neutralizing type I IFNs by sex and age in 13 cohorts from five countries.** (A) Prevalence of individuals with auto-Abs neutralizing at least one type I IFN at a concentration of 100 pg/ml (IFN- α 2 and IFN- ω) or 1 ng/ml (IFN- β) in the three groups of individuals infected with WNV (WNVIC, WNVF, and WNVD), by sex. (B) Prevalence of individuals with auto-Abs neutralizing at least one type I IFN at a concentration of 100 pg/ml (IFN- α 2 and IFN- ω) or 1 ng/ml (IFN- β) in the three groups of WNV-infected individuals (WNVIC, WNVF, and WNVD), by age class.

additional WNVD patients, sex was reported but not age; and for 1/61 WNVF patients, age was reported but not sex. In the WNVD group, the prevalence of auto-Abs was higher in male subjects than in female subjects in the new cohorts (60/147 [41%] vs. 19/76 [25%], odds ratio [OR] = 2.07, 95% CI: 1.13–3.89, $P = 0.02$), and in the total cohort of 664/981 patients with WNVD and for whom sex data were available (172/433 [40%] vs. 63/231 [27%], OR = 1.75, 95% CI: 1.24–2.48, $P = 1.7 \times 10^{-3}$). A similar trend was observed in the neuroinvasive disease group in the new cohorts (53/130 [41%] male patients vs. 17/68 [25%] female patients, OR = 2.06, 95% CI: 1.09–4.04, $P = 0.03$) and in the total cohort of 546/981 patients with neuroinvasive disease and available sex data (149/362 [41%] male patients vs. 56/184 [30%] female patients, OR = 1.59, 95% CI: 1.09–2.33, $P = 0.02$). By contrast, no difference in the prevalence of the auto-Abs between the sexes was observed in WNVIC (1/27 [4%] male patients vs. 0/5 female patients, $P = 1.0$) or in the WNVF group (1/33 [3%] male patients vs. 0/27 female patients, $P = 1.0$) in the seven new cohorts, and in WNVIC (3/114 [3%] male patients vs. 0/32 female patients, $P = 1$) or the WNVF group (11/88 [13%] male patients vs. 5/80 [6%] female patients, $P = 0.17$) in the total cohort (Fig. 4 A and Table S2).

Higher prevalence of auto-Abs in individuals over the age of 65 years

The individuals with neutralizing auto-Abs were significantly older than those without them in the seven new cohorts (mean age [SD], 70 [15] vs. 61 [15] years; $P = 2.43 \times 10^{-5}$) and in the total cohort of 977/981 individuals for whom age data were available (mean age [SD], 71 [15] vs. 60 [17] years; $P = 2.18 \times 10^{-19}$). This difference was driven primarily by the older age of the individuals with auto-Abs in the WNVD group (mean age [SD], 72 [14] vs. 65 [16] years; $P = 1.9 \times 10^{-7}$ in the group of 662 WNVD patients with available age data). By contrast, mean age did not differ significantly between individuals with and without auto-

Abs in the group of 168 individuals with WNVF and available age data (mean age [SD], 57 [18] vs. 54 [18] years; $P = 0.41$). Stratification by age group (≤ 40 , 41–65, and > 65 years) in the new cohorts confirmed our previous observations [19], with auto-Abs neutralizing low concentrations of IFN- α 2 and/or IFN- ω and/or intermediate concentrations of IFN- β being more frequent in patients with severe disease, particularly older individuals. In individuals aged ≤ 40 years, no auto-Abs were detected in the WNVIC group, whereas auto-Abs were present in 3/45 (7%) WNVF patients and 8/42 (19%) WNVD patients. The prevalence of auto-Abs increased with age. In the 41–65 years age group, auto-Abs were detected in 3/105 (3%) WNVIC, 8/83 (10%) WNVF patients, and 45/203 (22%) WNVD patients. The prevalence of the auto-Abs increased sharply after the age of 65 years, with auto-Abs detected in 0/14 WNVIC, 5/41 (12%) WNVF, and 181/417 (43%) WNVD cases (Fig. 4 B). Thus, the risk of carrying auto-Abs increased with age and was higher in subjects > 65 years old than in subjects ≤ 65 years old in the WNVD group (181/417 [43%] vs. 53/245 [22%], OR = 2.78, 95% CI: 1.95–4.01; $P = 2.77 \times 10^{-8}$), in patients with neuroinvasive disease (162/362 [45%], vs. 42/182 [23%] OR = 2.7, 95% CI: 1.82–4.07; $P = 1.30 \times 10^{-6}$), and in patients with encephalitis (108/262 [41%] vs. 25/111 [23%], OR = 2.78, 95% CI: 1.95–4.01; $P = 2.77 \times 10^{-8}$). The prevalence of these auto-Abs was highest in males > 65 years of age with encephalitis (81/173 [47%]).

Auto-Abs neutralizing type I IFNs are as rare in WNVIC and patients with WNVF as in the general population

For estimation of the risk of clinical disease conferred by the presence and nature of auto-Abs neutralizing type I IFNs, we compared the proportions of subjects carrying various types and combinations of neutralizing auto-Abs with the proportions in individuals carrying the corresponding neutralizing auto-Abs among 34,159 healthy men and women aged 18–100 years from the French general population, after adjustment for age and sex

(23). We found no significant difference in the prevalence of neutralizing auto-Abs between WNVIC and the general population, regardless of the type or combination of auto-Abs considered, in both a separate analysis of the new cohorts ($n = 32$ WNVIC) and a combined analysis of these and the previously described cohorts ($n = 146$ WNVIC) (1/32 [3%] vs. 295/13,570 [2%], OR = 1.97, 95% CI: 0.39–9.94, $P = 0.46$; and 3/146 [2%] vs. 295/13,570 [2%], OR = 1.04, 95% CI: 0.36–3.01, $P = 0.95$, for auto-Abs neutralizing 100 pg/ml IFN- $\alpha 2$ and/or IFN- ω for the new cohorts and the total cohort, respectively), consistent with the absence of clinical disease despite documented WNV infection in this group (Fig. 3, C and D; and Fig. 5). Similarly, we found no significant difference in the prevalence of neutralizing auto-Abs between individuals with WNVF in the new cohorts ($n = 61$ WNVF) and individuals in the general population, regardless of the type or combination of auto-Abs considered (1/61 [2%] vs. 295/13,570 [2%], OR = 1.12, 95% CI: 0.22–5.63, $P = 0.89$ for auto-Abs neutralizing 100 pg/ml IFN- $\alpha 2$ and/or IFN- ω) (Fig. 5, Fig. 6 C, and Table 3). This finding contrasts with the higher prevalence of auto-Abs neutralizing various combinations of type I IFNs in patients with WNVF in our previous study (15/108 [14%]) (20), and therefore in the combined analysis of the new and previously reported cohorts ($n = 169$ WNVF), resulting in an ~5–20 times increase in the risk of WNVF depending on the type and nature of the neutralizing auto-Abs (16/169 [10%] vs. 295/13570 [2%], OR = 4.7, 95% CI: 2.8–7.9, $P = 1.37 \times 10^{-6}$ for auto-Abs neutralizing 100 pg/ml IFN- $\alpha 2$ and/or IFN- ω), (Fig. 5, Fig. 6 D, and Table 3).

Risk of WNVF conferred by the auto-Abs in the new and total cohorts

The criteria for assignment to the WNVF group were identical for this and the initial study, but for several of the newly recruited cases, we obtained access to longitudinal documentation, making it possible to distinguish accurately between true WNVF cases (that is, individuals never hospitalized during the course of WNV infection) and patients initially assigned to the WNVF group but subsequently displaying progression to more severe disease. A longitudinal analysis of the clinical data showed that 17/78 [22%] of the patients initially referred for WNVF were later hospitalized (and were therefore reassigned to the WNVD group), with clinical signs of neuroinvasive disease reported in 12/17 (71%) cases. Strikingly, samples from one of these 17 patients (6%) neutralized both high and low concentrations of IFN- $\alpha 2$, and samples from another patient (6%) neutralized both high and low concentrations of IFN- ω . This two-step course of the disease is reminiscent of that observed in severe SARS-CoV-2 infection, for which a pathogenetic model has been proposed (40). These observations also highlight the need for consistent definitions of WNVD categories, particularly for WNVF, as considerable heterogeneity is observed between studies (41, 42, 43), with the definition of WNVF ranging from self-reported, mild clinical disease (14) to severe clinical disease requiring hospitalization in individuals not undergoing lumbar puncture (44). Overall, the study of seven new cohorts of patients with WNV infection modulates our previous findings, suggesting that the risk of mild WNVF is increased little, if at all, by the presence

of auto-Abs neutralizing type I IFNs, at least under the experimental conditions used here to test for type I IFN-neutralizing activity in patient samples.

Risk of WNVD in individuals with auto-Abs neutralizing type I IFNs

The prevalence of auto-Abs, both overall and in various combinations, was significantly higher in WNVD patients than in the general population. The presence of auto-Abs neutralizing at least low concentrations of IFN- $\alpha 2$ and/or IFN- ω , whether considering or not neutralization of IFN- β , was associated with an ~15–20 times higher risk of WNVD in both the new cohorts and in the overall study population consisting of the new and previously studied cohorts analyzed together (Fig. 5; Fig. 6, C and D; and Table 3). A combination of auto-Abs neutralizing at least low concentrations of both IFN- $\alpha 2$ and IFN- ω was associated with a 40–50 times increase in the risk of WNVD, both in the new cohorts and in the overall study population (Fig. 5; Fig. 6, C and D; and Table 3). The risk of clinical disease was higher in individuals carrying auto-Abs able to neutralize high concentrations (10 ng/ml) of IFN- $\alpha 2$, IFN- β , or IFN- ω . The presence of auto-Abs neutralizing combinations of high concentrations of IFN- $\alpha 2$ and/or IFN- β and/or IFN- ω resulted in an ~30–80-fold increase in the risk of severe WNVD (Table 3). The risk was higher in individuals with auto-Abs neutralizing a combination of high concentrations (10 ng/ml) of both IFN- $\alpha 2$ and IFN- ω , with an ~100–120 times increase in the risk for auto-Abs neutralizing high concentrations of both IFN- $\alpha 2$ and IFN- ω regardless of the presence of auto-Abs neutralizing IFN- β in the new and total cohorts (Fig. 6, C and D; and Table 3). Individuals with a combination of auto-Abs neutralizing high concentrations (10 ng/ml) of IFN- $\alpha 2$, IFN- β , and IFN- ω had the highest risk of severe clinical disease, with an ~170-fold (overall cohorts) to 350-fold (new cohorts) increase in the risk of WNVD (Table 3).

Risk of neuroinvasive disease in individuals with auto-Abs neutralizing type I IFNs

Among WNVD patients, the prevalence of auto-Abs, overall and in various combinations, was significantly higher in the subgroup of individuals with neuroinvasive disease than in the general population. The presence of auto-Abs neutralizing at least low concentrations of IFN- $\alpha 2$ and/or IFN- ω , whether considering or not neutralization of IFN- β , was associated with an ~15–20 times higher risk of neuroinvasive disease, encephalitis in particular, in both the new cohorts (OR = 18.8; 95% CI: 13.1–27, $P < 10^{-16}$ for encephalitis) and in the overall study population consisting of the new and previously studied cohorts analyzed together (OR = 17.7; 95% CI: 13.8–22.8, $P < 10^{-16}$ for encephalitis) (Fig. 5, Fig. 7, A–D, and Table S3). A combination of auto-Abs neutralizing at least low concentrations of both IFN- $\alpha 2$ and IFN- ω was associated with an ~40–50 times increase in the risk of neuroinvasive disease, encephalitis in particular, in both the new cohorts and in the overall study population (Fig. 7, A–D; and Table S3). The presence of auto-Abs neutralizing combinations of high concentrations of IFN- $\alpha 2$ and/or IFN- β and/or IFN- ω resulted in ~30–80-fold increase in the risk of neuroinvasive disease, encephalitis in particular (Fig. 7, A–D; and Table S3). The




Figure 5. Frequency of auto-Abs against type I IFNs in the WNVIC, WNVF, and WNVND groups and the two WNVND subgroups (WNVD without evidence of neuroinvasive disease and WNND) relative to the general population in the seven new cohorts and the overall study population consisting of all 13 cohorts. Horizontal bars represent the 95% CI limits. IFN- α , auto-Abs neutralizing IFN- α 2 (regardless of effects on other IFNs); IFN- ω , auto-Abs neutralizing IFN- ω ; IFN- $\alpha \pm \omega \pm \beta$, auto-Abs neutralizing IFN- α 2 and/or IFN- ω and/or IFN- β ; IFN- $\alpha + \omega$, auto-Abs neutralizing both IFN- α 2 and IFN- ω ; WNND, WNV neuroinvasive disease. ns, nonsignificant; ****P < 10⁻⁴.

presence of auto-Abs neutralizing a combination of high concentrations (10 ng/ml) of both IFN- α 2 and IFN- ω resulted in an ~100- to 200 times increase in the risk of neuroinvasive disease, encephalitis in particular, in both the new cohorts and the overall study population (Fig. 7, A-D; and Table S3). Individuals

with a combination of auto-Abs neutralizing high concentrations (10 ng/ml) of IFN- α 2 and IFN- β and IFN- ω had the highest risk of neuroinvasive disease, encephalitis in particular, with a >100-fold (overall cohorts) to >200-fold (new cohorts) increase in the risk (Table S3).




Figure 6. **ORs for the presence of auto-Abs in the WNV groups relative to the general population. (A–D)** ORs for the presence of auto-Abs in the WNVIC group relative to the general population, adjusted for age and sex with Firth’s bias-corrected logistic regression, in the seven new cohorts (A) and the overall study population consisting of 13 cohorts (B). ORs for the presence of auto-Abs in individuals with WNVF and WNVD in the seven new cohorts (C) and the overall study population consisting of 13 cohorts (D) relative to the general population, also adjusted for age and sex via logistic regression. Horizontal bars represent the 95% CI limits. IFN- α , auto-Abs neutralizing IFN- α 2 (regardless of effects on other IFNs); IFN- ω , auto-Abs neutralizing IFN- ω ; IFN- α \pm ω \pm β , auto-Abs neutralizing IFN- α 2 and/or IFN- ω and/or IFN- β ; IFN- α + ω , auto-Abs neutralizing both IFN- α 2 and IFN- ω . ns, nonsignificant; **** P < 10^{-4} .

Risk of WNVD due to auto-Abs neutralizing type I IFNs in subjects ≤ 65 years old

The calculated risk of WNVD conferred by the auto-Abs was greatest in subjects ≤ 65 years old for all combinations of auto-Abs neutralizing low, intermediate, or high concentrations of the three type I IFNs tested (Fig. 8, A–D and Table 3). In subjects ≤ 65 years old, the presence of auto-Abs neutralizing high concentrations (10 ng/ml) of IFN- α 2 combined with auto-Abs neutralizing high (10 ng/ml) or low (100 pg/ml) concentrations of IFN- ω resulted in an ~ 500 to $\sim 1,500$ increase in the risk of WNVD in the new cohorts and the overall study population (Fig. 8, A–D and Table 3). The calculated risk was similarly high in the subgroup of patients with neuroinvasive disease in the new cohorts and the total cohort (Table S3). In the subgroup of patients with encephalitis, the most severe manifestation of WNV infection, auto-Abs neutralizing at least 100 pg/ml of both IFN- α 2 and IFN- ω conferred an ~ 400 to ~ 700 times increase in

risk for the new and total cohorts, whereas auto-Abs neutralizing both high concentrations of IFN- α 2 and any concentration tested (high or low) of IFN- ω were associated with the highest risk, $>2,000$ times higher than for the reference group (OR = 2,597.5; 95% CI: 117.1–57597.3, $P = 3.93 \times 10^{-12}$ and OR = 2,218.4; 95% CI: 125.1–39337.7, $P < 10^{-16}$ for the new and total cohorts, respectively) (Table S3).

Risk of WNVD due to auto-Abs neutralizing type I IFNs in subjects >65 years old

In subjects >65 years old, a combination of auto-Abs neutralizing a combination of high concentrations of both IFN- α 2 and IFN- ω , regardless of the neutralization of IFN- β , resulted in an ~ 80 times higher risk of WNVD (Fig. 8, A–D and Table 3), neuroinvasive disease, and encephalitis in particular (Table S3), both in the new and total cohorts. A combination of auto-Abs neutralizing high concentrations of the three auto-Abs tested

Table 3. Risk of WNVF or WNVD in subjects carrying auto-Abs neutralizing specific sets of type I IFNs, relative to the general population, with adjustment for age and sex and risk of WNVD by age group

Anti-type I IFN auto-Ab (amount of type I IFN neutralized, in plasma diluted 1:10)	WNV group	New cohort		Overall cohort	
		OR [95% CI]	P value	OR [95% CI]	P value
Anti-IFN- ω (100 pg/ml) regardless of other type I IFNs neutralization	WNVF	1.6 [0.3–7.8]	0.6	5.8 [3.3–10.1]	5.0×10^{-7}
Anti-IFN- $\alpha 2$ (100 pg/ml) regardless of other type I IFNs neutralization	WNVF	0.7 [0.1–11.8]	0.8	5 [2.3–9.9]	1.5×10^{-4}
Anti-IFN- $\alpha 2$ (100 pg/ml) and/or anti-IFN- ω (100 pg/ml) and/or anti-IFN- β (10 ng/ml)	WNVF	1.0 [0.2–5.2]	1.0	4.3 [2.6–7.3]	4.1×10^{-6}
Anti-IFN- $\alpha 2$ (100 pg/ml) and/or anti-IFN- ω (100 pg/ml) regardless of anti-IFN- β	WNVF	1.12 [0.2–5.6]	0.9	4.7 [2.8–7.9]	1.4×10^{-6}
Anti-IFN- $\alpha 2$ (100 pg/ml) and anti-IFN- ω (100 pg/ml) regardless of anti-IFN- β	WNVF	1.6 [0.1–26.4]	0.8	8.7 [4–19.1]	2.1×10^{-5}
Anti-IFN- ω (10 ng/ml) regardless of other type I IFNs neutralization	WNVF	2.5 [0.2–39.8]	0.6	12.8 [6–27.5]	1.2×10^{-6}
Anti-IFN- $\alpha 2$ (10 ng/ml) regardless of other type I IFNs neutralization	WNVF	2.4 [0.1–39.1]	0.6	12.1 [5.6–26.2]	2×10^{-6}
Anti-IFN- $\alpha 2$ (10 ng/ml) and/or anti-IFN- ω (10 ng/ml) and/or anti-IFN- β (10 ng/ml)	WNVF	0.8 [0.1–13.4]	0.9	5.7 [2.9–11.3]	5.3×10^{-5}
Anti-IFN- $\alpha 2$ (10 ng/ml) and/or anti-IFN- ω (10 ng/ml) regardless of anti-IFN- β	WNVF	1.5 [0.1–24.4]	0.8	10.0 [5.1–19.8]	4.4×10^{-7}
Anti-IFN- $\alpha 2$ (10 ng/ml) and anti-IFN- ω (10 ng/ml) regardless of anti-IFN- β	WNVF	6.2 [0.4–101.9]	0.3	22.5 [9–56.1]	1.8×10^{-6}
Anti-IFN- $\alpha 2$ (10 ng/ml) and anti-IFN- ω (100 pg/ml) regardless of anti-IFN- β	WNVF	3.0 [0.2–50.6]	0.5	14.0 [5.7–34]	6.1×10^{-6}
Anti-IFN- ω (100 pg/ml) regardless of other type I IFNs neutralization	WNVD (all)	22.6 [16.2–31.2]	$<10^{-16}$	23.4 [18.8–29.3]	$<10^{-16}$
	WNVD ≤ 65	32 [17–59.9]	$<10^{-16}$	26 [17–39.6]	$<10^{-16}$
	WNVD >65	27.7 [18.3–41.9]	$<10^{-16}$	26.9 [20.4–35.5]	$<10^{-16}$
Anti-IFN- $\alpha 2$ (100 pg/ml) regardless of other type I IFNs neutralization	WNVD (all)	24.5 [17.3–34.5]	$<10^{-16}$	27.2 [21.5–34.4]	$<10^{-16}$
	WNVD ≤ 65	89.2 [43.9–181.3]	$<10^{-16}$	66.3 [38.1–115.4]	$<10^{-16}$
	WNVD >65	21.8 [14.4–32.8]	$<10^{-16}$	24.7 [18.9–32.4]	$<10^{-16}$
Anti-IFN- $\alpha 2$ (100 pg/ml) and/or anti-IFN- ω (100 pg/ml) and/or anti-IFN- β (10 ng/ml)	WNVD (all)	17.4 [12.7–23.7]	$<10^{-16}$	17.4 [14.1–21.4]	$<10^{-16}$
	WNVD ≤ 65	27.9 [16.3–47.8]	$<10^{-16}$	19.2 [13.2–27.9]	$<10^{-16}$
	WNVD >65	17.8 [11.9–26.4]	$<10^{-16}$	19 [14.5–24.6]	$<10^{-16}$
Anti-IFN- $\alpha 2$ (100 pg/ml) and/or anti-IFN- ω (100 pg/ml) regardless of anti-IFN- β	WNVD (all)	18.7 [13.8–23.4]	$<10^{-16}$	18.9 [15.4–23.1]	$<10^{-16}$
	WNVD ≤ 65	36.6 [21.1–63.6]	$<10^{-16}$	24.5 [16.7–36.0]	$<10^{-16}$
	WNVD >65	19.8 [13.4–29.3]	$<10^{-16}$	20.8 [16.2–26.9]	$<10^{-16}$
Anti-IFN- $\alpha 2$ (100 pg/ml) and anti-IFN- ω (100 pg/ml) regardless of anti-IFN- β	WNVD (all)	41.7 [27.9–62.2]	$<10^{-16}$	49.7 [37–67.1]	$<10^{-16}$
	WNVD ≤ 65	426.4 [114.5–1,587.9]	$<10^{-16}$	300 [106.5–845]	$<10^{-16}$
	WNVD >65	38.3 [24.3–60.2]	$<10^{-16}$	40.5 [29.3–56]	$<10^{-16}$
Anti-IFN- ω (10 ng/ml) regardless of other type I IFNs neutralization	WNVD (all)	52.9 [36–77.5]	$<10^{-16}$	57.1 [43.5–75]	$<10^{-16}$
	WNVD ≤ 65	162 [75.2–348.8]	$<10^{-16}$	138.9 [81.7–236]	$<10^{-16}$
	WNVD >65	43.7 [28.3–67.4]	$<10^{-16}$	42 [31–57]	$<10^{-16}$
Anti-IFN- $\alpha 2$ (10 ng/ml) regardless of other type I IFNs neutralization	WNVD (all)	51.1 [35–74.4]	$<10^{-16}$	63.1 [48.6–81.8]	$<10^{-16}$
	WNVD ≤ 65	300.2 [137.4–665.1]	$<10^{-16}$	213.7 [117.6–388.6]	$<10^{-16}$
	WNVD >65	37.4 [24.5–57.2]	$<10^{-16}$	46.9 [35.3–62.3]	$<10^{-16}$
Anti-IFN- $\alpha 2$ (10 ng/ml) and/or anti-IFN- ω (10 ng/ml) and/or anti-IFN- β (10 ng/ml)	WNVD (all)	28.8 [20–41.4]	$<10^{-16}$	31.2 [23.9–40.7]	$<10^{-16}$
	WNVD ≤ 65	60.3 [30.1–120.8]	$<10^{-16}$	44.4 [26.7–74]	$<10^{-16}$
	WNVD >65	25.8 [16.8–39.9]	$<10^{-16}$	29 [21.2–39.6]	$<10^{-16}$

Downloaded from http://rupress.org/jhi/article-pdf/23/3/e20250189/2028254/jhi_20250189.pdf by guest on 16 March 2026

Table 3. Risk of WNVF or WNVD in subjects carrying auto-Abs neutralizing specific sets of type I IFNs, relative to the general population, with adjustment for age and sex and risk of WNVD by age group (Continued)

Anti-type I IFN auto-Ab (amount of type I IFN neutralized, in plasma diluted 1:10)	WNV group	New cohort		Overall cohort	
		OR [95% CI]	P value	OR [95% CI]	P value
Anti-IFN- α 2 (10 ng/ml) and/or anti-IFN- ω (10 ng/ml) regardless of anti-IFN- β	WNVD (all)	38.4 [27.0–54.6]	$<10^{-16}$	45.6 [36.1–57.6]	$<10^{-16}$
	WNVD ≤ 65	172.1 [88.9–332.9]	$<10^{-16}$	120.7 [75.5–192.9]	$<10^{-16}$
	WNVD >65	28.8 [19.1–43.2]	$<10^{-16}$	34.8 [26.7–45.3]	$<10^{-16}$
Anti-IFN- α 2 (10 ng/ml) and anti-IFN- ω (10 ng/ml) regardless of anti-IFN- β	WNVD (all)	104.6 [66.5–164.4]	$<10^{-16}$	119.7 [83.8–171]	$<10^{-16}$
	WNVD ≤ 65	548.2 [173.3–1,733.9]	$<10^{-16}$	538.2 [208.1–1,391.8]	$<10^{-16}$
	WNVD >65	80 [49.1–130.5]	$<10^{-16}$	80.8 [55.4–117.8]	$<10^{-16}$
Anti-IFN- α 2 (10 ng/ml) and anti-IFN- ω (100 pg/ml) regardless of anti-IFN- β	WNVD (all)	57.2 [34.1–95.8]	$<10^{-16}$	77.5 [50.5–118.9]	$<10^{-16}$
	WNVD ≤ 65	1,730.9 [92.4–32,443.2]	2.2×10^{-16}	1,517 [91.5–25,181.1]	$<10^{-16}$
	WNVD >65	45.2 [26.2–78.1]	$<10^{-16}$	55.3 [35.7–85.5]	$<10^{-16}$
Anti-IFN- α 2 (10 ng/ml) and anti-IFN- ω (10 ng/ml) and anti-IFN- β (10 ng/ml)	WNVD (all)	347.7 [18.9–6,381.4]	5.9×10^{-9}	173.0 [10.0–3,002.8]	4.9×10^{-10}
	WNVD ≤ 65	200.2 [9.4–4,270.8]	7.9×10^{-5}	74.0 [3.5–1,559.6]	9×10^{-4}
	WNVD >65	177.2 [9.0–3,497.7]	1.1×10^{-5}	116.8 [6.6–2,063.7]	1.1×10^{-7}

≤ 65 and >65 indicate age cutoffs, in years; anti-IFN- ω and anti-IFN- α 2 indicate auto-Abs neutralizing IFN- ω or IFN- α 2, respectively, regardless of their effects on other type I IFNs.

conferred the highest risk in this age group, with a >100 times increase in the risk of WNVD and neuroinvasive disease (Table 3 and Table S3). Overall, the presence of auto-Abs neutralizing type I IFNs confers a very high risk of WNVD in all age groups. The biological and medical impact of various combinations of auto-Abs was considerable in both the under- and over-65s. The lower ORs calculated for individuals aged 65 years or older reflect both a higher baseline risk of severe disease in the absence of auto-Abs and the higher baseline prevalence of auto-Abs in the elderly (45). These results are consistent with previous reports on anti-type I IFN auto-Abs in individuals <65 years old (or 70 years old as a cutoff, in some studies) with hypoxemic COVID-19 pneumonia (45), critical influenza pneumonia (46), or WNVD (20). Overall, the data for the seven new cohorts of WNVD patients from five countries confirm the strong effect of auto-Abs neutralizing type I IFNs on the risk of developing severe disease following exposure to WNV and demonstrate that impaired type I IFN immunity is a major determinant of WNVD and neuroinvasive disease. Auto-Abs neutralizing type I IFNs underlie life-threatening WNVD in a significant proportion of cases, considerably increasing the likelihood of neuroinvasive disease and encephalitis in particular.

Discussion

These findings extend those of our previous report of seven cohorts (20) and three recent reports (24, 25, 26). They confirm that auto-Abs neutralizing type I IFNs underlie almost half of all cases of WNV encephalitis. These auto-Abs have now been found

in 13 unrelated cohorts from nine centers in five countries in Europe, the Middle East, and North America, in samples collected from 2002 to 2024, and in other patients from Italy, Germany, and Israel (24, 25, 26). When analyzed collectively, the ORs for encephalitis were found to be very high, ranging from ~ 20 for auto-Abs neutralizing low concentrations of IFN- α 2 and/or IFN- ω to $>2,000$ for auto-Abs neutralizing high concentrations of IFN- α 2 together with IFN- ω in the under-65s. The proportion of cases with neuroinvasive disease carrying auto-Abs neutralizing one or more type I IFN was also very high, and equally so across cohorts, with a mean frequency of 38% and a range of 17–50% of cases. These variations may depend on differences in disease group assignment, suggesting that consistent definitions of disease severity and neurological involvement and follow-up data for infected individuals are required.

Such high ORs and proportions are unprecedented for human infectious diseases (47, 48). The ORs were calculated by comparing patients with WNV encephalitis relative to individuals untested for WNV infection in a large sample of the general population. In our cohort, the proportion of individuals with silent WNV infection carrying the auto-Abs did not significantly differ from that in the general population. Yet, our sample of WNVIC is much smaller (146 WNVIC vs. $>10,000$ individuals of the corresponding age groups) and includes only 14 subjects >65 years old, none of whom carried the auto-Abs. It is reasonable to predict that the proportion of individuals carrying the auto-Abs among those with silent WNV infection, selected as appropriate controls on the basis of their resistance to severe disease, could be even smaller than that in the general population,




Figure 7. ORs for the presence of auto-Abs in the WNVD group relative to the general population. (A–D) ORs for the presence of auto-Abs in the WNVD group and the two WNVD subgroups (WNVD without evidence of neuroinvasive disease and WNV neuroinvasive disease, WNND) relative to the general population, adjusted for age and sex with Firth’s bias-corrected logistic regression, in the seven new cohorts (A) and the overall study population consisting of 13 cohorts (B). ORs for the presence of auto-Abs in the WNND group and the four WNND subgroups (WNE, WNV encephalitis; WNM, WNV meningitis; AFP, acute flaccid paralysis; UNS, unspecified neurological syndrome) in the seven new cohorts (C) and in the overall study population consisting of 13 cohorts (D) relative to the general population, also adjusted for age and sex via logistic regression. Horizontal bars represent the 95% CI limits. IFN- α , auto-Abs neutralizing IFN- α 2 (regardless of effects on other IFNs); IFN- ω , auto-Abs neutralizing IFN- ω ; IFN- α \pm ω \pm β , auto-Abs neutralizing IFN- α 2 and/or IFN- ω and/or IFN- β ; IFN- α + ω , auto-Abs neutralizing both IFN- α 2 and IFN- ω . ns, nonsignificant; ****P < 10⁻⁴.




Figure 8. ORs for the presence of auto-Abs in the WNV groups relative to the general population, by age group. (A–D) ORs for the presence of auto-Abs in patients with WNV relative to the general population, as determined by logistic regression, stratified for age group, for auto-Abs neutralizing different combinations of high concentrations (A) and low concentrations (B) of type I IFNs in the seven new cohorts, and for auto-Abs neutralizing different combinations of high concentrations (C) and low concentrations (D) of type I IFNs in the overall study population consisting of 13 cohorts. ORs were calculated separately for patients aged ≤ 65 and >65 years. Horizontal bars represent the 95% CI limits. IFN- α , auto-Abs neutralizing IFN- $\alpha 2$ (regardless of effects on other IFNs); IFN- ω , auto-Abs neutralizing IFN- ω ; IFN- $\alpha \pm \omega \pm \beta$, auto-Abs neutralizing IFN- $\alpha 2$ and/or IFN- ω and/or IFN- β ; IFN- $\alpha + \omega$, auto-Abs neutralizing both IFN- $\alpha 2$ and IFN- ω ; ORs, odds ratios. ns, nonsignificant; **** $P < 10^{-4}$.

in particular among individuals >65 years old, resulting in higher ORs for WNV and encephalitis. This is also suggested by our observation that, in the new cohorts, no samples from WNVIC and mild WNVF patients neutralized IFN- $\alpha 2$, and there were no subjects with auto-Abs neutralizing more than one type I IFN, while 1.4% of individuals in the general population carry auto-Abs neutralizing IFN- $\alpha 2$ and 0.6% carry auto-Abs neutralizing two or more type I IFNs (23).

The cells infected with WNV in which type I IFN activity is blocked by these auto-Abs, resulting in encephalitis, remain to be identified (22). Candidates include leukocytes, cells at the blood–brain barrier, and cells within the central nervous system (22). This study has important clinical implications. People at risk of having auto-Abs against type I IFNs, including individuals

with a history of autoimmunity or viral disease or over the age of 70 years, should be screened. Patients with auto-Abs neutralizing type I IFNs should be considered at risk and should avoid inhabiting or traveling to areas in which WNV is endemic. If this cannot be avoided or if they are outdoor workers living in endemic areas (49, 50), they should protect themselves against mosquitoes and, if bitten, should seek medical attention before the possible development of clinical disease. The development of a vaccine against WNV is a public health priority, and screening for these auto-Abs might guide future vaccine-based prevention strategies, as live-attenuated WNV vaccines may be harmful in patients with auto-Abs against type I IFNs (29, 30). Patients hospitalized for WNV encephalitis should be tested for auto-Abs, as should patients diagnosed with WNV infection in other

contexts. Treatment with a type I IFN not neutralized by the auto-Abs, such as IFN- β (51), or, in the future, with decoys that neutralize the auto-Abs but not the IFNs and their receptors (35, 52), or with chimeric antigen receptor (CAR)-T cells (53), may also be considered.

In 2020, we concluded that auto-Abs against type I IFNs were causal for critical COVID-19 for several individually indirect but collectively compelling reasons (34), as summarized by Manry et al. (45). This interpretation was supported by the longitudinal follow-up of a Swiss cohort of patients living with human immunodeficiency virus infection, which showed that these auto-Abs occur before severe viral diseases and that they later diversify and persist for life (52, 54). The production of auto-Abs against type I IFNs results from a failure of tolerance to self, which can be driven by monogenic inborn errors of immunity affecting central tolerance in the thymus (55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66). Once these auto-Abs appear, they do not disappear.

auto-Abs neutralizing type I IFN can underlie at least three severe viral respiratory diseases—COVID-19, influenza, and MERS—in 5–20% of cases (23, 34, 46, 67, 68), fulminant viral hepatitis caused by herpes simplex virus (HSV)-1 and HSV-2 (69), and severe adverse reactions to live-attenuated vaccines against YFV (YFV-17D) (29) and CHIKV (VLA1553) (30). They also underlie a case of lethal avian H5N1 influenza (70). We show here that they are the major determinants of WNV encephalitis in North America, Europe, and the Middle East. They also underlie 10% of cases of TBE (27) and the rarer cases of encephalitis due to the Powassan and Usutu viruses (28). They were also found in the most severe case of Ross River disease, which is caused by an alphavirus (28). They should be sought in patients with other types of encephalitis, particularly due to arboviruses, which are typically harmless in most infected individuals but may cause severe disease in a minority of cases (31). There are over 150 known human-pathogenic arboviruses, and studies to assess the contribution of auto-Abs against type I IFNs to the pathogenesis of each arboviral disease are warranted.

Materials and methods

Patients

We enrolled an international cohort of 318 individuals aged 10–99 years with documented WNV infection, 65.1% of whom were male and 33.9% female, living in Austria, Hungary, Israel, Italy, or the USA (Fig. 1 and Fig. S1), and a cohort of 663 individuals described in a previous study (20), living in Hungary, Italy, or the USA. In the new cohorts, neither sex nor age was reported for 2/318 individuals, age was not reported for 2/318 individuals, and sex was not reported for 1/318 individuals. In total, there were 981 individuals aged 9–99 years with documented WNV infection, 64.7% of whom were male and 34.9% female, living in Austria, Hungary, Israel, Italy, or the USA (Fig. S5). For all the enrolled individuals from the new and previously described cohorts, written informed consent was obtained in the country of residence of each patient, unless samples and anonymized medical information could be processed without prior written informed consent, in accordance with local regulations and with

institutional review board (IRB) approval. The Center for Virology at the Medical University of Vienna, Austria, serves as the national reference center for arboviruses and has the legal permission to process anonymized patient information on notifiable viral diseases. WNV infection was diagnosed on the basis of the serological demonstration of WNV-specific IgM or seroconversion to IgG, WNV neutralization assays (32), and/or RT-PCR on serum, plasma, or cerebrospinal fluid samples. Individuals were stratified according to the presence and/or severity of clinical manifestations, as defined by the need for hospitalization. Life-threatening WNV was defined as WNV infection requiring hospitalization. WNVF was defined as WNV infection not requiring hospitalization in patients reporting a febrile illness requiring outpatient care. The WNVIC were blood donors with documented WNV infection, diagnosed on the basis of the detection of WNV RNA in blood during screening at the time of blood donation, who remained asymptomatic or paucisymptomatic (headache) during follow-up. WNVIC patients included individuals with confirmed neurological disease (WNV neuroinvasive disease) and individuals without clinical evidence of neuroinvasive disease. The individuals in the neuroinvasive disease group were reported to have encephalitis (fever, acute signs of central or peripheral neurologic dysfunction, including altered mental status and neurological deficits), meningitis (fever, pleocytosis, headache, and nuchal rigidity), acute flaccid paralysis (poliomyelitis-like syndrome or Guillain-Barré-like syndrome), or other neurological syndromes. The experiments were conducted in Italy, France, and the USA, in accordance with local regulations and guidance from the Italian National Data Protection Authority, the French Ethics Committee (Comité de Protection des Personnes), the French National Agency for Medicine and Health Product Safety, the Institut National de la Santé et de la Recherche Médicale in Paris, France, and with the approval of the IRB of the Italian institutions (San Matteo Research Hospital in Pavia [Comitato Etico Territoriale Lombardia 6 - Policlinico San Matteo], the University Hospital of Padua, and the University Hospital of Bologna, Amedeo di Savoia Hospital-ASL Città in Torino in Turin), the Medical University of Vienna in Austria, the Tel-Aviv Sourasky Medical Center in Israel, the National Public Health Center in Budapest, and the Rockefeller University in New York, USA, respectively.

ELISA

ELISAs were performed as previously described (33). In brief, 96-well ELISA plates (MaxiSorp; Thermo Fisher Scientific) were coated by overnight incubation at 4°C with 1 μ g/ml rhIFN- α (ref. number 130-108-984; Miltenyi Biotec), rhIFN- ω (ref. number 300-02J; PeproTech), or rhIFN- β (ref. number 300-02BC; PeproTech). The plates were washed (PBS/0.005% Tween), blocked by incubation with the same buffer supplemented with 2% BSA, washed, and incubated with 1:50 dilutions of plasma samples from the patients or positive and negative controls for 2 h at room temperature. Each sample was tested once. Plates were thoroughly washed (PBS/0.005% Tween), and horseradish peroxidase-conjugated Fc-specific IgG fractions from polyclonal goat antiserum against human IgG (Nordic Immunological Laboratories) were added to a final concentration of 1 μ g/ml. Plates were

incubated for 1 h at room temperature and washed. The substrate was added, and OD was measured (450 nm). All the incubation steps were performed with gentle shaking (600 rpm).

Luciferase reporter assay

The blocking activity of anti-IFN- α 2, anti-IFN- ω , and anti-IFN- β auto-Abs was determined with a luciferase reporter assay, as previously described (23). Briefly, HEK293T cells were transfected with a plasmid encoding the firefly luciferase gene under the control of the human ISRE promoter in the pGL4.45 backbone and a plasmid constitutively expressing the *Renilla* luciferase as a control for transfection (pRL-SV40). Cells were transfected in the presence of the X-tremeGene9 transfection reagent (ref. number 6365779001; Sigma-Aldrich). After 24 h, cells in Dulbecco's modified Eagle medium (Thermo Fisher Scientific) supplemented with 2% fetal calf serum and 10% control or patient serum/plasma/whole blood (after heat inactivation at 56°C for 20 min) were either left unstimulated or were stimulated with unglycosylated rhIFN- α 2 (ref. number 130-108-984; Miltenyi Biotec), unglycosylated rhIFN- ω (ref. number 300-02J; PeproTech) at a concentration of 10 ng/ml or 100 pg/ml, or glycosylated rhIFN- β (ref. number 300-02BC; PeproTech) at a concentration of 10 or 1 ng/ml for 16 h at 37°C under an atmosphere containing 5% CO₂. Finally, the cells were lysed by incubation with a lysis buffer (provided in ref. number E1980; Promega) for 20 min at room temperature, and luciferase levels were measured with the Dual-Luciferase Reporter 1000 assay system (ref. number E1980; Promega) according to the manufacturer's protocol. Luminescence intensity was measured with a VICTOR-X Multilabel Plate Reader (PerkinElmer Life Sciences). Firefly luciferase activity values were normalized against *Renilla* luciferase activity values. The resulting values (luciferase induction) were then normalized against the median level of induction for non-neutralizing samples and expressed as a percentage (relative luciferase activity [RLA] ratio, %). Samples were considered to have neutralizing activity if the RLA ratio was below 15% of the median value for controls tested on the same day.

Statistical analysis

Odds ratios and P values for the effect of auto-Abs neutralizing each type I IFN in WNV patients relative to healthy individuals from the general population, adjusted for age in years and sex, were estimated by means of Firth's bias corrected logistic regression, as implemented in the logistf package of R software. Where relevant, statistical test results are indicated in the corresponding figures. ns, not significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001.

Online supplemental material

Fig. S1 shows demographic characteristics of individuals from seven new cohorts enrolled in the study. Fig. S2 shows correlation between luciferase-based IFN neutralization assay and ELISA results for the detection of auto-Abs to type I IFNs. Fig. S3 shows detection of auto-Abs against type I IFN by ELISA in patients with WNVD from seven new cohorts. Fig. S4 shows detection of auto-Abs neutralizing type I IFNs in the WNVD subgroups of patients from seven new cohorts. Fig. S5 shows

demographic characteristics of 981 individuals from 13 cohorts. Table S1 shows prevalence of WNVIC, WNVF, WNVD, or WNND of subjects carrying auto-Abs neutralizing at least 1 IFN-I by WNV lineage (WNV-1 or WNV-2). Table S2 shows risk of WNVIC, WNVF, WNVD, or WNND of subjects carrying auto-Abs neutralizing at least one IFN-I by sex. Table S3 shows risk of neuroinvasive disease and encephalitis for subjects carrying auto-Abs neutralizing specific sets of type I IFNs, relative to the general population, with adjustment for age and sex and risk by age group.

Data availability

All data supporting the findings of this study are available within the main text and supplemental material and from the corresponding authors upon request.

Acknowledgments

We thank the patients and their families for participating in our research. We thank all members of both branches of the Laboratory of Human Genetics of Infectious Diseases for discussions and technical and administrative support. Alessandro Borghesi wishes to thank Giampaolo Merlini, University of Pavia, for thoughtful discussions on the methods for anti-cytokine auto-Ab discovery in samples from patients with bacterial and viral infections. We thank Lazaro Lorenzo, Laboratory of Human Genetics of Infectious Diseases, Necker Branch, Institut National de la Santé et de la Recherche Médicale (INSERM) U1163, Necker Hospital for Sick Children, Paris, France, EU.

The Laboratory of Human Genetics of Infectious Diseases is supported by the Howard Hughes Medical Institute, The Rockefeller University, the St. Giles Foundation, the Stavros Niarchos Foundation (SNF) as part of its grant to the SNF Institute for Global Infectious Disease Research at The Rockefeller University, the National Institutes of Health (NIH) (R01AI163029), the National Center for Advancing Translational Sciences, the NIH Clinical and Translational Science Award program (UL1TR001866), the French Agence Nationale de la Recherche (ANR) under the France 2030 program (ANR-10-IAHU-01), the HORIZON-HLTH-2024-DISEASE-08-20 program under GA 101191725 (InFlaMe), the Agence Nationale de Recherches sur le Sida et les Hépatites Virales (ANRS) projects DéméléJEV (ANRS0629), IFNArboVE (ANRS00869-R-PR), and LSDengue (ANRS-23-PEPR-MIE-0007), the Integrative Biology of Emerging Infectious Diseases Laboratory of Excellence (ANR-10-LABX-62-IBEID), the French Foundation for Medical Research (FRM) (EQU202503020018), ANR GENVIR (ANR-20-CE93-003), and ANR AI2D (ANR-22-CE15-0046) projects, the HORIZON-HLTH-2021-DISEASE-04 program under grant agreement 101057100 (UNDINE), the ANR-RHU COVIFERON Program (ANR-21-RHUS-0008), the Square Foundation, Grandir - Fonds de solidarité pour l'enfance, the Fondation du Souffle, the SCOR Corporate Foundation for Science, the Battersea & Bowery Advisory Group, William E. Ford, General Atlantic's Chairman and Chief Executive Officer, Gabriel Caillaux, General Atlantic's Co-President, Managing Director and Head of Business in EMEA, and the General Atlantic Foundation, the French Ministry of Higher

Education, Research, and Innovation (MESRI-COVID-19), INSERM, REACTing-INSERM, Paris Cité University, and the Imagine Institute. P. Bastard was supported by a “Poste CCA-INSERM-Bettencourt” (with support from the Bettencourt-Schueller Foundation) and the FRM (EA20170638020). A. Gervais was supported by a French National Agency for Research grant as part of the “Investissement d’Avenir” program (ANR-10-LABX-62-01). The Host-pathogen group (A. Borghesi), Fondazione IRCCS Policlinico San Matteo, Pavia, Italy, is supported by grants from the Italian Ministry of Health, RC08061819 and RC08061822, and by the 5X1000 grant 08061821 from the San Matteo Hospital. The Rheumatology Unit (V. Codullo and F. Trespidi), Fondazione IRCCS Policlinico San Matteo, Pavia, Italy, is supported by grant RC08073123 from the Italian Ministry of Health. The Microbiology and Virology Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy, is supported by grants from the Ministry of Health, RC80206, and by the European Union under grant agreement 101191725 (InFlaMe project, HORIZON-HLTH-2024-DISEASE-08-20 program). A. Ferrari, at the Microbiology and Virology Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy, is affiliated with and supported by the National PhD Programme in One Health—Approaches to Infectious Diseases and Life Science Research, at the Department of Public Health, Experimental and Forensic Medicine, University of Pavia, Pavia, Italy, and by grant from the Italian Ministry of Health RC8048424. The Cell Factory (P. Comoli, M.A. Avanzini, and S. Croce), at Fondazione IRCCS Policlinico San Matteo, Pavia, Italy, is supported by grant RC8053724 from the Italian Ministry of Health. The Department of Molecular Medicine, University of Padua, Italy, is supported by the European Union’s Horizon 2020 research and innovation program (VEO, grant agreement 874735) and the EU4Health program (DURABLE, grant agreement 101102733). The Department of Clinical, Surgical, Diagnostic, and Pediatric Sciences, University of Pavia, Italy, and the Department of Molecular Medicine, University of Padua, Italy, are supported by EU funding within the Next-Generation EU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (project no. PE00000007, INF-ACT). The Medical University of Vienna, Austria is supported by the European Union under grant agreement 31101191725 (InFlaMe project, HORIZON-HLTH-2024-795 DISEASE-08-20 program). The Laboratory of Virology and Infectious Disease (M.R. MacDonald and C.M. Rice) is supported by NIH grants R01AI124690 and R01AI091707. The Murray laboratory is supported by NIH grants R01AI091816 and U19AI089992.

Author contributions: Adrian Gervais: conceptualization, investigation, methodology, validation, visualization, and writing—original draft, review, and editing. Francesca Trespidi: data curation, formal analysis, investigation, visualization, and writing—original draft, review, and editing. Alessandro Ferrari: methodology and resources. Francesca Rovida: resources. Astrid Marchal: formal analysis, Stefania Croce: investigation. Irene Cassaniti: methodology. Mattia Moratti: data curation, resources, and writing—review and editing. Jennifer L. Uhrlaub: resources. David M. Florian: resources and writing—review and editing. Karin Stiasny: resources and writing—review and editing. Elisa Burdino: investigation. Micol Angelini: data curation, investigation, and writing—review and editing. Lucy Bizien: investigation. Daniele

Lilleri: supervision, validation, and writing—review and editing. Veronica Codullo: data curation, investigation, resources, and writing—review and editing. Tal Freund: investigation, resources, and writing—review and editing. Yael Paran: data curation. Avi Gadoth: conceptualization, data curation, investigation, and project administration, supervision. Roni Biran: conceptualization. Alessandro Mancon: investigation and writing—review and editing. Camilla Lucca: data curation, investigation, and methodology. Stefania Vogiatzis: data curation, investigation, and methodology. Monia Pacenti: validation, visualization, and writing—original draft, review, and editing. Mélodie Aubart: writing—review and editing. Marco Zecca: investigation. Patrizia Comoli: data curation, investigation, methodology, and visualization. Jacques Fellay: supervision, validation, and writing—review and editing. Antonio Piralla: investigation and resources. Francesca Conti: conceptualization, methodology, project administration, supervision, and writing—review and editing. Alberto Dolci: supervision. Luisa Barzon: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, resources, and writing—review and editing. Valeria Ghisetti: conceptualization. Tiziana Lazzarotto: data curation, resources. Danilo Cereda: validation. Alessandro Aiuti: resources. Emmanuelle Jouanguy: resources and writing—review and editing. Paul Bastard: investigation, supervision, validation, and writing—review and editing. Margaret R MacDonald: supervision and writing—review and editing. Charles M. Rice: conceptualization and writing—review and editing. Anne Puel: funding acquisition, resources, and writing—review and editing. Laurent Abel: formal analysis and writing—review and editing. Giada Rossini: resources and writing—review and editing. Davide Mileto: resources, and visualization. Yannick Simonin: resources and writing—review and editing. Anna Nagy: resources. David Hagin: investigation, resources, and writing—review and editing. Kristy O. Murray: conceptualization, data curation, funding acquisition, investigation, and writing—review and editing. Fausto Baldanti: writing—review and editing. Judith H. Aberle: resources and writing—review and editing. Aurélie Cobat: data curation, formal analysis, funding acquisition, supervision, and writing—original draft, review, and editing. Shen-Ying Zhang: conceptualization, funding acquisition, investigation, resources, supervision, validation, and writing—review and editing. Jean-Laurent Casanova: conceptualization, funding acquisition, project administration, resources, supervision, validation, and writing—original draft, review, and editing. Alessandro Borghesi: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, validation, visualization, and writing—original draft, review, and editing.

Disclosures: K.O. Murray reported personal fees from Valneva, grants from NIH, and grants from CDC outside the submitted work. J.H. Aberle reported grants from Pfizer Austria outside the submitted work. J.-L. Casanova reported a patent to PCT/US2021/042741 pending. No other disclosures were reported.

Submitted: 1 September 2025

Revised: 23 January 2026

Accepted: 26 January 2026

References

- Young, J.J., J.M. Haussig, S.W. Aberle, D. Pervanidou, F. Riccardo, N. Sekulić, T. Bakonyi, and C.M. Gossner. 2021. Epidemiology of human West Nile virus infections in the European Union and European Union enlargement countries, 2010 to 2018. *Euro Surveill.* 26:2001095. <https://doi.org/10.2807/1560-7917.ES.2021.26.19.2001095>
- Postler, T.S., M. Beer, B.J. Blitvich, J. Bukh, X. de Lamballerie, J.F. Drexler, A. Imrie, A. Kapoor, G.G. Karganova, P. Lemey, et al. 2023. Renaming of the genus Flavivirus to Orthoflavivirus and extension of binomial species names within the family Flaviviridae. *Arch. Virol.* 168:224. <https://doi.org/10.1007/s00705-023-05835-1>
- Petersen, L.R., P.J. Carson, B.J. Biggerstaff, B. Custer, S.M. Borchardt, and M.P. Busch. 2013. Estimated cumulative incidence of West Nile virus infection in US adults, 1999-2010. *Epidemiol. Infect.* 141:591-595. <https://doi.org/10.1017/S0950268812001070>
- Chancey, C., A. Grinev, E. Volkova, and M. Rios. 2015. The global ecology and epidemiology of West Nile virus. *Biomed. Res. Int.* 2015:376230. <https://doi.org/10.1155/2015/376230>
- Chowdhury, P., and S.A. Khan. 2021. Global emergence of West Nile virus: Threat & preparedness in special perspective to India. *Indian J. Med. Res.* 154:36-50. https://doi.org/10.4103/ijmr.IJMR_642_19
- Mencattelli, G., M.H.D. Ndione, A. Silverj, M.M. Diagne, V. Curini, L. Teodori, M. Di Domenico, R. Mbaye, A. Leone, M. Marcacci, et al. 2023. Spatial and temporal dynamics of West Nile virus between Africa and Europe. *Nat. Commun.* 14:6440. <https://doi.org/10.1038/s41467-023-42185-7>
- European Centre for Disease Prevention and Control (ECDC). 2023. West Nile virus transmission season in Europe. <https://www.ecdc.europa.eu/en/news-events/epidemiological-update-west-nile-virus-transmission-season-europe-2023-0> (accessed August 5, 2024).
- European Centre for Disease Prevention and Control (ECDC). 2024. West Nile virus transmission season in Europe. <https://www.ecdc.europa.eu/archive/wnv-2024.html> (accessed June 26, 2025).
- Ministry of Health (Israel). 2025. West Nile virus disease in Israel. <https://www.gov.il/en/pages/disease-west-nile-fever?chapterIndex=8> (accessed June 26, 2025).
- Barzon, L., M. Pacenti, F. Montarsi, D. Fornasiero, F. Gobbo, E. Quaranta, I. Monne, A. Fusaro, A. Volpe, A. Sinigaglia, et al. 2022. Rapid spread of a new West Nile virus lineage 1 associated with increased risk of neuro-invasive disease during a large outbreak in northern Italy, 2022: One health analysis. *J. Trav. Med.* 31:taac125. <https://doi.org/10.1093/jtm/taac125>
- European Centre for Disease Prevention and Control (ECDC). 2024. West Nile virus seasonal surveillance - Historical data by year. <https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-data/historical> (accessed August 5, 2024).
- Centers for Disease Control and Prevention (CDC). 2026. West Nile virus - Historic data (1999-2023). https://www.cdc.gov/west-nile-virus/data-maps/historic-data.html?CDC_AAref_Val=https://www.cdc.gov/westnile/statsmaps/historic-data.html (accessed January 9, 2026).
- Gould, C.V., J.E. Staples, S.A.J. Guagliardo, S.W. Martin, S. Lyons, S.L. Hills, R.J. Nett, and L.R. Petersen. 2025. West Nile virus: A review. *JAMA.* 334:618-628. <https://doi.org/10.1001/jama.2025.8737>
- Mostashari, F., M.L. Bunning, P.T. Kitsutani, D.A. Singer, D. Nash, M.J. Cooper, N. Katz, K.A. Liljebjelke, B.J. Biggerstaff, A.D. Fine, et al. 2001. Epidemic West Nile encephalitis, New York, 1999: Results of a household-based seroepidemiological survey. *Lancet.* 358:261-264. [https://doi.org/10.1016/S0140-6736\(01\)05480-0](https://doi.org/10.1016/S0140-6736(01)05480-0)
- Busch, M.P., D.J. Wright, B. Custer, L.H. Tobler, S.L. Stramer, S.H. Kleinman, H.E. Prince, C. Bianco, G. Foster, L.R. Petersen, et al. 2006. West Nile virus infections projected from blood donor screening data, United States, 2003. *Emerg. Infect. Dis.* 12:395-402. <https://doi.org/10.3201/eid1205.051287>
- Lindsey, N.P., J.E. Staples, J.A. Lehman, M. Fischer, and Centers for Disease Control and Prevention (CDC). 2010. Surveillance for human West Nile virus disease - United States, 1999-2008. *MMWR Surveill. Summ.* 59:1-17.
- Carson, P.J., S.M. Borchardt, B. Custer, H.E. Prince, J. Dunn-Williams, V. Winkelman, L. Tobler, B.J. Biggerstaff, R. Lanciotti, L.R. Petersen, and M.P. Busch. 2012. Neuroinvasive disease and West Nile virus infection, North Dakota, USA, 1999-2008. *Emerg. Infect. Dis.* 18:684-686. <https://doi.org/10.3201/eid1804.111313>
- Ladbury, G.A.F., M. Gavana, K. Danis, A. Papa, D. Papamichail, S. Mourelatos, S. Gewehr, G. Theocharopoulos, S. Bonovas, A. Benos, and T. Panagiotopoulos. 2013. Population seroprevalence study after a West Nile virus lineage 2 epidemic, Greece, 2010. *PLoS One.* 8:e80432. <https://doi.org/10.1371/journal.pone.0080432>
- Lindsey, N.P., J.E. Staples, J.A. Lehman, and M. Fischer. 2012. Medical risk factors for severe West Nile virus disease, United States, 2008-2010. *Am. J. Trop. Med. Hyg.* 87:179-184. <https://doi.org/10.4269/ajtmh.2012.12-0113>
- Gervais, A., F. Rovida, M.A. Avanzini, S. Croce, A. Marchal, S.C. Lin, A. Ferrari, C.W. Thorball, O. Constant, T. Le Voyer, et al. 2023. Autoantibodies neutralizing type I IFNs underlie West Nile virus encephalitis in ~40% of patients. *J. Exp. Med.* 220:e20230661. <https://doi.org/10.1084/jem.20230661>
- Ferrari, A., I. Cassaniti, F. Rovida, D. Lilleri, S. Croce, F. Trespidi, S. Ghirardello, A. Gervais, S.Y. Zhang, J.L. Casanova, et al. 2025. Human type I interferons protect Vero E6 and ARPE-19 cells against West Nile virus and are neutralized by pathogenic autoantibodies. *Sci. Rep.* 15:11271. <https://doi.org/10.1038/s41598-025-89312-6>
- Lin, S.C., F.R. Zhao, H. Janova, A. Gervais, S. Rucknagel, K.O. Murray, J.L. Casanova, and M.S. Diamond. 2023. Blockade of interferon signaling decreases gut barrier integrity and promotes severe West Nile virus disease. *Nat. Commun.* 14:5973. <https://doi.org/10.1038/s41467-023-41600-3>
- Bastard, P., A. Gervais, T. Le Voyer, J. Rosain, Q. Philippot, J. Manry, E. Michailidis, H.H. Hoffmann, S. Eto, M. Garcia-Prat, et al. 2021. Autoantibodies neutralizing type I IFNs are present in ~4% of uninfected individuals over 70 years old and account for ~20% of COVID-19 deaths. *Sci. Immunol.* 6:eabl4340. <https://doi.org/10.1126/sciimmunol.abl4340>
- Barzaghi, F., C. Visconti, G.B. Pipitone, S. Bondesan, G. Molli, S. Giannelli, C. Sartirana, V. Lampasona, E. Bazzigaluppi, C. Brigatti, et al. 2025. Severe West Nile virus and severe acute respiratory syndrome coronavirus 2 infections in a patient with thymoma and anti-type I interferon antibodies. *J. Infect. Dis.* 231:e206-e212. <https://doi.org/10.1093/infdis/jiae321>
- Braune, J., L. Pechstein, C. Meisel, T. Meyer, J. Melchert, V.M. Corman, C. Franke, and T. Schneider. 2025. Detection of autoantibodies against interferon type I in patients with autochthonous West Nile virus encephalitis. *Brain Behav. Immun. Health.* 48:101085. <https://doi.org/10.1016/j.bbih.2025.101085>
- Roitblat, I.R., L. Hejla, Y. Moalem, E.H. Sklan, and E. Goldberg. 2026. Association of anti-interferon antibodies with severe clinical outcomes in West Nile virus -results of a recent outbreak. *Int. J. Infect. Dis.* 163:108282. <https://doi.org/10.1016/j.ijid.2025.108282>
- Gervais, A., A. Marchal, A. Fortova, M. Berankova, L. Krbkova, M. Pychova, J. Salat, S. Zhao, N. Kerrouche, T. Le Voyer, et al. 2024. Autoantibodies neutralizing type I IFNs underlie severe tick-borne encephalitis in ~10% of patients. *J. Exp. Med.* 221:e20240637. <https://doi.org/10.1084/jem.20240637>
- Gervais, A., P. Bastard, L. Bizien, C. Delifer, P. Tiberghien, C. Rodrigo, F. Trespidi, M. Angelini, G. Rossini, T. Lazzarotto, et al. 2024. Auto-Abs neutralizing type I IFNs in patients with severe Powassan, Usutu, or Ross River virus disease. *J. Exp. Med.* 221:e20240942. <https://doi.org/10.1084/jem.20240942>
- Bastard, P., E. Michailidis, H.H. Hoffmann, M. Chbihi, T. Le Voyer, J. Rosain, Q. Philippot, Y. Seeluthner, A. Gervais, M. Materna, et al. 2021. Auto-antibodies to type I IFNs can underlie adverse reactions to yellow fever live attenuated vaccine. *J. Exp. Med.* 218:e20202486. <https://doi.org/10.1084/jem.20202486>
- Gervais, A., P. Bastard, Q. Zhang, M.C. Jaffar-Bandjee, L. Bizien, L. Dahmane, M.P. Moiton, J. Jabot, R. Cally, A. Maillard, et al. 2026. Type I IFN autoantibodies underlie chikungunya live-attenuated vaccine encephalitis. *Proc. Natl. Acad. Sci. USA.* 123:e2532212123. <https://doi.org/10.1073/pnas.2532212123>
- Gervais, A., A. Borghesi, J.L. Casanova, and S.Y. Zhang. 2025. Auto-Abs against type I IFNs: Strong, common, and global determinants of severe arboviral diseases. *J. Hum. Immun.* 2:e20250090. <https://doi.org/10.70962/jhi.20250090>
- Percivalle, E., I. Cassaniti, A. Sarasini, F. Rovida, K.M.G. Adzeshoun, I. Colombini, P. Isernia, I. Cuppari, and F. Baldanti. 2020. West Nile or Usutu virus? A three-year follow-up of humoral and cellular response in a group of asymptomatic blood donors. *Viruses.* 12:157. <https://doi.org/10.3390/v12020157>
- Puel, A., C. Picard, M. Lorrot, C. Pons, M. Chrabieh, L. Lorenzo, M. Mamani-Matsuda, E. Jouanguy, D. Gendrel, and J.L. Casanova. 2008. Recurrent staphylococcal cellulitis and subcutaneous abscesses in a child with autoantibodies against IL-6. *J. Immunol.* 180:647-654. <https://doi.org/10.4049/jimmunol.180.1.647>
- Bastard, P., L.B. Rosen, Q. Zhang, E. Michailidis, H.H. Hoffmann, Y. Zhang, K. Dorgham, Q. Philippot, J. Rosain, V. Béziat, et al. 2020. Autoantibodies against type I IFNs in patients with life-threatening COVID-19. *Science.* 370:eabd4585. <https://doi.org/10.1126/science.abd4585>
- Groen, K., R. Kuratli, J. Enkelmann, S. Fernbach, P.D. Wendel-Garcia, W.I. Staiger, M. Lejeune, E. Sauras-Colón, F. Roche-Campo, P. Filippidis, et al. 2025. Type I interferon autoantibody footprints reveal neutralizing mechanisms and allow inhibitory decoy design. *J. Exp. Med.* 222:e20242039. <https://doi.org/10.1084/jem.20242039>

36. Groen, K., R. Kuratli, L. Sar, A. Vasou, M. Huber, D.J. Hughes, and B.G. Hale. 2024. Highly sensitive reporter cell line for detection of interferon types I-III and their neutralization by antibodies. *Eur. J. Immunol.* 54: e2451325. <https://doi.org/10.1002/eji.202451325>
37. Gervais, A., C. Le Floch, T. Le Voyer, L. Bizien, J. Bohlen, F. Celmeli, F. Al Qureshah, C. Masson, J. Rosain, M. Chbihi, et al. 2024. A sensitive assay for measuring whole-blood responses to type I IFNs. *Proc. Natl. Acad. Sci. USA.* 121:e2402983121. <https://doi.org/10.1073/pnas.2402983121>
38. Honda, N., U. Lindberg, P. Andersson, S. Hoffmann, and H. Takei. 2005. Simultaneous multiple immunoassays in a compact disc-shaped microfluidic device based on centrifugal force. *Clin. Chem.* 51:1955–1961. <https://doi.org/10.1373/clinchem.2005.053348>
39. Pons, S., L. Generenaz, A. Gervais, A. Puel, P. Bastard, N. Renard, V. Guyot, C. Vinit, F. Zheng, K. Brengel-Pesce, et al. 2025. Rapid detection of anti-IFN- α autoantibodies using a new automated VIDAS assay prototype. *Eur. J. Immunol.* 55:e202451516. <https://doi.org/10.1002/eji.202451516>
40. Zhang, Q., P. Bastard, A. Bolze, E. Jouanguy, S.Y. Zhang, COVID Human Genetic Effort, A. Cobat, L.D. Notarangelo, H.C. Su, L. Abel, and J.L. Casanova. 2020. Life-threatening COVID-19: Defective interferons unleash excessive inflammation. *Med.* 1:14–20. <https://doi.org/10.1016/j.medj.2020.12.001>
41. Patel, H., B. Sander, and M.P. Nelder. 2015. Long-term sequelae of West Nile virus-related illness: A systematic review. *Lancet Infect. Dis.* 15: 951–959. [https://doi.org/10.1016/S1473-3099\(15\)00134-6](https://doi.org/10.1016/S1473-3099(15)00134-6)
42. Petersen, L.R., A.C. Brault, and R.S. Nasci. 2013. West Nile virus: Review of the literature. *JAMA.* 310:308–315. <https://doi.org/10.1001/jama.2013.8042>
43. Bigham, A.W., K.J. Buckingham, S. Husain, M.J. Emond, K.M. Bofferding, H. Gildersleeve, A. Rutherford, N.M. Astakhova, A.A. Perelygin, M.P. Busch, et al. 2011. Host genetic risk factors for West Nile virus infection and disease progression. *PLoS One.* 6:e24745. <https://doi.org/10.1371/journal.pone.0024745>
44. Watson, J.T., P.E. Pertel, R.C. Jones, A.M. Siston, W.S. Paul, C.C. Austin, and S.I. Gerber. 2004. Clinical characteristics and functional outcomes of West Nile fever. *Ann. Intern. Med.* 141:360–365. <https://doi.org/10.7326/0003-4819-141-5-200409070-00010>
45. Manry, J., P. Bastard, A. Gervais, T. Le Voyer, J. Rosain, Q. Philippot, E. Michailidis, H.H. Hoffmann, S. Eto, M. Garcia-Prat, et al. 2022. The risk of COVID-19 death is much greater and age dependent with type I IFN autoantibodies. *Proc. Natl. Acad. Sci. USA.* 119:e2200413119. <https://doi.org/10.1073/pnas.2200413119>
46. Zhang, Q., A. Pizzorno, L. Miorin, P. Bastard, A. Gervais, T. Le Voyer, L. Bizien, J. Manry, J. Rosain, Q. Philippot, et al. 2022. Autoantibodies against type I IFNs in patients with critical influenza pneumonia. *J. Exp. Med.* 219:e20220514. <https://doi.org/10.1084/jem.20220514>
47. Casanova, J.L., and L. Abel. 2022. From rare disorders of immunity to common determinants of infection: Following the mechanistic thread. *Cell.* 185:3086–3103. <https://doi.org/10.1016/j.cell.2022.07.004>
48. Casanova, J.L., and L. Abel. 2024. The microbe, the infection enigma, and the host. *Annu. Rev. Microbiol.* 78:103–124. <https://doi.org/10.1146/annurev-micro-092123-022855>
49. Loconsole, D., F. Centrone, A. Sallustio, D. Casulli, V. Colella, O. Mongelli, G. Venturi, A. Bella, L. Marino, D. Martinelli, and M. Chironna. 2023. Abrupt increase in detection of locally acquired West-Nile-virus-lineage-2-mediated neuroinvasive disease in a previously non-endemic area of Southern Italy (2023). *Viruses.* 16:53. <https://doi.org/10.3390/v16010053>
50. Stufano, A., V. Schino, G. Sacino, R. Ravallesse, R. Ravallesse, L. De Benedictis, A. Morea, R. Iatta, S. Giannecchini, M.A. Stincarelli, et al. 2025. West Nile virus seroprevalence among outdoor workers in Southern Italy: Unveiling occupational risks and public health implications. *Viruses.* 17:310. <https://doi.org/10.3390/v17030310>
51. Tel-Aviv Sourasky Medical Center. 2024. Early interferon-beta treatment for West-Nile virus infection. <https://clinicaltrials.gov/study/NCT06510426> (accessed November 3, 2024).
52. Groen, K., and B.G. Hale. 2026. Human autoantibodies against type I interferons in severe viral disease. *Curr. Opin. Virol.* 74:101511.101511. <https://doi.org/10.1016/j.coviro.2026.101511>
53. Peng, J.J., J.Y. Ding, Y. Xu, H.P. Shih, Y.N. Lin, T.Y. Wu, Y.F. Lo, C.C. Lo, C.F. Yeh, C.Y. Kuo, et al. 2025. Chimeric autoantibody receptor T cells clonally eliminate B cells producing autoantibodies against IFN- γ . *Sci. Immunol.* 10:eadm8186. <https://doi.org/10.1126/sciimmunol.adm8186>
54. Fernbach, S., N.K. Mair, I.A. Abela, K. Groen, R. Kuratli, M. Lork, C.W. Thorball, E. Bernasconi, P. Filipkiddis, K. Leuzinger, et al. 2024. The Swiss HIV cohort study. Loss of tolerance precedes triggering and lifelong persistence of pathogenic type I interferon autoantibodies. *J. Exp. Med.* 221:e20240365. <https://doi.org/10.1084/jem.20240365>
55. Meager, A., K. Visvalingam, P. Peterson, K. Möll, A. Murumägi, K. Krohn, P. Eskelin, J. Perheentupa, E. Husebye, Y. Kadota, and N. Willcox. 2006. Anti-interferon autoantibodies in autoimmune polyendocrinopathy syndrome type 1. *Plos Med.* 3:e289. <https://doi.org/10.1371/journal.pmed.0030289>
56. Walter, J.E., L.B. Rosen, K. Csomos, J.M. Rosenberg, D. Mathew, M. Keszei, B. Ujhazi, K. Chen, Y.N. Lee, I. Tirosh, et al. 2015. Broad-spectrum antibodies against self-antigens and cytokines in RAG deficiency. *J. Clin. Invest.* 125:4135–4148. <https://doi.org/10.1172/JCI80477>
57. Oftedal, B.E., A. Hellesen, M.M. Erichsen, E. Bratland, A. Vardi, J. Perheentupa, E.H. Kemp, T. Fiskerstrand, M.K. Viken, A.P. Weetman, et al. 2015. Dominant mutations in the autoimmune regulator AIRE are associated with common organ-specific autoimmune diseases. *Immunity.* 42: 1185–1196. <https://doi.org/10.1016/j.immuni.2015.04.021>
58. Meyer, S., M. Woodward, C. Hertel, P. Vlaicu, Y. Haque, J. Kärner, A. Macagno, S.C. Onuoha, D. Fishman, H. Peterson, et al. 2016. AIRE-deficient patients harbor unique high-affinity disease-ameliorating autoantibodies. *Cell.* 166:582–595. <https://doi.org/10.1016/j.cell.2016.06.024>
59. Maccari, M.E., A. Scarselli, S. Di Cesare, M. Floris, A. Angius, A. Deodati, M. Chiriaco, P. Cambiaso, S. Corrente, G.S. Colafati, et al. 2017. Severe Toxoplasma gondii infection in a member of a NFKB2-deficient family with T and B cell dysfunction. *Clin. Immunol.* 183:273–277. <https://doi.org/10.1016/j.clim.2017.09.011>
60. Rosenberg, J.M., M.E. Maccari, F. Barzaghi, E.J. Allenspach, C. Pignata, G. Weber, T.R. Torgerson, P.J. Utz, and R. Bacchetta. 2018. Neutralizing anti-cytokine autoantibodies against interferon- α in immunodysregulation polyendocrinopathy enteropathy X-linked. *Front. Immunol.* 9:544. <https://doi.org/10.3389/fimmu.2018.00544>
61. Hetemäki, I., S. Laakso, H. Välimaa, I. Kleino, E. Kekäläinen, O. Mäkitie, and T.P. Arstila. 2021. Patients with autoimmune polyendocrine syndrome type 1 have an increased susceptibility to severe herpesvirus infections. *Clin. Immunol.* 231:108851. <https://doi.org/10.1016/j.clim.2021.108851>
62. Hetemäki, I., M. Kaustio, M. Kinnunen, N. Heikkilä, S. Keskitalo, K. Nowlan, S. Miettinen, J. Sarkkinen, V. Glumoff, N. Andersson, et al. 2021. Loss-of-function mutation in IKZF2 leads to immunodeficiency with dysregulated germinal center reactions and reduction of MAIT cells. *Sci. Immunol.* 6:eabe3454. <https://doi.org/10.1126/sciimmunol.abe3454>
63. Bastard, P., E. Orlova, L. Sozaeva, R. Lévy, A. James, M.M. Schmitt, S. Ochoa, M. Kareva, Y. Rodina, A. Gervais, et al. 2021. Preexisting autoantibodies to type I IFNs underlie critical COVID-19 pneumonia in patients with APS-1. *J. Exp. Med.* 218:e20210554. <https://doi.org/10.1084/jem.20210554>
64. Rosain, J., T. Le Voyer, X. Liu, A. Gervais, L. Polivka, A. Cederholm, L. Berteloot, A.V. Parent, A. Pescatore, E. Spinosa, et al. 2024. Incontinentia pigmenti underlies thymic dysplasia, autoantibodies to type I IFNs, and viral diseases. *J. Exp. Med.* 221:e20231152. <https://doi.org/10.1084/jem.20231152>
65. Materna, M., O.M. Delmonte, M. Bosticardo, M. Momenilandi, P.E. Conrey, B. Charmetieu-De Muyllder, C. Bravetti, R. Bellworthy, A. Cederholm, F. Staels, et al. 2024. The immunopathological landscape of human pre-TCR α deficiency: From rare to common variants. *Science.* 383:eadh4059. <https://doi.org/10.1126/science.adh4059>
66. Le Voyer, T., A.V. Parent, X. Liu, A. Cederholm, A. Gervais, J. Rosain, T. Nguyen, M. Perez Lorenzo, E. Rackaityte, D. Rinchai, et al. 2023. Autoantibodies against type I IFNs in humans with alternative NF- κ B pathway deficiency. *Nature.* 623:803–813. <https://doi.org/10.1038/s41586-023-06717-x>
67. Bastard, P., A. Gervais, M. Taniguchi, L. Saare, K. Särekannu, T. Le Voyer, Q. Philippot, J. Rosain, L. Bizien, T. Asano, et al. 2024. Higher COVID-19 pneumonia risk associated with anti-IFN- α than with anti-IFN- ω auto-Abs in children. *J. Exp. Med.* 221:e20231353. <https://doi.org/10.1084/jem.20231353>
68. Alotaibi, F., N.K. Alharbi, L.B. Rosen, A.Y. Asiri, A.M. Assiri, H.H. Balkhy, M. Al Jeraisy, Y. Mandourah, S. Al Johani, S. Al Harbi, et al. 2023. Type I interferon autoantibodies in hospitalized patients with Middle East respiratory syndrome and association with outcomes and treatment effect of interferon beta-1b in MIRACLE clinical trial. *Influenza Other Respir. Viruses.* 17:e13116. <https://doi.org/10.1111/irv.13116>
69. Gervais, A., A. Marchal, S. Boucherit, A. Abi Haidar, L. Bizien, A. Yalcinkaya, E. Sandström, X.F. Kong, E. Jacquemin, O. Bernard, et al. 2026. Autoantibodies neutralizing type I IFNs in patients with fulminant herpes simplex virus hepatitis. *J. Exp. Med.* 223:e20251760. <https://doi.org/10.1084/jem.20251760>
70. Zhang, Q., T.S. Conrad, M. Moncada-Velez, K. Jiang, A. Cupic, J. Eaton, K. Hutchinson, A. Gervais, R. Chen, A. Puel, et al. 2026. Autoantibodies neutralizing type I IFNs in a fatal case of H5N1 avian influenza. *J. Exp. Med.* 223:e20251962. <https://doi.org/10.1084/jem.20251962>

Supplemental material




Figure S1. **Demographic characteristics of individuals from seven new cohorts enrolled in the study.** (A) Number of individuals from each WNV group enrolled per center. (B) Number of individuals from each WNVD group enrolled per center. (C) Age distribution of the individuals enrolled, by WNV group. (D) Age distribution of the WNVD individuals, by center. (E) Sex distribution of the individuals enrolled, by WNV group. (F) Sex distribution of the individuals enrolled, by center. WNND, WNV neuroinvasive disease, a subset of WNVD; WNE, WNV encephalitis; WNM, WNV meningitis; AFP, acute flaccid paralysis; UNS, unspecified neurological syndrome (subsets of the WNND group).




Figure S2. **Correlation between luciferase-based IFN neutralization assay and ELISA results for the detection of auto-Abs to type I IFNs.** (A) Detection of auto-Abs against IFN- α 2 by ELISA versus neutralizing capacity for concentrations of 10 ng/ml and 100 pg/ml. (B) Detection of auto-Abs against IFN- β by ELISA versus neutralizing capacity for concentrations of 10 ng/ml and 1 μ g/ml. (C) Detection of auto-Abs against IFN- ω by ELISA versus neutralizing capacity for concentrations of 10 ng/ml and 100 pg/ml. The panels show concordance between luciferase-based neutralization assay and ELISA in the upper-left (samples with OD <0.5 on ELISA not displaying IFN neutralizing activity) and lower-right quadrants (samples with OD >0.5 with IFN neutralizing activity) and discordance between luciferase-based IFN neutralization assay and ELISA in the upper-right (samples with OD >0.5 on ELISA but not displaying IFN neutralizing activity) and lower-left quadrants (samples with IFN neutralizing activity despite OD <0.5 on ELISA). Values below 15% are considered to be associated with the presence of neutralizing auto-Abs.




Figure S3. **Detection of auto-Abs against type I IFN by ELISA in patients with WNVD from seven new cohorts. (A)** Detection, by ELISA, of auto-Abs against IFN- α 2, IFN- β , and IFN- ω for the WNVD group and the two subgroups of WNVD patients: non-neuroinvasive WNVD and neuroinvasive WNVD (WNND). **(B)** Detection, by ELISA, of auto-Abs against IFN- α 2, IFN- β , and IFN- ω in the subgroups of WNND patients: WNV encephalitis (WNE), WNV meningitis (WNM), acute flaccid paralysis (AFP), and unspecified neurological syndrome (UNS).




Figure S4. **Detection of auto-Abs neutralizing type I IFNs in the WNVD subgroups of patients from seven new cohorts. (A and B)** Luciferase-based neutralization assays for detecting auto-Abs neutralizing 10 ng/ml IFN- α 2, IFN- ω , or IFN- β and 100 pg/ml IFN- α 2 or IFN- ω , or 1 ng/ml IFN- β in the WNVD group and in the two WNVD subgroups of hospitalized patients with neuroinvasive WNVD (WNND) and without neuroinvasive disease. Neutralizing activity was defined as a luciferase signal below 15%. **(C and D)** Percentages of individuals with auto-Abs neutralizing type I IFNs at a concentration of 10 ng/ml or 100 pg/ml for IFN- α 2 or IFN- ω , or 1 ng/ml for IFN- β in the WNVD group and in the two subgroups of WNVD patients hospitalized with WNND and without neuroinvasive disease. IFN- α , auto-Abs neutralizing IFN- α 2 (regardless of their effects on other IFNs); IFN- ω , auto-Abs neutralizing IFN- ω (regardless of their effects on other IFNs); IFN- $\alpha \pm \omega \pm \beta$, auto-Abs neutralizing IFN- α 2 and/or IFN- ω and/or IFN- β ; IFN- $\alpha + \omega$, auto-Abs neutralizing both IFN- α 2 and IFN- ω .

Downloaded from <http://rupress.org/jhi/article-pdf/12/3/e20250189/20250189.pdf> by guest on 16 March 2026




Figure S5. **Demographic characteristics of 981 individuals from 13 cohorts.** (A) Number of individuals from each WNV group enrolled per center. (B) Number of individuals from each WNVD group enrolled per center. (C) Age distribution of the individuals enrolled, by WNV group. (D) Age distribution of the WNVD individuals, by center. (E) Sex distribution of the individuals enrolled, by WNV group. (F) Sex distribution of the individuals enrolled, by center. WNND, WNV neuroinvasive disease, a subset of WNVD; WNE, WNV encephalitis; WNM, WNV meningitis; AFP, acute flaccid paralysis; UNS, unspecified neurological syndrome (subsets of WNND group).

Provided online is Table S1, Table S2, and Table S3. Table S1 shows prevalence of WNVIC, WNVF, WNVD, or WNND of subjects carrying auto-Abs neutralizing at least 1 IFN-I by WNV lineage (WNV-1 or WNV-2). Table S2 shows risk of WNVIC, WNVF, WNVD, or WNND of subjects carrying auto-Abs neutralizing at least one IFN-I by sex. Table S3 shows risk of neuroinvasive disease and encephalitis for subjects carrying auto-Abs neutralizing specific sets of type I IFNs, relative to the general population, with adjustment for age and sex and risk by age group.